

Supporting Information for

Inter-comparison study of atmospheric ^{222}Rn and ^{222}Rn progeny monitors

Claudia Grossi^{1,2}, Olivier Llido³, Felix R. Vogel⁴, Victor Kazan³, Alessandro Capuana⁵, Scott D. Chambers⁶, Sylvester Werczynski⁶, Roger Curcoll^{7,8}, Marc Delmotte³, Arturo Vargas¹, Josep-Anton Morguí^{7,9}, Ingeborg Levin⁵, Michel Ramonet³.

¹ Institut de Tècniques Energètiques (INTE), Universitat Politècnica de Catalunya (UPC), Barcelona, Spain;

² Physics Department, Universitat Politècnica de Catalunya (UPC), Barcelona, Spain;

³ Laboratoire des Sciences du Climat et de l'Environnement, Université Paris-Saclay (LSCE/IPSL, CEA-CNRS-UVSQ), Gif-sur-Yvette, France;

⁴ Climate Research Division, Environment and Climate Change Canada, Toronto, Canada;

⁵ Institut für Umweltphysik (IUP), Heidelberg University, Heidelberg, Germany;

⁶ Environmental Research, ANSTO, Lucas Heights, Australia;

⁷ Institut de Ciència i Tecnologia Ambientals (ICTA), Universitat Autònoma de Barcelona (UAB), Cerdanyola del Vallès, Spain;

⁸ Chemical Department, Universitat Politècnica de Catalunya (UPC), Barcelona, Spain;

⁹ Departament Biologia Evolutiva, Ecologia i Ciències Ambientals, Universitat de Barcelona (UB), Barcelona, Spain.

Correspondence to: Claudia Grossi (claudia.grossi@upc.edu)

Contents of this file

S1

S2

S3

S4

Introduction

Here we show the plots of the linear regression fits calculated during the Phase I and II of our intercomparison study.




Figure S1. **Left panel:** Linear regression fits between the hourly atmospheric ^{214}Po activity concentrations measured by the LSCE and the ^{222}Rn activity concentrations measured by ARMON during the Phase I at the ODM site. **Right panel:** Linear regression fits between the hourly atmospheric ^{214}Po activity concentrations measured by the LSCE and the HRM during the Phase I at the ODM site.




Figure S2. **Left panel:** Linear regression fits between the hourly atmospheric ^{222}Rn activity concentrations measured by the ANSTO_ODM and by ARMON during the Phase I at the ODM site. **Right panel:** Linear regression fits between the hourly atmospheric ^{222}Rn activity concentrations measured by the ANSTO_ODM and the hourly atmospheric ^{214}Po activity concentrations measured by the HRM during the Phase I at the ODM site.




Figure S3. **Left panel:** Linear regression fits between the hourly atmospheric ^{214}Po activity concentrations measured by the HRM and the hourly atmospheric ^{222}Rn activity concentration

measured by the ARMON during the Phase I at the ODM site. **Right panel:** Linear regression fits between the hourly atmospheric ^{214}Po activity concentrations measured by the HRM and the hourly atmospheric ^{222}Rn activity concentration measured by the ARMON during the Phase II at the SAC site.




Figure S4. **Left panel:** Linear regression fits between the hourly atmospheric ^{222}Rn activity concentrations measured by the ANSTO_SAC and by ARMON during the Phase II at the SAC site. **Right panel:** Linear regression fits between the hourly atmospheric ^{222}Rn activity concentrations measured by the ANSTO_SAC and the hourly atmospheric ^{214}Po activity concentrations measured by the HRM during the Phase II at the SAC site.