

In this study, an experimental version of the model was used for simulations (Makar et al., 2018; 2017; 2015a; 2015b; Akingunola et al., 2018), designed for better process representation where processing time is not as crucial a consideration as for the operational code, and where more detailed science representations are desired. Some of the main differences between the experimental and operational versions of GEM-MACH are given in Table S1, below.

Table S1: Process Description / Features for the Operational versus Experimental GEM-MACH

<b>Feature</b>	<b>Operational</b>	<b>Experimental</b>
Particle size distribution	2-bin sectional approach, with rebinning to 12 bins in key particle microphysical processes	12-bin sectional approach applied throughout.
Cloud microphysics package	Kain-Fritsch (convective), Sundqvist (all other cloud)	P3 cloud microphysics package
Treatment of ammonia	One-way emissions and deposition	Bidirectional NH <sub>3</sub> fluxes (Whaley et al., 2018)
Anthropogenic plume rise formula	Briggs top-of-stack stability function formulation.	Residual buoyancy calculation between individual model layers (Akingunola et al., 2018)
Leaf Area Index information	BELD3 database	Monthly satellite-derived LAI values (Makar et al., 2017)
Methane treatment	Constant background values	Emitted, transported, reacted species.
Gas-phase chemistry solver	Young and Boris	KPP / RODAS3
Meteorological modulation of fugitive dust	Applied to all particulate species emitted as fugitive dust	Applied only to the “crustal material” fraction of fugitive dust emissions.
Particle Deposition	As per Gong et al. (2003): calculation of mass flux leaving layer and transfer to destination layer or surface.	1D semi-Lagrangian advection in the vertical, to prevent discontinuities in the resulting vertical profile of particle mass.
Forest Canopy Shading and	None	As per Makar et al. (2017):

Turbulence Treatment		three additional layers are added to the model to explicitly simulate the attenuation of light and the profile of vertical diffusion within plant canopies.
Coupling with meteorology	One-way: meteorological fields modify the chemical fields.	Two-way (fully coupled): as per Makar et al. (2015); particle direct and indirect effects are incorporated into the modelling framework, allowing meteorology to perturb the chemistry.
Forest Fire Emissions Module	<p>CFFEPSv2.03 (Chen et al., 2019): Off-line calculations using near-real-time satellite hotspot data, a priori meteorological forecast temperature profiles at 5 levels, and 3 coupled models (a fire growth model, a fire emissions model, and a column radiative transfer model; (Anderson et al., 2011).) to determine the mass of emissions and vertical height reached by smoke plumes. The resulting emissions data into the column are input into GEM-MACH along with other emissions fields.</p> <p>Near-real time fire hotspot information is obtained from three satellite sensors: MODIS, the Advanced Very High Resolution Radiometer (AVHRR), and Visible Infrared Imaging Radiometer Suite (VIIRS)</p>	CFFEPSv4.0. Key differences from v2.03 include: (a) The original C algorithms of v2.03 were converted to Fortran90 (CFFEPSv3.0) and then incorporated into GEM-MACH as an on-line subroutine package; (b) plume rise is calculated internally using GEM's full vertical resolution for temperature profiles (greater resolution in plume rise calculations); (c) particulate emissions are subdivided into all 12 particle bins; (d) due to the use of the on-line paradigm, forest fire emissions may influence the weather, in turn modifying the predicted plume height on subsequent calculations.

	<p>processed through the Canadian Wildland Fire Information System operated by the Canadian Forest Service, Natural Resources Canada. Further details describing the implementation of the GEM-MACH wildfire component within the model can be found in e.g. Munoz-Alpizar et al. (2017); Pavlovic et al. (2016); Chen et al. (2019).</p>	
--	---	--

## References

- Akingunola, A., Makar, P. A., Zhang, J., Darlington, A., Li, S.-M., Gordon, M., Moran, M. D., and Zheng, Q.: A chemical transport model study of plume-rise and particle size distribution for the Athabasca oil sands, *Atmos. Chem. Phys.*, 18, 8667–8688, <https://doi.org/10.5194/acp-18-8667-2018>, 2018.
- Chen, J., Anderson, K., Pavlovic, R., Moran, M. D., Englefield, P., Thompson, D. K., Munoz-Alpizar, R., and Landry, H.: The FireWork v2.0 air quality forecast system with biomass burning emissions from the Canadian Forest Fire Emissions Prediction System v2.03, *Geosci. Model Dev.*, 12, 3283–3310, <https://doi.org/10.5194/gmd-12-3283-2019>, 2019.
- Gong, S.L., Barrie, L.A., Blanchet, J.-P., von Salzen, K., Lohmann, U., Lesins, G., Spacek, L., Zhang, L.M., Girard, E., Lin, H., Leaitch, R., Leighton, H., Chylek, P., and Huang, P., Canadian Aerosol Module: A size-segregated simulation of atmospheric aerosol processes for climate and air quality models 1. Module development, *J. Geophys. Res.*, 108, D1, 4007, doi:10.1029/2001JD002002, 2003.
- Makar, P.A., Staebler, R.M., Akingunola, A., Zhang, J., McLinden, C., Kharol, S.K., Pabla, B., Cheung, P. and Zheng, Q., The effects of forest canopy shading and turbulence on boundary layer ozone, *Nature Communications*, 8, art. no. 15243, doi: 10.1038/ncomms15243, 2017.
- Makar, P.A., Akingunola, A., Aherne, J., Cole, A.S., Aklilu, Y.-A., Zhang, J., Wong, I., Hayden, K., Li, S.-M., Kirk, J., Scott, K., Moran, M.D., Robichaud, A., Cathcart, H., Baratzedah, P., Pabla, B., Cheung, P., Zheng, Q., Jeffries, D.J., Estimates of exceedances of critical loads for acidifying deposition in Alberta and Saskatchewan, *Atmos. Chem. Phys.*, 18, 9897-9927, 2018
- Makar, P., Gong, W., Hogrefe, C., Zhang, Y., Curci, G., Zabkar, R., Milbrandt, J., Im, U., Balzarini, A., Baró, R., Bianconi, R., Cheung, P., Forkel, R., Gravel, S., Hirtl, M., Honzak, L., Hou, A., Jiménez-Guerrero, P., Langer, M., Moran, M., Pabla, B., Pérez, J., Pirovano, G., José, R. S., Tuccella, P., Werhahn, J., Zhang, J., and Galmarini, S.: Feedbacks between air pollution and weather, part 2: Effects on chemistry, *Atmospheric Environment*, 115, 499 – 526, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2014.10.021>, <http://www.sciencedirect.com/science/article/pii/S1352231014008103>, 2015a.

Makar, P., Gong, W. F., Milbrandt, J., Hogrefe, C., Zhang, Y., Curci, G., Zabkar, R., Im, U., Balzarini, A., Baró, R., Bianconi, R., Cheung, P., Forkel, R., Gravel, S., Hirtl, M., Honzak, L., Hou, A., Jimenez-Guerrero, P., Langer, M., and Galmarini, S.: Feedbacks between air pollution and weather, Part 1: Effects on weather, *Atmospheric Environment*, 115, 2015b.

Munoz-Alpizar, R., Pavlovic, R., Moran, M. D., Chen, J., Gravel, S., Henderson, S. B., Ménard, S., Racine, J., Duhamel, A., Gilbert, S., Beaulieu, P.-A., Landry, H., Davignon, D., Cousineau, S., and Bouchet, V.: Multi-Year (2013–2016) PM<sub>2.5</sub> Wildfire Pollution Exposure over North America as Determined from Operational Air Quality Forecasts, *Atmosphere*, 8, <https://doi.org/10.3390/atmos8090179>, <http://www.mdpi.com/2073-4433/8/9/179>, 2017.

Pavlovic, R., Chen, J., Anderson, K., Moran, M. D., Beaulieu, P.-A., Davignon, D., and Cousineau, S.: The FireWork air quality forecast system with near-real-time biomass burning emissions: Recent developments and evaluation of performance for the 2015 North American wildfire season, *Journal of the Air & Waste Management Association*, 66, 819–841, <https://doi.org/10.1080/10962247.2016.1158214>, <https://doi.org/10.1080/10962247.2016.1158214>, PMID: 26934496, 2016.

Pendlebury, D., Gravel, S., Moran, M. D., and Lupu, A.: Impact of chemical lateral boundary conditions in a regional air quality forecast model on surface ozone predictions during stratospheric intrusions, *Atmospheric Environment*, 174, 148 – 170, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2017.10.052>, <http://www.sciencedirect.com/science/article/pii/S1352231017307185>, 2018.

Whaley, C. H., Makar, P. A., Shephard, M. W., Zhang, L., Zhang, J., Zheng, Q., Akingunola, A., Wentworth, G. R., Murphy, J. G., Kharol, S. K., and Cady-Pereira, K. E.: Contributions of natural and anthropogenic sources to ambient ammonia in the Athabasca Oil Sands and north-western Canada, *Atmos. Chem. Phys.*, 18, 2011–2034, <https://doi.org/10.5194/acp-18-2011-2018>, 2018.