Authors responses to Atmos. Meas. Tech. Discuss., https://doi.org/10.5194/amt-2019-504

Establishment of AIRS Climate-Level Radiometric Stability using Radiance Anomaly Retrievals of Minor Gases and SST

by L. Larrabee Strow and Sergio DeSouza-Machado

Our responses are given below. For ease of review, we type-faced the reviewers questions in blue. At the bottom of our responses we include a copy of the new manuscript with removals in red and additions in blue with an underline. The main changes to the manuscript are:

- 1. Additional material on our clear selection has been added to Section 3 to show that we are not "matching" to ERA-I and indeed have objectively chosen the clear scenes.
- 2. Enhanced justification for using ERA-I model fields for evaluation of the anomaly retrieval Jacobians. This is done in a general manner in new Section 4.3. In Section 5 we include a detailed numerical analysis of any possible errors in our trend and anomaly results due to inaccurate ERA-I model fields, esp. see Table 4. (The blue underlining denoting new material in this table is overlapping the Jacobian denominators in Columns 1 and 2, but that will not show up in the final manuscript.)
- 3. The description of how we handle the co-linearity of the CO₂ and temperature Jacobians in Sect. 4.4 has been considerably lengthened. In addition, our main results shown in Table 3 now include (second row) an estimate of the AIRS stability using ESRL CO₂ trends as the standard peformed **without** applying our correction for the co-linearity of these Jacobians.
- 4. Section 6 now includes new material, as suggested by Reviewer 1, that illustrates how the AIRS BT anomaly trends are modified by individual gases, temperature, and water vapor.
- 5. In several places in this manuscript, such as Eq. 1 and 2, we have rewritten terms and tried to clarify the language so that the reader understands that our "measurement" is BT anomalies, not BT absolute spectra. This fact removes many concerns about biases in our data that seemed to trip people up.

Reply to: Anonymous Referee 2

Note: Many comments refer the reviewer to changes in the manuscript that are included at the end of this file.

1 General Comments

The manuscript introduces and discusses important methodology and results to the suitability and utilisation of AIRS for climate applications, showing also a way for other hyperspectral sounder products (e.g. IASI, CrIS...). Based on a 16-year series, it indicates that AIRS radiance measurements and retrieved quantities match stability and sensitivity requirements for climate trend studies, as evaluated indirectly by Obs fit computations and direct intercomparisons to external reference measurements. This is found in line wit the scope of the journal and expected scientific novelty.

I find the manuscript overall very well structured and written, providing sufficient results and discussions, with clear illustrations and appropriate references.

I recommend the publication of the manuscript pending few clarifications listed below.

2 Use of ERA for Jacobians

"However, ERA-I is so accurate, that is not necessary" and similar other statement, sounds too absolute statement. The "so accurate" sould be elaborated a bit more, especially in view of some non-negligible biases seen in Fig.2.

Indeed, we just stated in the paper that ERA-Interim (ERA-I) was accurate enough. It is important to remember that the data being retrieved, BT anomalies, are very small, which implies that the Jacobians used in the retrieval do not have to be terribly accurate. This was also noticed by the other reviewers, therefore we put a significant effort into addressing the concerns about the accuracy of ERA-I for Jacobian evaluations.

The biases in Fig. 2 are actually not that big. Note that we are showing a slightly different bias in the revised paper. Since our RTA (SARTA) has a default CO2 amount of 385 ppm we selected a time period where the atmosphere had the same CO2 amount as well. That made the biases in the CO2 sensitive region from 700 to 750 cm⁻¹ even smaller, around 0.2-0.3K on average. The AIRS radiometry may not be that accurate. The water region biases (1300-1615 cm⁻¹) are larger, but we do not heavily rely on accurate water vapor in the mid-troposphere for this work. Even if the Jacobians valuels are slightly incorrect, our retrieval still removes their effect in terms of interference with N_{2} and CH_{4} . We are somewhat sensitive to the column amount of water vapor for window region channels that are used to fit for the SST anomalies. However, as noted below, we added quite a bit of material to justify the use of ERA for Jacobian evaluation and find that this introduces extremely small inaccuracies that can be ignored. Some details of what we did are discussed below.

Figure 2 has been enhanced to also show the single-footprint standard deviation between the AIRS observations and our ERA-I simulated BTs. We also include the AIRS noise in the new bottom panel of this figure, which shows that in the CO2 sounding region, the AIRS noise is barely smaller then the standard deviation of the biases, showing the ERA-I temperature fields are very close to what is observed.

A new section, 4.3 "Construction of Jacobians" has been added to address this issue qualitatively, introducing some estimates of ERA accuracy and AIRS radiometric accuracy (which would limit accuracy of standard retrievals used to generate the atmospheric state Jacobians, if we had done so.).

In Section 5.4 (CO2 anomalies) and 5.7 (SST anomalies) we now include a very detailed analysis of how potential uncertainties in the Jacobians generated using ERA-I would affect our anomaly retrievals and trends. These are summarized in a new Table (#4), which shows that using ERA-I for the Jacobians is likely to introduce errors that are far far below our statistical uncertainties. This is mainly due to the fact that the anomalies are quite small, so extremely accurate Jacobians are not needed.

I content that the ERA-I fields, once heavily averaged as in this paper, are more accurate than any 1D-var retrievals (like those done in the AIRS Level 2 product), since the assimilation includes many instruments, including AIRS, IASI, and CrIS.

3 Specific Comments

• L38: has it ever been considered to use AMSU in combination to disentangle T/CO2 signals? Like e.g. in Crevoisier et al. 2011 (TBC). would independence be more useful to climate studies, as oppoosed to

using climatological CO2?

This is certainly a valid idea and many people use it. We did not want to invoke another instrument in these anomaly retrievals since that introduces uncertainties in say, the AMSU radiometric stability, which could greatly complicate the analysis. Our temperature trends are quite similar to those in ERA, which lends credibility that we have indeed separated CO_2 from temperature. In addition, we added quite a bit of material to the new Section 4.4 to justify our approach on separating CO_2 from temperature. We believe the results shown show that this was successful. Please note that the RHS plot in Fig. 7 had a labelling error that may have introduced some confusion on our mitigation of co-linearities in the CO_2 and temperature Jacobians.

• L56: not sure what the retrieval residuals can tell us really. The fit, if minimisation well programmed, will always come down to about the observation error in the end.

Exactly. And what this paper is after **is** the observation error, ie how stable is AIRS, what channels are mis-behaving and by how much. This is discussed in Section 6.2.

• L65: how about any bias correction prior to the 1D-Var? NWP DA for instance need BC in variational minimisation to fit OBS with CALC. Has it been ever considered in AIRS L2 retrieval?

That is the whole point of retrieving from BT anomalies. AIRS calibration errors (that do not change in time) are removed when forming the BT anomaly. And, that also means we only use relative changes in the RTA to determine the anomalies, so RTA bias is removed as well. That is why this approach works so well. In order to minimize confusion on this point we have added a sentence at the end of Sect. 4.1 stating this more explicitly, and have changed Eqs. 1 and 2 slightly to make it a bit more transparent that we are retrieving from a BT anomaly. In addition, we also clarified our approach by explicitly defining the observation y in Eqs. 4 and 5 in terms of the BT anomaly derived in Eq. 2.

• L115: over year+? clarify editorial

We changed the working to "mult-year".

• L161: by by (or bye bye typo)

Fixed.

• L162: stddev in window may be due also to uncertainties in the forward modelling, including RTM/spectro as well as input SST/H2O profiles. eq(5): explicit L?

Yes, we didn't explicitly define *L*, now fixed. The stddev here would be extremely insensitive to forward modelling errors and spectroscopy because these are mean anomalies, and only represent small changes in the atmospheric state, not how well we can simulate the observed BT. Agreed that SST/H2O profile errors could cause some of the window STD, now stated in the paper.

• L191: why are forward model uncertainties not included? The rationale (and consequences) should be discussed. Any bias correction?

There is no concept of a bias correction here, since we are retrieving anomalies. Time independent bias errors in the AIRS calibration are removed in forming the anomaly, and the retrieval of anomalies does not need to compute the absolute BT, only it's variations. The forward model is fixed throughout this time period so there was no need to put in any RTA uncertainties into the retrieval. There **are** second order RTA errors that come into the retrieval via the Jacobians, and these are discussed in detail in the revised paper.

Please note that our a-priori values for the temperature and water profiles is "zero", as well as our linearization point. That's because these are anomalies, not absolute BTs. Granted, we need the atmospheric state for computing the Jacobians, but ERA-I serves that purpose with high accuracy for our purposes.

• L194: typo "more layers thAn"

Fixed.

• L206: needs a little more explanation how the 0.004K and even 0.001K extremely low noise values were found. I assumed simple signal/noise enhancements resulting from massive averaging. However it is difficult to believe that one can fit the observation down to that level, usually the RTM uncertainties combined with the effect of state vector not varied in the retrieval are larger than the instrument noise.

We tried to clarify this, but it is indeed due to massive averaging. I agree, it was hard for us to believe we needed to faithfully use these low errors in our OE retrieval, but as stated in the paper, until we did that we could not retrieve the large CO_2 anomalies near the end of the time series. Again, we are retrieving BT anomalies, and they are very small (not much larger than single footprint noise). And, as stated above, using anomalies avoids calibration and RTA bias errors, resulting in very robust retrievals.

• 238, 264: incomplete ref (Aumann)

Fixed.

• 240-242: the DoF for O3 and H2O appears quite large compared to what is commonly accepted, as pointed out (usually \approx 3 for O3 and 6-8 for H2O). I think this is more directly due to the massive averaging which effectively results in lowered instrument noise. 321 H2O channels on a single pixel would not bring such a high DoF, would it? Temeprature is a little under what is commonly expected of hyperspectral sounders \approx 10-12 DoFs. But in this case, the channel pruning might be responsible for the signal loss.

Yes, we agree. It's the low noise that does this. Generally you would expect 5 DOFS or so for H2O. Indeed the high number of H_2O channels contributed to this DOF, and we ignored all sorts of correlations that might exist. We didn't pursue this in detail because it is not all that relevant for this study, where we are not that interested in the final temperature and H_2O profiles, as long as their trends are reasonable. We also used a very simple approach for the DOF measurements, using only the mean profile. I strongly suspect that that these averaged anomalies, and associated trends don' have enough variability to need more than a few DOFs. These issues will be pursued in another study where we are not concentrating just on highly averaged clear scene anomalies.

• 254: typo to to (two to)

Fixed.

• 271: complete ref (Tans and Keeling)

I am doing what their web site suggests, and they don't give a date. Maybe the AMT editors/proofreaders can help me on this.

• L312: section reference broken

Fixed.

• Fig.11: isn't it possible to plot break-down of ESRL components in their different latitudes location?

That would involve using a much more complicated ESRL data set that is not gridded by latitude, but just provides point sources. Or, it would involve using CarbonTracker, which is an assimilated product that I though was not appropriate. What ESRL does provide in simple form are a few high-quality stations (MLO and CGRIM) and a "global" CO_2 product which is easy to use. Since we are after AIRS stability in this paper, using the "global" product seemed appropriate. A scientific study of CO_2 spatial variability using this appproach would need more complete in-situ CO_2 data, but that is for another study.

• Explain Lag-1 autocorrelations

We have removed several reference to lag-1 correlations in Section 3.3 since they are not needed at that point. Later in the paper, when we do discuss "lag-1 autocorrelations" we clarify that these are corrections to least-squares uncertainty estimates for the serial correlations in the anomaly time series.

The Santer reference is very widely used in climate research and it details how the lag-1 autocorrelation of a time series can be used to empirically correct least-squares uncertainty estimates that assume that the time series (residuals) will have uncorrelated gaussian noise (which is rarely true). An explanation of the lag-1 approach would be too detailed for this paper and should be unnecessary given its popularity in the climate community.

• 5.5 For clarity, move Table 4 and Fig. 12 in section 5.5.

I can't do that with latex submissions. Hopefully the AMT typesetting will fix that problem.

• The larger departure AIRS - ESRL for CH4 and N2O over time is interesting, yet unexplained. Seems noticeable enough in Climate app context.

I do not think that departure is correct, it is due to some shifts in the AIRS calibration due to AIRS hardware "events". See the discussion in Section 5.5 for the details on this.

• 5.7: I understand that OSTIA provides the foundation SST (Merchant et al. 2014, Corlette et al., GHRSST website...), which is physically different to the radiative skin SST which is accessible to AIRS. In that respect, I find the agreement rather impressive with nearly no biases, while one could expect some given the different SST quantities. The authors should confirm the respective intrinsic nature of the SST datasets (model and retrieved) and possibly discuss the agreement accordingly. A correction of e.g. skin-to-bulb bias of 0.17K may be necessary in absolute term, it would however not impact the relative variation over time.

We are only measuring OSTIA (via ERA-I) and OISST SST trends and anomallies. So, any biases in these products relative to AIRS radiances would not show up in our analysis.

Establishment of AIRS Climate-Level Radiometric Stability using Radiance Anomaly Retrievals of Minor Gases and SST

L. Larrabee Strow and Sergio DeSouza-Machado

Dept of Physics and JCET, University of Maryland Baltimore County, Baltimore, Maryland **Correspondence:** L. Larrabee Strow (strow@umbc.edu)

Abstract. Temperature, H_2O , and O_3 profiles, as well as CO_2 , N_2O , CH_4 , CFC12, and SST scalar anomalies are computed using a clear subset of AIRS observations over ocean for the first 16-years of NASA's EOS-AQUA AIRS operation. The AIRS <u>Level 1c Level-1c</u> radiances are averaged over 16 days and 40 equal-area zonal bins and then converted to brightness temperature anomalies. Geophysical anomalies are retrieved from the brightness temperature anomalies using a relatively

- 5 standard optimal estimation approach. The CO₂, N₂O, CH₄, and CFC12 anomalies are derived by applying a vertically uniform multiplicative shift to each gas in order to obtain an estimate for the gas mixing ratio. The minor gas anomalies are compared to the NOAA ESRL in-situ values and used to estimate the radiometric stability of the AIRS radiances. Similarly the retrieved SST anomalies are compared to the SST values used in the ERA-Interim reanalysis and to NOAA's OISST SST product. These inter-comparisons strongly suggest that many AIRS channels are stable to better than 0.02 to 0.03 K/Decade, well
- 10 below climate trend levels, indicating that the AIRS blackbody is not drifting. However, detailed examination of the anomaly retrieval residuals (observed minus computed) show various small unphysical shifts that correspond to AIRS hardware events (shutdowns, etc.). Some examples are given highlighting how the AIRS radiances stability could be improved, especially for channels sensitive to N_2O and CH_4 . The AIRS short wave channels exhibit larger drifts that make them unsuitable for climate trending, and they are avoided in this work. The AIRS Level 2 surface temperature retrievals only use short wave channels. We
- 15 summarize how these short wave drifts impacts recently published comparisons of AIRS surface temperature trends to other surface climatologies.

1 Introduction

The Atmospheric Infrared Sounder (AIRS) on NASA's AQUA satellite platform (Aumann et al., 2003) measures 2378 high-spectral resolution infrared radiances between 650 and 2665 cm⁻¹ with a resolving power ($\lambda/\Delta\lambda$) of ~1200. Launched in 2002

20 into a sun-synchronous polar orbit with a 13:30 ascending node equator crossing time, AIRS now has been operating almost continuously for 17+ years.

The long record of AIRS allows measurements of short-term climate trends that are especially useful given it's global coverage. Nominal decadal climate temperature trends are in the 0.1-0.2K/decade range. For example a recent Intergovernmental Panel on Climate Change (IPCC) report (Masson-Delmotte et al., 2018) suggests 20th century trends (2000-2017) of about

25 0.17K/decade. If AIRS is to contribute to climate-level trend measurements, uncertainty estimates for the time stability of the

AIRS radiances are a pre-requisite before using AIRS Level 2/3 products for climate level trending. Estimating the level of any instrument-related trends, for a wide range of AIRS channels, is the subject of this work.

A recent study (Aumann et al., 2019) addressed the stability of a single AIRS channel by comparisons to sea surface temperatures (<u>SST</u>). Some limitations of this that study are addressed below, but the major limitation of this work is that its

30 <u>major limitation is that is evaluates only one channel</u>. AIRS retrievals use 400+ AIRS channels, and there is no guarantee that the AIRS stability in one channel applies to all channels, as acknowledged in (Aumann et al., 2019).

AIRS is sensitive to a host of atmospheric and surface variables, including atmospheric temperature (via CO_2 emissions), humidity, surface temperature, O_3 , CH_4 , N_2O , carbon monoxide, clouds, coarse mode aerosols and other minor gases. 1D-var retrievals such as the AIRS Level 2 products (Susskind et al., 2014) attempt to retrieve all relevant atmospheric and surface

- variables in order to produce the most accurate temperature and H_2O profiles. The atmospheric CO_2 concentration is especially important for AIRS retrievals since most of the radiance measured in the temperature sounding channels is due to CO_2 emission. However, it is difficult to separate the CO_2 concentration from variations in the temperature profile due to co-linearity of their Jacobians. Consequently, the AIRS Level 2 retrievals instead vary CO_2 in the forward model to account for CO_2 growth during the mission (Blaisdell, 2019).
- The largest radiance trends seen by AIRS are due to the growth rate of CO_2 in the atmosphere. Assuming a nominal growth rate of 2 ppm/year, and max sensitivity of AIRS channels of CO_2 of 0.03K/ppm, the brightness temperature (BT) shift in AIRS over 16-years is ~1K, or 0.06K/year. Concentrations of atmospheric carbon dioxide have been measured worldwide for many years with extremely high accuracy (Masarie and Tans, 1995; Tans and Keeling) by NOAA Earth System Research Laboratory (ESRL). Averaged yearly, CO_2 concentrations are highly uniform globally, with little latitudinal variation in growth rates.
- 45 Similarly NOAA ESRL also provides a wide network of measurements of N_2O and CH_4 , which are also relatively uniformly mixed over yearly time-periods. Here we use the high accuracy of the trends in these in-situ measurements of minor gases to determine the stability of a large number of AIRS channels.

Sea surface temperature (SST) SST trends are also extremely well measured and generally referenced to the in-situ ARGO (Argo, 2019) buoy network but interpolated to a full grid using instruments such as the AVHRR. Two SST products referenced
to the buoy network are compared to AIRS trends here: (1) NOAA's Optimum Interpolation SST (version 2) (OISST) (Banzon et al., 2016), and (2) the Operational Sea Surface Temperature and Ice Analysis (OSTIA) (Stark et al., 2007), which has

been used in the ERA-I Interim Reanalysis (ERA-I) since 2009 (Dee et al., 2011). Prior to Feb. 2009 ERA-I used the NCEP

- Real-Time Global SST (RTG) product, a precursor to OISST.
- AIRS stability is referenced to trends in these minor gases and SST by performing 1D-var retrievals of clear scene radiance anomalies averaged into 40 equal-area latitude bins and 16-day time periods. Comparisons of the retrieved gas concentrations and SST trends, combined with examination of the retrieval residuals, provides a number of powerful tests of AIRS radiometric stability as well as detailed information on AIRS performance changes due to several minor instrument shutdowns that took place occasionally over the mission.

After a summarizing the characteristics of the AIRS instrument, and the data used in this work, the retrieval methodology 60 is reviewed with a short discussion of the retrieved temperature profile time series. We follow with stability estimates derived from the anomaly spectra retrievals of CO_2 , N_2O , CH_4 , and SST. Although AIRS is most sensitive to the two best in-situ data sets, CO_2 and SST, we also compare to retrievals of N_2O and CH_4 since they are also relatively well measured and help test the AIRS performance in spectral regions not covered by CO_2 and SST. Finally we examine the time series of the anomaly retrieval residuals (BT observed - fit) time series since, together with the anomaly geophysical retrievals, they provide detailed information on AIRS radiances over time, especially the instrument response to various short shutdowns that occurred during the mission.

2 AIRS Instrument and Data

65

80

Several details of the AIRS instrument design are relevant to the processing performed here and are needed to understand some of the results. AIRS has 2378 spectral channels divided up into 17 different detector arrays. Appendix A gives the nominal

- 70 wavenumber boundaries of these arrays. Arrays M-11 and M-12 are linear arrays of single photoconductive HgCdTe detectors. The other AIRS arrays are photovoltaic detectors, and each reported detector output is actually some linear combination of two detectors offset from each other in the vertical (not dispersive) direction. The photovoltaic detectors for each AIRS channels channel are labeled "A" and "B". The relative contributions of A and B detectors can be changed by command to the spacecraft. The majority of these detectors are wired for equal contributions by the A and B detectors, which we denote as A+B detectors.
- 75 However, some detectors have always been inoperable, or their performance characteristics changed in orbit, so there are a number of A-only and B-only detectors.

The radiometric and spectral characteristics of the A versus B detectors can be slightly different. During the mission, good A+B detectors can suddenly exhibit greatly increased noise when one or the other of the two detectors fails or degrades. In many circumstances the AIRS Project has changed A+B detectors to be either A-only or B-only in order to recover that particular channel, albeit at slightly lower noise levels than if both detectors were working properly. Fortunately, many of the

A-only and B-only detectors are in the window regions where AIRS has tremendous redundancy. Unfortunately, the M-10 array which covers the tropospheric CO_2 sounding channels also has a good number of A-only, B-only detectors.

Here we avoid any photovoltaic channel that is not A+B, and any channel with a state change during the mission. Although A-only and B-only channels may perform well, many of these single detector channels exhibit drifts over the mission for colder

85 scenes. This is especially apparent in time series of cold scene observations (deep convective clouds) by comparison to similar time series derived from IASI on METOP-1. In addition, we avoid any channels with detector noise above 0.5K NEDT (for a 250K scene). As discussed below in more detail, we also avoid all short wave AIRS channels, meaning channels past 2000 cm⁻¹ for our final trend measurements, since we find that the short wave is drifting slightly.

3 Radiance(BT)/Brightness Temperature Anomalies

90 3.1 Clear Selection

The new AIRS Level-1c radiance product (Aumann et al., 2020) is used in this work rather than the standard L1b product. The Level-1c product provides single-footprint radiance estimates for channels in Level-1b that are not functional, or are extremely noisy. Even high quality Level-1b channels can sometimes "pop" or experience radiation hits that invalidate the measurement. In these extremely rare cases the Level-1c algorithm substitutes an estimated radiance using a principal-components approach.

95 These corrections are rare enough that they have no effect on the long-term trends under study in this work. Level-1c also includes some channels (in-between detector arrays) that do not exist.

More importantly for this work, the radiances in Level-1c have been corrected for small drifts in the channel center frequencies. These drifts are small, but are large enough to have some minor impact on radiance trends. We emphasize that the channels selected for the anomaly retrievals are all valid Level-1b channels, and most have undergone no corrections other the redication there are also been as found for small drifts are small.

- 100 than adjusting the radiances back to a fixed frequency scale.
 - AIRS L1c clear scenes are primarily detected using a uniformity filter. (Throughout this paper the term "scene" refers to a single AIRS nominal 12-by-12 km. footprint or field-of-view.) The BT of each AIRS ocean scene is subtracted from the BT of each of its 8 neighbors for two window channels at 819.3 and 961.1 cm⁻¹. A scene is <u>initially</u> deemed clear only if the absolute value of all of these differences, averaged over the two channels, is less than 0.4K. The selected scenes are matched
- to ERA-I model fields and a simulated clear BT for the 961.1 cm⁻¹ channel is computed using a stand-alone version of the AIRS radiative transfer algorithm (Strow et al., 2003) called SARTA, implemented using HITRAN 2008 line parameters. If the difference between the observed and computed clear scene BT values is more than \pm 4K the scene is discarded from the clear list. This test mostly removes colder scenes made up of very uniform marine boundary layer stratus clouds. The clear yield and mean zonal radiances are quite insensitive to the exact value of this threshold. The uniformity test is not performed
- 110 on the first and last of the 135 along-track scans in each AIRS granule since they do not have 8 neighbors and we wanted to avoid cross-granule processing. The total number of clear scenes is limited to ~2040,000 daily clear scenes by randomly sub-setting the detected clear scenes, however this daily limit is almost never reached. In this work we only use descending node observations in order to avoid solar and nonLTE contributions to the AIRS radiances in the short wave. After subsetting for descending (ocean) only the total number of clear scenes detected is ~10,000 per day.
- 115 The 4K (observed minus computed) BT test removes ~20% of the scenes detected with the uniformity filter. A map of these deleted scenes very clearly shows that they are almost all located along the west coasts of the Americas and Africa, where marine bounday layer stratus clouds commonly occur. The (observed minus computed) BT values for the 1231 cm⁻¹ window channel have a nearly gaussian distribution with a width of ~0.6K. Note that this distribution of biases is well separated from the 4K cutoff used to remove marine bounday layer stratus clouds.
- Another important characteristic of this clear subset is the stability of the observing times. If the mean observing time changes during this 16-year time period, trends in the SST could be confused with the diurnal cycle of the SST. Due to the high stability of the AQUA orbit, this is not an issue. The short term day-to-day variations in the mean clear subset times can vary by several hours. In addition, there is a seasonal variation of several hours in the clear subset. But, these variation are extremely stable, and the total linear drift of the clear subset over the 16-year observing period, for any given latitude bin in the tropics, is ~20 ± 40 (2-σ) seconds per year, effectively zero.
 - 4

All observing parameters, on a footprint basis, are saved, such as satellite viewing zenith angle and noise (converted to BT units).

The In addition, the ERA-I model parameters (temperature, H₂O, and O₃ profiles, and surface temperature with a spatial resolution of approximately 80 km on 60 levels in the vertical from the surface up to 0.1 hPa-) are matched to each clear scene
are also and saved along with their associated simulated L1c radiances. This allows our processing to use simulated rather than observed radiances for testing. The ERA-I profiles are also used to compute the anomaly Jacobians , as discussed below. All of our processing can be tested using the simulated radiances rather than the observed radiances as input used in the retrievals, and are discussed in detail in Sects. 4.3 and 5.

3.2 Clear Scene Characteristics

Figure 1 illustrates the density and location of the clear ocean dataset, averaged over 2012. Retrievals are only performed on



Figure 1. Density of AIRS clear ocean scenes for calendar years 2012.

135

140

zonally averaged data, which translates into ~44/25 observations/day at $-50^{\circ}/+50^{\circ}$ latitude respectively, with a maximum of 200 observations per day at -0.5° latitude. The non-uniform nature of this sampling should be kept in mind when examining temperature, H₂O, or O₃ trends in that this data set is not necessarily representative of global/zonal climate trends. However, we do assume that the minor gas anomaly trends we are retrieving are uniformly mixed over year+multi-year time scales. Our anomaly retrieval results show uniform mixing is generally quite accurate over even 16-day time scales.

The ERA-I model fields, suitably converted to temperature, humidity, and ozone profile anomalies are used to compute the BT profile Jacobians needed for the anomaly retrievals. In principle we could have performed 1D-var retrievals on each 16-day averaged BT spectrum in each latitude zone. However, ERA-I is so accurate, that is not necessary. Figure 2 illustrates the accuracy of ERA-I for this dataset , where we show by plotting the (observed - ERA-I) based simulated BT bias for 28.4°N.

= 0 (Sept. 2002). The time series fit are discussed in Sect. 3.3 simulated BT used our SARTA RTA, which has a default value of 385 ppm for CO₂. This CO₂ value is matched in the observations by comparing to AIRS observations for the time period centered around June 2008 when the nominal global CO₂ amount was 385 ppm. The window regions (800-1000 cm⁻¹) exhibit a bias of ~-0.5K, which is quite small and likely some combination of instrument bias, evaporative cooling of the ocean surface

relative to the ERA-I SST, incorrect ERA-I water vapor column affecting the H₂O continuum, and some cloud-contamination. 150 Sampling errors may contribute to the larger biases in the water region beyond 1300 cm⁻¹. A zoom of the bias in the bottom panel of Fig. 2 highlights the low bias in the 700-750 cm⁻¹ region which is sensitive to tropospheric CO_2 , with a mean of ~0.2-0.3K.



Figure 2. Top Panel: (AIRS - ERA-I simulated) BT bias for 28.4°N in Sept. 2002. The for a time period centered around June 2008 when the global CO₂ feature in amount was ~385 ppm. Bottom Panel: Zoom of top panel. Shows that the mid-wave region near 700-760 cm⁻¹, which is due our use of a constant most sensitive to CO₂ concentration, has a mean bias of 385 ppm for the ERA-I simulations~0.2-0.3K, rather than and a value single-footprint standard deviation of ~370 ppm more appropriate for this date0.3K. Also shown is the AIRS NEDT, which is barely smaller than the bias standard deviation.

The CO₂ tropospheric sounding region in the 700-780 cm⁻¹ region shows a sharp increase in the bias of about 0.6K (depending on channel) that is due to the fact that the single-footprint standard deviation of the bias is also shown in Fig. 155 2, bottom panel along with the average AIRS NEDT for these footprints. The ERA-I bias standard deviation is barely larger than the AIRS noise in this spectral region, indicating that ERA-I based radiance simulations used a constant amount of CO₂ set at 385 ppm. This feature will go through zero bias and become negative by approximately the same BT amount at the end of our 16-year test period temperatures in the mid-troposphere track the AIRS observations very closely with a standard deviation considerably smaller than the AIRS noise. This makes a strong case for the accuracy of the BT Jacobians computed 160 from ERA-I temperature fields.

Figure 3 shows the linear trend for the clear dataset averaged over $\pm 50^{\circ}$ latitude.



Figure 3. Mean BT trends (a_1 in Eq 1) averaged over $\pm 50^{\circ}$ in Δ BT/year units. The 2- σ uncertainty shown has been corrected for serial correlations in the BT time series. Channels used in the anomaly retrievals are denoted in red, and the BT trend uncertainty is in yellow.

Mean radiance trends for $\pm 50^{\circ}$ in Δ BT/year units. The 2- σ uncertainty shown is lag-1 corrected for time series correlations. Channels used in the anomaly retrievals are denoted in red, and the BT trend uncertainty is in yellow.

165

These BT trends prominently exhibit the growth in CO_2 in the tropospheric channels from 700 to 750 cm⁻¹, which results in a negative change in the observed BT since increasing CO_2 shifts the emission to higher and therefore colder regions of the atmosphere. The growth in CO_2 in the stratospheric channels (a positive BT change) below 700 cm⁻¹ is roughly cancelled by cooling in the stratosphere. All window channels exhibit warming, with larger values in the shortwave past 2450 cm⁻¹. The non-uniform spatial sampling of these clear scenes precludes any general statements about climate warming, although for these observations we clearly see warmingsurface warming in the 800-1250 cm⁻¹ region, if the AIRS radiometry is stable. In

170 these observations we clearly see warmingsurface warming in the 800-1250 cm⁻¹ region, if the AIRS radiometry is stable. In addition, the effects of much stronger water vapor absorption in the long wave compared to the short wave windows makes definitive inter-comparisons of the BT trends complicated, which is addressed below by doing retrievals on these data.

3.3 Construction of Anomalies

The clear scene radiance subset is sorted into 40 equivalent area latitude bins that cover the full -90° to 90° latitude range and are averaged over every 16 days. This results in a data set for the first 16-years of AIRS that has the size a size of (40 x 2645 x 365denoting) latitude bins, AIRS L1c channels, and the total number of 16-day averages. The following time-series function was fit to these averaged radiances, $r_{obs}(t)$, for each latitude and AIRS L1c channel,

$$r_{\text{fit}}(t) = r_o + a_1 t + \sum_{\underline{i} \in \underline{1}} {}^4 c_i \sin(2\pi n t + \phi_i) \tag{1}$$

where t is AIRS mission times in years, r_o is a constant, c_i are the amplitudes of the season cycle and three harmonics, and the 180 ϕ_i are their associated phases. The function models periodic variations in the radiances using an annual term and the first three harmonics. At 28°N, for example, the annual amplitude relative to the mean radiance, c_1/r_o , has a median value (taken over channel) of 4.2%. The median amplitudes of the three harmonics terms, c_2, c_3, c_4 relative to r_o is 0.32%, 0.45% and 0.23% respectively, all with 2- σ uncertainties of ~0.05%.

For the retrievals performed we created the radiance anomaly by inserting the linear trend into the residuals of the above 185 time series fits.

$r_a(t) = r_{\rm obs} - r(t) + a_1 t.$

The linear term The linear trends a_1 represents the linear-part of the minor gas signals we aim to measure. The a_1 terms are included in the anomaly time-series fits since they are a useful way to quickly understand AIRS trends and because this allows us to measure a more correct lag-1 auto-correlation of the time series noise, which is used to estimate corrections for the uncertainties in the time-series parameters using the approach popularized in (Santer et al., 2000). The radiances anomalies for simple diagnostic purposes, and are not used directly in the anomaly retrievals.

190

200

210

The radiance anomalies, $r_a(t)$ are formed by removing the constant r_o , and the sinusoidal terms in Eq. 1, from the observed radiance time series $r_{obs}(t)$. This can be expressed as

$$r_{a}(t) = r_{obs} - \left(r_{o} + \sum_{i=1}^{4} c_{i} \sin(2\pi nt + \phi_{i})\right)$$
(2)

195 The radiance anomalies $r_a(t)$ were converted to brightness temperature units using

$$y(\nu) \equiv BT_a(\nu, t) = \frac{r_a(\nu, t)}{\frac{\partial r(\nu)}{\partial BT(\nu)}}.$$
(3)

for our retrievals. The 40 x 2645 x 365 array of BT_a vectors are the retrieval inputs y as denoted in the retrieval formation discussed in Sect. 4.

All uncertainties quoted in this paper derived from time series are for 95% uncertainty levels and are lag-1 corrected for correlations in the time series residuals.

The anomaly BT time series mean BT spectra and their standard deviations are shown in Figure 4 for the 28.4°N latitude bin. The BT anomaly is set to zero at the mission start, therefore the mean BT in the CO_2 region channels sensitive to tropospheric CO_2 between 700 and 750 cm⁻¹ is -0.5Kgiven that it changes by by ~-1K, which then increases by ~1K during the mission. The standard deviation indicates that the SST (and H₂O continuum) vary by ~1K during this time period. Some of this is likely

205 due to changes in sampling from day to day . The upper-tropospheric water vapor and ERA-I errors in SST and column H_2O . Upper-tropospheric water vapor, which dominates the spectral region between 1350-1615 cm⁻¹, has the highest variability, which is expected due to both the high temporal variability of water vapor in time, and our non-uniform sampling.

An example radiance BT anomaly for the 710.141 cm⁻¹ channel is shown in Fig. 5, for the same latitude bin. This channel is heavily influence influenced by the CO_2 growth, so the AIRS observed trends are becoming more negative, although there is considerable noise, again due to weather and sampling. For comparison we also plot the ERA-I simulated BT anomaly, which



Figure 4. Mean and standard deviation of the AIRS BT anomalies for the zonal bin centered at 28.3°N.

does not contain the CO_2 growth, since it is set to a fixed value of 385 ppm in the simulations. The difference between these two BT anomalies will primarily be due to CO_2 growth, and is shown in black. Note that since the ERA-I tracks the atmospheric state quite accurately most of the time-series "noise" is removed. This helps lend credence to our use of the ERA-I model fields for Jacobian evaluation.



Figure 5. Sample AIRS observed and ERA-I simulated BT anomalies for the zonal bin centered at 28.3° N for the AIRS channel centered at 710.14 cm^{-1} . The differences in the AIRS and ERA-I anomalies is are plotted in black. Note, this difference anomaly is not used in the anomaly retrievals.

215 4 Retrieval Methodology

4.1 Approach

Geophysical retrievals are derived from BT anomalies the BT spectral anomalies $y(\nu)$, defined in Eq. 3. Using standard retrieval notation the atmospheric state x is derived from the observations y, $y(\nu) \equiv BT_a(\nu,t)$, by minimizing the cost function J

$$J = (y - F(x))^T \mathbf{S}_{\epsilon}^{-1} (y - F(x)) + (x - x_a)^T \mathbf{R}(x) (x - x_a)$$
(4)

where S_{ϵ} is a diagonal observation error covariance matrix containing the square of the BT noise, K are the anomaly Jacobians, and R is a regularization matrix. The retrieved atmospheric state *x* (the geophysical anomalies) are given by

$$x = x_a + (\mathbf{K}^T \mathbf{S}_{\epsilon}^{-1} \mathbf{K} + \mathbf{R})^{-1} (\mathbf{K}^T \mathbf{S}_{\epsilon}^{-1} (y - F(x_n)), \text{ where } \mathbf{R} = \mathbf{S}_a^{-1} + \alpha \mathbf{L}^T \mathbf{L}_{\underline{\cdot}},$$
(5)

where S_{α} is the a-priori covariance matrix, and αL is an empirical regularization constraint using Tikhonov L1-type derivative smoothing. This retrieval approach is standard Optimal Estimation (OE) (Rodgers, 1976) enhanced to include both covariance and empirical Tikhonov regularization in **R** (Steck, 2002). Forward model uncertainty is not included in the measurement error

- 225 and empirical Tikhonov regularization in R (Steck, 2002). Forward model uncertainty is not included in the measurement error covariance. The mathematical approach is very similar to the author's single-footprint AIRS retrieval algorithm (DeSouza-Machado et al., 2)
 - *

230

245

A-priori estimates for $x_a(t) \equiv 0$ for T(z), O₃(z), H₂O(z) and T_{SST} were set to zero. Two approaches were used for the minor gas a-priori estimates. The first approach set $x_a(t) = x_a(t-1)$ where $x_a(t=0) = 0$ for the minor gases, iteratively increasing the a-priori gas amount in time based on the previous 16-day retrieval.

Another approach used the known growth rates in the minor gases (from ESRL) by setting $x_a(t) = g * (t - t_o)$ for the apriori minor gas amount, where g is the nominal yearly growth rate for each gas from the NOAA ESRL atmospheric gas trends. For both approaches we set the a-priori covariance to g times one year, the yearly variation in that gas. Nearly identical results are obtained if we used g times five years. The iterative approach for setting the minor gas a-priori produces noisier retrieval anomalies. However, if our retrievals are averaged over $\pm 50^{\circ}$ latitude, both approaches produced identical differences

retrieval anomalies. However, if our retrievals are averaged over $\pm 50^{\circ}$ latitude, both approaches produced identical differences compared to in-situ measurements, including error uncertainties. The figures shown here use the a-priori ramp from the ESRL data, although the figures for the iterative ramp are only distinguishable from what is shown for single zonal retrievals (such as the Mauna Loa and Cape Grim comparisons).

The retrieval approach is standard Optimal Estimation (OE) (Rodgers, 1976) enhanced to include both covariance and
 empirical Tikhonov regularization in R (Steck, 2002). Here we use Tikhonov L1-type derivative smoothing. Forward model uncertainty is not included in the measurement error covariance.

The temperature, H_2O , and O_3 profile retrievals use 20 atmospheric layers, selected from the AIRS standard 100-layer pressure grid (Strow et al., 2003) by accumulating five of the standard AIRS layers at a time. The lowest layer is about 1.5 km thick, with increasingly wider layers as you go higher in the atmosphere. This layering scheme allows more layers then than degrees-of-freedom (DOFs) although it does limit retrievals in the upper-stratosphere. We wish to minimize our sensitivity to the upper-stratosphere since our comparisons to in-situ measurements are made in the troposphere. Consequently we removed all channels peaking above 10 hPa.

Most of the regularization in the retrieval comes from the Tikhonov terms, since we do not want to invoke climatology too strongly for a climate level measurement. Appendix B discusses the profile retrievals, and simulations of these retrievals, in

- 250 more detail. In summary, after experimentation with Tikhonov regularization we added some a-priori covariance uncertainties in temperature and water vapor of 2.5K and 60% respectively. These are extremely large values for a-priori covariance uncertainty terms improved simulated retrievals and uncertainties compared to the anomaly variations. For example, the retrieved 400 HPa temperature anomalies shown in Fig. B3 are all less then ± 3K, indicating that the temperature a-priori covariance uncertainty is providing very minimal regularization. This means that almost all of the retrieved temperature 255 variability is coming from the data, and is not damped by the a-priori estimates, a desirable situation for the measurement
- of climate trends. These a-priori covariance uncertainty terms did, however, improve the profile trends generated from these retrievals by in simulation by a slight amount, 3-10%, and thus were retained in our retrieval.

The observation error covariances were derived by averaging the noisefrom each observations contributing to the averaged anomaly being retrieved(noise) are the mean AIRS NEDT for each channel, averaged over 16-days, and then divided by the

- square root of N, the number of scenes averaged. Originally a fixed value of 0.01K observation noise was used, but we found that this noise value depressed the CO_2 anomaly retrievals as they grew in size over time. This problem disappeared once we switched to the true measurement noise values, which are in the range of noise equivalent brightness temperature (NEDT) equal to 0.004K for long wave CO_2 channels from 700-750 cm⁻¹, about 0.001K in window regions between 800-1250 cm⁻¹, and 0.001 to 0.002K in the water bands, band that covers the 1300-1615 cm⁻¹ spectra region. These are externely low noise
- 265 values, which help explain why the anomaly retrievals have a relatively high number of degrees-of-freedom.
 As stated earlier, the profile Jacobians used the ERA-I profiles, which were converted to anomaly profiles for each pressure

layer. The minor gas Jacobians were computed using our pseudo line-by-line kCARTA radiative transfer algorithm (Strow et al., 1998; DeSouza-Machado et al., 2019). kCARTA allows for extremely accurate Jacobian calculations, including analytic trace gas and temperature Jacobians. Initial retrievals used a fixed value for the minor-gas Jacobians. However, given the large

- 270 increase in the minor gases (10% for CO_2), we determined that the minor-gas Jacobians need to be updated as the gas amounts increase. Therefore we used finite-difference Jacobians, computed using the minor gas amount retrieved from the previous time-step during the anomaly retrievals (or from the gas amount estimated using NOAA ESRL in-situ gas amount data). The minor gas profiles used in the Jacobian calculations are from (Anderson et al., 1986). The CO_2 profile is essentially constant in ppm until you reach the highest atmospheric layer.
- There exists a weak dependence of these retrievals on the ERA-I model fields since we use the ERA-I model fields for the temperature, H₂O, and O₃ profiles in the profile Jacobians, **K**. While we could retrieve the atmospheric profiles from the full radiance at each time step and latitude zone, ERA-I is so accurate we do not believe this is needed. Moreover, we *do* retrieve the profile anomaly in each step, so we are not dependent on the actual Section 5.4 discusses potential errors introduced by using ERA-I model values for the retrieval other than a weak dependence via the Jacobians. In the results shown later the
- 280 impact of attempting to use the ERA-I profile anomalies , rather than retrieved profile anomalies, is presented. The end result

show conclusively that we did need to retrieve the true profile anomalies, using ERA-I profile anomalies, for example, results in increased errors in the trace gas retrievals for Jacobian evaluation, where they are shown to be extremely small and unimportant.

The direct retrieval of anomalies from the BT anomaly spectra represents a very different approach than normally used in infrared remote sounding. Although the mathematical approach is the same as in single-fooprint retrievals (DeSouza-Machado et al., 2018), the often troublesome problem of static measurement and RTA bias errors is largely removed here since instrument calibration and/or absolute RTA biases do not appear in the retrieval process.

4.2 Channel Selection

As discussed in Section 2, only channels that remain A+B throughout the mission are used, noting that the designation A+B does not apply to detectors in the M-11 and M-12 long wave detector arrays. Initial retrievals showed that the AIRS short wave detectors are drifting slightly, so these channels are also excluded from the anomaly fits (except for demonstration tests as discussed below). Unfortunately, the use of only A+B detectors greatly restricts the number of available channels in the important long wave CO₂ temperature sounding region from 710-780 cm⁻¹, where many channels are either A-only or B-only. It is important to weight these channels relatively strongly in the retrieval minimization. Since we also wish to de-emphasize stratospheric contributions to the minor-gas rates only every 5th channel from 650-720 cm⁻¹ was included in the retrieval. In

addition, any channels in this range with Jacobians that peaked above 10 hPa were excluded.

All channels in the M-5 array were excluded since they have relatively poor radiometric stability (as will be shown later). Several window channels that are sensitive to CFC11 were excluded, although many channels sensitive to CFC12 were included, and CFC12 trends were retrieved. Many H₂O channels were included, since they are mostly A+B and have been stable

300

throughout the mission. After some experimentation, four channels sensitive to N_2O were also excluded since they appear to be behaving significantly out-of-family. Three of these channels are located near the end of the M-4c array, which also exhibits some anomalous frequency shifting behavior (Aumann et al., 2020).

305

A total of 470 channels remained after this pruning process. These channels are nicely distributed throughout the AIRS spectrum and are easily sufficient for 1D-var retrievals. The nominal number of DOFs for tropical scenes for this channel set are ~6 ozone DOFs, ~8 temperature DOFs, and 12 H₂O DOFs. The larger number of H₂O DOFs is likely due to the large number of H₂O channels used (321 out of 470 channels).

The overall sensitivity of the anomaly retrievals to CO_2 is shown is shown in Figure 6 where the mean CO_2 Jacobian, averaged over all channels, is plotted. The CO_2 sensitivity peaks around 400 hPa, and drops to near zero at the surface. There is some dependence on stratospheric CO_2 , but stratospheric CO_2 trends, especially in the lower stratosphere, should track the

310 tropospheric trends, albeit with growth rates that are slightly influenced by previous years due to age-of-air. This figure also shows the mean CO_2 Jacobian is if all channels below 700 cm⁻¹ are removed (all sensitive to the stratosphere). Retrieval tests using this these restrictions are discussed later.

4.3 Construction of Jacobians

The relatively high accuracy of ERA-I temperature fields was highlighted previously in Fig. 5 which plots the time dependence
of the bias between the observed and simulated BT for this channel. This bias, in black, has very little variability (other than the smooth decrease due to increasing CO₂) compared to either the observed or simulated BT values due to the high accuracy of the ERA-I temperature profiles. This is not unexpected in a reanalysis product that assimilates a wide range of in-situ measurements (radiosondes) and satellite measurements (microwave and infrared sounders, including AIRS). In principal we could use the AIRS Level-2 atmospheric state for generating the Jacobians for the anomaly retrievals. However, for the large-scale averaging used in this work errors introduced by the relatively large ERA-I spatial grid compared to AIRS are minimized.

- Moreover ERA-I is constrained by a large number of instruments and in-situ measurements for the temperature profile. Monthly mean ERA-I observation minus analysis differences for radiosonde temperatures are below 0.2K throughout the troposphere, rising to 0.3K in the lower stratosphere (Simmons et al., 2014). We note that the statistical accuracy of the AIRS Level-2 algorithm is mainly verified by inter-comparisons with ECMWF forecast/analysis fields (Susskind et al., 2014), which
- are likely even more stable in a reanalysis product. The AIRS Level-2 retrieved temperature and H₂O global biases relative to ECMWF are very small, well below 0.5K for temperature and 5% for water vapor.

In principle we could have performed 1D-var retrievals on each 16-day averaged BT spectrum in each latitude zone, but given the relatively small biases between ERA-I and AIRS shown in Fig. 2 retrievals will produce minimal improvements to the ERA-I fields. Note that the ERA-I bias in the 700-750 cm⁻¹ region with the most sensitivity to tropospheric CO_2 is only

in the 0-0.5K range. Moreover, 1D-var retrievals using AIRS will also be limited by uncertainties in the AIRS radiometric calibration, which is estimated to be in the 0.2K range (Pagano and Broberg, 2016).

More importantly, since we are only retrieving anomalies, highly accurate Jacobians are unnecessary since the BT variations in the anomalies are so small, especially when applied to trends. A quantitative assessment of errors in our measured anomaly trends from using using ERA-I for Jacobian evaluations is presented in Sections 5.4 and 5.7.

335 4.4 Temperature and Minor Gas Jacobian Co-linearity

A non-standard "correction" is made to the minor gas retrievals that attempts to correct for the co-linearity of the temperature and minor gas Jacobians. COWe demonstrate that this new approach clearly removes un-physical variability in the CO₂ anomaly retreivals. Co-linearity of the temperature and minor gas Jacobians makes it difficult for the retrieval to separate temperature profile variations from variations in CO₂retrievals using hyperspectral infrared are difficult because of

this co-linearity. As stated above, the , CH_4 , and N_2O . Usually this is managed by constraining the retrievals with accurate a priori estimates that have small enough covariances to allow some separation of T(z) and CO_2 variability. Kulawik et.al. Kulawik et al. (2010) discuss this problem in the context of CO_2 retrievals using the NASA EOS-AQUA TES instrument, where they describe the selection of constraints as a way to "determine the partitioning of shared degrees of freedom between CO_2 and temperature".

- 345 Here we take a different approach based on the fact that we have highly accurate simulated anomalies computed from the ERA-I model fieldsare matched one-to-one with each clear observations, followed by a RTAcomputation of a simulated radiance, using constant amounts of . The simulated anomalies are derived using our SARTA radiative transfer algorithm (RTA) and were generated using constant values for the minor gases throughout the mission. Since the 16-year time period. Except for the minor gas signatures, the ERA-I spectral anomalies are very similar to to observed AIRS anomalies, we can
- 350 partially evaluate the observed anomalies since both AIRS calibration errors and RTA errors are largely removed when forming the anomalies. Fig. 5 shows the excellent agreement between the observed and simulated BT anomalies for the 710.14 cm⁻¹ channel. The only major difference in these anomalies is the downward drift in the observations primarily due to the growth of CO₂. Note that almost all the high frequency variability in the observed and simulated anomalies is removed when taking their difference, shown in black in Fig. 5, indicating that the ERA-I temperature fields match the AIRS observations very closely.
- 355 Given that the ERA-I spectral anomalies are very similar to observed anomalies we can largely determine the effect of the Jacobian co-linearities by retrieving the minor gas amounts on the observed CO₂ anomaly retrievals by retrieving a (fictitious, or non-existing) CO₂ anomaly from the ERA-I simulated anomalies that contain no minor gas variations. We illustrate this with CO₂ retrievals in Figure 7BT anomalies using an identical retrieval algorithm. Since the simulated anomalies have a constant value for each minor gas the variations in the retrieved CO₂ (or other minor gases) is a measure of the inability of the retrieval 360 to separate the minor gas anomalies from the temperature profile.



Figure 6. Mean of CO_2 Jacobians for all channels used in the anomaly retrievals, and the same if all channels below 700 cm⁻¹ (stratospheric channels) are excluded.

While the long-term trends are not very sensitive to this removal of co-linearity, the lowered noisethis approach affords is extremely useful for detecting and understanding shifts in the AIRS radiometry due to various instrument shutdowns that occurred over the mission. Figure 7 illustrates this process for; (1) a single latitude bin near -55° lat (with a width of \sim 4° latitude) in the left hand panel and (2) the average of 30 latitude bins covering ± 50° latitude in the right hand panel. The



Figure 7. Illustration of "noise" removal in the CO₂ anomaly retrievals by subtracting the CO₂ retrieved from ERA-I simulations from the observed CO₂ retrievals. Left: -55° latitude CO₂ retrieval. Right: $\pm 50^{\circ}$ latitude average CO₂ retrieval.

- 365 yellow curve (labelled Simulated) is the retrieved CO₂ anomaly derived from the ERA-I simulated anomalies. Although close to zero in the mean, this retrieved CO₂ anomaly varies considerably by up to ± 15 ppm. The red curve (labelled AIRS Raw) shows the CO₂ anomaly retrieved from the AIRS observations, which has similar variability superimposed on a linear ramp of ~2 ppm/year. Our adjusted observed CO₂ anomaly is generated by subtracting the simulated from the observerd CO₂. This is shown in blue (labelled AIRS Adjusted), where we see that most of the "noise" has been removed resulting in a very smooth
- CO_2 anomaly curve.

The right hand panel of Fig. 7 shows similar results but using the average of all latitude bins between $\pm 50^{\circ}$ latitude. The co-linearity of the temperature and CO₂ Jacobians apparently changes randomly enough with latitude that the simulated CO₂ has far less variability than in the left panel. The utility of this approach is nicely illustrated by examining the dip of about 7 ppm in the "Simulated" CO₂ retrieval in early 2010 for the $\pm 50^{\circ}$ latitude bin. This dip is also visible in the observed anomaly

curve (AIRS Raw). The "Simulated" CO₂ anomaly is subtracted from the "AIRS Raw" curve to obtain the final observed CO₂

375

380

anomaly (AIRS adjusted) and it is quite evident that the dip in early 2010 has cancelled out, as desired.

The above adjustments to the CO_2 anomaly retrievals have little effect on estimates of AIRS stability over 16 years, as outlined later in Sect. 5.4, although it does increase the statistical uncertainty in the AIRS BT trends by a factor 2.4. More importantly, the application of these adjustments greatly reduces the apparent noise in the derived CO_2 trends, making the detection of instrument shifts in the AIRS BT time series much more sensitive.

5 Anomaly Retrievals

5.1 AIRS Events

Evaluation of the anomaly retrievals requires some knowledge of the AIRS mission events. Table 1 summarizes the major events during the AIRS mission that had thermal consequences for either the spectrometer or the focal plane arrays. While
most of these events were minor, recent measurements of the AIRS frequency shifts (Aumann et al., 2020) highlight that these

events are associated with small shifts in the AIRS frequency scale. These shifts are indicative of very small movements of the detectors relative to the instrument spectrometer axis, and therefore they could also and could, for example, slightly alter the detector's view of the blackbody and cold scene. Any small non-uniformities in these calibration looks could affect the absolute radiometrybecause viewing angles to the Earth and cold scenes might change every so slightly. We will refer to these events during discussions of the anomaly retrieval results.

390

Table 1. Summary of AIRS events that had a thermal impact on either the spectrometer, the focal plane, or both.

Date Event		Event
	10/29/03	AQUA shutdown lasting for several weeks (solar flare)
	01/09/10	Single event upset, focal plane temperature cycling
	03/28/14	Single event upset, small focal plane cooler variation
	09/25/16	Single event upset, one cooler restart

5.2 Truth Anomalies

The retrieved minor gas anomalies are compared to the NOAA Earth System Research Laboratories (ESRL) monthly mean data derived from in-situ measurements (Tans and Keeling). We chose the ESRL Mauna Loa, Cape Grim, and Global mean data for CO₂, N₂O, and CH₄. Monthly anomalies for these in-situ datasets were computed using the same methods used to compute the BT anomalies for consistency. We focus mainly on the global CO₂ ESRL anomalies since they are derived from a

395

wide geographical range and sites and carefully merged to avoid local sources. The N₂O ESRL anomalies provide information on AIRS channels in the 1250 -1310 cm⁻¹ region that are distinct from the main CO₂ channels below 780 cm⁻¹. (There are also strong N₂O channels in the short wave band of AIRS.) The CH₄ anomalies mostly probe AIRS channels from 1230 to 1360 cm^{-1} . There is some concern that CH₄ anomaly trends may have more spatial variability than CO₂ and N₂O, however we find good overall agreement with the ESRL global CH₄ trends, and CH₄ provides some sensitivity to channels that overlap with

400

N₂O, but extend a bit further into the water band.

We focus mostly on the use of CO_2 for AIRS stability estimations since CO_2 is so well measured and has the largest BT signal in the AIRS spectrum (relative to N₂O and CH₄). In addition, the N₂O and CH₄ spectra overlap strongly in the AIRS BT spectrum, possibly introducing some retrieval uncertainty relative to CO₂. Absolute errors in the ESRL CO₂ data are estimated

405 to be ~0.2 ppm (https://www.esrl.noaa.gov/gmd/ccl/ccl uncertainties co2.html), with yearly growth rate uncertainties of ~0.07 ppm/year (https://www.esrl.noaa.gov/gmd/ccgg/trends/gl_gr.html). Anomaly growth rate errors averaged over 16 years are likely much lower since yearly sampling errors should diminish over time. Moreover, most absolute errors will not be applicable to the CO₂ anomaly, which is a relative measurement. Therefore it is difficult to definitively estimate the ESRL anomaly trend uncertainty. If the yearly growth rate uncertainties of 0.07 ppm/year are random, then the average of 16 of these growth rates

would be 0.018 ppm/year, which corresponds to a percentage uncertainty of 0.8% in the anomaly trend. 410

Estimates for N_2O and CH_4 anomaly trend uncertainties using the ESRL stated uncertainties in yearly growth rates, and assuming these are random errors each year, are 3.5% and 2.4%. These larger uncertainties, and the smaller total impact of these two gases on the AIRS BT anomalies, suggest that the best estimates for AIRS stability are likely derived from the CO_2 anomalies.

415 5.3 Short Wave Trends



Figure 8. Anomaly fit residual, averaged over all 365 16-day time steps for $\pm 30^{\circ}$ latitude. The L1c fill channels have no L1b counterparts and are simulated in the production of L1c. Note the offset in the short wave.

Most of the anomaly retrievals performed here only included AIRS channels in located below 1615 cm⁻¹, avoiding the short wave channels in the 2181 to 2665 cm⁻¹ region. Early retrievals showed that the AIRS short wave channels exhibit a positive trend compared to the longer wave channels. Moreover, anomaly fits to just the short wave channels return SST trends that are significantly larger than both the long wave channels and both the ERA-I (OSTIA) and OISST SST products.

420

The behavior of the AIRS short wave channel relative to the long wave is easily seen in the anomaly retrieval fit residuals. Figure 8 shows the mean value (taken over the 365 16-day time steps for \pm 30° latitude) for the residuals. All AIRS L1c channels are plotted, which includes many bad channels, and channels that do not exist but are filled during L1c creation (Aumann et al., 2020). The channels selected for the anomaly fits (see Sect. 4.2) are shown in red circles. The fit residuals for channels used in these retrievals are almost all well below 0.02K. However, the short wave channels show anomalies inconsistent with the long wave of up to ~0.07K in the window channels past 2450 cm⁻¹.

425

The anomaly retrievals can respond to drifts/offsets in the AIRS radiances by retrieving geophysical variables (CO_2 , temperature, etc.) that vary incorrectly in time. Alternatively, un-physical changes in the radiances could also be reflected in larger non-zero fit residuals. This could happen when the forward model Jacobians cannot model time-dependent radiance errors, especially for jumps in the radiometric calibration that happen due to AIRS events (shutdowns). One way to examine this



Figure 9. Linear trends in the anomaly fit residuals, averaged over all 365 16-day time steps for $\pm 30^{\circ}$ latitude. Note the linear trend in the short wave in these fit residuals. Also shown is the trend difference (ERA-I SST - AIRS SST) for these data.

430 possibility is to look for any remaining trends in the anomaly fit residuals. These are shown for the same data set used in Fig. 8 in Fig. 9. Most of the channels used in the anomaly fits have residual slopes below 0.002K/year, although careful examination of the residual time series for particular channels can exhibit jumps associated with AIRS shutdowns.



Figure 10. Retrieved CO_2 anomalies compared to ESRL global in-situ data. The CO_2 anomaly difference between AIRS and ESRL is shown in yellow. The magenta curve is that difference converted into BT units.

The main observation in Fig. 9 is a clear positive trend in the short wave relative to the longer wave channels used in the retrievals. The (AIRS - ERA) SST trend plotted as a solid horizontal line in this figure (discussed in Sect. {sec:sst}5.7) shows that the AIRS short wave trends are more different from the ERA-I SST trends than the long wave channels. Most of the

short wave channels, including those in the mid-troposphere, exhibit positive trends relative to the long wave, except for some channels that are peaking very high in the stratosphere, below 10 hPa, that are marked in gray.

Consequently, unless otherwise noted, all the remaining results presented here <u>use avoid the avoid short</u> wave channels, and use the channel set (470 channels) denoted in these figures.

440 5.4 CO₂ Anomaly Retrievals

Retrieved CO₂ anomalies compared to ESRL global in-situ data. The CO₂ anomaly difference between AIRS and ESRL is shown in yellow. The magenta curve is that difference converted into BT units.

Figure 10 shows the retrieved CO_2 anomalies averaged over $\pm 50^{\circ}$ latitude in blue and the ESRL global anomaly product in red. The correspondence over time is excellent. The AIRS minus ESRL anomaly differences are shown in yellow.



Figure 11. Observed linear trend in the AIRS CO₂ anomalies versus latitude, compared to NOAA ESRL Mauna Loa (MLO), ESRL Cape Grim (GCRIM), and the ESRL global CO₂ product trends (black line).

In order to convert the variation in the gas anomalies to an equivalent AIRS BT anomaly temperature we computed anomaly retrievals with the observed AIRS BT anomaly spectra modified by a 0.01K/year ramp, for all channels. This 0.01K/year ramp is divided by the resulting changes in the CO₂ anomaly linear trends (ppm/year) to obtain the sensitivity of the retrieval to a trend in the AIRS radiances, in K/ppm. For CO₂ this sensitivity is 0.073-0.073K/ppm. This is about 2X larger than the largest column Jacobians in the AIRS spectra, which have a value of ~0.030K/ppm. This is not unexpected, since the CO₂ column measurement is partially a relative measurement, especially for weak CO₂ channels in the window region where the absolute BT errors are mostly accounted for by (incorrect) adjustments in the SST that minimize the effect of the 0.01K/year applied ramp. It is also possible that the temperature profile could also adjust to minimize sensitivity of the ramp on the CO₂ ppm values. In addition, this sensitivity estimate assumes all AIRS channels are drifting, which is clearly an approximation given the results shown here.

Table 2. Slope of the (AIRS - ESRL) CO₂ anomalies in ppm/year units.

Data Set	Mean Trend Difference	Uncertainty in Trend
CO_2	(ppm/year)	(ppm/year)
Global	0.032	0.012
Mauna Loa	0.033	0.023
Cape Grim	0.056	0.020

The magenta curve in Fig. 10 is the (AIRS minus ESRL) anomaly differences converted to BT units using the 0.073-0.073K/ppm sensitivity factor. This curve has been slightly smoothed for clarity. The right-hand side vertical axis shows the variations in this curve in BT units. Most of the BT variability is within ± 0.05K, however a transition in BT in late 2003 is larger. This larger transition is likely due to the Nov 2003 shutdown of the AQUA spacecraft. The AIRS channel center frequencies were shifted due to this shutdown (Strow et al., 2006) and were subsequently corrected in the AIRS L1c product (Aumann et al., 2020; Manning et al., 2019). In addition, as reported in (Strow et al., 2006) interference fringes in the AIRS entrance filters shifted after the Nov. 2003 AQUA shutdown because AIRS was restarted at a slightly different spectrometer temperature. The

fringes change the AIRS spectral response functions, which has not yet been corrected in the AIRS L1c product radiances. Figure 11 illustrates the differences between the AIRS and ESRL CO₂ linear growth rates. The growth rate were computed

using rates for both our CO₂ retrievals and the ESRL CO₂ time series were computed by re-using the fitting function in Eq.

465 1where the input data is the CO₂ ppm anomaly rather than a radiance. Figure 11 shows the a_1 term in this equation, but now applied to the retrieved CO₂ anomalies, ie.

$$CO_{2}(t) = CO_{2}(t=0) + b_{1}t + \sum_{i=1}^{4} d_{i}\sin(2\pi nt + \phi_{i})$$
(6)

where b_1 are the CO₂ trends in ppm/year. Later this equation will be used to fit the N₂O, CH₄, and SST anomalies, instead of the CO₂ anomalies as shown here.

Figure 11 plots the fitted values for the AIRS growth rates (the b1 term in Eq. 6), computed as a function of latitude. The CO2 growth rates are not completely uniform from year-to-year, so Eq. 1-6 cannot perfectly fit the trend data. However, it provides a convenient metric for inter-comparing these two CO2 anomalies. Note that the error bars shown for AIRS are slightly over-estimated because of the fact that Eq. 1-6 does not perfectly fit the slightly non-linear anomaly curve. The error estimates are for 95% confidence intervals and have been they have been corrected for serial correlations in the anomaly time series using the popular lag-1 auto-correlation corrected using the auto-correlation approach detailed in (Santer et al., 2000).

The Mauna Loa and Cape Grim growth rates are also shown, also derived using Eq. 16, as is the ESRL global rate, indicated by the dark black horizontal line. If the 16-year in-situ rates indeed have an estimated error of 0.018 ppm/year (assuming the 0.07 ppm/year uncertainties in the ESRL rates are random), then AIRS is in close agreement with ESRL averaged over latitude. The latitude dependence of the AIRS derived rates appear to have clear latitudinal dependencies, with lower rates near

480 the ITCZ and higher rates in regions of descending air. We do not examine this latitude dependence in this work, not only is it small, it could also be related to small inaccuracies in our retrieval algorithm.

CO. Tost

CO_2 lest	Medil Helid Difference	Oncertainty in Trend
	(K/Decade)	(K/Decade)
Global		
Baseline	-0.023	0.009
Baseline (no CO ₂ adjustment)	+0.019	0.022
No Strat	-0.034	0.008
No Cov Reg.	-0.043	0.009
No ν Cal.	-0.059	0.010
Shortwave Only	+0.070	0.009
ERA-I T(z)	+0.060	0.035
Mauna Loa		
Baseline	-0.024	0.017
Cape Grim		
Baseline	-0.040	0.020

Table 3. Slope of the (AIRS - ESRL) CO_2 anomalies in K/Decade units. Trend differences for various modifications of our retrieval algorithm are shown, see the text for details. Note that Baseline is the algorithm configuration detailed in the text and used for inter-comparisons.

Mean Trend Difference Uncertainty in Trend

Since the CO₂ linear growth rate measurements are not sensitive to year-to-year variability in the CO₂ anomaly, we instead use the (AIRS - ESRL) global anomaly differences shown in Fig. 10 to quantify the AIRS stability. Any linear trend differences between the AIRS and ESRL CO₂ in Fig. 10 are quantified by fitting the (AIRS - ESRL) CO₂ anomaly differences to Eq. 16. Table 2 summarizes any trend in AIRS relative to ESRL by tabulating the $a_{T}b_{L}$ terms from the fit for the ESRL global, Mauna Loa, and Cape Grim sites. The uncertainties are as before, 95% confidence intervals corrected for lag-1 auto-correlations. As one might expect, the global trends agree the best, and Cape Grim the worst. The higher errors for Cape Grim may be related to our clear subset having fewer samples at -40° latitude relative to the 20° latitude zone occupied by Mauna Loa. These mean differences are extremely small, corresponding, for global, to 1.5 ± 0.6% trend differences.

- Table 3 shows the conversion of the CO₂ ppm trend differences to equivalent BT differences using the 0.073-0.073 K/ppm sensitivity conversion. The baseline entry, first line of the table, represents the final configuration for the anomaly retrievals and represents is our best estimate for the true differences between the ESRL and AIRS CO₂ anomaly trends, namely -0.023 ± 0.009 K/decade. This is an exceedingly small trend difference. While suggesting that AIRS is extremely stable, for channels sensitive to CO₂ and temperature, systematic errors may be larger than the differences reported here. Our optimistic estimate
- 495 of the ESRL global anomaly uncertainty trend uncertainty discussed in Section 5.2, 0.8%, is equivalent to 0.017 ppm/yearor 0.27 ppm over 16 years. From Table tableco2ppmthe. The AIRS minus ESRL global trend differences are difference shown in Table 2 is about 2X times this optimistic larger than this estimate for the ESRL uncertainty. This translates, in Table 3 to an

estimate for AIRS stability of -0.023 ± 0.009 and slightly larger than the statistical uncertainty in this trend difference. In BT units, this potential uncertainty in the ESRL global CO₂ anomaly trend is -0.012 K/decadeDecade.

500

505

In addition to the possible 0.8% The sensitivity of these results to uncertainties in the Jacobians are derived from the second partials derivative of BT as follows.

$$M_{unc} = \frac{\partial}{\partial X} \left(\frac{\partial BT}{\partial Y} \right) \times X_{unc} \times Y_{meas} = \left(\frac{\partial^2 BT}{\partial X \partial Y} \right) \times X_{unc} \times Y_{meas}$$
(7)

where M is the quantity being measured (here, CO₂ anomalies and trends), X_{unc} is the uncertainty in the ESRL global trends, our AIRS anomaly retrieval could be in error if profile variables used to compute the Jacobians, and Y_{meas} is either the maximum anomaly or the mean trend we measured for Y. These are quantified in Table 4.

The first entry accounts for errors in $\partial BT/\partial CO_2$ due to uncertainties in the CO₂ Jacobians are inaccuratespectroscopy. The HITRAN database (Gordon et al., 2017) reports uncertainties in the CO₂ line strengths of 1-2%. These uncertainties would translate into the same percentage error in the Jacobians. In addition, atmospheric spectra are sensitive to line widths, line shape, line mixing, often at temperatures that are not measured in laboratory spectra. Characterizing the combination of these

- 510 errors is essentially impossible, so here we assume a 1% uncertainty in the CO₂ Jacobians, using the line strength uncertainty only. The maximum CO₂ anomaly error occurs at the end of the time series when the CO₂ anomaly is highest (35 ppm). Therefore the max anomaly error is $1\% \times 35$ ppm = 0.35 ppm. Using our retrieval sensitivity of -0.073K/ppm, this translates into an effect max error in the BT anomaly error of 0.026K. Dividing this anomaly uncertainty by the 16 year time period under study gives a trend uncertainty due to CO₂ spectroscopy errors of 0.016 K/decade as shown in Table 4. This value is slightly
- 515 larger than the statistical uncertainty in the baseline CO₂ trend shown in Table 3, and slightly smaller than the derived trend differences versus ESRL CO₂.

The second entry in Table 4 lists estimated uncertainties in the CO₂ anomalies and trends (converted to BT units) that could arise due to errors in the ERA-I temperature profile. The second parital derivative was computed with finite differences using a fixed temperature offset for all levels and then summed over all levels, a worst case scenario. The mean of these second order

520 derivatives, taken over the retrieval channels in the ~700-750 cm⁻¹ spectral region that has high sensitive to CO₂, represents an effective scalar value for $\partial^2(BT)/(\partial X \partial Y)$ in Eq. 7. This term is multiplied by an assumed uncertainty in the ERA-I temperature of 0.5K and by the maximum anomaly value of 35 ppm to obtain a maximum uncertainty of 0.0035K in the CO₂ anomaly. The maximum effect on the CO₂ trend is again this value divided by 16 years giving an uncertainty of 2.2 × 10^{-3} K/decade, an insignificant uncertainty. Note that our assumed uncertainty of 0.5K is higher than ERA-I error estimates

```
525 discussed in Section 4.3.
```

Clearly the estimated 1% uncertainty in the CO₂ spectroscopy is the dominant source of error in our CO₂ retrievals. If the ESRL 0.8% uncertainty is combined in quadrature with the 1% HITRAN uncertainty, a total minimum expected uncertainty in the CO₂ anomaly trends is 1.3%. This translates to a BT uncertainty of 0.02 K/decade, close to our <u>derived</u> mean trend difference between AIRS and ESRL based on the CO₂ anomaly measurements. This may be a more accurate uncertainty

530 estimate for this measurement rather than the 0.009 <u>K/Decade</u> statistical uncertainty derived from fitting the AIRS minus ESRL anomalies.

Table 3

Table 4. Anomaly and trend error estimates for CO_2 and SST due to uncertainties in BT Jacobians via their second derivatives with respect to possible ERA-I uncertainties. As noted, the maximum effect on the CO_2 anomalies would be at the time of the largest anomaly, which is at the end of our time series in Aug. 2019. See the text for details.

Jacobian	Sensitivity	Uncertainty	Max Effect on Anomaly	Effect on Trend
$\frac{\partial BT}{\partial CO_{2}}$	$rac{\partial^2 BT}{\partial C O_2^2} \sim$	<u>1% CO₂ Spectroscopy</u>	0.026K (in Aug. 2019)	0.016 K/decade
	$rac{\partial^2 BT}{\partial T_{air}\partial CO_2}$ ~	0.5K T profile	0.0035K (in Aug. 2019)	2.2×10^{-3} K/decade
$rac{\partial BT}{\partial T_{SST}} \sim$	$rac{\partial^2 BT}{\partial T_{SST}^2}$	0.5K T _{SST}	4.0×10 ⁻⁴	9.6×10 ⁻⁵ K/decade
	$rac{\partial^2 BT}{\partial T_{air}\partial T_{SST}} \sim$	0.5K T profile	<u>8×10⁻⁴K</u>	1.8×10^{-4} K/decade
	$rac{\partial^2 BT}{\partial T_{H_2O}\partial T_{SST}} \sim$	10% H ₂ O column	<u>0.02K</u>	4.5×10^{-3} K/decade

The second entry in Table 3 lists the mean trend difference and its uncertainty if the adjustment for co-linearity discussed in Sect. 4.4 is not applied, which leads to a larger trend uncertainty by a factor of 2.4. The resulting trend difference is somewhat

535 smaller, but with a different sign. The baseline retrievals with and without the co-linear CO₂ adjustments do not quite overlap within their respective 2σ uncertainties, missing statistical agreement by 0.013 K/Decade, which is relatively small. However, based on the discussion in Sect. 4.4 we believe that the application of the co-linear CO₂ adjustment improves the accuracy of the AIRS CO₂ anomaly.

Table 3 also shows the results of a number of fit testing the sensitivity of the retrievals to various retrieval alternatives. The "No Strat" entry removed all channels that primarily sense the stratosphere by removing all channels below 700 cm⁻¹. Fig. 6 shows how this modifies the mean CO₂ Jacobian used in the retrieval, essentially removing all sensitivity to CO₂ above 60 hPa. Unfortunately channels above 700 cm⁻¹ have some residual sensitivity to CO₂ in the stratosphere, and removing channels below 700 cm⁻¹ may make it more difficult to properly minimize the retrieval residuals for some channels above 700 cm⁻¹. If S_a is completely removed, removing a-priori profile regularization, the CO₂ anomaly trend difference increases by a factor of two. Removing the L1c frequency calibration adjustments increases the anomaly trend differences by nearly a factor of three, and changes their sign. If only short wave channels are fit (excluding channels that peak above 10 hPA, and some channels sensitive to both carbon monoxide), the mean trend differences are more than three times larger than the baseline, again with a

sign change.

The last test, labeled "ERA-I T(z)", examines the impact of performing simultaneous retrievals of temperature profiles while retrieving the CO_2 anomalies by using the ERA-I temperature profiles anomalies instead of fitting for them. This test increased the anomaly differences between AIRS and ESRL by almost a factor of three, with a significant increase in the uncertainty of

the trend, giving 0.35 K/decade instead of close to 0.009 K/decade for the baseline.

23



Figure 12. Retrieved N_2O anomalies compared to ESRL global in-situ data. The N_2O anomaly difference between AIRS and ESRL is shown in yellow. The magenta curve is that difference converted into BT units.

Table 3 also shows the Mauna Lao anomaly difference, which is close to the global result, although accompanied by a higher uncertainty of 0.017 K/decade compared to the 0.009 K/decade for the global anomaly. Cape Grim anomaly differences are almost two times higher than the global trend differences, but this is not surprising given the much lower number of observations at that latitude.

555

The retrieved AIRS global CO_2 anomalies did detect a small seasonal pattern in the anomaly for latitudes above 40° N of with an amplitude of ~0.5 ppm. This is due to the residual of the seasonal cycle of CO_2 that is not completely removed when constructing the BT anomalies.

Table 5. Slope of the (AIRS - ESRL) N₂O anomalies in K/Decade units.

Data Set	Mean Trend Difference	Uncertainty in Trend
N_2O	(K/Decade)	(K/Decade)
Global	-0.141	0.012
Mauna Loa	-0.200	0.030
Cape Grim	-0.080	0.033

Note that radiometric shifts or drifts in the AIRS BT time series could be either reflected in incorrect geophysical trends, or partially buried in the anomaly fit residuals. The high quality of the anomaly retrievals for CO_2 and the small fit residuals for CO_2 channels strongly suggest that the AIRS blackbody is extremely stable, at least for long and mid wave A+B channels. The SST retrievals discussed later reinforce this conclusion. However, we do see evidence of radiometric shifts due to discrete AIRS events (especially for N_2O and CH_4) that might be amenable to correction. Future work will include careful examination of

⁵⁶⁵ both the anomaly retrievals and their residuals, likely in an iterative fashion, in order to determine what channels are responsible for unphysical shifts in the anomaly products.

5.5 N₂O Anomaly Retrievals

The N₂O retrieved anomaly time series is shown in Fig. 12 and primarily senses the 1240-1325 cm⁻¹ spectral region. Clearly the observed N₂O anomaly is growing slightly faster than the ESRL values. The N₂O anomalies are converted to equivalent
BT variations just as for CO₂, but with a derived sensitivity of 0.140 K/ppb. Table 5 tabulates the derived trend for the (AIRS minus ESRL) anomaly by fitting the difference to Eq. 46, and then converting to BT units.

The trend differences here are much larger than for CO_2 . Examination of either the AIRS minus ESRL anomalies in ppb, or their equivalent in BT units (left hand y-axis) suggest that two unphysical steps might be present in the time series, one in mid-2005 and another on one in mid-to-late 2010. Unfortunately, these steps do not closely coincide with AIRS events, possibly appearing more than one year after the Nov 2003 event and and slightly less than one year after the Ian. 2010 event

575 possibly appearing more than one year after the Nov. 2003 event and and slightly less than one year after the Jan. 2010 event.



Figure 13. Retrieved CH₄ anomalies compared to ESRL global in-situ data. The CH₄ anomaly difference between AIRS and ESRL is shown in yellow. The magenta curve is that difference converted into BT units.

Table 6. Slope of the (AIRS - ESRL) CH₄ anomalies in K/Decade units.

Data Set	Mean Trend Difference	Uncertainty in Trend
CH_4	(K/Decade)	(K/Decade)
Global	-0.107	0.024
Mauna Loa	-0.062	0.039
Cape Grim	-0.100	0.037

To illustrate the effect of these two discrete shifts on the anomaly trend differences we empirically introduce a step in our retrieved N₂O time series of -0.6 ppb on July 1, 2005 and another step on Jan. 18, 2010 of -0.5 ppb. The trend difference between this empirically modified time series and ESRL, in BT units, becomes -0.022 ± 0.009 K/decade, very similar to the CO₂ trend differences. The main point of this exercise is to illustrate that just two two-small discrete radiometric shifts could be responsible for the higher trend differences between AIRS and ESRL for N₂O. More work is needed to map these discrete

580

non-physical events in the retrieved N_2O anomaly time series back into steps in the AIRS BT time series. The hope is that careful examination of the anomaly time series residuals during this process would highlight specific channels (or cluster of channels) that are behaving non-physically.

5.6 CH₄ Anomaly Retrievals

- The CH₄ retrieved anomalies have some similarities to the N₂O anomalies, since the spectra of both gases occur in the same general spectral region. The CH₄ the region of sensitivity is ~1210-1380 cm⁻¹. Figure 13 shows the CH₄ results using the same approach as for CO₂ and N₂O. The ppb to BT conversion for CH₄ was measured to be 0.023 K/ppb, significantly lower than for CO₂ or N₂O, although total BT trend due to CH₄ is only marginally lower than CO₂ and N₂O.
- The high variability of atmospheric CH₄ growth is well known, as can be seen in the ESRL curve in Fig. 13. The AIRS derived anomalies follow that variable growth rate quite nicely overall. It should be noted that the ESRL CH₄ curve is more variable than CO₂ and N₂O, and may be less uniform globally, making CH₄ a less ideal gas for testing AIRS stability. However, the AIRS minus ESRL anomaly differences are valuable in that they, like N₂O, highlight discrete jumps that can often be identified with AIRS events, such as late 2003 (biggest jump), early 2010, and possibly in early 2014. The positive jump in the CH₄ anomaly difference near March 2014 also coincides with a jump in the N₂O anomaly difference, both taking place after
- 595 the March 2014 event. However, this apparent jump seems to fade within one year for both gases. We believe this might be caused by AIRS frequency shifts that occurred in the M-4a and M-4c detector modules after this event. Those frequency shifts appeared to disappear within one year, and at present they are not corrected for in the AIRS L1c product.

Table 6 lists the trend differences between AIRS and ESRL for CH_4 , showing trends differences that similar to those for N_2O , presumably since both gases occur in the same spectral region.

600 5.7 SST Retrievals

The SST anomaly retrievals are compared to the ERA-I supplied SST (mostly OSTIA) and to NOAA's OISST operational SST product. Although both of these SST products are tied to the ARGO floating buoy network, they are gridded SST products using interpolation derived from satellite data such as AVHRR.

A recent study (Fiedler et al., 2019) compared various SST products to the buoy network and found differences for OSTIA of 1.1 mK/year, and 7.8 mK/Year for OISST. This establishes a rough estimate of the differences in these products when evaluating them relative to our retrieved SST anomalies.

Figure 14 plots time series of our retrieved SST anomaly and the co-located ERA-I SST (mostly OSTIA) anomaly, averaged over $\pm 30^{\circ}$ latitude, where these products are expected to be most accurate since most buoy's are located in the tropics. The AIRS SST trend derived from this time series is 0.096 \pm 0.046 K/decade. The AIRS and ERA-I 16-day averaged anomalies

610 agree very closely, their difference is shown in black. A zoom of the AIRS minus ERA-I SST anomaly is shown in Fig. 15 to highlight their differences. Steps in these differences are possibly evident near the end of 2003 and especially near the end of September 2016 when AIRS had an a cooler-restart.



Figure 14. Tropical (\pm 30°) SST anomalies retrieved from AIRS compared to the ERA-I anomalies. The black curve is the difference between the AIRS and ERA-I anomalies.



Figure 15. Zoom of Fig. 14 that highlights the shift in the AIRS - ERA-I SST anomaly presumably due to the AIRS Sept. 25, 2016 cooler restart. A small shift is also seen at the date of the Nov. 2003 AQUA shutdown.

Table 7 summarizes the AIRS minus (ERA-I and OISST) anomaly trend differences, computed using Eq. 16. The trend differences are quite small for both SST products. The (AIRS minus ERA-I) trend has the same magnitude as the trend derived

615 using CO_2 , but with the opposite sign. Overlap of the CO_2 and ERA-I SST within their stated uncertainty estimates is missed by 0.01K/decade, which is very small. The CO_2 and OISST trend estimates miss overlap by slightly more, 0.02K/decade. However, this overlap difference is small compared to the differences between OISST and the buoy network reported by (Fiedler et al., 2019).

Overall the excellent agreement of these two extremely independent assessments (CO_2 versus SST) to within 0.02K/decade is very encouraging given the complexity of the CO_2 measurement and the uncertainties in the SST product trends.

Comparisons between AIRS-derived SST and ERA-I or OISST products will contain biases due to time aliasing between the AIRS observations and daily means used in the SST products. Although these time dependent biases can have random and seasonal variations of several hours the observed linear drift in the AIRS local observing time over the 16-year observation period was less than one minute/year, far too small to introduce any drifts in the AIRS SST relative to the ERA-I or OISST daily averages

625 <u>daily averages</u>.

620

Uncertainties in the SST anomaly retrievals due to our use of ERA-I fields for the evaluation of the SST Jacobians were estimated using the same approach for the CO₂ anomaly retrievals. The BT Jacobians (dBT/dSST) for channels sensitive to SST depend on accurate values for the SST itself, the air temperature profile, and most importantly the H₂O profile, especially in the lower troposphere. We computed the partial derivatives of the BT Jacobians with respect to all three of these variabiles.

- 630 again using finite differences and a constant offset for the air temperature profile and constant percentage offsets for the H₂O profile. The partial derivatives were averaged for all AIRS channels used in our retrievals in the 800-1235 cm⁻¹ region that is sensitive to surface temperature. The uncertainties assumed in the ERA model fields (X_{unc} in Eq. 7) are listed in column three of Table 4 and are likely higher than the estimated uncertainties summarized in Section 4.3. The uncertainties in the BT Jacobians are then multiplied by Y_{meas} in Eq. 7 which is either 0.4 K (the maximum SST anomaly, see Fig. 14), or
- 635 0.0096K/year (our retrieved trend in SST).

The results shown in columns four and five of Table 4 clearly indicate that using ERA profile fields for estimated BT surface temperature Jacobian is extremely accurate. The highest uncertainties are due to H_2O , but even these are far below the statistical uncertainties shown in Table 7.

Table 7. Slope of the (AIRS - (ERA/OISST)) SST anomaly differences.

Data Set	Mean Trend Difference	Uncertainty in Trend
	(K/Decade)	(K/Decade)
(AIRS - ERA-I)	0.022	0.012
(AIRS - OISST)	0.034	0.021

Aumann (Aumann et al., 2019) recently compared the 1231 cm^{-1} AIRS channel trends to RTGSST, a precursor to OISST. 640 He used a statistical approach to remove trends in water vapor that affect the 1231 cm^{-1} channel radiances, which he concedes could introduce artifacts if there is a shift in the mean vertical distribution of water vapor. Our approach does not contain this limitation in principle, although we have not carefully examined the retrieved water vapor trends, mainly because there is no truth for comparison. An intercomparison of our results to his are not strictly possible since we used different SST products for truth and our SST anomalies used many channels. However, the trend of the 1231 cm⁻¹ channel in our retrievals can be

- 645 derived by adding the slope of our fit residual for the 1231 cm⁻¹ channel (-0.7 mK/year) to our derived SST trends for ERA-I and OISST. Using Aumann's units of mK/year, the result is a trend of 1.5 mK/year and 2.7mK/year for ERA-I and OISST respectively, with respective uncertainties of 1.2 and 2.1 mK/year. These two trends compare favorably with Aumman's night trend for 1231 cm⁻¹ of +2.9 \pm 0.4 mK/year. It is interesting that our OISST trend differences agrees more closely with his RTGSST trend difference since these two data sets have similar heritage. Of course the extremely low statistical errors reported
- 650 by Aumann do not allow overlap of these two results, but that is not necessarily expected since we use different SST products. Agreement for AIRS radiometric trends at the several mK/year level for at least a single channel should be considered quite remarkable.

We also derived AIRS minus (ERA-I, OISST) SST trend differences using AIRS short wave only anomaly retrievals. For tropical latitudes, ± 30°, the (AIRS - ERA-I) trend is 0.078 ± 0.040 K/decade and 0.065 ± 0.09 K/decade for OISST. These represent significantly higher trend than observed using long and mid wave channels only. The trend difference between (AIRS long wave minus AIRS short wave) anomaly fits is -0.058 ± 0.026 K/decade, clearly indicating the short wave positive drift relative to the long wave.

The latitude dependence of the AIRS derived SST trends versus ERA-I and OISST may eventually help determine the source of some of these differences. Figure 16 shows these trends between \pm 60° latitude. The uncertainties in these trends are

- 660 ~0.005K/year, but are not shown since these uncertainties are primarily geophysical in nature (how linear is the SST trend) and affect each SST product identically. Agreement is quite good among all products in the northern hemisphere, while OISST is systematically lower than AIRS and ERA-I in the southern hemisphere. Also shown are the AIRS SST trends using only the short wave channels (gray curve), which are always higher than the long wave AIRS trends except at the highest latitudes and near the equator.
- 665 Unfortunately, the AIRS Level 2 retrieval algorithm only uses short wave channels for surface temperature retrievals (Susskind et al., 2014). A recent inter-comparison of surface temperature trends from the AIRS Level 2 retrievals to three established surface temperature climate products (Susskind et al., 2019) concluded that the AIRS surface temperature trends were 0.24 K/decade, slightly higher than GISTEMP's (Hansen et al., 2010) value of 0.22 K/decade, and significantly higher than the HadCRUT4 (Morice et al., 2012) and Cowtan and Way (Cowtan et al., 2015) values of 0.17 and 0.19 K/decade respectively.
- The results presented here conclude that the AIRS short wave channels are drifting positive by about 0.058 K/decade relative to the long wave channels, which appear to be in extremely good agreement with established SST climate products as discussed above. If we subtract this 0.058 K/decade AIRS short wave drift from the the AIRS 0.24 K/decade trend presented in (Susskind et al., 2019) we obtain a corrected AIRS trend of 0.18 K/decade, much more in line with the HadCRUT4 and C+W values. In this case GISTEMP is now the only outlier. A more straightforward way to validate the reported AIRS Level 2 surface trends

675 reported by (Susskind et al., 2019) would be to directly compare them to other SST products such as OISST, but unfortunately this was not part of the (Susskind et al., 2019) analysis.



Figure 16. Latitude dependence of the linear trend in the AIRS retrieved SST, OISST, and ERA-I SST. Also shown are the SST trends when only the AIRS short wave channels are used to compute the anomalies.

5.8 CFC12 Retrieval

All anomaly retrievals presented here included CFC12 retrievals. Although these are not used for quantitative assessments of AIRS radiometric stability, the retrieved CFC12 anomaly is shown in Fig. 17 for completeness. Excellent agreement is found between the AIRS observed CFC12 and the ESRL Northern Hemisphere measurements (ESRL). The linear trends derived from these two curves are -2.94 ± 0.04 ppt/year for AIRS, and -2.93 ± 0.02 ppt/year for ESRL, nearly perfect agreement. These results give us confidence that the SST retrievals have not been compromised by CFC12 contamination, since there are a number of channels sensitive to both. Note that the trend of ~40 ppt of CFC12 derived here from AIRS is equivalent to only ~0.11K in BT!

685 6 Retrieval BT Breakouts and Residuals

The anomaly fit residuals provide a wealth of information on the behavior of each AIRS channel versus time. As stated earlier, unphysical shifts in the AIRS radiance time series can be reflected in either the retrieved geophysical anomalies or in the fit residuals. Jumps in the fit residuals will generally take place when the shifted radiances cannot be "adjusted away" by the BT Jacobians, which require a reasonably accurate physical response to radiance jumps. We believe that the anomaly retrieval approach presented here will allow objective corrections to AIRS radiances, especially for radiance jumps that can be tied to

690



Figure 17. AIRS CFC12 retrieved anomaly compared to the NOAA ESRL Northern Hemisphere anomaly. Note that a 40 ppt trend in CFC12 corresponds to about 0.11K in brightness temperature for the channel with the highest CFC12 Jacobian.

instrument events. The excellent agreement between the CO_2 and SST anomalies and in-situ data strongly suggests that the AIRS blackbody is very stable, which is key to climate-level trend measurements.

There are several likely causes for some of the differences seen here between our observed anomalies and the N₂O and CH₄ truth anomalies from ESRL. Shifts in the frequency calibration of AIRS (Strow et al., 2006; Manning et al., 2019) have largely been removed in the AIRS L1c product, although some transient shifts in the AIRS M-4a and M-4c arrays (that cover N₂O and CH₄ channels) have not yet been corrected in L1c (see (Aumann et al., 2020)). The AIRS frequency shifts imply that detector views of the blackbody and cold scene targets have also shifted during the mission. While these shifts are very small, radiometric drifts/shifts could arise from these focal plane movements if the blackbody and cold scene targets are not perfectly uniform. As mentioned in Sec. 5.4, shifts of interference fringes in some of the AIRS entrance filters when AQUA

Aqua was restarted in Nov. 2003 may also contribute to the observed anomaly shifts. These fringe shifts have been modeled by the authors and future work may include modification of AIRS radiances before Nov. 2003 to remove the effects of these small shifts in the instrument spectral response function.

Here we present several views of the AIRS anomaly fit fits and their residuals as examples on how future work might proceed to potentially correct the AIRS radiances for small remaining radiometric drifts/shifts.

705 6.1 Retrieved Anomalies in BT Units

First to provide some context, Figure 18 shows the contribution of the various geophysical trends to the observed BT anomalies for channels sensitive to different geophysical variables. This is done by multiplying the BT Jacobian for some particular geophysical variable by its retrieved anomaly over time. For illustration purposed we averaged the trends over the latitude bins from $\pm 50^{\circ}$ latitude.



Figure 18. Contribution to the observed BT anomalies caused by the retrieved geophysical anomalies. These are simply the BT Jacobian multiplied by the time-dependent retrieved geophysical anomalies. The BT anomalies in the bottom panel are multiplied by the sum, over all layers, of the retrieved profile anomalies.

The upper panel shows that the retrieved CO_2 anomaly translates into a BT trend for the 722.1 cm⁻¹ channel of more than -1K. Channels very sensitive to the retrieved CH_4 and N_2O anomalies have BT trends that are lower than CO_2 . The anomaly for a channel sensitive to SST in this panel has an upward trend due to increasing SST values, but these are quite small compared to the minor gas trends.

The bottom panel of Fig. 18 plots the BT anomalies due to the retrieved temperature, H_2O , and O_3 anomalies. The profile anomalies have been summed over all levels for this figure. The same channel chosen to illustrate the BT anomaly due to the CO₂ anomaly, 722.1 cm⁻¹, is also used to illustrate the contribution of the temperature anomaly. The BT trend for the 722.1 cm⁻¹ channel due to the temperature anomaly is far smaller than for CO₂, is slightly noiser, and has a small positive trend that mostly occurs after 2014. This would be expected since there is also a positive trend for SST with the same general time dependence. The BT trend due to the retrieved H₂O anomaly is plotted for the 1418.6 cm⁻¹ channel sensitive to mid-tropospheric H₂O. This

720 BT anomaly moves in the opposite direction to the BT anomaly due to temperature, which is expected since on a large scale increasing temperatures raise H₂O amounts, which leads to lower BT values.

Spectra illustrating how the various geophysical anomalies contribute to the BT anomalies are constructed by multiplying the BT Jacobians times the 16-year mean of the retrieved geophysical anomalies. Since these are computed quantities, all channels can be included. These are plotted in Fig. 19, where we separate the geophysical contributions just as in Fig. 18. If

725 the trends are linear in time, the 16-year mean anomalies represent the anomalies for year eight. Divided these by eight gives the nominal BT trend in K/year.

This figure clearly shows that CO_2 dominates the changes in the longwave region, as expected. The N₂O and CH₄ BT anomalies are conentrated in the 1230-1400 cm⁻¹ region with significant overlap, which is largely separable in the retrieval.



Figure 19. Contribution to the observed BT anomalies caused by the retrieved geophysical anomalies. These are simply the BT Jacobian multiplied by the mean, over time, of the 16-year record of geophysical anomalies. The BT anomalies in the bottom panel are multiplied by the sum, over all layers, of the retrieved profile anomalies.

On this scale the BT changes due to SST are quite small. In the lower panel the temperature, H_2O , and O_3 BT anomaly 730 trends are derived from the sum of the profile Jacobians over all layers. In many regions of the spectrum the temperature and H₂O BT anomaly trends are dominant, an indication that our anomaly retrievals successfully accounted for variability in those parameters. BT trends in the channels sensitive to tropospheric temperature (700-750 cm⁻¹) are in the range of 0.01-0.02K/year (after dividing the plotted mean anomaly by eight), nominally consistent with global warming during this period.

The H₂O greenhouse effect is clearly seen in the bottom panel of Fig. 19. The increased emission in the water band 735 (1200-1615 cm⁻¹) due to higher atmospheric temperatures is largely negated by the decrease in emission due to increasing amounts of H_2O , which shifts the emission in any given channel to higher altitudes where the temperature is lower.

Also note that channels sensitive to stratospheric temperatures in the $650-690 \text{ cm}^{-1}$ region have a negative trend, indicating stratospheric cooling. This is also an expected result for global warming, but great care should be taken in using this data set for general conclusions since the sampling is non-uniform, and the trend standard deviation (Fig. 20) is almost 2X larger than

740 the trend.

> For completeness the standard deviation of the nominal linear anomaly trends shown in Fig. 19 are plotted in Fig. 20 using the same breakouts of geophysical anomalies. The CO_2 BT anomaly trend maximum standard deviation of ~0.1 K near 730 cm^{-1} is only slightly higher than the standard deviation expected if it was solely due to a linear trend in CO₂, 0.08K. The air temperature stratospheric standard deviation is large, as previously noted, presumably due to the effects of the quasi-biennial

745 oscillation (QBO) and possibly ENSO variability. The variability due to air temperature and H₂O produces standard deviations in the water region (1250-1615 cm⁻¹) that is generally larger than variability due to trends in CH_4 and N_2O , but apparently our retrieval successfully removeds those interferences. Note the relatively high O₃ variability, which we do retrieve but have not



Figure 20. Standard deviation for the contribution to the observed BT anomalies caused by the mean retrieved geophysical anomalies shown in Fig. 19.

examined carefully. It is important to remember that these are anomaly standard deviations, so they do not include seasonal variability.

750 6.2 Anomaly BT Residuals

765

The anomaly fits shown above are summed and then subtracted from the observed BT anomalies to obtain the the BT anomaly fit residuals. Any trends in these residuals can also be examined to search for channels that changed characteristics during the 16-year time period.

Figure 21 shows the BT anomaly fit residual slopes for A+B, A-only, and B-only channels separately. Most of the A+B
channels shown, all of which were used in the anomaly retrievals, are within ± 0.004 K/year of zero. While a large number of A-only and B-only channel are in agreement, there are a number of cases where they exhibit significant slopes (trends) that are not in agreement with the A+B channels. Module M-05 channels near 1100 cm⁻¹ are clearly drifting differently than the other channels (we did not use any A+B M-05 channels in the retrievals since they are also in error). Module M-08 channels near 851 cm⁻¹ show a clear separation between A+B channels and A-only, B-only. Clearly, the opposite sign of the A-only versus B-only drifts are largely cancelled when A+B channels are used. Since the SST retrievals are quite good, and because the surface channels near 1200 cm⁻¹ agree with the A+B channels, we conclude that the A-only and B-only drifts are real, and possibly due to drifts, or offsets, in the exact part of the blackbody and/or cold target scenes observed by these detectors.

Since the N_2O retrieved anomalies exhibit some small unphysical behaviors, we examine the fit residuals for the 24 channels (used in the retrievals) that are most sensitive to N_2O . Visual inspection of these channel's residual time series clearly indicated that 12 of them had easily identifiable features due to AIRS events. Figure 22 shows three different averages of these residual

time series; (a) 12 good channels, with no strong evidence of AIRS events, (b) 12 bad channels which clearly exhibit jumps at



Figure 21. Slope of the AIRS anomaly residuals separated by A+B (Fit channels), A-Only, and B-Only. This illustrates trends in the A-only and B-only channels relative to A+B channels in some modules. The A-only and B-only channels were not used in the fitting, so they are not strictly residuals, but Observed - Computed differences.

the time of AIRS events, and (c) the mean time series for all 24 channels used in the anomaly fits. We see that the good channel mean (blue) is very flat, with a slight indication of a jump near the Nov. 2003 event. The bad channel curve (red) shows a large jump near Nov. 2003, possibly some longer-term drifts, and a feature in March 2014 that seems to last for 1 to 1 1/2 years. This last feature can change sign depending on which bad channel is observed, making it very likely that this is due to the

M-4a/M-4c frequency calibration shift that is not yet corrected in the L1c product.

770

A new set of anomaly retrievals were produced, but with the 12 bad N_2O channels removed. When compared with the ESRL N_2O anomalies, this change produced slightly better agreement with ESRL after Nov. 2013. The slope of the (AIRS - ESRL) anomaly difference curve was reduced from -0.141 K/Decade (as reported in Table 5) to -0.113 K/Decade, a slight

- 775 improvement. This drift relative to ESRL reduces to -0.069 K/Decade if anomaly data before Nov. 2013 is ignored. This illustrates that improvements to the AIRS products can be improved achieved by removing channels with residuals that have non-physical jumps. If the Nov. 2013 radiometric jumps can be removed (whether due to frequency shifts, fringe shifts, or pure radiometric jumps) even higher stability is possible. However, one could presently begin the AIRS time series, say on Jan. 1, 2004 and retain a stability approximately 2X better than climate trends.
- These results illustrate a simple case for how the anomaly fit residuals can be used to improve AIRS trend products. In this work we have not looked for non-physical jumps in the retrieved temperature, H_2O , and O_3 profile anomalies. These products likely exhibit some of these behaviors and need to be included in any comprehensive study to further improve the AIRS radiance stability. Some sort of iterative approach will likely be needed in order to ensure that these small remaining radiometric jumps become undetectable in both the retrieved anomalies and in the anomaly residuals.



Figure 22. Anomaly fit residual time series for various combinations of 24 channels sensitive to N_2O in the long wave. The bad N_2O channels have easily visible jumps at times corresponding to AIRS hardware events.

785 7 Conclusions

A framework for establishing stability of the AIRS radiances has been introduced that uses retrievals of minor gas and SST trends from BT anomaly spectra. Extremely good agreement between retrieved CO_2 trends (or anomalies) and in-situ trends from NOAA ESRL illustrates that a large fraction of AIRS channels are extremely stable, well below climate trends, where agreement with ESRL CO_2 anomalies is -0.023 ± 0.009 K/Decade. The SST anomaly retrievals also compare favorably to the

790 ERA-I reanalysis and to NOAA's OISST SST product, with differences of less than 0.022 K/Decade, and slightly higher values for comparisons to OISST. Such good agreement for a wide range of detectors strongly suggests that the AIRS blackbody is very stable.

Unphysical radiometric jumps are observed in the all all the retrieved anomaly time series, but especially for N_2O and CH_4 . These jumps can largely be related to AIRS events, and we illustrate how the anomaly fit residuals, combined with inter-comparisons to truth anomaly trends such as N_2O may provide a way to correct small remaining jumps in some AIRS

795

channels.

This work emphasizes that users of AIRS data-radiances (both Level-1b and Level-1c) for climate applications must pay careful attention to channel selection, since certain detector arrays and channels are presently not suitable for climate trending, including all of the AIRS short wave channels. However, establishment of such a high level of stability for so many remote

800 sensing observations/channels is highly unusual, and should lead to a high level of trust in AIRS climate trends that pay careful attention to only using validated climate-level channels.

Array Name	Start ν (cm ⁻¹)	End ν (cm ⁻¹)
1a	2552	2677
2a	2432	2555
1b	2309	2434
2b	2169	2312
4a	1540	1614
4b	1460	1527
3	1337	1443
4c	1283	1339
4d	1216	1273
5	1055	1136
6	973	1046
7	910	974
8	851	904
9	788	852
10	727	782
11	687	729
12	649	682

Table A1. The wavenumber ranges covered by each of the 17 AIRS arrays.

Acknowledgement

The authors thank Steve Broberg, NASA JPL AIRS Project Office, for supplying us with a table of AIRS events. We also thank Steven Buczkowski at UMBC/JCET for the extensive data handling and production needed for this work. The hardware used for this work is part of the UMBC High Performance Computing Facility (HPCF). The facility is supported by the U.S. National Science Foundation through the MRI program (grant nos. CNS-0821258, CNS-1228778, and OAC-1726023) and the SCREMS program (grant no. DMS-0821311), with additional substantial support from the University of Maryland, Baltimore County (UMBC).

Appendix A: AIRS Detector Array Wavenumbers

810 Table A1 shows the wavenumber ranges covered by each of the 17 AIRS arrays.

Appendix B: Anomaly and Profile Trend Retrievals

A complete simulated BT anomaly dataset was generated using ERA-I model fields, by matching each AIRS clear observation to ERA-I and generating a simulated radiance. This simulated dataset was used to set the regularization parameters for the profile inversions. The measurement of anomalies largely removes systematic errors in both the radiance observations (radio-metric accuracy) and in the RTA (spectroscopy errors). We believe that these two factors helped make the retrieval inversions

quite stable, requiring only minimal regularization.

Since our interest is mainly in the minor-gas profile offsets we used 20 atmospheric layers for the retrievals (20 each for temperature, H_2O , and O_3), created by concatenating layers from the 100-layer atmospheric profile model in (Strow et al., 2003). This choice, coupled with our regularization, provided more layers than degrees of freedom, as desired. We found that low noise of the AIRS zonally averaged 16-day anomalies (see Sect. 4.1 coupled with low bias errors in the measurement covariances allowed us to use only minimal regularization.

820

We first adapted Tikhonov-only first-derivative (L1-type) regularization which mostly removed obvious outliers, mostly in the higher latitudes in the stratosphere. This gave averaged linear-trend accuracies in our simulations of -0.03 ± 0.07 K/year compared to the ERA-I model field trends used to generate the anomaly data set. (This degrades to -0.05 ± 0.08 K/year if the regularization is lowered by a factor of 10X.) A reasonable goal is to achieve trends in simulation accurate to 0.01K/year, averaged over the troposphere. We then added a-priori uncertainties to the temperature and H₂O profiles of 2.5K and 60% respectively, which are roughly the maximum variation in these quantities over time for $\pm 50^{\circ}$ latitude. These covariances are not very restrictive given that measurement uncertainties are so low. It appears that their main impact is again for high latitudes under conditions where we have higher noise due to low number of clear samples.



Figure B1. Temperature kernels for the anomaly retrievals. These are taken from a random day for the zonal bin centered at 28.3°N.

⁸¹⁵



Figure B2. H₂O kernels for the anomaly retrievals. These are taken from a random day for the zonal bin centered at 28.3°N.

830 The temperature and water vapor retrieval kernels are shown in Figs. B1,B2. They exhibit a very regular spacing in the troposphere with roughly 12 well-separated kernels.



Figure B3. Retrieved 400 hPa temperature anomalies versus latitude. Top: Our retrievals from the AIRS observations. Bottom: ERA-I anomalies.

Figure B3 illustrates the 400 hPa temperatures we retrieved from the AIRS data (top panel) along with the ERA-I anomalies computed directly from the model fields. We do not expect these two data sets to compare perfectly, since for example, the ERA-I anomalies are from relatively large gridded data and the AIRS measurement are from a nominal 15 x 15 km field-of-view. Given the non-uniform sampling of our data set we do not think detailed examination of the observed versus ERA-I

835



Figure B4. Temperature trends from the 16-year data period studied here. Left: ERA-I trends derived directly from the model temperature fields. Middle: Simulated retrievals of the ERA-I trends using radiance anomalies created from the ERA-I fields and our SARTA RTA. Right: Temperature profile trends retrieved from the AIRS observed anomalies. The middle panel simulation assumes that RTA is perfectly accurate.

anomalies is warranted. We do note the there are many similarities in time and latitude that give some measure of validation to our profile retrievals. Similar results are seen with water vapor profiles.

Figure B4 summaries the temperature trend simulations and comparisons between ERA-I trends, our anomaly retrievals from the ERA-I generated radiances, and those observed with the AIRS clear subset. The trends are computed from the anomaly retrievals (or model fields) using Eq. 16, where the input is the layer temperature instead of a radianceCO₂ amount.

These results have been slightly smoothed to make visual inter-comparisons easier. The left panel shows the vertical trends versus latitude directly computed from the ERA-I temperature fields. The middle panel shows our simulated temperature trend retrievals. These simulations agree quite well with the ERA-I model fields, the largest differences are seen in the lower troposphere at the higher latitudes, and near the boundary layer in the tropics. The simulated retrievals are also placing the tropopause too high, not surprising given the lack of sensitivity of the infrared to the tropopause height and our limited number of vertical layers. The right panel are the temperature enomely trends retrieved from the AIPS observed enomalies. Clearly

of vertical layers. The right panel are the temperature anomaly trends retrieved from the AIRS observed anomalies. Clearly there are significant differences between the ERA-I temperature profile trends and those we retrieved from AIRS, although the basic structure is relatively similar. Note that the uncertainties in these trends are quite high in the stratosphere (not shown) due to variations in the quasi-biennal oscillation (QBO), especially in the tropics, with errors larger than the observed trends in the vicinity of the tropopause. However, these uncertainties are largely present in both ERA-I and the AIRS observations.

The AIRS observed anomalies may also be impacted by errors in the BT Jacobians. The middle panel in Fig. B4 used similar RTAs for both simulations and the retrieval. The version of SARTA used for the radiance simulations is based on HITRAN2008 while the Jacobians used in the retrieval used kCARTA which is based on HITRAN2016 and a slightly modified version of CO_2 line-mixing. We expect that these spectroscopy differences have little impact since the CO_2 line strengths for the strong 15 μ m bands have not changed between HITRAN versions. In addition, no noise was added to the simulated anomalies.

855

840

845

We believe that these results show that the anomaly retrievals used for measuring minor-gas trends exhibit realistic behavior and given our simulation testing this retrieval approach is likely to give accurate minor-gas trends. The impact of some of the regularization choices are discussed in Sect.5.4. *Author contributions.* LLS led the study and made the comparisons between the anomaly fits and in-situ data. SDM developed the anomaly retrieval algorithm. LLS and SDM together optimized the anomaly retrieval regularization.

Competing interests. The authors declare that they have no conflict of interest.

References

890

- Anderson, G. P., Clough, S. A., Kneizys, F. X., Chetwynd, J. H., and Shettle, E. P.: AFGL atmospheric constituent profiles (0.120km), Tech. rep., 1986.
- 865 Argo: (2019). Argo float data and metadata from Global Data Assembly Centre (Argo GDAC)., SEANOE, 2019.

Aumann, H., Broberg, S., Manning, E., Pagano, T., Sutin, B., and Strow, L.: AIRS Level 1C Algorithm Theoretical Basis Document, Version 6.7 AIRICRADATBDand TestDocuments, ., available as https://docserver.gesdisc.eosdis.nasa.gov/public/project/AIRS/AIRICRAD.v6.7. ATBD.pdf, 2020.

- 870 Staelin, D. H., Strow, L. L., and Susskind, J.: AIRS/AMSU/HSB on the Aqua Mission, IEEE Transactions on Geoscience and Remote Sensing, 41, 253, https://doi.org/10.1109/TGRS.2002.808356, https://doi.org/10.1109/TGRS.2002.808356, 2003.
 - Aumann, H. H., Broberg, S., Manning, E., and Pagano, T.: Radiometric Stability Validation of 17 Years of AIRS Data Using Sea Surface Temperatures, Geophysical Research Letters, 46, 12504–12510, https://doi.org/10.1029/2019GL085098, https://doi.org/10.1029/2019GL085098, 2019.
- 875 Banzon, V., Smith, T. M., Chin, T. M., Liu, C., and Hankins, W.: A long-term record of blended satellite and in situ sea-surface temperature for climate monitoring, modeling and environmental studies, Earth System Science Data, 8, 165–176, https://doi.org/10.5194/essd-8-165-2016, https://doi.org/10.5194/essd-8-165-2016, 2016.

Blaisdell, J. M.: private communication, 2019.

- Cowtan, K., Hausfather, Z., Hawkins, E., Jacobs, P., Mann, M. E., Miller, S. K., Steinman, B. A., Stolpe, M. B., and Way, R. G.: Robust
 comparison of climate models with observations using blended land air and ocean sea surface temperatures, Geophysical Research Letters,
 42, 6526–6534, https://doi.org/10.1002/2015gl064888, 2015.
 - Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., Berg, L. v. d., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.,
- 885 Park, B., Peubey, C., Rosnay, P. d., Tavolato, C., Thépaut, J., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, Quarterly Journal of the Royal Meteorological Society, 137, 553–597, https://doi.org/10.1002/qj.828, https://doi.org/10.1002/qj.828, 2011.
 - DeSouza-Machado, S., Strow, L. L., Tangborn, A., Huang, X., Chen, X., Liu, X., Wu, W., and Yang, Q.: Single-footprint retrievals for AIRS using a fast TwoSlab cloud-representation model and the SARTA all-sky infrared radiative transfer algorithm, Atmospheric Measurement
- DeSouza-Machado, S., Strow, L., Motteler, H., and Hannon, S.: kCARTA: A fast pseudo line-by-line radiative transfer algorithmwith analytic Jacobians, fluxes, Non-Local ThermodynamicEquilibrium and scattering for the infrared, Atmos. Meas. Tech., https://doi.org/10.5194/amt-2019-282, accepted for publication, 2019.

ESRL: Combined Chloroflurocarbon-12 data from the NOAA/ESRL Global Monitoring Division.

Techniques, 11, 529–550, https://doi.org/10.5194/amt-11-529-2018, 2018.

895 Fiedler, E. K., McLaren, A., Banzon, V., Brasnett, B., Ishizaki, S., Kennedy, J., Rayner, N., Roberts-Jones, J., Corlett, G., Merchant, C. J., and Donlon, C.: Intercomparison of long-term sea surface temperature analyses using the GHRSST Multi-Product Ensemble (GMPE) system, Remote Sensing of Environment, 222, 18–33, https://doi.org/10.1016/j.rse.2018.12.015, https://doi.org/10.1016/j.rse.2018.12.015, 2019.

Aumann, H. H., Chahine, M. T., Gautier, C., Goldberg, M. D., Kalnay, E., McMillin, L. M., Revercomb, H., Rosenkranz, P. W., Smith, W. L.,

Gordon, I. E., Rothman, L. S., Hill, C., Kochanov, R. V., Tan, Y., Bernath, P. F., Birk, M., Boudon, V., Campargue, A., Chance, K. V., Drouin, B. J., Flaud, J. M., Gamache, R. R., Hodges, J. T., Jacquemart, D., Perevalov, V. I., Perrin, A., Shine, K. P., Smith, M. A. H., Tennyson, J.,

- Toon, G. C., Tran, H., Tyuterev, V. G., Barbe, A., Császár, A. G., Devi, V. M., Furtenbacher, T., Harrison, J. J., Hartmann, J. M., Jolly, A., Johnson, T. J., Karman, T., Kleiner, I., Kyuberis, A. A., Loos, J., Lyulin, O. M., Massie, S. T., Mikhailenko, S. N., Moazzen-Ahmadi, N., Müller, H. S. P., Naumenko, O. V., Nikitin, A. V., Polyansky, O. L., Rey, M., Rotger, M., Sharpe, S. W., Sung, K., Starikova, E., Tashkun, S. A., Auwera, J. V., Wagner, G., Wilzewski, J., Wcisło, P., Yu, S., and Zak, E. J.: The HITRAN2016 molecular spectroscopic database, Journal of Quantitative Spectroscopy and Radiative Transfer, 203, 3–69, https://doi.org/10.1016/j.jqsrt.2017.06.038, 2017.
- 905 Hansen, J., Ruedy, R., Sato, M., and Lo, K.: GLOBAL SURFACE TEMPERATURE CHANGE, Reviews of Geophysics, 48, https://doi.org/10.1029/2010RG000345, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2010RG000345, 2010.
 - Kulawik, S. S., Jones, D. B. A., Nassar, R., Irion, F. W., Worden, J. R., Bowman, K. W., Machida, T., Matsueda, H., Sawa, Y., Biraud, S. C., Fischer, M. L., and Jacobson, A. R.: Characterization of Tropospheric Emission Spectrometer (TES) CO2 for carbon cycle science, Atmospheric Chemistry and Physics, 10, 5601–5623, https://doi.org/10.5194/acp-10-5601-2010, 2010.
- 910 Manning, E. M., Strow, L. L., and Aumann, H. H.: AIRS version 6.6 and version 7 level-1C products, in: Earth Observing Systems XXIV, edited by Butler, J. J., Xiong, X. J., and Gu, X., vol. 11127, pp. 247 253, International Society for Optics and Photonics, SPIE, https://doi.org/10.1117/12.2529400, https://doi.org/10.1117/12.2529400, 2019.
 - Masarie, K. A. and Tans, P. P.: Extension and integration of atmospheric carbon dioxide data into a globally consistent measurement record, Journal of Geophysical Research: Atmospheres, 100, 11 593–11 610, https://doi.org/10.1029/95JD00859, https://doi.org/10.1029/
- 915 95JD00859, 1995.
- Masson-Delmotte, V., Zhai, P., Pörtner, H. O., Roberts, D., Skea, J., Shukla, P. R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., Connors, S., Matthews, J. B. R., Chen, Y., Zhou, X., Gomis, M. I., Lonnoy, E., Maycock, T., Tignor, M., and (eds.), W.Waterfield): IPCC, 2018: Summary for Policymakers. In: Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the
- 920 threat of climate change, sustainable development, and efforts to eradicate poverty, Tech. rep., World Meteorological Organization, 2018. Morice, C. P., Kennedy, J. J., Rayner, N. A., and Jones, P. D.: Quantifying uncertainties in global and regional temperature change
 - using an ensemble of observational estimates: The HadCRUT4 data set, Journal of Geophysical Research: Atmospheres, 117, https://doi.org/10.1029/2011JD017187, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011JD017187, 2012.
- Pagano, T. S. and Broberg, S. E.: Recent checks on the radiometric and spatial calibration of AIRS in-orbit, in: Earth Observing Systems
 XXI, edited by Butler, J. J., Xiong, X. J., and Gu, X., vol. 9972, pp. 68 76, International Society for Optics and Photonics, SPIE, https://doi.org/10.1117/12.2238765, https://doi.org/10.1117/12.2238765, 2016.
 - Rodgers, C. D.: Retrieval of atmospheric temperature and composition from remote measurements of thermal radiation, Reviews of Geophysics, 14, 609–624, https://doi.org/10.1029/RG014i004p00609, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/RG014i004p00609, 1976.
- 930 Santer, B. D., Wigley, T. M. L., Boyle, J. S., Gaffen, D. J., Hnilo, J. J., Nychka, D., Parker, D. E., and Taylor, K. E.: Statistical significance of trends and trend differences in layer-average atmospheric temperature time series, Journal of Geophysical Research: Atmospheres, 105, 7337–7356, https://doi.org/10.1029/1999JD901105, https://doi.org/10.1029/1999JD901105, 2000.
- Simmons, A., Poli, P., Dee, D., Berrisford, P., Hersbach, H., Kobayashi, S., and Peubey, C.: Estimating low-frequency variability and trends in atmospheric temperature using ERA-Interim, Quarterly Journal of the Royal Meteorological Society, 140, 329–353, https://doi.org/10.1002/qj.2317, 2014.

- Stark, J. D., Donlon, C. J., Martin, M. J., and McCulloch, M. E.: Ostia: An operational, high resolution, real time, global sea surface temperature analysis system, OCEANS 2007 - Europe, pp. 1-4, https://doi.org/10.1109/OCEANSE.2007.4302251, https://doi.org/10. 1109/OCEANSE.2007.4302251, 2007.
- Steck, T.: Methods for determining regularization for atmospheric retrieval problems., Applied Optics, 41, 1788–97, 2002.
- 940 Strow, L., Motteler, H. E., Benson, R. G., Hannon, S. E., and Souza-Machado, S. D.: Fast computation of monochromatic infrared atmospheric transmittances using compressed look-up tables, Journal of Quantitative Spectroscopy and Radiative Transfer, 59, 481-493, https://doi.org/10.1016/S0022-4073(97)00169-6, https://doi.org/10.1016/S0022-4073(97)00169-6, 1998.
 - Strow, L. L., Hannon, S. E., Souza-Machado, S. D., Motteler, H. E., and Tobin, D.: An Overview of the AIRS Radiative Transfer Model, IEEE Transactions on Geoscience and Remote Sensing, 41, 303, https://doi.org/10.1109/TGRS.2002.808244, https://doi.org/10.1109/ TGRS.2002.808244, 2003.
- 945
 - Strow, L. L., Hannon, S. E., Machado, S. D., Motteler, H. E., and Tobin, D. C.: Validation of the Atmospheric Infrared Sounder radiative transfer algorithm, Journal of Geophysical Research: Atmospheres (1984–2012), 111, https://doi.org/10.1029/2005JD006146, https://doi. org/10.1029/2005JD006146, 2006.
 - Susskind, J., Blaisdell, J. M., and Iredell, L.: Improved methodology for surface and atmospheric soundings, error estimates, and quality
- 950 control procedures: the atmospheric infrared sounder science team version-6 retrieval algorithm, Journal of Applied Remote Sensing, 8, 084 994-084 994, https://doi.org/10.1117/1.JRS.8.084994, https://doi.org/10.1117/1.JRS.8.084994, 2014.
 - Susskind, J., Schmidt, G. A., Lee, J. N., and Iredell, L.: Recent global warming as confirmed by AIRS, Environmental Research Letters, 14, 044 030, https://doi.org/10.1088/1748-9326/aafd4e, https://doi.org/10.1088/1748-9326/aafd4e, 2019.

Tans, P. and Keeling, R.: Trends in Atmospheric Carbon Dioxide, http://www.esrl.noaa.gov/gmd/ccgg/trends/.