Supplementary material to TROPOMI tropospheric ozone column data : Geophysical assessment and comparison to ozonesondes, GOME-2B and OMI

Daan Hubert¹, Klaus-Peter Heue^{2,3}, Jean-Christopher Lambert¹, Tijl Verhoelst¹, Marc Allaart⁴, Steven Compernolle¹, Patrick D. Cullis⁵, Angelika Dehn⁶, Christian Félix⁷, Bryan J. Johnson⁵, Arno Keppens¹, Debra E. Kollonige^{8,9}, Christophe Lerot¹, Diego Loyola², Matakite Maata¹⁰, Sukarni Mitro¹¹, Maznorizan Mohamad¹², Ankie Piters⁴, Fabian Romahn², Henry B. Selkirk^{9,13}, Francisco R. da Silva¹⁴, Ryan M. Stauffer^{9,15}, Anne M. Thompson⁹, J. Pepijn Veefkind⁴, Holger Vömel¹⁶, Jacquelyn C. Witte¹⁶, and Claus Zehner⁶

¹Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Ringlaan 3, 1180 Uccle (Brussels), Belgium

²German Aerospace Centre (DLR), Münchener Straße 20, 82234 Weßling, Germany

³Technische Universität München, Arcisstrasse 21, 80333 München, Germany

⁴Royal Netherlands Meteorological Institute (KNMI), Utrechtseweg 297, 3730 AE De Bilt, The Netherlands

⁵NOAA Global Monitoring Laboratory (NOAA/ESRL/GML), 1325 Broadway, Boulder 80305-3337, CO, USA

⁶European Space Agency/Centre for Earth Observation (ESA/ESRIN), Largo Galileo Galilei 1, 00044 Frascati (Roma), Italy

⁷Federal Office of Meteorology and Climatology, MeteoSwiss, Payerne, Switzerland

⁸Science Systems and Applications, Inc., Lanham, MD, USA

⁹Atmospheric Chemistry and Dynamics Lab, NASA Goddard Space Flight Center, Greenbelt, MD, USA

¹⁰School of Biological and Chemical Sciences, University of the South Pacific, Fiji

¹¹Meteorological Service of Suriname, Paramaribo, Suriname

¹²Atmospheric Science and Cloud Seeding Division, Malaysian Meteorological Department, Petaling Jaya, Selangor, Malaysia

¹³Universities Space Research Association, Columbia, MD, USA

¹⁴Laboratory of Environmental and Tropical Variables, Brazilian Institute of Space Research, Natal, Brazil

¹⁵Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD, USA

¹⁶National Center for Atmospheric Research, Boulder, CO, USA

Correspondence: daan.hubert@aeronomie.be



Figure S1. Sampling statistics of two years of screened TROPOMI tropospheric O3 column data (1 May 2018 – 30 Apr 2020): (top) number of TrOC values per year, (bottom) mean day of year. In the outer tropics, winter / early spring data are often screened due to the absence of thick convective clouds in the stratospheric reference region. Grid cells with a uniform sampling time distribution are whited out. Red markers show the location of the SHADOZ sites considered in this study. Red isolines trace the 500, 1000 and 2000 m surface elevation levels. This figure supplements the annual statistics of TrOC in Fig. 1 in manuscript.



Figure S2. Skipped Pearson correlation coefficients for TROPOMI versus OMI (left) or GOME-2B (right) TrOC time series; white contours trace isolines of spread in TROPOMI TrOC (Fig. 1 centre in manuscript).



Figure S3. Like Fig. 4 in manuscript. Time series of the difference between spatially and temporally co-located OMI and ozonesonde tropospheric ozone column data over nine SHADOZ sites. Positive values indicate that OMI overestimates the ozonesonde value. Blue line and area show median and 68% interpercentile over the entire period.



Figure S4. Like Fig. 4 in manuscript. Time series of the difference between spatially and temporally co-located GOME-2B and ozonesonde tropospheric ozone column data over nine SHADOZ sites. Positive values indicate that GOME-2B overestimates the ozonesonde value. Blue line and area show median and 68% interpretentile over the entire period.



Figure S5. Statistics of local solar time of ozonesonde launches at each SHADOZ station in comparison to the fixed overpass time by the three satellite sensors (horizontal lines). The box shows the 25%, 50% and 75% quantiles, whiskers cover the 99.3% interpercentile, red markers represent individual flights. Stations are ordered West to East.



Figure S6. Like Fig. 12 in manuscript. Median biweekly OMI (top) and GOME-2B (bottom) TrOC over Atlantic basin between early July and mid December in 2018 (left) and 2019 (right). Grid cells with sparse or inhomogeneous temporal sampling or a value below 30 DU are blanked. Contours indicate the 35 DU (dashed), 40 DU (solid) and 45 DU (red) isolines. As a result of a \sim 2 DU low bias of GOME-2B with respect to OMI and TROPOMI the area above 30 DU is considerably smaller.