



A new measurement approach for validating satellite-based above cloud aerosol optical depth

Charles K. Gatebe^{1,2}, Hiren Jethva^{1,2}, Ritesh Gautam³, Rajesh Poudyal^{2,4} and Tamas Várnai^{2,5}

¹Universities Space Research Association (USRA), Columbia, MD, 21046, USA

5 ²NASA Goddard Space Flight Center, Greenbelt, Maryland, 20771, USA

³Environmental Defense Fund, Washington, DC 20009, USA

⁴Science Systems and Applications, Inc. (SSAI), Lanham, MD 20706, USA

⁵University of Maryland, Baltimore County, Baltimore, MD 21250, USA

10 *Correspondence to:* Charles K. Gatebe (charles.k.gatebe@nasa.gov)

Abstract. The retrieval of aerosol parameters from passive satellite instruments in cloudy scenes is challenging partly because clouds and cloud-related processes may significantly modify aerosol optical depth (AOD) and particle size, a problem that is further compounded by the 3D radiative processes. Recent advances in retrieval algorithms such as the “color ratio” method which utilizes the measurements at a shorter (470 nm) and a longer (860 nm) wavelength have demonstrated the simultaneous
15 derivation of AOD and cloud optical depth (COD) for scenes where absorbing aerosols are found to overlay low-level cloud decks. This study shows simultaneous retrievals of above-cloud aerosol optical depth (ACAOD) and aerosol-corrected cloud optical depth (COD) from airborne measurements of cloud-reflected and sky radiances using the color ratio method. These airborne measurements were taken over marine stratocumulus clouds with NASA’s Cloud Absorption Radiometer (CAR) during SAFARI 2000 field campaign offshore of Namibia. The ACAOD is partitioned between the AOD below aircraft
20 (AOD_{cloudtop}) and above aircraft AOD (AOD_{sky}). The results show good agreement between AOD_{sky} and sunphotometer measurements of the above aircraft AOD. The results also show that the use of aircraft-based sunphotometer measurements to validate satellite retrievals of the ACAOD is complicated by the lack of information on AOD below aircraft. Specifically, the CAR-retrieved AOD_{cloudtop} captures this “missing” aerosol layer caught between the aircraft and cloud top, which is required to quantify above cloud aerosol loading and effectively validate satellite retrievals. In addition, the
25 study finds a strong anticorrelation between the AOD_{cloudtop} and COD for cases where COD <10 and a weaker anticorrelation for COD >10, which may be associated with the uncertainties in the color ratio method at lower AODs and CODs. The influence of 3D radiative effects on the retrievals is examined and the results show that at cloud troughs, 3D effects increase retrieved ACAOD by about 3-10% and retrieved COD by about 25%. The results show that the color ratio method has little sensitivity to 3D effects at overcast stratocumulus cloud decks. These results demonstrate a novel airborne
30 measurement approach for assessing satellite retrievals of aerosols above clouds, thereby filling a major gap that exists in the global aerosol observations.



1 Introduction

The uncertainties of aerosols measurements in the vicinity of clouds has implication for the direct shortwave radiative aerosol effect and forcing on the climate system. Also, aerosols are known to exert an indirect forcing on climate by altering cloud properties and precipitation. According to the last Assessment Report of the Intergovernmental Panel on Climate Change (Boucher et al., 2013), the interactions between clouds and aerosols remain among the largest sources of uncertainty, pointing to a lack of good understanding of the aerosol-cloud system, and holding back progress in the enhancement of Earth system predictions/projections.

Space-based retrievals of aerosol optical properties in the vicinity of clouds is complex because of the difficulty in distinguishing the contributions from aerosols and clouds in top of the atmosphere (TOA) reflectance measurements. However, in the last two decades, several studies have demonstrated new approaches for aerosol retrievals in the vicinity of clouds. The absorbing aerosols such as smoke plumes, desert dust, and volcanic ash have been monitored from satellite observations in the presence of clouds using the ultraviolet measurements of Total Ozone Mapping Spectrometer (TOMS)/ Nimbus 7 (Herman et al., 1997; Torres et al., 1998), Ozone Monitoring Instrument (OMI)/Aura (Torres et al., 2012), and the Scanning Imaging Absorption Spectrometer for Atmospheric Chartography (SCIAMACHY) (De Graaf et al. 2007). The near-UV retrieval approach was extended to the visible and near-infrared spectral regions for the simultaneous derivation of aerosol optical depth and cloud optical depth based on Moderate Resolution Imaging Spectroradiometer (MODIS) measurements in regions where light-absorbing carbonaceous and dust aerosols overlay low-level clouds (c.f. Jethva et al. 2013, Sayer et al. 2016). Similarly, Waquet et al (2009) developed a method based on multiangle polarization measurements at visible and near-infrared wavelengths to retrieve aerosol properties over clouds and successfully applied it to measurements of the Polarization and Directionality of Earth Reflectances (POLDER)–Polarization and Anisotropy of Reflectances for Atmospheric Sciences Coupled with Observations from a Lidar (PARASOL) instrument. These advancements have provided hope for realizing global scale monitoring of aerosol properties over clouds, thereby filling a major gap that exists in the global aerosol observations, but significant challenges remain in the validation of above cloud aerosol products (Shinozuka et al. 2019). There is no question that the above-cloud aerosol retrievals need to be validated with airborne measurements.

This study demonstrates the applicability of the color ratio method (Jethva et al., 2013; 2016), which utilizes the measurements at a shorter (470 nm) and a longer (860 nm) wavelength for the simultaneous derivation of aerosol optical depth (AOD) and cloud optical depth (COD), to airborne observations. The study uses airborne data taken over marine stratocumulus clouds by the NASA's Cloud Absorption Radiometer (CAR) during SAFARI 2000 field campaign offshore of Namibia. The CAR instrument provides unique views of the cloud-aerosol system, from far, from close, or even from inside clouds—and from all the viewing directions (c.f. King et al. 1986; Gatebe et al. 2012; Gautam et al. 2016; Varnai et al. 2019; Gatebe and King 2016; Melnikova and Gatebe, 2018). The area selected has unique and reliable juxtaposition of regional and temporal patterns of meteorological conditions that are conducive to persistent low level clouds as seen from satellite imagery over the southeastern Atlantic region (cf. Figure 1); a region known to be impacted by optically thick smoke from intense biomass



65 burning activities (agriculture crop residue burning in central and southern Africa) (Das et al. 2020). The primary objective of this study is to retrieve aerosol optical depth above clouds using a novel airborne measurement approach of simultaneously measuring scattered radiation above and below the aircraft, and thereby demonstrate an effective observational tool to validate satellite-based aerosol retrievals above clouds.

2 Instruments and Methods

70 The Southeast Atlantic is widely used to study aerosol direct and indirect radiative effects because of the presence of stratiform marine clouds over ocean and the annual recurrence of very high concentrations of biomass burning aerosols between June and September (cf. Das et al., 2020; De Graaf et al. 2007; De Graaf et al. 2012; Keil and Haywood, 2003; Meyer et al., 2013; Sayer et al., 2016). The measurements analyzed here were taken aboard the University of Washington's Convair-580 research aircraft. During several portions of the flight analyzed here, the aircraft followed a circular flight track (Figure 1) at a near-
 75 constant distance from the cloud top (~650 m) occurring below ~1 km altitude (Gatebe et al 2003; Sinha et al. 2003). The image acquired by MODIS/Terra on the same day at about 09:25 UTC (see Figure 1, map inset), shows widespread clouds over the entire Namibian coast. There were reports during the Southern African Regional Science Initiative's (SAFARI 2000) dry season campaign (Swap et al., 2002)) that optically thick smoke that originated from intense biomass burning activities was advected over to the marine stratiform clouds off Namibian coast. The CV-580 flight began just prior to 10:00 UTC and
 80 ended at about 13:00 UTC. Table 1 summarizes the times and locations of the cases analysed, which are labelled alphabetically, a-p, based on the time of observations.

2.1 Aircraft & Sensors

The CAR instrument flew aboard the UW CV-580 research aircraft (Figure 2a), and obtained the bidirectional reflectance distribution function (BRDF) over an extensive and persistent stratocumulus cloud deck with an overlaying smoke aerosol
 85 layer. The aircraft was also equipped with other instruments to measure gases, aerosols and radiation (see Appendix A by P. V. Hobbs in the work of Sinha et al. 2003). Figure 2b shows a cutaway drawing of CAR. The instrument is approximately 72 cm long, 41 cm wide, and 39 cm deep and weighs 42 kg. CAR was designed primarily to image the sky and surface at an instantaneous field of view (IFOV) of 1° through a 190° wide plane as shown in Figure 2c. CAR measures both transmitted and reflected radiances at 14 narrow spectral bands located in the ultraviolet, visible and near-infrared (0.340-2.301 μm; Figure
 90 2d). This combination provides a convenient and efficient means of obtaining complete BRDFs for any surface type at a landscape level and ensures that surface albedo, which is an angular-weighted integration of the reflection function over a hemisphere, can be derived from these measurements covering the required angular range (Nicodemus et al., 1977; Kimes et al., 1987).

During the BRDF measurements over the marine stratiform clouds, the instrument obtained unique views of the
 95 cloud-aerosol system, scanning from zenith to the horizon and then from the horizon to nadir, and covering the entire 360°



range of azimuthal directions as the aircraft flew in a circular flight track (c.f. Gatebe et al. 2003; Fig. 3). The quicklook RGB image in Figure 3 ($R=1.04\ \mu\text{m}$, $G=0.87\ \mu\text{m}$, and $B=0.47\ \mu\text{m}$) illustrates measurements taken from 12:27 UTC to 12:54 UTC. The sun can be seen in the sky at about 33° view zenith angle, which also corresponds to the solar zenith angle, and a bright cloud system is seen on the image from view zenith angles 90° - 180° . The horizon coincides with the 90° view zenith angle, which is easily identified by the contrast between the sky and surface. In this image, the principal plane is defined by the vertical plane containing the sun and the plane that is equidistant between two solar disks.

Note that the circular flight track during the BRDF measurements above the clouds ($\sim 650\ \text{m}$) is about $4\ \text{km}$ in diameter, and with an aircraft bank angle of 20° - 30° , which is compensated by CAR to help maintain the full 180° view from zenith to nadir, the plane took ~ 3 minutes to complete an orbit. The marine stratiform clouds are generally characterized by a well-defined cloud top height corresponding to a strong boundary layer inversion. Given this viewing geometry of the cloud-aerosol system, the CAR measurements permit the retrieval of aerosol optical properties above clouds separated into above and below the aircraft, plus the cloud optical properties, using the color ratio method. These measurements provide the best data for validating above cloud aerosol retrieved from satellite measurements, analogous to the validation of cloud-free aerosol retrievals from satellites, which is typically done with observations from the AErosol RObotic NETwork (AERONET) ground-based sunphotometer network (Holben et al. 1998).

2.2 The color ratio method and its application to airborne observations

The color ratio (CR) method has been used to retrieve simultaneously the above-cloud aerosol optical depth (ACAOD) and aerosol-corrected cloud optical depth (COD) from OMI (Torres et al. 2012) and MODIS observations (Jethva et al. 2013; 2016). The technique is physically based on the reduction of the ultraviolet (UV), visible (VIS), and near-infrared (NIR) radiation reaching the top of atmosphere, due to particle absorption above cloud. The effects of aerosol absorption have a spectral signature, in which the absorption strength is found to be stronger at shorter wavelengths than at longer wavelengths. This produces a strong color effect in spectral measurements, hence, the name color ratio method. The near-UV based color ratio algorithm has been applied to the long-term record of OMI to derive a global product of ACAOD (Jethva et al., 2018). The ACAOD product has been validated against airborne measurements taken from HSRL-2 lidar operated during the ORACLES campaign conducted over the south eastern Atlantic Ocean. On the other hand, the ACAOD derived from the visible/near-IR observations of MODIS was validated against the direct AOD measurements acquired from airborne AATS and 4STAR sunphotometers operated during different field campaigns (Jethva et al. 2016). In both OMI and MODIS validation studies, the satellite-retrieved ACAOD product was found to agree well with the airborne measurements within the expected uncertainty limits associated with the inversion technique, which mainly arises from the chosen aerosol model and its absorption properties.

Here, the CR method was applied to CAR observations, which include direct and diffuse solar radiance (or sky radiance), at nine spectral channels (see Fig. 4.). The direct solar component is given by the extra-terrestrial solar radiance attenuated by atmospheric absorption and scattering. On the other hand, sky radiance results from single and multiple scattering



processes due to interaction of sunlight with aerosols and gas molecules. Atmospheric gas molecules (e.g. nitrogen, oxygen,
 carbon dioxide, ozone, water vapor, etc.) and aerosols are likely to strongly affect the solar radiance in the visible and near
 infrared regions. The attenuation (scattering and/or absorption) by each atmospheric constituent is strongly dependent on
 wavelength and can be determined through the optical thickness using simple parametric models (e.g. Zibordi and Voss, 1989).
 In the case of CAR measurements close to the sun (solar aureole), the signal from the direct solar radiance measurements
 saturate the detectors and therefore pixels that are especially close to the solar direction (scattering angles are $\leq 10^\circ$) should be
 excluded from any retrieval (Gatebe et al. 2010). The sky radiance distribution seen here is typical of clear skies (cloud free),
 where the radiance of a point in the sky depends both on its position relative to the sun (i.e., azimuth angle) and on its airmass
 number (i.e., zenith angle). The sky radiance distribution is generally symmetrical about the principal plane, where the
 maximum value of the sky radiance for each wavelength is observed. This is illustrated in Fig. 4e: $45^\circ \approx 315^\circ$; $90^\circ \approx 270^\circ$;
 $135^\circ \approx 225^\circ$ for $\lambda > 0.4 \mu\text{m}$. The minimum values of sky radiance are found to be in the area directly opposite to the sun's
 position.

The CAR observations are indicative of the presence of absorbing aerosols above the clouds due to apparent
 brightening/darkening, which is evident when looking at the measured sky radiances /cloud bidirectional reflectance factor
 (BRF) (cf. Fig. 4). Aerosol loading has a strong influence, especially in the forward scattering directions (relative azimuth
 angle (ϕ) $< 90^\circ$ and $\phi > 270^\circ$), with reflectances in the shorter wavelengths (e.g. $0.38 \mu\text{m}$) larger by a factor of > 3 relative to
 the longer wavelengths (e.g. $1.22 \mu\text{m}$). This asymmetry is largely attributed to aerosol scattering and not to Rayleigh scattering,
 as the latter is expected to exhibit symmetrical distribution in either scattering directions. More interestingly, there seems to
 be a strong aerosol absorption signal above clouds. It is well known that clouds reflect uniformly across the visible-near-IR
 spectrum, however, the presence of absorbing aerosols above clouds (in this case smoke transported from southwestern Africa)
 induces an overall absorption or darkening in the UV and shorter visible wavelengths, thus resulting in a strong reflectance
 gradient from UV to blue to near-IR spectrum, $\sim 35\%$ reduced reflectance at $0.34 \mu\text{m}$ compared to that at $1.04 \mu\text{m}$, as seen in
 Fig. 4f. Overall, the positive spectral gradient seen in Fig. 4f, is normally associated with cloud darkening at the shorter
 wavelengths (cf. Gautam et al. 2016).

3 Results

3.1 The observations

Figures 5 and 6 show the full BRF of low stratiform clouds at selected wavelengths of $0.472 \mu\text{m}$ and $0.870 \mu\text{m}$, respectively,
 from each of the 16 different cases described in Section 2. The two wavelengths form the basis of the “color ratio” method for
 the simultaneous retrieval of above-cloud aerosol optical depth (ACAOD) and cloud optical depth (COD). The spectral BRF
 of stratiform clouds observed in the 16 cases is highly anisotropic due to a combination of factors ranging from cloud
 heterogeneity (including sub-pixel heterogeneity), solar illumination geometry, sensor viewing geometry, and cloud
 parameters such as optical thickness and effective radius (cf. Cornet et al. 2018). The 16 cases have a range of solar zenith



angles ($23^\circ < \text{SZA} < 34^\circ$). Measurements span an area of ~ 55 km (N-S) \times ~ 12 km (E-W), with most cases (9 cases: cases h-p) concentrated over a much smaller area (~ 8 km \times ~ 4 km) (cf. Fig. 1). The observations were taken at approximately the same altitude (Table 1: cases a-d: 1420 -1541 m above mean sea level (AMSL) & cases h-p: 1608-1616 m AMSL), implying that corresponding pixels for the different cases have similar measurement scale. The only exceptions (cases e-f) were taken at different altitudes during the aircraft spiral from 1814 m to 3369 m AMSL. The cloud top height was ~ 1000 m AMSL (Sinha et al. 2003) and the cloud geometrical thickness was at most 300 m (cf. Melnikova and Gatebe 2019, subsection 2.2). Based on these characteristics, the 16 cases may be grouped into three groups (cf. Table 1): Group 1: cases a-d ($\text{SZA} \approx 24^\circ$, measurements were taken close to each other in time, $\Delta t < 16$ minutes, altitude ≈ 1508 m and the location is about the same as shown in Figure 1). Group 2: cases e-g ($\text{SZA} \approx 24^\circ$, $\Delta t < 6$ minutes, altitude \approx variable from low to high, and same location near the Namibian coastline as shown in Figure 1. Group 3 cases h-p ($\text{SZA} \approx 34^\circ$, $\Delta t < 23$ minutes, altitude ≈ 1614 m, and the location is about the same as shown in Figure 1). Since the stratiform clouds are formed and maintained by a balance of various marine boundary layer processes (cf. Duynkerke and Teixeira 2001; Woods 2012; Feingold et al. 2017), the variations in the BRF patterns with time, especially where other parameters are similar, are possibly linked to formation of open cells caused by the drizzle-cloud dynamical interactions and inevitably leading to changes in the cloud liquid water path and BRF. The pronounced circular brightness feature (see cases h-p, Fig. 5, $\lambda = 0.470 \mu\text{m}$, or Fig. 6, $\lambda = 0.870 \mu\text{m}$) shows a cloud bow (or primary rainbow), which is typical of water clouds (cf. Gatebe et al. 2003, where case h was analysed in detail). Figure 7 shows the derived spectral albedo (with atmosphere) for all the 16 cases at $\lambda = 0.470 \mu\text{m}$ and $\lambda = 0.870 \mu\text{m}$ (see Table 2 for the spectral albedo (with atmosphere) for all the wavelengths). Clearly, Group 3 cases had higher spectral albedo and was optically thicker, while Group 2 cases from near the Namibian coastline had the lowest spectral albedo (with atmosphere). It is interesting to note that the spectral albedo remains almost constant in Group 2 cases, despite the change in measurement scale during the spiral.

In the following subsections, we will examine how the surface reflectance anisotropy impacts the retrievals of the optical depth (both clouds and aerosols) using the color ratio method.

3.2 The retrieved ACAOD and COD

Figure 8 shows the retrieved AOD for aerosol layers located above the aircraft-level (AOD_{sky}) derived from the observed diffuse sky radiance by CAR. The retrievals were performed using a single-channel fit at 470 nm between the observed sky radiance aerosol look-up table accounting for the variations in AOD and geometry. Note that the aerosol model used for AOD_{sky} retrievals was the same for the inversion of AOD below aircraft (AOD_{cloudtop}). It is complicated to characterize and model the anisotropic effects of reflecting clouds with varying optical depths on the hemispherical diffuse sky radiances measured by CAR. Therefore, we adopted a simple approach to account for these effects, at least partially, by retrieving AOD above the aircraft assuming an averaged underneath cloud optical depth field retrieved from the



AOD_{cloudtop} inversion for each CAR BRDF case. For the most part the hemispherical distribution of retrieved AOD_{sky} along the azimuth direction is found to be smooth and near-uniform suggesting that the sky retrievals of AOD aren't significantly affected by the cloud anisotropy and that the simple approach of assuming an averaged value of COD for the full azimuthal scan works reasonably well in capturing the cloud effects on the sky radiances. The angular pattern in cases a-d is similar and in good agreement with the airborne direct sunphotometer measurements as discussed later (Figure 12 and Table 1).

The retrieved AOD below the aircraft (AOD_{cloudtop}) for all the 16 CAR BRDF cases are shown in Figure 9. The white areas in each polar plot are devoid of AOD_{cloudtop} retrievals either due to no cloud detection and/or the observations fall outside the color ratio vs. reflectance look-up table domain including extreme viewing geometry. In almost all cases (a-p), the retrieved AOD_{cloudtop} shows a dependence on viewing zenith angle, where lower (higher) AOD_{cloudtop} values are associated with slant (near-nadir) viewing angles (see also Figure 11 – scatter plots of AOD_{cloudtop} vs COD). Such gradient in the retrieved AOD_{cloudtop} can result from the limitations of the radiative transfer calculations at slant angles and the fact that CAR observations are interpreted within the look-up table after linearly interpolating between aerosol geometry nodes. The nodes in geometry used in the RT calculations include solar view zenith angles (sza_{nodal}: 0°, 10°, 20°, 30°, 40°, 50°, and 60°), view zenith angles (vza_{nodal}: 0°, 6°, 12°, 18°, 24°, 30°, 36°, 42°, 48°, 54°, 60°, 66°, 72°, and 80°), and relative azimuthal angles (raa_{nodal}: 0°, 20°, 40°, 60°, 80°, 100°, 120°, 140°, 160°, and 180°). Another salient feature of the retrieved AOD_{cloudtop} field is the intermittent patches of high AODs that extend in the viewing zenith direction along an azimuthal plane. A careful correlation of this feature with BRFs measured at 0.47 μm (see Fig. 5) and 0.87 μm (Fig. 6) reveals that the higher AODs are spatially collocated with relatively lower values of BRF, indicating that these observations belonged to either clear-sky or partially cloudy sky or thin heterogeneous scenes, for which the assumption of fully overcast thick homogeneous pixels made in the CR algorithm breaks down. Under such situations, it is expected that the uncertainty in the retrieved AOD_{cloudtop} would be larger than the expected errors due to other algorithmic assumptions. This issue is explored further in subsection 3.4 under the influence of 3D effects on the retrieved AOD_{cloudtop} and COD.

Another important observation in Fig. 9 is the increasing magnitudes of AOD above cloud for the cases e, f, and g. Table 1 shows that the altitude of aircraft for these three cases was recorded as 1533±2, 1814±259, 2646±223 meters above mean



sea level. It is expected that as the aircraft altitudes moves higher in the atmosphere, the CAR sensor would see an aerosol layer of greater geometrical thickness, thereby resulting in greater aerosol extinction and AOD. The retrieved AOD_cloudtop for these cases precisely demonstrates this effect by showing increasing magnitudes for higher aircraft altitudes.

220 The color ratio algorithm, along with the above-cloud AOD, also co-retrieves aerosol-corrected cloud optical depth, which is shown in Figure 10. Unlike aerosol fields, both seen above and below the aircraft level show more homogeneous distributions, the cloud optical depth fields retrieved from most of the cases show a great deal of variability along the azimuthal plane. Except for the cases m, n, o, and p, all other cases (a through l) show overall higher cloud optical depth in the back scattering directions shown in the bottom hemisphere opposite to the Sun and between the azimuth angle 90° and 270° . Unlike
 225 polar orbiting satellite observations at a fixed geometry for a given overpass, the CAR measurements offer a complete picture over all the viewing directions relative to the sun direction. This unique observational geometry provides increased information content that would allow quantification of the effects of angular reflectance distribution in remote sensing retrieval algorithms.

3.3 The relationship between AOD_cloudtop and COD

Figure 11 shows the scatter plots of AOD_cloudtop vs COD for view zenith angles 0° - 30° (blue color), 30° - 60° (green
 230 color), and 60° - 90° (red color), which shows very interesting patterns. The retrievals of AOD_cloudtop are found to exhibit a systematic dependence on COD (similar to an exponential decay function), especially the blue color and green color dots, where larger values of AOD_cloudtop correspond to lower values of COD and crawling along the x-axis on the right as COD increases. An exception to this rule are the retrievals made at the higher view zenith angles, 60° - 90° (red color), where the retrieved ACAOD remains low (<0.2) despite an increase in the COD, which seems unrealistic and confirms some of the
 235 limitations of the color ratio method. Another exception is seen in cases e, f and g, where AOD_cloudtop vs COD show no clear dependence on viewing zenith angle and COD was around 5, indicating that these observations belonged to either clear-sky or partially cloudy sky or thin heterogeneous scenes, for which the assumption of fully overcast thick homogeneous pixels made in the CR algorithm breaks down. The relationship between the two retrieved quantities appears to be confined for $COD < 10$, after which both retrievals are found to be not correlated with each other. Such observed dependence was expected
 240 as noticed in the color ratio algorithm introduced in Jethva et al. (2013). The uncertainties in satellite ACAOD inversion is known to be larger at lower CODs. This is because the retrieval domain space, i.e., color ratio versus reflectance at a longer wavelength, at lower CODs becomes narrower with steep changes in the color ratio, especially at $COD < 10$. Therefore, any



uncertainty in the assumptions made in the retrieval algorithm, i.e., single-scattering albedo, an assumption of fully overcast pixels, and linear interpolation between the nodes where reflectances and its ratio of a joint aerosol-cloud scene behaves non-linearly would result in the amplification of the error in the retrieved ACAOD. These artifacts are more pronounced at lower values of both ACAOD and COD, where uncertainties in the retrieved ACAOD could reach 40% to 80% at $COD < 10$ and $ACAOD < 0.5$ typically observed in the present CAR AOD retrievals (Jethva et al., 2013, Table II).

Figure 12 shows the two main aerosol-above-cloud retrieved parameters, namely AOD_{sky}, when CAR views upward flying above the cloud field, and the AOD below aircraft (AOD_{cloudtop}), when CAR views downward measuring the cloud field averaged over all the viewing directions (see also, Table 1, columns 6-9). The summation of AOD_{sky} and AOD_{cloudtop} provides the column AOD above the stratocumulus cloud fields (ACAOD), as retrieved from CAR measurements over marine stratus clouds during SAFARI 2000 in the southeast Atlantic region. In addition to the two aerosol-above-cloud parameters retrieved from CAR, Fig. 12 also shows simultaneous COD retrievals using CAR measurements as well as AOD retrievals from the AATS sunphotometer that made coincident measurements of AOD on the UW CV-580 flights. The AOD retrievals from AATS are based on direct Sunphotometer measurements and therefore represent aerosol loading above the aircraft-level.

In the case of flight transects shown in Fig. 1, the AATS AOD retrievals were largely obtained above the marine stratocumulus clouds. However, when the cloud top is well separated from the aircraft, i.e., the altitude of aircraft is higher than that of the cloud tops; the AATS measurements do not capture the aerosol layer below the aircraft as the instrument is always pointing upwards, toward the Sun. Therefore, the reported AOD data from AATS is not representative of the total column AOD above clouds, unless the aircraft is flying at the same altitude where cloud top is located. Often, the altitude difference is not negligible, for example, during the SAFARI flights shown in Fig. 3, there was a clear separation of ~600 m between the aircraft and cloud top. Specifically, the CAR-retrieved AOD_{cloudtop} captures this “missing” aerosol layer caught between the aircraft and cloud top, which is in addition to the AOD_{sky} retrieved above the aircraft level. The latter quantity is equivalent to that retrieved by AATS, whereas AOD_{cloudtop} is the remainder of the column AOD that we retrieve from CAR in this study. For these reasons, Jethva et al. (2016) in validating MODIS-retrieved ACAOD for the same September 13, 2000 AATS flight extrapolated the airborne measurements from the respective altitudes to cloud-top using a detailed profile



measurements and associated altitude-AOD polynomial in order to make the comparisons between satellite and airborne measurements consistent.

270 To illustrate the various retrievals, we consider flight measurements from cases h-p. The COD associated with the marine stratocumulus clouds appear relatively uniform, varying between 15 - 20 for measurements from cases h-m (Fig. 12). These retrievals (for cases h-m) are based on some of the most homogeneous clouds observed during the three separate circular measurements obtained from transects a-d, e-g and h-p. These relatively homogeneous and similar sets of circular transects are also noted in the BRF polar plots shown in Fig. 6h-m. The simultaneous retrievals of Sky_AOD show moderately high
 275 aerosol loading, $AOD = 0.5$ across circles h-m, which is in very close agreement with the AATS_AOD retrievals. The consistency in AOD retrievals (above the aircraft level) between the two disparate measurement approaches, i.e. AATS and CAR, is generally found throughout data obtained from the 16 cases (a-p), as indicated by the high correlation ($R^2 = 0.92$) between the two retrievals shown in Table 1. However, the central distinction here is that the CAR approach also allows us to directly retrieve aerosols above clouds that are present below the aircraft level ($AOD_{cloudtop}$). For instance, in cases h-m,
 280 the $AOD_{cloudtop}$ is 0.2, implying the total above-cloud column AOD is 0.7 or 40% higher relative to the above aircraft-level AOD. Overall, we find $AOD_{cloudtop}$ ranging between 0.17 and 0.40 from the 16 cases shown in Fig. 12, indicating a notable enhancement of the overall presence of aerosols above clouds. These observations show that a significant aerosol layer is not captured in previous measurements, indicating the strength and effectiveness of near-simultaneous multiangular measurements scanning the sky and surface, as demonstrated in this study using CAR measurements.

285 3.4 The influence of 3D effects on the retrieved ACAOD and COD

Numerous earlier studies indicate that passive remote sensing of both cloud and aerosol properties can be significantly impacted by three-dimensional (3D) radiative processes (e.g., Marshak and Davis, 2005; Wen et al., 2006; <http://i3rc.gsfc.nasa.gov/Publications.html>). Since the impact of 3D effects is different for different observations and retrieval algorithms (e.g., Cornet et al., 2018), we next examine the impact of 3D effects on the CAR aerosol and cloud retrievals
 290 discussed above. Our goal does not lie in providing quantitative estimates of 3D effects; instead we examine whether 3D effects are likely to play a substantial role in shaping the behavior of CAR-retrieved cloud and aerosol optical depths.

Our tests consider the scene shown in Figures 5k, 6k, 9k and 10k as a representative of heterogeneous areas with potentially significant 3D effects. The figures show that around 60° azimuth angle, CAR observed a roughly 300 m wide and very long



trough in which the retrieved COD drops by roughly 50% while the retrieved ACAOD increases by roughly 50%. Figures 9, 10, and 11 show that this behaviour is not unique, and that in many cases with COD values below 10 or sometimes even 20, the retrieved AOD values increase sharply as COD decreases. In principle, this behaviour appears consistent with earlier findings that showed 3D effects to increase retrieved AOD values for pixels that were surrounded by brighter (thick-cloud-covered) areas (e.g., Wen et al., 2013).

To examine 3D influences in CAR retrievals, we performed 1D and 3D radiative transfer simulations using the Monte Carlo model that powers the online simulator of 3D radiative processes that was created as part of the I3RC (Intercomparison of 3D Radiation Codes) project and is publicly available at <http://i3rcsimulator.umbc.edu/>. The key simulation parameters are listed in Table 3 and the simulation results are shown in Table 4.

In Table 4, each row, the left column indicates whether or not below-CAR aerosols (BCA) were considered, what the cloud optical depth was at the trough center, and whether the simulations considered 1D or 3D radiative processes. The indicated uncertainties come from Monte Carlo simulation noise.

Since COD retrievals are shaped mainly by the $0.87 \mu\text{m}$ reflectance values, 3D BRFs exceeding 1D BRFs by about 25% for COD=7 indicates that 3D radiative processes significantly enhance CAR BRFs and thus the COD values retrieved in the center of the trough—which means that 3D effects make the COD drop in the trough appear less deep than it really is. This behavior is consistent with earlier studies showing that radiative smoothing (caused by the diffusion of photons scattered from thick to thin areas) make horizontal cloud variability appear less strong than it really is. Several studies proposed counteracting this effect by artificially roughening the retrieved COD fields (e.g., Marshak et al., 1998; Zinner et al., 2006), but these methods are yet to gain wide usage. By performing additional simulations, we found that if we decreased COD at the center of the trough from 7 to 4.7, 3D simulations would yield $0.87 \mu\text{m}$ BRF values around 0.32—thus resulting in hypothetical retrievals yielding COD=7 (similar to the actual CAR retrievals). We note, however, that the value of 4.7 depends on our assumption of cloud base altitude (hence cloud geometrical thickness), and so it is somewhat uncertain.

Regarding aerosol retrievals, the results in Table 4 imply that 3D effects enhance by about 3-10% the impact of below-CAR aerosols (BCAs) on the $\text{BRF}_{0.47 \mu\text{m}} / \text{BRF}_{0.87 \mu\text{m}}$ color ratio (CR) values:

$$((\text{CR}_{3\text{D}, \text{COD}=4.7(\text{BCA})} - \text{CR}_{1\text{D}(\text{no BCA})}) / (\text{CR}_{1\text{D}(\text{BCA})} - \text{CR}_{1\text{D}(\text{no BCA})})) = 1.075 \pm 0.02$$



$$((CR_{3D, COD=7.0(BCA)} - CR_{1D(no\ BCA)}) / (CR_{1D(BCA)} - CR_{1D(no\ BCA)})) = 1.052 \pm 0.02.$$

320 Since CR is the key signal in our ACAOD retrievals, this implies that 3D effects are likely to increase retrieved ACAOD values by 3-10%.

To understand this result, we need to consider both the radiative smoothing discussed above for COD retrievals, and the 3D process often called “bluing” (e.g., Marshak et al., 2008). Bluing occurs when nearby thick clouds reflect more sunlight than the clouds in the field-of-view do, and some of the extra reflection is then scattered into the instrument field-of-view by
 325 air molecules and aerosol particles that reside between the cloud and the sensor. As expected, Table 4 reveals that 3D processes do indeed enhance BRFs: For COD=7, BRF_{3D} values exceed the corresponding BRF_{1D} values at both 0.47 μm and 0.87 μm . However, the table also reveals that given a certain 0.87 μm BRF value, 3D and 1D processes yield fairly similar 0.47 μm BRFs and thus color ratios: $BRF_{0.47\ \mu m, COD=4.7, 3D} \approx BRF_{0.47\ \mu m, COD=7, 1D}$ and $CR_{3D, COD=4.7} \approx CR_{1D, COD=7}$.

The weak impact of 3D effects on CR is likely due to two factors. First, while the bluing process implies a larger molecular
 330 and aerosol scattering enhancement at 0.47 μm than at 0.87 μm (i.e., a higher CR), this is partially compensated by the aerosol absorption cross section being larger at 0.47 μm than at 0.87 μm . Second, much of the 3D effects that cause the enhancements apparent in Table 4 are likely caused by the in-cloud radiative smoothing process discussed above, which causes similar relative enhancements in trough BRFs at 0.47 μm and 0.87 μm : Cloud droplets, which cause radiative smoothing through multiple scattering, have similar scattering properties at 0.47 μm and 0.87 μm .

335 We note that simulations (not shown) indicate that 3D effects would have similar or even weaker influence on ACAOD retrievals over the linear trough if the measurements were taken not by CAR flying only 600 m above the clouds, but by a satellite passing overhead. This is because the compensating effect of aerosol scattering and absorption and the spectrally neutral in-cloud radiative smoothing cause 3D relative enhancements that are spectrally quite neutral.

Overall, the results discussed above imply that 3D radiative processes had a significant impact on retrieved cloud optical
 340 depths, but also that the 3D impacts on retrieved ACAOD values is fairly small and is not the main reason for the retrieved ACAOD values increasing over thin clouds.



4. Conclusion

In conclusion, the study accomplished the simultaneous retrieval of above cloud total aerosol optical depth (ACAOD) and aerosol-corrected Cloud Optical Depth (COD) from airborne CAR measurements of cloud-reflected and sky radiances using the color ratio method. The ACAOD is partitioned between the AOD below the aircraft (AOD_cloudtop) and the AOD above the aircraft (AOD_sky) with full angular coverage provided by the CAR measurements. The study demonstrates a novel measurement approach for retrieving and quantifying aerosols above clouds, in particular recovering the aerosol layer between cloud tops and aircraft level that is missed in typical airborne sunphotometer measurements of above-cloud-aerosols. Overall, this work provides a path forward for filling a critical gap in aircraft-based sunphotometer measurement strategies that are currently used to validate satellite retrievals of the ACAOD.

The results show a strong anticorrelation between the AOD_cloudtop and COD for cases where COD <10, and a weaker anticorrelation for COD >10. The impact of 3D radiative effects on the retrievals is examined and the results show that at cloud troughs, 3D effects increase retrieved ACAOD by about 3-10% and retrieved COD by about 25%. This indicates that the color ratio method has little sensitivity to 3D effects at overcast stratocumulus cloud decks. The results also display good agreement between CAR and sunphotometer measurements of the above aircraft AOD. However, the results also show that the use of aircraft-based sunphotometer measurements to validate satellite retrievals of the ACAOD is complicated by the lack of information on AOD below the aircraft, indicating the strength and effectiveness of near-simultaneous multiangular measurements scanning the sky and surface, as demonstrated in this study using CAR measurements.

Acknowledgements

We are grateful to all our colleagues who helped in many ways and made it possible to collect the analyzed observations. This research was primarily funded by NASA's Atmospheric Composition Campaign Data Analysis and Modeling solicitation, ACCDAM, which is managed by Hal Maring. CAR SAFARI 2000 data acquisition was funded by the EOS Project Science Office led by Michael King.

References

- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B. and Zhang, X.Y.: Clouds and Aerosols. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.



Cornet, C., Labonnote, L. C., Waquet, F., Szczap, F., Deaconu, L., Parol, F., Vanbauce, C., Thieuleux, F., and Riedi, J.: Cloud heterogeneity on cloud and aerosol above cloud properties retrieved from simulated total and polarized reflectances, *Atmos. Meas. Tech.*, 11, 3627–3643, 2018.

375

Das, S., Harshvardhan, H., and Colarco, P. R.: The influence of elevated smoke layers on stratocumulus clouds over the SE Atlantic in the NASA Goddard Earth Observing System (GEOS) model, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031209, <https://doi.org/10.1029/2019JD031209>, 2020

380 De Graaf, M., Stammes, P., Torres, O., and Koelemeijer, R. B. A.: Absorbing Aerosol Index: Sensitivity analysis, application to GOME and comparison with TOMS, *J. Geophys. Res.*, 110, D01201, <https://doi.org/10.1029/2004JD005178>, 2005.

De Graaf, M., Stammes, P., and E. Aben, A.A.: Analysis of reflectance spectra of UV-absorbing aerosol scenes measured by SCIAMACHY, *J. Geophys. Res.*, 112, D02206, <https://doi.org/10.1029/2006JD007249>, 2007.

385

Feingold, G., Balsells, J., Glassmeier, F., Yamaguchi, T., Kazil, J., and McComiskey, A.: Analysis of albedo versus cloud fraction relationships in liquid water clouds using heuristic models and large eddy simulation, *J. Geophys. Res. Atmos.*, 122, 7086–7102, <https://doi.org/10.1002/2017JD026467>, 2017.

390 Gatebe, C. K., King, M. D., Platnick, S., Arnold, G. T., Vermote, E. F., and Schmid, B.: Airborne spectral measurements of surface-atmosphere anisotropy for several surfaces and ecosystems over southern Africa, *Journal of Geophysical Research*, 108, 8489, <https://doi.org/10.1029/2002JD002397>, 2003.

Gatebe, C. K., Dubovik, O., King, M. D., and Sinyuk, A.: Simultaneous retrieval of aerosol and surface optical properties from
 395 combined airborne- and ground-based direct and diffuse radiometric measurements, *Atmos. Chem. Phys.*, 10, 2777–2794, <https://doi.org/10.5194/acp-10-2777-2010>, 2010.

Gatebe, C. K., Varnai, T., Poudyal, R., Ichoku, C., and King, M. D.: Taking the pulse of pyrocumulus clouds, *Atmospheric Environment*, 52, 121–130, <https://doi.org/10.1016/j.atmosenv.2012.01.045>, 2012.

400

Gatebe, C. K., and King, M. D.: Airborne spectral BRDF of various surface types (ocean, vegetation, snow, desert, wetlands, cloud decks, smoke layers) for remote sensing applications, *Remote Sensing of Environment*, 179, 131–148, <https://doi.org/10.1016/j.rse.2016.03.029>, 2016.



- 405 Gautam, R., Gatebe, C. K., Singh, M. K., Várnai, T., and Poudyal, R.: Radiative characteristics of clouds embedded in smoke derived from airborne multiangular measurements, *Journal of Geophysical Research: Atmospheres*, 121(15), 9140–9152, 2016.
- Jethva, H., Torres, O., Remer, L., and Bhartia, P.: A Color Ratio Method for Simultaneous Retrieval of Aerosol and Cloud Optical Thickness of Above-Cloud Absorbing Aerosols From Passive Sensors: Application to MODIS Measurements, *IEEE T. Geosci. Remote*, 51, 3862–3870, <https://doi.org/10.1109/TGRS.2012.2230008>, 2013.
- 410 Jethva, H., Torres, O., Remer, L., Redemann, J., Livingston, J., Dunagan, S., Shinozuka, Y., Kacenelenbogen, M., Rosenheimer, M. S., and Spurr, R.: Validating MODIS above-cloud aerosol optical depth retrieved from “color ratio” algorithm using direct measurements made by NASA’s airborne AATS and 4STAR sensors, *Atmos. Meas. Tech.*, 9, 5053–5062, <https://doi.org/10.5194/amt-9-5053-2016>, 2016.
- 415 Jethva, H., Torres, O., and Ahn, C.: A 12-year long global record of optical depth of absorbing aerosols above the clouds derived from the OMI/OMACA algorithm, *Atmos. Meas. Tech.*, 11, 5837–5864, <https://doi.org/10.5194/amt-11-5837-2018>, 2018.
- 420 King, M. D., Strange, M. G., Leone, P., and Blaine, L. R.: Multiwavelength scanning radiometer for airborne measurements of scattered radiation within clouds, *Journal of Atmospheric and Oceanic Technology*, 3, 513–522, [https://doi.org/10.1175/1520-0426\(1986\)003<0513:MSRFAM>2.0.CO;2](https://doi.org/10.1175/1520-0426(1986)003<0513:MSRFAM>2.0.CO;2), 1986.
- 425 Herman, J. R., Bhartia, P. K., Torres, O., Hsu, C., Seftor, C., and Celarier, E.: Global distribution of UV-absorbing aerosols from Nimbus 7/TOMS data, *J. Geophys. Res.*, 102, 16911–16922, <https://doi.org/10.1029/96JD03680>, 1997.
- Levy, R. C., Remer, L. A., and Dubovik, O.: Global aerosol optical properties and application to Moderate Resolution Imaging Spectroradiometer aerosol retrieval over land, *J. Geophys. Res.*, 112, D13210, <https://doi.org/10.1029/2006JD007815>, 2007.
- 430 Marshak, A., Davis, A., Cahalan, R., and Wiscombe, W.: Nonlocal Independent Pixel Approximation, direct and inverse problems, *IEEE Trans. Geosci. Remote Sens.*, 36, 192–205, <https://doi.org/10.1109/36.655329>, 1998.
- Marshak, A., A. B. Davis (Eds.): *3D Radiative Transfer in Cloudy Atmospheres*, Springer, Heidelberg, 686 pp., 2005.
- 435 Marshak, A., Wen, G., Coakley, J., Remer, L., Loeb, N. G., and Cahalan, R. F.: A simple model for the cloud adjacency effect and the apparent bluing of aerosols near clouds. *J. Geophys. Res.*, 113, D14S17, <https://doi.org/10.1029/2007JD009196>, 2008.



- Melnikova, I., and Gatebe, C. K.: Vertical Profile of Cloud Optical Parameters Derived from Airborne Measurements Above,
 440 Inside and Below Clouds, *J. Quant Spectrosc. Radiat. Transf.*, 214:39-60. <https://doi.org/10.1016/j.jqsrt.2018.04.005>, 2018.
- Meyer, K., Platnick, S., Oreopoulos, L., and Lee, D.: Estimating the direct radiative effect of absorbing aerosols overlying
 marine boundary layer clouds in the southeast Atlantic using MODIS and CALIOP, *J. Geophys. Res. Atmos.*, 118, 4801–4815,
<https://doi.org/10.1002/jgrd.50449>, 2013.
- 445 Roy, D., and Zhang, H.: NASA Global Web-Enabled Landsat Data Annual Global 30 m V031 [Data set]. NASA EOSDIS
 Land Processes DAAC. Accessed 2020-06-23 from <https://doi.org/10.5067/MEaSUREs/GWELD/GWELDYR.031>, 2019.
- Sayer A. M., Hsu, N. C., Bettenhausen, C., Lee, J., Redemann, J., Schmid, B., and Shinozuka, Y.: Extending “Deep Blue”
 450 aerosol retrieval coverage to cases of absorbing aerosols above clouds: Sensitivity analysis and first case studies, *J. Geophys.
 Res. Atmos.*, 121, 4830–4854, <https://doi.org/10.1002/2015JD024729>, 2016.
- Schmid, B., Redemann, J., Russell, P. B., Hobbs, P. V., Hlavka, D. L., McGill, M. J., Holben, B. N., Welton, E. J., Campbell,
 J. R., Torres, O., Kahn, R. A., Diner, D. J., Helling, M. C., Chu, D. A., Robles-Gonzalez, C., and de Leeuw, G.:
 455 Coordinated airborne, spaceborne, and ground-based measurements of massive thick aerosol layers during the dry season in
 southern Africa, *J. Geophys. Res.*, 108, 8496, <https://doi.org/10.1029/2002JD002297>, 2003.
- Shinozuka, Y., Saide, P. E., Ferrada, G. A., Burton, S. P., Ferrare, R., Doherty, S. J., Gordon, H., Longo, K., Mallet, M.,
 Feng, Y., Wang, Q., Cheng, Y., Dobracki, A., Freitag, S., Howell, S. G., LeBlanc, S., Flynn, C., Segal-Rosenhaimer, M.,
 460 Pistone, K., Podolske, J. R., Stith, E. J., Bennett, J. R., Carmichael, G. R., da Silva, A., Govindaraju, R., Leung, R., Zhang,
 Y., Pfister, L., Ryoo, J.-M., Redemann, J., Wood, R. and Zuidema, P.: Modeling the smoky troposphere of the southeast
 Atlantic: a comparison to ORACLES airborne observations from September of 2016, *Aerosols/Field
 Measurements/Troposphere/Chemistry (chemical composition and reactions)*, doi:10.5194/acp-2019-678, 2019.
- Torres, O., Bhartia, P. K., Herman, J. R., Ahmad, Z., and Gleason, J.: Derivation of aerosol properties from satellite mea-
 465 surements of backscattered ultraviolet radiation: Theoretical basis, *J. Geophys. Res.*, 103, 17099–17110,
<https://doi.org/10.1029/98JD00900>, 1998.
- Torres, O., Jethva, H., and Bhartia, P. K.: Retrieval of Aerosol Optical Depth above Clouds from OMI Observations: Sensitivity
 470 Analysis and Case Studies, *J. Atmos. Sci.*, 69, 1037–1053, <http://doi.org/10.1175/JAS-D-11-0130.1>, 2012.



- Várnai, T., Gatebe, C., Gautam, R., Poudyal, R., & Su, W. (2019). Developing an Aircraft-Based Angular Distribution Model of Solar Reflection from Wildfire Smoke to Aid Satellite-Based Radiative Flux Estimation. *Remote Sensing*, 11(13), 1509.
- 475 Waquet, F., Riedi, J., Labonnote, L. C., Goloub, P., Cairns, B., Deuzeand, J. -L., and Tanre, D.: Aerosol Remote Sensing over Clouds Using A-Train Observations, *J. Atmos. Sci.*, 66, 2468– 2480, <https://doi.org/10.1175/2009JAS3026.1>, 2009.
- 480 Waquet, F., Cornet, C., Deuzé, J.-L., Dubovik, O., Ducos, F., Goloub, P., Herman, M., Lapyonok, T., Labonnote, L. C., Riedi, J., Tanré, D., Thieuleux, F., and Vanbauce, C.: Retrieval of aerosol microphysical and optical properties above liquid clouds from POLDER/PARASOL polarization measurements, *Atmos. Meas. Tech.*, 6, 991–1016, <https://doi.org/10.5194/amt-6-991-2013>, 2013.
- Wen, G., Marshak, A. and Cahalan, R. F.: Impact of 3D clouds on clear sky reflectance and aerosol retrieval in a biomass burning region of Brazil, *IEEE Geo. Rem. Sens. Lett.*, 3, 169-172, <https://doi.org/10.1109/LGRS.2005.861386>, 2006.
- 485 Wen, G., Marshak, A., Levy, R. C., Remer, L. A., Loeb, N. G., Várnai, T., and Cahalan, R. F.: Improvement of MODIS aerosol retrievals near clouds. *J. Geophys. Res.*, 118, 9168-9181, <https://doi.org/10.1002/jgrd.50617>, 2013.
- Wood, R.: Stratocumulus clouds, *Mon. Weather Rev.*, 140, 2373–2423, <https://doi.org/10.1175/MWR-D-11-00121.1>, 2012
- 490 Zinner, T., Mayer, B., and Schröder, M.: Determination of three-dimensional cloud structures from high-resolution radiance data, *J. Geophys. Res.*, 111, D08204, <https://doi.org/10.1029/2005JD006062>, 2006.



495 Table 1: Retrieved parameters from a total of 16 CAR BRDF cases taken on September 13, 2000 during
 SAFARI-2000 campaign.

Case	Location (°S, °E)	Time (UTC) HH: MM: SS	Solar Zenith (°)	Mean Aircraft Alt. , m, (AMSL)	Retrieved COD	Retrieved ACAOD (below aircraft)	Retrieved ACAOD (above aircraft)	AATS AOD
a	20.67, 13.13	10:44:51	(24.67) 24.36	1420±40	12±5	0.25±0.14	0.56±0.09	0.56±0.01
b	20.62, 13.12	10:50:47	24.11	1540±2	12±5	0.24±0.16	0.55±0.09	0.55±0.00
c	20.62, 13.12	10:53:21	24.04	1541±2	11±5	0.27±0.18	0.55±0.09	0.55±0.01
d	20.61, 13.13	11:01:13	23.95	1533±2	8±3	0.34±0.18	0.58±0.09	0.55±0.00
e	20.24, 13.20	11:18:00	23.94	1814±259	7±2	0.32±0.21	0.55±0.10	0.55±0.01
f	20.24, 13.20	11:21:00	24.09	2646±223	7±3	0.33±0.19	0.45±0.11	0.48±0.01
g	20.25, 13.20	11:23:47	(24.25)	3369±250	7±4	0.41±0.17	0.32±0.14	0.40±0.01
h	20.26, 13.22	12:28:07	(31.70) 31.88	1608±3	19±7	0.18±0.10	0.50±0.11	0.52±0.00
i	20.48, 13.10	12:30:34	(32.10) 32.28	1613±2	19±7	0.19±0.10	0.49±0.10	0.51±0.00
j	20.47, 13.10	12:33:00	32.69	1614±3	18±6	0.19±0.10	0.50±0.10	0.51±0.01



k	20.47, 13.11	12:35:30	33.11	1616±3	17±5	0.19±0.10	0.52±0.10	0.52±0.01
l	20.47, 13.11	12:37:58	33.54	1615±3	16±4	0.19±0.10	0.52±0.10	0.51±0.01
m	20.47, 13.11	12:40:28	33.97	1614±3	17±6	0.19±0.10	0.52±0.10	0.52±0.00
n	20.47, 13.11	12:45:25	34.85	1615±1	25±10	0.17±0.08	0.47±0.10	0.51±0.00
o	20.46, 13.12	12:47:55	35.30	1614±2	28±11	0.17±0.08	0.45±0.11	0.50±0.00
p	20.46, 13.13	12:50:23	35.76	1614±2	29±10	0.17±0.08	0.44±0.11	0.50±0.00



Table 2: Measured spectral albedo (with atmosphere) for each BRDF case.

Case	Wavelength (μm)							
	0.340	0.381	0.472	0.682	0.870	1.036	1.219	1.273
a	0.32	0.38	0.41	0.45	0.46	0.47	0.41	0.40
b	0.34	0.40	0.44	0.48	0.49	0.51	0.44	0.43
c	0.32	0.38	0.41	0.45	0.46	0.47	0.41	0.40
d	0.25	0.30	0.31	0.33	0.34	0.35	0.31	0.30
e	0.22	0.26	0.28	0.30	0.31	0.32	0.28	0.27
f	0.23	0.26	0.27	0.30	0.31	0.31	0.27	0.27
g	0.23	0.27	0.27	0.31	0.31	0.32	0.27	0.27
h	0.42	0.51	0.54	0.60	0.62	0.64	0.55	0.53
i	0.40	0.48	0.52	0.57	0.58	0.61	0.52	0.50
j	0.40	0.47	0.51	0.56	0.57	0.60	0.51	0.49
k	0.39	0.47	0.50	0.55	0.56	0.58	0.50	0.49
l	0.39	0.47	0.50	0.55	0.57	0.59	0.50	0.49
m	0.40	0.48	0.51	0.57	0.58	0.60	0.52	0.50
n	0.45	0.55	0.59	0.65	0.68	0.70	0.59	0.57



o	0.47	0.57	0.62	0.69	0.71	0.73	0.61	0.59
p	0.49	0.59	0.64	0.71	0.73	0.75	0.62	0.61

500

Table 3. Key parameters of the performed radiative transfer simulations.

Parameter	Value
Aircraft altitude	1.6 km
Cloud base and top altitudes	0.5 km, 1 km
Base and top altitudes of homogeneous aerosol layer	1 km, 2.5 km
Cloud optical depth (COD)	Linear decrease from the edge to the center line of a 300 m wide and infinitely long trough. Outside trough: COD = 17; center line of trough: COD = 7 or 4.7.
Cloud droplet effective radius	10 μm
Aerosol optical depth at 0.5 μm	Above CAR: 0.5; below CAR: 0.35 (0 in some tests)
Aerosol size distribution	Small mode of MODIS absorbing smoke model in Levy et al. (2007)
Aerosol absorption	Refractive index: $1.5 + i \cdot 0.033$. Resulting single scattering albedos: 0.85 at 0.47 μm and 0.79 at 0.87 μm
Surface albedo	0.05
Solar zenith angle	33°
Viewing zenith angle	0°

Table 4: Table 4. Simulated CAR BRFs at the center of a hypothetical trough.

	BRF _{0.47 μm}	BRF _{0.87 μm}	BRF _{0.47 μm} / BRF _{0.87 μm}
No BCA, COD=7.0, 1D	0.28861 \pm 0.00007	0.34162 \pm 0.00007	0.84483 \pm 0.00038
No BCA, COD=7.0, 3D	0.35663 \pm 0.00008	0.42296 \pm 0.00008	0.84318 \pm 0.00035
No BCA, COD=4.7, 3D	0.28829 \pm 0.00008	0.34243 \pm 0.00008	0.84189 \pm 0.00044
Yes BCA, COD=7.0, 1D	0.25203 \pm 0.00004	0.32416 \pm 0.00006	0.77749 \pm 0.00027
Yes BCA, COD=7.0, 3D	0.31018 \pm 0.00006	0.40075 \pm 0.00007	0.77400 \pm 0.00028



Yes BCA, COD=4.7, 3D	0.25037 ± 0.00005	0.32414 ± 0.00006	0.77241 ± 0.00030
----------------------	-----------------------	-----------------------	-----------------------

505



Figure 1: Location of the measurements. On 13 September 2000, the NASA's Cloud Absorption Radiometer (CAR) on board the University of Washington Convair-580 research aircraft obtained measurements over marine stratocumulus offshore of Namibia at several locations marked by the aircraft ground track on the map inset. The aircraft completed multiple circular flight tracks (>16) at different locations, shown on the enlarged map of the rectangular box area, and labelled alphabetically, a-p, based on the time of observations (see Table 1). The circular flight tracks were performed primarily for the airborne measurements of bidirectional reflectance distribution function (BRDF) (cases a-d and h-p), and in a few instances (cases e-g) represent vertical profiles for physical and chemical measurements. The marine stratus clouds were extensive as seen by the MODIS/Terra instrument on the same day around 09:25 UTC (see the map inset). The CV-580 flight began just prior to 10:00 UTC and ended at 13:00 UTC. The enlarged map is derived from GWELD product generated browse image (Roy and Zhang 2019).

Figure 2: (a). The University of Washington's Convair-580 research aircraft in Pietersburg, South Africa, for SAFARI 2000. (b) Schematic of NASA's Cloud Absorption Radiometer (CAR), which was mounted in the nose of the CV 580 aircraft. (c) A cumulonimbus cloud observed with CAR during Flight No. 2034 on September 14, 2011, 18:35 - 18:40 UTC, in Florida to illustrate the kind of images acquired by CAR. (d) Specifications for the CAR, which contains 14 narrow spectral bands between 0.34 and 2.30 μm .

Figure 3: CAR quicklook image (constructed from three bands at 1.04, 0.87, and 0.47 μm) obtained over the marine stratocumulus clouds. The circular flight track by the aircraft allows the CAR to image the sky and surface in all viewing zenith and azimuthal angles, and covering an area defined by a diameter of about 4 km on the surface (assuming the aircraft is flying 600 m above the surface). The unique feature of these measurements is the solar disks, which define the start and end point for each circle. A prominent feature of the marine stratocumulus clouds is the presence of the cloud bow ring associated with scattering by water droplets and with a peak at $\sim 75^\circ$ zenith angle in the antisolar direction.

Figure 4: Measured angular distribution of sky radiance (a & c) and cloud reflected radiance (b & d) at selected wavelengths ($\lambda=0.682 \mu\text{m}$ and $\lambda=0.874 \mu\text{m}$) obtained at about 12:47:55 UTC with a solar zenith angle of $\sim 35.30^\circ$ (Table 1: case o). The view zenith angle (θ) on the polar plots is represented as the radial distance from the center (0°) towards the periphery (90°) and the azimuthal angle (ϕ) as the arc length from the solar principal plane ($0^\circ \leq \phi \leq 360^\circ$). The principal plane is within the $0^\circ - 180^\circ$ azimuthal plane (the vertical plane passing through the solar position). Figures 4e & 4f show measured radiance at eight CAR spectral bands (0.34-1.27 μm) (sky and clouds) at a constant view zenith angle (50°) at different azimuthal planes angled 0° , 45° , 90° , 135° , 180° , 225° , 270° and 315° .

Figure 5: BRDF at $\lambda=0.472 \mu\text{m}$ for different solar zenith angles ($23^\circ < \text{SZA} < 34^\circ$) and cloud optical thickness. The marine stratocumulus are often extensive and flat, but contain areas that have thinner clouds or even open cells that allows radiation to penetrate through and therefore have lower BRDF values as shown by the blue colors. A prominent feature of the marine stratocumulus clouds is the presence of the cloud bow ring associated with scattering by water droplets and with a peak at $\sim 75^\circ$ zenith angle in the antisolar direction.

Figure 6: BRDF at 0.874 μm obtained at different solar zenith angles ($23^\circ < \text{SZA} < 34^\circ$) and locations over the marine stratocumulus off the Skeleton coastline in Namibia for the 16 cases described in Table 1. A prominent feature of the marine stratocumulus clouds is the presence of the cloud bow ring associated with scattering by water droplets and with a peak at $\sim 75^\circ$ zenith angle in the antisolar direction.

Figure 7: Spectral albedo (with atmosphere) for all the 16 cases at $\lambda=0.470 \mu\text{m}$ and $\lambda=0.870 \mu\text{m}$.

Figure 8: Retrieved aerosol optical depth above clouds and the aircraft, obtained from the CAR sky radiance measurements. Pixels without valid retrievals are shaded white.

Figure 9: Retrieved aerosol optical depth above clouds and below the aircraft (ACAOD). Pixels without valid retrievals are shaded white.

Figure 10: Retrieved cloud optical depth. Pixels without valid retrievals are shaded white.

Figure 11: Scatter plot ACAOD vs COD for view zenith angles 0° - 30° (blue color dots), 30° - 60° (green color dots), and 60° - 90° (red color dots).

Figure 12: Comparison of the retrieved parameters averaged over all the viewing directions for each case (a-p).




Figure 1: Location of the measurements. On 13 September 2000, the NASA's Cloud Absorption Radiometer (CAR) on board the University of Washington Convair-580 research aircraft obtained measurements over marine stratocumulus offshore of Namibia at several locations marked by the aircraft ground track on the map inset. The aircraft completed multiple circular flight tracks (>16) at different locations, shown on the enlarged map of the rectangular box area, and labelled alphabetically, a-p, based on the time of observations (see Table 1). The circular flight tracks were performed primarily for the airborne measurements of bidirectional reflectance distribution function (BRDF) (cases a-d and h-p), and in a few instances (cases e-g) represent vertical profiles for physical and chemical measurements. The marine stratus clouds were extensive as seen by the MODIS/Terra instrument on the same day around 09:25 UTC (see the map inset). The CV-580 flight began just prior to 10:00 UTC and ended at 13:00 UTC. The enlarged map is derived from GWELD product generated browse image (Roy and Zhang 2019).




Figure 2: (a). The University of Washington’s Convair-580 research aircraft in Pietersburg, South Africa, for SAFARI 2000. (b) Schematic of NASA’s Cloud Absorption Radiometer (CAR), which was mounted in the nose of the CV 580 aircraft. (c) A cumulonimbus cloud observed with CAR during Flight No. 2034 on September 14, 2011, 18:35 - 18:40 UTC, in Florida to illustrate the kind of images acquired by CAR. (d) Specifications of CAR, which contains 14 narrow spectral bands between 0.34 and 2.30 μm.






Figure 3: CAR quicklook image (constructed from three bands at 1.04, 0.87, and 0.47 μm) obtained over the marine stratocumulus clouds. The circular flight track by the aircraft allows the CAR to image the sky and surface in all viewing zenith and azimuthal angles, and covering an area defined by a diameter of about 4 km on the surface (assuming the aircraft is flying 600 m above the surface). The unique feature of these measurements is the solar disks, which define the start and end point for each circle. A prominent feature of the marine stratocumulus clouds is the presence of the cloud bow ring associated with scattering by water droplets and with a peak at $\sim 75^\circ$ zenith angle in the antisolar direction.




(a) sky radiance ($0.682 \mu\text{m}$)




(b) cloud reflected radiance ($0.682 \mu\text{m}$)




(c) sky radiance ($0.874 \mu\text{m}$)



(d) cloud reflected radiance ($0.874 \mu\text{m}$)



(e) sky spectral radiance ($\theta=50^\circ$)



(f) cloud spectral reflected radiance ($\theta=50^\circ$)






Figure 4: Measured angular distribution of sky radiance (a & c) and cloud reflected radiance (b & d) at selected wavelengths ($\lambda=0.682 \mu\text{m}$ and $\lambda=0.874 \mu\text{m}$) obtained at about 12:47:55 UTC with a solar zenith angle of $\sim 35.30^\circ$ (Table 1: case o). The view zenith angle (θ) on the polar plots is represented as the radial distance from the center (0°) towards the periphery (90°) and the azimuthal angle (ϕ) as the arc length from the solar principal plane ($0^\circ \leq \phi \leq 360^\circ$). The principal plane is within the $0^\circ - 180^\circ$ azimuthal plane (the vertical plane passing through the solar position). Figures 4e & 4f show measured radiance at eight CAR spectral bands ($0.34\text{-}1.27 \mu\text{m}$) (sky and clouds) at a constant view zenith angle (50°) at different azimuthal planes angled $0^\circ, 45^\circ, 90^\circ, 135^\circ, 180^\circ, 225^\circ, 270^\circ$ and 315° .




Figure 5: BRF at $\lambda=0.472 \mu\text{m}$ for different solar zenith angles ($23^\circ < \text{SZA} < 34^\circ$) and cloud optical thickness. The marine stratocumulus are often extensive and flat, but contain areas that have thinner clouds or even open cells that allows radiation to penetrate through and therefore have lower BRF values as shown by the blue colors. A prominent feature of the marine stratocumulus clouds is the presence of the cloud bow ring associated with scattering by water droplets and with a peak at $\sim 75^\circ$ zenith angle in the antisolar direction.




Figure 6: BRF at $0.874 \mu\text{m}$ obtained at different solar zenith angles ($23^\circ < \text{SZA} < 34^\circ$) and locations over the marine stratocumulus off the Skeleton coastline in Namibia for the 16 cases described in Table 1. A prominent feature of the marine stratocumulus clouds is the presence of the cloud bow ring associated with scattering by water droplets and with a peak at $\sim 75^\circ$ zenith angle in the antisolar direction.

600




Fig. 7: Spectral albedo (with atmosphere) for all the 16 cases at $\lambda=0.470 \mu\text{m}$ and $\lambda=0.870 \mu\text{m}$.

605




Figure 8: Retrieved aerosol optical depth above clouds and the aircraft, obtained from the CAR sky radiance measurements. Pixels without valid retrievals are shaded white.





Figure 9: Retrieved aerosol optical depth above clouds and below the aircraft (ACAOD). Pixels without valid retrievals are shaded white.



615 Figure 10: Retrieved cloud optical depth. Pixels without valid retrievals are shaded white.





Figure 11: Scatter plot ACAOD vs COD for view zenith angles 0°-30° (blue color dots), 30°-60° (green color dots), and 60°-90° (red color dots).



Comparison of the retrieved parameters averaged over all the viewing directions for each case (a-p).