

We provide here again our answer to Referee #2, where we have explained all changes that were made to the original manuscript version. Starting on page 4, we have then attached the manuscript in the “tracked-changes” version.

Answer to Anonymous Referee #2

- 5 Thank you very much for your positive evaluation of our manuscript and the helpful comments. Below, we address your individual comments and describe the associated changes made in the revised manuscript version.

Anonymous Referee #2

Spectroscopic methods are often use to probe regions of the atmosphere that are not easy to get to, so to speak. The ability to use these methods to obtain information on the composition, number, size and shape of atmospherically important compounds
10 *relies on the availability of high-quality, wavelength-dependent complex refractive indices. This manuscript contributes nicely to the library of such data, this time for ammonium nitrate (AN) at low temperature. The authors have carried out a carefully planned experiment and have followed it up with a detailed optical analysis that sets some basic limits on the applicability of the refractive indices with respect to the value of the real index at high wavenumber as well as shape factor and aspect ratio. To no one’s surprise, the optical constants at low temperature are strikingly different from those at room temperature. The*
15 *authors demonstrate the utility of using temperature-appropriate indices in the workup of measurements taken in the field. I for one appreciated the discussion in Appendix B regarding the shifting of peaks in the fingerprint region below 1500 cm⁻¹.*

Now for some general comments...

- 1.) Can the authors provide a sense of how many spectra were recorded and how long the scans were? Although I do not think this is a problem when using AIDA, I was looking for verification that the size distributions of the AN particles were stable*
20 *over the data collection period.*

Yes, in the AIDA chamber the number concentration of the AN aerosol particles was very stable during the period where the size distribution measurements and FTIR scans were conducted. Due to sedimentation and sampling losses, the number concentration of the crystallized AN particles decreased by about 5% per hour. A combined size distribution measurement with the SMPS and APS instruments took about 6 min, so the number concentration of the AN particles only varied by about
25 0.5% within that period. For the FTIR measurements, we adjusted the number of scans such that the overall recording time was also about 6 min, overlapping with the period where the size distribution of the AN particles was measured. The spectrum shown in Fig. 2a was thus averaged over 500 scans at 0.5 cm⁻¹ resolution. We will add this information to the revised manuscript version:

Line 114: “500 individual scans were averaged for each spectrum.”

- 30 Line 140: “Note that these two measurements were carried out in an overlapping time range and lasted about 6 min. During this time, N_{aer} varied by less than 0.5%.”

2.) With regard to the real index at the anchor point, presumably if the density of AN was available as a function of temperature, the Lorentz-Lorenz transform could be used to to get a good estimate of n at 223 K. Nevertheless, the small range used by the authors provides a general sense of the sensitivity of these types of retrievals to the anchor point value.

35 Yes, that would be a good way to better define the value for the anchor point. However, we must also consider the polymorphic phase change of the solid AN particles from phase IV to V when going to lower temperatures, which might also affect the density and the refractive index (but we did not find any measurements in the literature). Since we could not limit this value any better at this time, we thought it best to define a reasonable range of uncertainty for the anchor point value and examine its impact on the retrieval results.

40 3.) *In Appendix A, the authors note that they extended their computation below 800 cm⁻¹ which is typical in such work to avoid truncation errors in the finite evaluation of the Kramers-Kronig integral. How far was this extension and what was k set to?*

Yes, we will extend this discussion in the revised manuscript version. As said in line 315, we used the room-temperature k data from Jarzembki et al. (2003) for the extension. These data extend to 500 cm⁻¹, but only show one further, spectrally narrow absorption band at 717 cm⁻¹ with a maximum k value of about 0.1, thus ten times smaller than the $\nu_2(\text{NO}_3^-)$ mode of AN at 831 cm⁻¹. The influence of the non-measured range below 800 cm⁻¹ on the Kramers-Kronig integral was therefore rather small. For the extension, we considered the 800 – 690 cm⁻¹ range to fully capture the additional 717 cm⁻¹ band, interpolated the Jarzembki et al. (2003) k data to our wavenumber resolution, and added them to our measured spectrum below 800 cm⁻¹ prior to performing the Kramers-Kronig integration. We will extend our statement on line 315 as follows:

45 “These data are available down to 500 cm⁻¹, but just show one further, spectrally narrow absorption band at 717 cm⁻¹ with a maximum k value of about 0.1. To fully capture this mode, it was sufficient to consider the range from 800 to 690 cm⁻¹ for the extension. We interpolated the Jarzembki et al. (2003) k data in this range to the resolution of our measurements and added them to the $k(\tilde{\nu})$ spectrum below 800 cm⁻¹ before the Kramers-Kronig integration.”

50 4.) *Out of interest, have the authors done a retrieval for which k in the inter-band regions was not set to 0.002 (or zero beyond 3500 cm⁻¹), i.e., performed the retrieval with all of the baseline noise in place? If so, are the inter-band values of k warranted?*

55 We have first tried to perform the retrieval with the original data, but it has led to a considerable noise in the k data in the inter-band regions and was also extremely time-consuming because our approach requires that each wavenumber point is individually optimized (see also answer to point 5 below). Smoothing and/or setting the inter-band values to a small residual value was therefore the only practicable solution. For the wavenumber ranges between 3500 and 2800 cm⁻¹ and between 2000 and 800 cm⁻¹, we have included all relevant spectral subsections with distinct AN infrared bands (as also tabulated in Fernandes et al., 1979) in the full k retrieval, meaning that there should not be any unconsidered absorption in the inter-band regions where we have set k to 0.002. However, the situation is different in the range between 2800 and 2000 cm⁻¹, where absorption is generally very low, but some very weak infrared bands, mostly combination modes, have been assigned by Fernandes et al. (1979) (see Table 3 therein). Here, it is important to say that we do not resolve these modes with our approach, and we will do so in the revised manuscript version, line 369:

65 “Note that Fernandes et al. (1979) have assigned further, very weak infrared bands of AN, mostly combination modes, in the region between 2000 and 2800 cm⁻¹. We were not able to detect these signatures in our measured spectrum, hence, these bands are not represented in our retrieved $k(\tilde{\nu})$ spectrum.”

5.) Additionally, what is meant by the occurrence of "singular spikes" which required the application of a smoothing function. Did they originate in the spectrum from which the initial estimate of k was derived? Again, I congratulate the authors on a well-done study.

We assume that the occurrence of these "spikes" in the k spectrum results from the nature of our optimization method, where we had to include each individual wavenumber grid point in the retrieval. Sometimes the routine (optimization code) got stuck (i.e., found a local minimum) when there was still considerable noise, or even "spikes" in the retrieval result for k . A much better and faster convergence to the "global" minimum was obtained when applying a weak smoothing function. A better formulation than "to avoid the occurrence of singular spikes in the $k(\tilde{\nu})$ spectrum" (line 362) would therefore be "to improve the convergence behavior of the optimization algorithm" – we will change this in the revised manuscript version.

Ideally, as briefly outlined in Appendix A and elaborated in detail in some of the cited articles (e.g. Ossenkopf et al., 1992; Segal-Rosenhaimer et al., 2009), the retrieval of optical constants from aerosol infrared spectra would involve particles which are small enough to be treated with Rayleigh theory. In principle, smaller-sized AN crystals could easily be generated by reducing the solute concentration in the nebulizer that we used for aerosol generation. Only, the injection period to fill the huge volume of the AIDA chamber with such small particles and achieve a sufficient mass loading for a good signal-to-noise ratio in the infrared spectrum even for small absorption bands would just be too long. Hence, we had to use larger AN particles and the iterative approach to deduce the optical constants.

85 **High-resolution optical constants of crystalline ammonium nitrate for infrared remote sensing of the Asian Tropopause Aerosol Layer**

Robert Wagner¹, Baptiste Testa², Michael Höpfner¹, Alexei Kiselev¹, Ottmar Möhler¹, Harald Saathoff¹,
Jörn Ungermann³, and Thomas Leisner¹

¹Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

90 ²Department of Physics and Chemistry, University of Lyon, Lyon, France

³Institute of Energy and Climate Research, Stratosphere, Forschungszentrum Jülich, Jülich, Germany

Correspondence to: Robert Wagner (robert.wagner2@kit.edu)

Abstract. Infrared spectroscopic observations have shown that crystalline ammonium nitrate (AN) particles are an abundant
95 constituent of the upper tropospheric aerosol layer which is formed during the Asian summer monsoon period, the so-called
Asian Tropopause Aerosol Layer (ATAL). At upper tropospheric temperatures, the thermodynamically stable phase of AN is
different from that at 298 K, meaning that presently available room-temperature optical constants of AN, that is, the real and
imaginary parts of the complex refractive index, cannot be applied for the quantitative analysis of these infrared measurements.
In this work, we have retrieved the first low-temperature data set of optical constants for crystalline AN in the 800 – 6000
100 cm^{-1} wavenumber range with a spectral resolution of 0.5 cm^{-1} . The optical constants were iteratively derived from an infrared
extinction spectrum of $1 \mu\text{m}$ -sized AN particles suspended in a cloud chamber at 223 K. The uncertainties of the new data set
were carefully assessed in a comprehensive sensitivity analysis. We show that our data accurately fit aircraft-borne infrared
measurements of ammonium nitrate particles in the ATAL.

1 Introduction

105 The term “Asian Tropopause Aerosol Layer” (ATAL) has been established in 2011 when analyses of CALIPSO lidar
measurements have revealed the existence of a layer of enhanced aerosol concentrations at altitudes from 13 to 18 km during
the Asian summer monsoon (Vernier et al., 2011). The formation of the ATAL is related to the strong convection within the
Asian monsoon system, leading to the transport of boundary layer pollutants to the tropical tropopause layer (Vernier et al.,
2018). Since its original discovery, the ATAL has been investigated by a variety of balloon-borne, aircraft-borne, and long-
110 term satellite measurements (Höpfner et al., 2019; Lau et al., 2018; Thomason and Vernier, 2013; Vernier et al., 2018; Vernier
et al., 2015; Yu et al., 2017). Additionally, chemical transport models have been used to simulate the chemical composition
and concentration of aerosol particles lifted to and formed in the ATAL (Fairlie et al., 2020; Gu et al., 2016).

The model simulations predicted that nitrate aerosol is a prominent constituent of the ATAL. Ion chromatographic analyses of
115 filter samples collected during the BATAL campaigns (Balloon Measurements of the ATAL) (Vernier et al., 2018), as well as

in situ aerosol mass spectrometer measurements during high-altitude research aircraft flights within the StratoClim project (Stratospheric and upper tropospheric processes for better climate predictions) (Höpfner et al., 2019), confirmed the presence of nitrate aerosol particles. Another important step forward in understanding the chemical composition of the ATAL was made by the analysis of infrared limb observations from various satellite missions dating back as far as 1997 and from the above-
120 mentioned StratoClim research flights over Nepal and India in 2017 (Höpfner et al., 2019). A distinct infrared signature at 831 cm^{-1} with a width of about 3 cm^{-1} , consistently observed for spectra recorded in the ATAL during the Asian summer monsoon, was ascribed to the $\nu_2(\text{NO}_3^-)$ mode of crystalline ammonium nitrate (AN), suggesting that AN is an ubiquitous part of the ATAL. The presence of NH_3 in the upper troposphere during the Asian summer monsoon, as required for the neutralization of nitric acid and the formation of AN particles (Wang et al., 2020), had already been detected by previous satellite observations
125 (Höpfner et al., 2016).

To support the interpretation of the satellite and aircraft-based infrared measurements, laboratory infrared extinction spectra of crystalline AN particles and supercooled aqueous AN solution droplets have been recorded at temperature conditions of the upper troposphere in the aerosol and cloud simulation chamber AIDA (Aerosol Interaction and Dynamics in the Atmosphere)
130 (Höpfner et al., 2019). These measurements confirmed that the signature at 831 cm^{-1} is due to AN particles in the crystalline state, because the respective $\nu_2(\text{NO}_3^-)$ mode of supercooled aqueous AN solution droplets is shifted to 829 cm^{-1} and has a larger width of about 6 cm^{-1} (Höpfner et al., 2019). The crystallization of AN from aqueous AN solution droplets can be induced by various heterogeneous mechanisms, including contact efflorescence (Davis et al., 2015), inclusion of insoluble dust grains (Han et al., 2002), or admixing small amounts of ammonium sulfate (AS), which precipitates first and
135 triggers the crystallization of AN (Schlenker et al., 2004; Schlenker and Martin, 2005). The phase state of AN is crucial for predicting the particles' impact on cloud formation in the upper troposphere. Whereas supercooled aqueous AN solution droplets only favor cirrus formation by homogeneous ice nucleation (Cziczo and Abbatt, 2001; Koop et al., 2000), crystalline AN particles induce heterogeneous ice formation at much lower relative humidities compared to homogeneous freezing conditions (Shilling et al., 2006; Wagner et al., 2020).

140
To enable the quantitative analysis of the infrared limb observations with respect to the total mass of AN, the infrared $\nu_2(\text{NO}_3^-)$ mode recorded in the AIDA chamber has been scaled by the particles' mass concentration to yield mass-specific absorption coefficients (Höpfner et al., 2019). More versatile input parameters for the accurate retrieval of aerosol parameters from infrared remote sensing measurements are the so-called optical constants, that is, the real and imaginary parts of the complex refractive index (Bohren and Huffman, 1983; Zsazsky et al., 2007). To date, there exists only one data set of the
145 complex refractive index of solid ammonium nitrate at infrared wavelengths (Jarzembski et al., 2003). The data were derived from the analysis of room-temperature infrared spectra of AN compiled in a spectroscopic library, measured using powders dispersed in an inert paraffin oil. For two reasons, however, this data set cannot be applied for the analysis of the infrared limb observations of the ATAL. First, the optical constants are not provided at sufficient spectral resolution to mimic the relatively

150 narrow $\nu_2(\text{NO}_3^-)$ mode of AN at 831 cm^{-1} . Second, crystalline AN is known to undergo various thermal phase transitions (Chellappa et al., 2012; Herrmann and Engel, 1997). Specifically, there are different thermodynamically stable phases at room temperature (phase IV) and upper tropospheric temperatures (phase V, stable below 255 K), going along with marked changes of the habitus of the corresponding infrared spectra in certain wavenumber regimes, e.g. in the region between 1300 and 1500 cm^{-1} with contributions from the $\nu_4(\text{NH}_4^+)$ and $\nu_3(\text{NO}_3^-)$ modes (Fernandes et al., 1979; Koch et al., 1996; Shen et al., 1993; Théorêt and Sandorfy, 1964).

In this work, we have derived the first data set of complex refractive indices of crystalline AN at low temperature (223 K) with a sufficient spectral resolution (0.5 cm^{-1}) to resolve the $\nu_2(\text{NO}_3^-)$ mode. The optical constants were deduced for the wavenumber range from 800 to 6000 cm^{-1} , so that they can be employed for the analysis of measurements in any appropriate region in the mid-infrared not too strongly affected by trace gas signatures. The complex refractive indices were iteratively derived from an infrared extinction spectrum of about $1\text{ }\mu\text{m}$ -sized, almost pure crystalline AN particles suspended in the AIDA chamber at 223 K (Sects. 2 and 3.1). A comprehensive sensitivity analysis shows the effect of measurement uncertainties and different assumptions on the shape of the AN particles (spherical or aspherical particle habits) on the optical constants deduced (Sect. 3.2). We demonstrate that this new data set is appropriate to accurately fit infrared limb observations of AN in the ATAL (Sect. 3.3).

2 Methods

2.1 Experimental setup

The experimental setup has been described in detail in our preceding articles (Höpfner et al., 2019; Wagner et al., 2020) and will only briefly be summarized here. Figure 1 shows a scheme of the AIDA aerosol and cloud chamber facility with the instrumentation employed in the present study (Möhler et al., 2003; Wagner et al., 2006b; Wagner et al., 2009). The 84 m^3 sized aerosol vessel, placed inside a temperature controlled housing, was cooled to a temperature of 223 K to record the low-temperature infrared extinction spectrum of crystalline AN particles in their thermodynamic phase V. The mean gas temperature was recorded by averaging the measurements of four vertically oriented thermocouple sensors mounted at different heights of the chamber with an uncertainty of $\pm 0.3\text{ K}$. The RH_w (relative humidity with respect to supercooled water) was controlled to 22% by evaporating an equivalent amount of purified water (GenPure Pro UV ultrapure water system, Thermo Scientific) into the chamber. The resulting water vapor partial pressure, as measured by scanning the intensity of the rotational-vibrational water vapor absorption line at $1.37\text{ }\mu\text{m}$ with a tunable diode laser (TDL) (Fahey et al., 2014), was divided by the saturation water vapor pressure over liquid supercooled water to compute RH_w (Murphy and Koop, 2005).

After the AIDA chamber was set to the desired temperature and relative humidity conditions, the injection of the mixed AN/AS aerosol particles was started. As described above, a small admixture of AS was needed to catalyze the crystallization of AN.

We prepared an aqueous solution of 99.4 mol% AN and 0.6 mol% AS with an overall solute concentration of about 9 g/100 ml by dissolving respective amounts of AN (99%, VWR Chemicals) and AS (99.5% Merck) in purified water. The aqueous solution was aerosolized with an ultrasonic nebulizer (GA 2400, SinapTec). The particle flow of the mixed AN/AS solution droplets was first dried to $RH_w \leq 3\%$ with a set of silica gel diffusion dryers (Topas GmbH) and then injected into the cooled
185 AIDA chamber.

In our preceding studies, we have analyzed in detail the composition-dependent crystallization behavior of the mixed AN/AS solution droplets by varying the amount of AS and have found three different scenarios (Höpfner et al., 2019; Wagner et al., 2020). (i) Pure, 100 mol% AN solution droplets did not crystallize at all and could be maintained for at least 4 hr in the
190 supercooled liquid state at 223 K. (ii) Mixed AN/AS solution droplets with an AS admixture of ≥ 10 mol% already crystallized during the short transit time through the diffusion dryers and were immediately present as crystalline particles upon injection into the AIDA chamber. (iii) Mixed AN/AS solution droplets with smaller AS admixtures of 0.6 and 2.9 mol% gradually crystallized at low temperature inside the AIDA chamber, with the crystallization rate depending on RH_w . We used two in situ
195 extinction spectroscopy as well as laser light scattering and depolarization measurements. A Fourier Transform Infrared (FTIR) spectrometer (IFS66v, Bruker) was coupled to an internal multiple reflection cell to measure aerosol infrared extinction spectra over an optical path length of 166.8 m at wavenumbers between 800 and 6000 cm^{-1} with a resolution of 0.5 cm^{-1} (Wagner et al., 2006a). 500 individual scans were averaged for each spectrum. The gradual crystallization of the AN/AS solution droplets was indicated by the continuous decrease of the liquid water extinction band at about 3500 cm^{-1} . The near backscattering linear
200 depolarization ratio, δ , of the aerosol particles at a scattering angle of 178° and a wavelength of 488 nm was measured by the light scattering instrument SIMONE (Schnaiter et al., 2012). A continuous increase of δ over time was indicative of the formation of aspherical, crystalline AN particles and the concomitant loss of spherical, non-depolarizing solution droplets. The crystallization of the aerosol population was completed when a steady-state δ value was reached.

205 With regard to the crystallization experiment with the 99.4 mol% AN and 0.6 mol% AS solution droplets, the time series of δ indicated that the entire particle population had crystallized within a period of 1 hr after aerosol injection (see Fig. 2a in Wagner et al. (2020)). The infrared extinction spectrum of the almost pure, crystallized AN particles is shown in Fig. 2a of our article, including the AN vibrational assignment according to Fernandes et al. (1979). The minor fraction of AS likely crystallizes as one of the double salts $2\text{AN}\cdot\text{AS}$ or $3\text{AN}\cdot\text{AS}$ (Bothe and Beyer, 2007; Schlenker and Martin, 2005), but their specific extinction
210 signatures, e.g. the additional sulfate mode at about 1100 cm^{-1} (Wagner et al., 2020), are barely visible. We can therefore treat the infrared measurement to a good approximation as a pure AN spectrum.

Concomitantly with their infrared spectrum, we measured the particles' number size distribution by combining the size spectra from a scanning mobility particle sizer (SMPS, model 3071A, TSI, mobility diameter range 0.014 – 0.82 μm) and an aerodynamic particle spectrometer (APS, model 3321, TSI, aerodynamic diameter range 0.523 – 19.81 μm). To convert the mobility diameter of the crystalline AN particles from the SMPS measurement into the equal-volume sphere diameter, d_p , we adopted a dynamic shape factor, χ , of 1.1 to account for the slightly aspherical particle habits of the AN crystals (Hinds, 1999). The same value for χ as well as the particle density of AN, $\rho(\text{AN}) = 1.72 \text{ g cm}^{-3}$, were chosen to transform the aerodynamic diameter from the APS measurement into d_p (Hinds, 1999; Kelly and McMurry, 1992). The resulting number size distribution as a function of d_p is shown in Fig. 2b. The dominant particle mode mostly falls into the measurement range of the APS instrument and is centered at $d_p = 0.98 \mu\text{m}$. The integrated total number concentration of the AN particles, N_{aer} , is 985 cm^{-3} , in good agreement with an independent measurement of N_{aer} with a condensation particle counter (CPC, model 3010, TSI). The particle size measurement from Fig. 2b and the infrared spectrum from Fig. 2a are the basic input parameters for the retrieval scheme of the optical constants (Sect. 2.2). Note that these two measurements were carried out in an overlapping time range and lasted about 6 min. During this time, N_{aer} varied by less than 0.5%.

Another important parameter for the implementation of the retrieval approach is the shape of the crystalline AN particles. We have therefore recorded electron microscope images of filter-collected AN particles from the AIDA chamber (Fig. 2c). Note that these images were obtained from a different experiment with crystallized particles from 97.1 mol% AN and 2.9 mol% AS, but we do not expect that the slightly higher AS content significantly alters the particle shape. For particle sampling from the AIDA chamber, we have developed a method to collect aerosol particles in the cryo-preserved state (Wagner et al., 2017). For that purpose, we used a vacuum cryo-transfer system (EM VCT500, Leica) to ensure an unbroken chain of cryogenic transfer steps from particle sampling to electron microscopic analysis (Fig. 1). This avoids the polymorphic phase change of the solid AN particles from phase V to IV, which might induce a change of particle morphology. The cryo-transfer unit includes a liquid-nitrogen cooled transfer shuttle with a silicon substrate mounted on a copper sample holder. For particle sampling, the Si substrate was introduced through a pneumatic air lock into a sampling chamber located inside the isolating containment of the AIDA chamber. The sampling of the AN particles was electrostatically assisted by directing the aerosol flow from the AIDA chamber through a Polonium neutralizer (model 3077A, TSI) onto the sampling substrate connected to a high voltage power supply set to 2 kV. After sampling, the substrate was retracted into the pre-cooled shuttle without exposing it to the ambient air. Thereafter, the shuttle was detached from the AIDA air lock and transferred under continuous supply of liquid nitrogen to the ESEM (environmental scanning electron microscope) laboratory, where it was introduced through the air lock into the microscope (Quattro S, FEI ThermoFisher Scientific). The sample imaging was conducted in the N_2 atmosphere, thus avoiding sputtering of the particle surface normally required for high-resolution imaging in a SEM. The electron microscope images shown in Fig. 2c reveal that the crystalline AN particles are of rather compact shape with aspect ratios predominantly in the range from 0.80 to 1.25. Similar morphologies have been reported for crystallized AS particles (Earle et al., 2006).

2.2 Retrieval scheme for deriving the optical constants

The retrieval scheme, as sketched in Fig. 3, follows standard procedures for the derivation of optical constants from aerosol infrared extinction spectra (Dohm and Niedziela, 2004; Earle et al., 2006; Norman et al., 1999; Segal-Rosenheimer et al., 2009; Signorell and Luckhaus, 2002; Zsatsky et al., 2005). Briefly, an initial guess for the wavenumber-dependent spectrum of the imaginary part of the complex refractive index, $k(\tilde{\nu})$, was derived by subtracting the scattering contribution from the measured infrared extinction spectrum (step 1). Using the subtractive Kramers-Kronig transformation (Ahrenkiel, 1971; Milham et al., 1981), we then computed the real part of the complex refractive index, $n(\tilde{\nu})$, for each wavenumber grid point of the measured spectrum (step 2). In its subtractive form, the Kramers-Kronig relation needs a so-called anchor point, a known value for the real index of refraction at some reference wavenumber, $n(\tilde{\nu}_x)$. As a reasonable estimate for the anchor point, we employed a value of $n = 1.56$ at 4600 cm^{-1} from the tabulated room-temperature refractive indices by Jarzembki et al. (2003). The anchor point value was included as one of three parameters in our sensitivity analysis to investigate the variability of the retrieval results for the optical constants on the uncertainty ranges of the input parameters (Sect. 3.2). In this analysis, we also considered two other input values for $n(\tilde{\nu}_x)$, namely 1.52 and 1.60.

In step 3 of the retrieval scheme, the $n(\tilde{\nu})$ and $k(\tilde{\nu})$ spectra as well as the size distribution of the AN particles were fed into the optical model to compute the extinction spectrum. The size distribution measurement involved another sensitivity parameter, namely the dynamic shape factor χ (Hinds, 1999). As explained in Sect. 2.1, this parameter was needed to account for the effect of shape on particle motion when converting the measured mobility and aerodynamic diameters of the crystalline AN particles into equal-volume sphere diameters as input for the optical model (Fig. 2b). Our best estimate for χ was a value of 1.1, as representative for slightly aspherical particle habits (Hinds, 1999). For $\chi = 1.1$, we obtained a median particle diameter of $0.98 \text{ }\mu\text{m}$ and a total aerosol volume concentration, V_{aer} , of $410 \text{ }\mu\text{m}^3 \text{ cm}^{-3}$. In our sensitivity analysis (Sect. 3.2), we have repeated the retrieval procedure of the optical constants with two other χ values, namely 1.05 and 1.15, in order to test the impact of the uncertainty of χ on the deduced refractive indices. For $\chi = 1.05$, the median diameter of the dominant particle mode is shifted to $0.96 \text{ }\mu\text{m}$, decreasing V_{aer} by about 7% to $380 \text{ }\mu\text{m}^3 \text{ cm}^{-3}$, whereas for $\chi = 1.15$, it is shifted to $1.00 \text{ }\mu\text{m}$, increasing V_{aer} by about 7% to $440 \text{ }\mu\text{m}^3 \text{ cm}^{-3}$.

The third sensitivity parameter was the particle aspect ratio, ϕ , employed in the optical model. We first derived the optical constants using Mie theory under the assumption of spherical particles ($\phi = 1$). In the sensitivity analysis, we additionally used the T-matrix code to model the AN particles as randomly-oriented spheroids with six different aspect ratios between $\phi = 0.5$ and 2.0 (Mishchenko and Travis, 1998). After the computation of the extinction spectrum (step 4), we calculated the root-mean square deviation (RMSD) between calculated and measured infrared spectrum (step 5), and finished the loop by iteratively adjusting the initial guess $k(\tilde{\nu})$ spectrum to minimize the RMSD (step 6). The computational details of the retrieval

scheme are described in Appendix A. Figure A1 therein shows the good agreement between calculated and measured infrared spectrum after the minimization procedure.

280 3 Results and discussion

3.1 The new low-temperature data set of optical constants for crystalline AN

Figure 4 shows our newly derived, low-temperature complex refractive index data set for crystalline AN, as obtained when employing the parameter values $n(\tilde{\nu}_x) = 1.56$, $\chi = 1.1$, and $\phi = 1$ in the retrieval scheme (black line). In comparison with the room-temperature data from Jarzembki et al. (2003) (red line), the general magnitude of the imaginary indices in the various absorption regimes is similar, with maximum k values of about 0.5, 2, and 1 in the wavenumber regimes 2800 – 3500 cm^{-1} , 1250 – 1500 cm^{-1} , and 820 – 840 cm^{-1} , respectively. The low-temperature $k(\tilde{\nu})$ spectrum reveals more fine structure in the 2800 – 3500 cm^{-1} and 1250 – 1500 cm^{-1} absorption regions compared to the room-temperature data set. Very similar temperature-dependent spectral changes have been observed between the infrared spectra of AN films recorded at 270 and 90 K (Koch et al. (1996), see spectra B and C in their Fig. 4), and were attributed to the temperature-induced polymorphic phase change of AN between phase IV and phase V. The peak positions of the infrared absorption bands in our low-temperature $k(\tilde{\nu})$ spectrum, that is, 3235 cm^{-1} for $\nu_3(\text{NH}_4^+)$, 3062 cm^{-1} for $(\nu_2+\nu_4)(\text{NH}_4^+)$, 1760 cm^{-1} for $\nu_2(\text{NH}_4^+)$, 1055 cm^{-1} for $\nu_1(\text{NO}_3^-)$, and 831 cm^{-1} for $\nu_2(\text{NO}_3^-)$, are in good agreement with the tabulated values for ammonium nitrate films probed at 90 K (Koch et al., 1996). In contrast, we observed a different spectral pattern in the 1250 – 1500 cm^{-1} regime, with band maxima centered at 1356, 1418, and 1476 cm^{-1} , whereas peak positions at 1320, 1367, 1397, 1462, and 1492 cm^{-1} were reported by Koch et al. (1996). They emphasized that this particular wavenumber region was very sensitive to the procedure by which the AN films were deposited, arguing that the $\nu_4(\text{NH}_4^+)$ and $\nu_3(\text{NO}_3^-)$ modes are prone to site-sensitive coupling due to their similar frequencies. Regarding our $k(\tilde{\nu})$ spectrum, we will show in Sect. 3.2 that the 1250 – 1500 cm^{-1} wavenumber regime is also particularly sensitive to the variation of the parameter values $n(\tilde{\nu}_x)$, χ , and ϕ , which contributes to the observed frequency and intensity shifts with respect to the infrared spectrum of the AN film in that region. With regard to the $\nu_2(\text{NO}_3^-)$ mode of AN, Koch et al. (1996) did not observe any frequency shift upon cooling from 270 to 90 K. Also, room-temperature infrared measurements of solid AN particles pointed to the same band position at 831 cm^{-1} as in our low-temperature spectrum (Schlenker and Martin, 2005). The apparent shift of this band in the k spectrum reported by Jarzembki et al. (2003) (Fig. 4c) might therefore be related to the insufficient spectral resolution of the underlying infrared spectrum used for the retrieval of the optical constants.

305 3.2 Sensitivity analysis

An overview about our sensitivity analysis regarding the effect of different values for the parameters $n(\tilde{\nu}_x)$, χ , and ϕ on the retrieved $k(\tilde{\nu})$ spectrum of the AN particles is shown in Fig. 5. In the first part (panels a – c), we varied $n(\tilde{\nu}_x)$ between 1.52

and 1.60 while keeping χ and ϕ constant at 1.1 and 1, respectively. The associated changes in $k(\tilde{\nu})$ were small; most notably, an increasing value for the real refractive index at the anchor point of 4600 cm^{-1} led to slightly decreasing intensities of the $k(\tilde{\nu})$ spectrum in the N – H stretching mode regime between 2800 and 3500 cm^{-1} , whereas the k values of the $\nu_2(\text{NO}_3^-)$ mode at 831 cm^{-1} almost remained unaffected. Different values for χ (panels d – f) immediately affected the size distribution and thereby the overall volume concentration of the AN particles, V_{aer} , with lower values for χ giving rise to lower values for V_{aer} (see Sect. 2.2). Within the approximation that particle absorption is primarily governed by the magnitude of the imaginary index k , a lower value for χ , that is, V_{aer} , must be balanced by higher k values in the retrieved $k(\tilde{\nu})$ spectrum. Such behavior was indeed observed in the wavenumber regimes of the moderately intense absorption bands of AN between 2800 and 3500 cm^{-1} as well as 810 and 850 cm^{-1} . In particular, the k values were regularly increased or decreased over the whole wavenumber region of the absorption bands when χ was decreased or increased, respectively.

The above approximation breaks down in the regime of the very intense vibrational modes of AN between 1200 and 1500 cm^{-1} . With high values for the imaginary index ($k > 1$), going along with high-amplitude anomalous dispersion signals in the n spectrum (Fig. 4), the resonance condition for inducing so-called Fröhlich, or surface modes can be fulfilled, meaning that n approaches zero and k adopts a value of $\sqrt{2}$ (Bohren and Huffman, 1983; Clapp and Miller, 1993; Leisner and Wagner, 2011). In such a case, there will be an enhanced cross section in the absorption spectrum of small spherical particles and the spectral habitus of an absorption band (including e.g. band intensity and peak position) can strongly differ from the $k(\tilde{\nu})$ spectrum. Regarding our sensitivity analysis, this meant that the variation of χ , that is, V_{aer} , did not lead to a regular, predictable change in the retrieved $k(\tilde{\nu})$ spectrum as in the other wavenumber regimes with the less intense absorption bands. The variation of χ only affected a certain wavenumber region between 1320 and 1370 cm^{-1} in the retrieval result for $k(\tilde{\nu})$ (Fig. 5e). In Appendix B and the associated Figs. B1 and B2, we provide an extended description of the peculiar signature of the AN infrared spectrum in the $1200 - 1500\text{ cm}^{-1}$ regime.

Frequency and intensity of the surface modes are strongly dependent on the particle shape (Bohren and Huffman, 1983; Clapp and Miller, 1993). We therefore obtained pronounced, shape-dependent variations in the retrieval results for the $k(\tilde{\nu})$ spectrum in the $1200 - 1500\text{ cm}^{-1}$ regime when modifying the aspect ratio of the crystalline AN particles (Fig. 5g). For spherical particles with $\phi = 1$, the two most intense maxima in the retrieved $k(\tilde{\nu})$ spectrum were at 1418 and 1356 cm^{-1} , with the latter having the higher intensity. An increasing degree of asphericity inverted the intensities of these two maxima, with the 1418 cm^{-1} band, slightly shifted to higher wavenumbers, becoming the more intense one, and the 1356 cm^{-1} band, also slightly shifted to higher wavenumbers, becoming the less intense one. These shape-dependent absorption signatures induce a further degree of complexity to the infrared spectrum of AN in the $1200 - 1500\text{ cm}^{-1}$ wavenumber regime, in addition to the variability already highlighted by Koch et al. (1996) regarding the thin film AN spectra. In contrast, the shape-dependent variations of the retrieved $k(\tilde{\nu})$ spectrum in the regime of the $\nu_2(\text{NO}_3^-)$ mode were small (Fig. 5h). The maximum value for k at 831 cm^{-1} varied

by at most 6% between the retrieval results for the various aspect ratios, with a trend of decreasing maximum value with increasing degree of asphericity. The shape-related spectral changes in the 2800 – 3500 cm⁻¹ regime of the $k(\tilde{\nu})$ spectrum were insignificant and are therefore not shown.

3.3 Application of the new refractive index data set

345 We consider the refractive index data set derived for $n(\tilde{\nu}_x) = 1.56$, $\chi = 1.1$, and $\phi = 1$ as our currently best estimate for the optical constants of crystalline AN at 223 K (Fig. 4). However, we also provide the data sets retrieved for all sensitivity studies described in Sect. 3.2 and strongly encourage any modelers who apply our data to take into account these uncertainties of n and k in their own spectral analyses. Regarding the influence of particle asphericity investigated by the T-matrix calculations, we have considered aspect ratios partly exceeding the actual particles' eccentricity as revealed by the electron microscope
350 images (Fig. 2c) to ensure that we derived a valid upper estimate for the uncertainty of n and k related to the shape dependency of the infrared extinction signatures of AN. As each individual n and k data set relies on an idealized representation for the shape of the AN particles (either spherical or spheroidal with a fixed aspect ratio), modelers might also consider using a shape-averaged n and k data set in their analyses (Mishchenko et al., 1997).

355 We used the new refractive index data set of AN to re-analyze infrared limb observations of the ATAL with the airborne GLORIA (Gimballed Limb Observer for Radiance Imaging of the Atmosphere) instrument during the StratoClim research flight on 31 July 2017 (Fig. 6a). The AN mass concentrations derived with the $n(\tilde{\nu}_x) = 1.56$, $\chi = 1.1$, and $\phi = 1$ data set agree within 0.01 ± 0.06 (3σ) $\mu\text{g m}^{-3}$ with those previously estimated using the mass-specific absorption coefficients (see Fig. 3b in Höpfner et al., 2019). Maximum deviations of about ± 0.04 $\mu\text{g m}^{-3}$ for the retrieved AN mass concentrations are obtained in
360 the sensitivity analysis when considering the uncertainties of the n and k values. The measured infrared spectrum of the $\nu_2(\text{NO}_3^-)$ mode of the AN particles (blue line in Fig. 6b), showing the difference of mean GLORIA spectra recorded in periods of high and low AN mass concentrations, is accurately reproduced in the spectral fit with the new optical constants (orange line). Apart from the $\nu_2(\text{NO}_3^-)$ mode, our new data set of refractive indices also opens the possibility to exploit bands in other spectral regimes for the detection and quantification of AN in remote sensing observations of the atmosphere.

365 Appendix A – Computational details of the retrieval scheme

Here, we describe the computational details of the individual steps in the retrieval scheme for deriving the infrared complex refractive indices of AN (Fig. 3). The wavenumber-dependent complex refractive index, $N(\tilde{\nu})$, is given as $N(\tilde{\nu}) = n(\tilde{\nu}) + ik(\tilde{\nu})$, with the real (n) and imaginary (k) parts called the optical constants. Our retrieval approach takes advantage of the Kramers-Kronig relation (Bohren and Huffman, 1983), by which the real refractive index can be computed at any given
370 wavenumber $\tilde{\nu}_k$ from the full wavenumber spectrum of $k(\tilde{\nu})$ (Eq. A1).

$$n(\tilde{\nu}_k) - 1 = \frac{2}{\pi} P \int_0^{\infty} \frac{k(\tilde{\nu}) \tilde{\nu}}{\tilde{\nu}^2 - \tilde{\nu}_k^2} d\tilde{\nu} \quad (\text{A1})$$

In step 1 of our retrieval approach, where we derived an initial guess for the optical constants of AN from the measured infrared extinction spectrum, we made use of a different form of the Kramers-Kronig relation, linking the real and imaginary parts of the composite function $f = (N^2(\tilde{\nu}) - 1)/(N^2(\tilde{\nu}) + 2)$ (Rouleau and Martin, 1991).

$$\text{Re}\{f\}(\tilde{\nu}_k) = \frac{2}{\pi} P \int_0^{\infty} \frac{\text{Im}\{f\}(\tilde{\nu}) \tilde{\nu}}{\tilde{\nu}^2 - \tilde{\nu}_k^2} d\tilde{\nu} \quad (\text{A2})$$

For the initial estimate of $k(\tilde{\nu})$, we analyzed the extinction spectrum of the 1 μm -sized AN crystals in the framework of Rayleigh theory (Bohren and Huffman, 1983). The scattering contribution to extinction was subtracted from the measurement by assuming its intensity to be proportional to $\tilde{\nu}^4$ (Norman et al., 1999). In the Rayleigh limit, the so-derived absorption spectrum is directly proportional to the total aerosol volume concentration, V_{aer} , and to the imaginary part of the composite function f , $\text{Im}\{f\}$ (Ossenkopf et al., 1992). With V_{aer} given by the SMPS and APS size distribution measurements as discussed in Sect. 2.1, we could thus derive the full wavenumber spectrum of $\text{Im}\{f\}$ from the estimated absorption spectrum, perform the Kramers-Kronig integration in Eq. A2 to obtain the spectrum of $\text{Re}\{f\}$, and calculate the initial guess for $k(\tilde{\nu})$ from $\text{Re}\{f\}$ and $\text{Im}\{f\}$ (Leisner and Wagner, 2011; Segal-Rosenheimer et al., 2009).

The initial guess $k(\tilde{\nu})$ spectrum had to be iteratively adjusted using Mie theory or the T-matrix method because the generic requirement for the validity of the Rayleigh approximation, that is, the AN particles have to be very small compared to any of the wavelengths (Bohren and Huffman, 1983), was not fulfilled for the entire measurement range in the mid-infrared. The Kramers-Kronig integration in its form of Eq. A2 was therefore only applied once in step 1 of the retrieval scheme. In step 2, that was part of the loop for the iterative adjustment of $k(\tilde{\nu})$, we employed the direct relation between $n(\tilde{\nu})$ and $k(\tilde{\nu})$, implementing Eq. A1 in the version of the subtractive Kramers-Kronig transformation to minimize the effect of truncation errors due to the unknown behavior of $k(\tilde{\nu})$ beyond the experimentally accessible wavenumber range (Ahrenkiel, 1971; Milham et al., 1981; Segal-Rosenheimer and Linker, 2009).

$$n(\tilde{\nu}_k) = n(\tilde{\nu}_x) + \frac{2(\tilde{\nu}_k^2 - \tilde{\nu}_x^2)}{\pi} P \int_0^{\infty} \frac{k(\tilde{\nu}) \tilde{\nu}}{(\tilde{\nu}^2 - \tilde{\nu}_k^2)(\tilde{\nu}^2 - \tilde{\nu}_x^2)} d\tilde{\nu} \quad (\text{A3})$$

As described in Sect. 2.2, the anchor point value, $n(\tilde{\nu}_x)$, was set to 1.56 at 4600 cm^{-1} . Two other values for $n(\tilde{\nu}_x)$, namely 1.52
400 and 1.60, were adopted in our sensitivity analysis (Sect. 3.2). Prior to performing the Kramers-Kronig integration, we extended
the $k(\tilde{\nu})$ spectrum below the 800 cm^{-1} cutoff of our infrared measurements with the room-temperature k values reported by
Jarzembski et al. (2003). These data are available down to 500 cm^{-1} , but just show one further, spectrally narrow absorption
band at 717 cm^{-1} with a maximum k value of about 0.1. To fully capture this mode, it was sufficient to consider the range from
800 to 690 cm^{-1} for the extension. We interpolated the Jarzembski et al. (2003) k data in this range to the resolution of our
405 measurements and added them to the $k(\tilde{\nu})$ spectrum below 800 cm^{-1} before the Kramers-Kronig integration. The integral was
computed with Maclaurin's formula method (Ohta and Ishida, 1988).

For the Mie calculations, modeling the crystalline AN particles as spheres with an aspect ratio, ϕ , of one, we extended the Mie
code provided by Bohren and Huffman (1983) to average the computed extinction cross sections over the measured number
410 distribution of particle sizes. Due to their computational efficiency, the Mie computations were explicitly included in each
iteration, meaning that for each new adjustment of the $k(\tilde{\nu})$ and $n(\tilde{\nu})$ spectra, a new Mie calculation of the extinction spectrum
was executed. In order to investigate the effect of particle shape on the retrieval results for the optical constants, we also
modeled the AN particles as randomly-oriented spheroids, considering both oblate ($\phi > 1$) and prolate ($\phi < 1$) particle shapes.
To compute the extinction cross sections, we used the extended precision T-matrix code for randomly oriented particles by
415 Mishchenko and Travis (1998), choosing six different aspect ratios, namely $\phi = 1/2$, $\phi = 2/3$, $\phi = 4/5$, $\phi = 5/4$, $\phi = 3/2$, and $\phi = 2$.
Here, the computational burden would have been too high to perform a complete T-matrix computation of the size-averaged
extinction spectrum of the AN particles each time the $k(\tilde{\nu})$ and $n(\tilde{\nu})$ spectra were newly adjusted during the optimization
procedure. We therefore computed a priori for each aspect ratio an extinction cross section database on a three-dimensional
parameter space, which served as a look-up table in the retrieval procedure. Specifically, the extinction cross sections were
420 computed for

- 27 equal-volume sphere size parameters x_p ($x_p = \pi d_p / \lambda$) between 0.005 and 5 (0.005, 0.01, 0.03, 0.05, 0.1, 0.2 – 4.0
with $\Delta x_p = 0.2, 4.5$, and 5),
- 30 values for the real refractive index n between 0.4 and 3.3 with $\Delta n = 0.1$, and
- 29 values for the imaginary refractive index k between 0.0001 and 2.5 (0.0001, 0.025, 0.05, 0.075, 0.1 – 2.5 with Δk
425 = 0.1),

summing up to 23490 individual calculations for each aspect ratio. In the iterative loop of the retrieval scheme, the T-matrix
computed extinction spectrum for a new adjustment of $k(\tilde{\nu})$ and $n(\tilde{\nu})$ was then simply obtained by spline interpolation from
430 the pre-computed extinction cross section database. We have validated the accuracy of the interpolation scheme with the

denoted number of grid points in the (x_p, n, k) space by the comparison between directly computed and interpolated extinction cross sections.

435 To minimize the root-mean square deviation between measured and calculated extinction spectrum, we used the downhill simplex algorithm as the optimization technique in the iterative adjustment of the $k(\tilde{\nu})$ spectrum (Press et al., 1992). The measured spectrum is provided in a digital resolution of about 0.24 cm^{-1} , resulting from an approximately doubled size of the original interferogram due to zero-filling with a factor of two (Aroui et al., 2012). Overall, this amounts to 21572 wavenumber grid points for the $k(\tilde{\nu})$ spectrum in the wavenumber range between 800 and 6000 cm^{-1} . The $k(\tilde{\nu})$ spectrum of AN features regimes with vastly varying types of absorption signatures, including, for example, completely non-absorbing regions, regimes of spectrally broad but weak absorption, and regimes with strong absorption bands, either spectrally broad or narrow. In the retrieval scheme, the various spectral regimes were treated differently and not all wavenumber grid points were included as optimization parameters.

- 445 • Wavenumber region $800 - 1250 \text{ cm}^{-1}$. This regime includes the spectrally narrow $\nu_2(\text{NO}_3^-)$ and $\nu_1(\text{NO}_3^-)$ modes at 831 and 1055 cm^{-1} . Apart from these two pronounced bands, the absorption is very low. We therefore optimized the $k(\tilde{\nu})$ values in the regimes $826 - 836 \text{ cm}^{-1}$ and $1050 - 1060 \text{ cm}^{-1}$ with the full spectral resolution. All other k values were set to a small, constant background value of 0.002 in order to not transfer the noise from the baseline in the experimental spectrum to the $k(\tilde{\nu})$ spectrum.
- 450 • Wavenumber region $1250 - 1500 \text{ cm}^{-1}$. This regime comprises the intense, spectrally broad absorption bands due to the $\nu_4(\text{NH}_4^+)$ and $\nu_3(\text{NO}_3^-)$ modes. All $k(\tilde{\nu})$ values were included in the optimization, but we applied a weak smoothing function prior to the Kramers-Kronig transformation to improve the convergence behavior of the optimization algorithm to avoid the occurrence of singular spikes in the $k(\tilde{\nu})$ spectrum (Savitzky-Golay smoothing filter, quadratic polynomial fit with 5 data points in the moving window) (Press et al., 1992).
- 455 • Wavenumber region $1500 - 2800 \text{ cm}^{-1}$. This regime includes two very weak, but spectrally narrow absorption signatures between about 1750 and 1785 cm^{-1} , presumably due to the $\nu_2(\text{NH}_4^+)$ and $(\nu_1+\nu_4)(\text{NO}_3^-)$ modes (Fernandes et al., 1979; Koch et al., 1996). In that part, we optimized all $k(\tilde{\nu})$ values without any smoothing filter. Outside the $1750 - 1785 \text{ cm}^{-1}$ region, the residual absorption is spectrally broad and very small, so that we either used a strong smoothing filter or set the k values to a small, constant background value of 0.002 . Note that Fernandes et al. (1979) have assigned further, very weak infrared bands of AN, mostly combination modes, in the region between 2000 and 2800 cm^{-1} . We were not able to detect these signatures in our measured spectrum, hence, these bands are not represented in our retrieved $k(\tilde{\nu})$ spectrum.
- 460 • Wavenumber region $2800 - 3500 \text{ cm}^{-1}$. This regime comprises the intense N-H stretching modes and was treated in the same way as the $1250 - 1500 \text{ cm}^{-1}$ wavenumber region.

- Wavenumber region 3500 – 6000 cm⁻¹: In this non-absorbing regime, all $k(\tilde{\nu})$ values were set to zero.

465

A comparison between the measured (black lines) and the computed (red lines) infrared extinction spectrum of crystalline AN particles in five different wavenumber regimes after convergence of the optimization algorithm is shown in Fig. A1. The above description specifically relates to the retrievals performed with Mie theory. Regarding the T-matrix computations, the analysis of the extinction cross section data base showed that the wavenumber region above 1500 cm⁻¹ was only prone to minor shape-related changes for the considered range of aspect ratios. Therefore, the analysis of the shape-dependent variations in the retrieval results for the optical constants of AN was confined to the 800 – 1500 cm⁻¹ regime (see Figs. 5g and 5h).

470

Appendix B – Analysis of the infrared extinction signature of the AN particles between 1200 and 1500 cm⁻¹

In order to better understand the signature of the infrared extinction spectrum of the crystalline AN particles in the 1200 – 1500 cm⁻¹ wavenumber regime, we show in Fig. B1 the comparison between the measured extinction spectrum (panel a) and the retrieved optical constants for $n(\tilde{\nu}_\chi) = 1.56$, $\chi = 1.1$, and $\phi = 1$ (panel b). It is obvious that the spectral signature of the measured extinction bands of the AN particles is clearly different from that of the underlying $k(\tilde{\nu})$ spectrum, that is, different from a bulk absorption spectrum of AN whose intensity would be proportional to $k(\tilde{\nu}) \cdot \tilde{\nu}$ (Bohren and Huffman, 1983). The two most intense maxima in the $k(\tilde{\nu})$ spectrum at 1418 and 1356 cm⁻¹ (dashed red lines) are shifted to higher wavenumbers in the measured particle spectrum (1424 and 1373 cm⁻¹, dashed blue lines), accompanied by significant changes in their relative intensities. The intensity of the 1424 cm⁻¹ band is disproportionately high compared to that of the 1373 cm⁻¹ band, considering that the value of the imaginary refractive index is almost identical at both frequencies ($k \sim 1.4$). However, the real refractive index at 1424 cm⁻¹ is much lower than at 1373 cm⁻¹, thereby better fulfilling the resonance condition for the manifestation of a Fröhlich, or surface mode, in the particle spectrum ($n \approx 0$ and $k \approx \sqrt{2}$) (Bohren and Huffman, 1983; Clapp and Miller, 1993). The same reasoning explains the frequency shift of the particle extinction bands towards higher wavenumbers compared to the maxima in the $k(\tilde{\nu})$ spectrum. Although this frequency shift leads to a reduction of the magnitude of the imaginary index k , the intensity of the particle extinction bands is enhanced due to the simultaneous decrease of the value for the real refractive index n , yielding a higher cross section due to a better match with the resonance condition. For a less intense absorption band like the $\nu_2(\text{NO}_3^-)$ mode at 831 cm⁻¹, the amplitude of the anomalous dispersion feature in the $n(\tilde{\nu})$ spectrum is reduced and there are no explicit frequencies where the optical constants match the resonance condition. Therefore, the particle extinction band only experiences a minor frequency shift of about 0.4 cm⁻¹ compared to the location of the absorption peak in the $k(\tilde{\nu})$ spectrum (Fig. B2).

480

485

490

In our sensitivity analysis (Sect. 3.2), we have investigated the response of the retrieved $k(\tilde{\nu})$ spectrum for the AN particles to a change of the parameter χ , which controls the overall aerosol volume concentration, V_{aer} . We have observed that in the 1200

495 – 1500 cm^{-1} regime, the variation of χ did not induce a regular increase or decrease of the k values over the whole frequency range (Fig. 5e). Obviously, the peak at 1418 cm^{-1} in the retrieved $k(\tilde{\nu})$ spectrum is much less influenced by a change in V_{aer} than the absorption band at 1356 cm^{-1} . As a tentative explanation, one might argue that the particle extinction band at 1424 cm^{-1} , resulting from the k maximum at 1418 cm^{-1} , is less affected by V_{aer} because its intensity is primarily governed by the strongly enhanced cross sections resulting from the match of the optical constants with the resonance condition. At 1356
500 cm^{-1} , the resonance condition is of much less importance for the band intensity, meaning that the magnitude of the retrieved k values at this frequency are more directly influenced by a change of χ , that is, V_{aer} .

Data availability

Upon manuscript acceptance, we will archive all refractive index data sets derived in this work in the KITopen repository, the central publication platform for KIT (Karlsruhe Institute of Technology) scientists (Open Access, contact:
505 KITopen@bibliothek.kit.edu), and assign them a citable persistent identifier (DOI).

Author contributions

Conceptualization, methodology, and supervision: RW, MH. Formal analysis: BT, MH, RW. Investigation: RW, BT, AK, MH, HS, JU. Project administration: OM, TL. Software: BT, MH, RW. Visualization: RW, BT, AK, MH. Writing – original draft: RW, MH. Writing – review & editing: all authors.

510 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

We gratefully acknowledge the continuous support by all members of the Engineering and Infrastructure group of IMK-AAF, in particular by Olga Dombrowski, Rainer Buschbacher, Tomasz Chudy, Steffen Vogt, and Georg Scheurig. This work has
515 been funded by the Helmholtz-Gemeinschaft Deutscher Forschungszentren as part of the program “Atmosphere and Climate”. Additional funding has been received by the French Erasmus+ agency.

References

Ahrenkiel, R. K.: Modified Kramers-Kronig Analysis of Optical Spectra, J. Opt. Soc. Am., 61, 1651-1655, 1971.

- 520 Aroui, H., Orphal, J., and Tchana, F. K.: Fourier Transform Infrared Spectroscopy for the Measurement of Spectral Line Profiles, in: Fourier Transform - Materials Analysis. Ed.: S. M. Salih, InTech, New York, 69-102, 2012.
- Bohren, C. F., and Huffman, D. R.: Absorption and Scattering of Light by Small Particles, John Wiley & Sons, Inc., New York, 1983.
- 525 Bothe, J. R., and Beyer, K. D.: Experimental determination of the $\text{NH}_4\text{NO}_3/(\text{NH}_4)_2\text{SO}_4/\text{H}_2\text{O}$ phase diagram, *J. Phys. Chem. A*, 111, 12106-12117, 2007.
- Chellappa, R. S., Dattelbaum, D. M., Velisavljevic, N., and Sheffield, S.: The phase diagram of ammonium nitrate, *J. Chem. Phys.*, 137, 064504, 10.1063/1.4733330, 2012.
- 530 Clapp, M. L., and Miller, R. E.: Shape Effects in the Infrared Spectrum of Ammonia Aerosols, *Icarus*, 105, 529-536, 1993.
- Cziczo, D. J., and Abbatt, J. P. D.: Ice nucleation in NH_4HSO_4 , NH_4NO_3 , and H_2SO_4 aqueous particles: Implications for cirrus cloud formation, *Geophys. Res. Lett.*, 28, 963-966, 2001.
- 535 Davis, R. D., Lance, S., Gordon, J. A., Ushijima, S. B., and Tolbert, M. A.: Contact efflorescence as a pathway for crystallization of atmospherically relevant particles, *Proc. Natl. Acad. Sci. U.S.A.*, 112, 15815-15820, 2015.
- Dohm, M. T., and Niedziela, R. F.: Infrared complex refractive indices for nopinone, *Geophys. Res. Lett.*, 31, L14109, 10.1029/2004gl019737, 2004.
- 540 Earle, M. E., Pancescu, R. G., Cosic, B., Zsetsky, A. Y., and Sloan, J. J.: Temperature-dependent complex indices of refraction for crystalline $(\text{NH}_4)_2\text{SO}_4$, *J. Phys. Chem. A*, 110, 13022-13028, 2006.
- 545 Fahey, D. W., Gao, R. S., Möhler, O., Saathoff, H., Schiller, C., Ebert, V., Krämer, M., Peter, T., Amarouche, N., Avallone, L. M., Bauer, R., Bozóki, Z., Christensen, L. E., Davis, S. M., Durrý, G., Dyroff, C., Herman, R. L., Hunsmann, S., Khaykin, S. M., Mackrodt, P., Meyer, J., Smith, J. B., Spelten, N., Troy, R. F., Vömel, H., Wagner, S., and Wienhold, F. G.: The AquaVIT-1 intercomparison of atmospheric water vapor measurement techniques, *Atmos. Meas. Tech.*, 7, 3177-3213, 10.5194/amtd-7-3159-2014, 2014.
- 550 Fairlie, T. D., Liu, H. Y., Vernier, J. P., Campuzano-Jost, P., Jimenez, J. L., Jo, D. S., Zhang, B., Natarajan, M., Avery, M. A., and Huey, G.: Estimates of Regional Source Contributions to the Asian Tropopause Aerosol Layer Using a Chemical Transport Model, *J. Geophys. Res. (Atmos.)*, 125, e2019JD031506, 10.1029/2019JD031506, 2020.

- 555 Fernandes, J. R., Ganguly, S., and Rao, C. N. R.: Infrared Spectroscopic Study of the Phase-Transitions in CsNO₃, RbNO₃ and NH₄NO₃, *Spectrochim. Acta A*, 35, 1013-1020, 1979.
- Gu, Y. X., Liao, H., and Bian, J. C.: Summertime nitrate aerosol in the upper troposphere and lower stratosphere over the Tibetan Plateau and the South Asian summer monsoon region, *Atmos. Chem. Phys.*, 16, 6641-6663, 10.5194/acp-16-6641-560 2016, 2016.
- Han, J. H., Hung, H. M., and Martin, S. T.: Size effect of hematite and corundum inclusions on the efflorescence relative humidities of aqueous ammonium nitrate particles, *J. Geophys. Res. (Atmos.)*, 107, 4086, 10.1029/2001JD001054, 2002.
- 565 Herrmann, M. J., and Engel, W.: Phase transitions and lattice dynamics of ammonium nitrate, *Propellants Explosives Pyrotechnics*, 22, 143-147, 1997.
- Hinds, W. C.: *Aerosol Technology*, John Wiley & Sons, Inc., New York, 1999.
- 570 Höpfner, M., Volkamer, R., Grabowski, U., Grutter, M., Orphal, J., Stiller, G., von Clarmann, T., and Wetzell, G.: First detection of ammonia (NH₃) in the Asian summer monsoon upper troposphere, *Atmos. Chem. Phys.*, 16, 14357-14369, 2016.
- Höpfner, M., Ungerer, J., Borrmann, S., Wagner, R., Spang, R., Riese, M., Stiller, G., Appel, O., Batenburg, A. M., Bucci, S., Cairo, F., Dragoneas, A., Friedl-Vallon, F., Hünig, A., Johansson, S., Krasauskas, L., Legras, B., Leisner, T., Mahnke, C., 575 Möhler, O., Molleker, S., Müller, R., Neubert, T., Orphal, J., Preusse, P., Rex, M., Saathoff, H., Strohm, F., Weigel, R., and Wohltmann, I.: Ammonium nitrate particles formed in upper troposphere from ground ammonia sources during Asian monsoons, *Nature Geoscience*, 12, 608-612, 10.1038/s41561-019-0385-8, 2019.
- Jarzembski, M. A., Norman, M. L., Fuller, K. A., Srivastava, V., and Cutten, D. R.: Complex refractive index of ammonium 580 nitrate in the 2-20- μ m spectral range, *Appl. Opt.*, 42, 922-930, 10.1364/Ao.42.000922, 2003.
- Kelly, W. P., and McMurry, P. H.: Measurement of Particle Density by Inertial Classification of Differential Mobility Analyzer Generated Monodisperse Aerosols, *Aerosol. Sci. Tech.*, 17, 199-212, 10.1080/02786829208959571, 1992.
- 585 Koch, T. G., Holmes, N. S., Roddis, T. B., and Sodeau, J. R.: Low-temperature photochemistry of submicrometer nitric acid and ammonium nitrate layers, *J. Phys. Chem.*, 100, 11402-11407, 1996.

- Koop, T., Luo, B. P., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611-614, 2000.
- 590
- Lau, W. K. M., Yuan, C., and Li, Z. Q.: Origin, Maintenance and Variability of the Asian Tropopause Aerosol Layer (ATAL): The Roles of Monsoon Dynamics, *Scientific Reports*, 8, 3960, 10.1038/s41598-018-22267-z, 2018.
- Leisner, T., and Wagner, R.: Infrared Spectroscopy of Aerosol Particles, in: *Fundamentals and Applications in Aerosol Spectroscopy*, edited by: Signorell, R., and Reid, J. P., CRC Press, Boca Raton, 2011.
- 595
- Milham, M. E., Frickel, R. H., Embury, J. F., and Anderson, D. H.: Determination of optical constants from extinction measurements, *J. Opt. Soc. Am.*, 71, 1099-1106, 1981.
- 600
- Mishchenko, M. I., Travis, L. D., Kahn, R. A., and West, R. A.: Modeling phase functions for dustlike tropospheric aerosols using a shape mixture of randomly oriented polydisperse spheroids, *J. Geophys. Res. (Atmos.)*, 102, 16831-16847, 1997.
- Mishchenko, M. I., and Travis, L. D.: Capabilities and limitations of a current FORTRAN implementation of the T-matrix method for randomly oriented, rotationally symmetric scatterers, *J. Quant. Spectrosc. Radiat. Transfer*, 60, 309-324, 1998.
- 605
- Möhler, O., Stetzer, O., Schaefers, S., Linke, C., Schnaiter, M., Tiede, R., Saathoff, H., Krämer, M., Mangold, A., Budz, P., Zink, P., Schreiner, J., Mauersberger, K., Haag, W., Kärcher, B., and Schurath, U.: Experimental investigation of homogeneous freezing of sulphuric acid particles in the aerosol chamber AIDA, *Atmos. Chem. Phys.*, 3, 211-223, 2003.
- 610
- Murphy, D. M., and Koop, T.: Review of the vapour pressures of ice and supercooled water for atmospheric applications, *Q. J. R. Meteorol. Soc.*, 131, 1539-1565, 2005.
- Norman, M. L., Qian, J., Miller, R. E., and Worsnop, D. R.: Infrared complex refractive indices of supercooled liquid HNO₃/H₂O aerosols, *J. Geophys. Res. (Atmos.)*, 104, 30571-30584, 1999.
- 615
- Ohta, K., and Ishida, H.: Comparison among Several Numerical Integration Methods for Kramers-Kronig Transformation, *Appl. Spectrosc.*, 42, 952-957, 1988.
- Ossenkopf, V., Henning, T., and Mathis, J. S.: Constraints on Cosmic Silicates, *Astron. Astrophys.*, 261, 567-578, 1992.
- 620

- Press, W. H., Teukolsky, S. A., Vetterling, W. T., and Flannery, B. P.: Numerical Recipes in C: The Art of Scientific Computing, Cambridge University Press, Cambridge, 1992.
- 625 Rouleau, F., and Martin, P. G.: Shape and Clustering Effects on the Optical Properties of Amorphous Carbon, *Astrophys. J.*, 377, 526-540, 1991.
- Schlenker, J. C., Malinowski, A., Martin, S. T., Hung, H. M., and Rudich, Y.: Crystals formed at 293 K by aqueous sulfate-nitrate-ammonium-proton aerosol particles, *J. Phys. Chem. A*, 108, 9375-9383, 2004.
- 630 Schlenker, J. C., and Martin, S. T.: Crystallization pathways of sulfate-nitrate-ammonium aerosol particles, *J. Phys. Chem. A*, 109, 9980-9985, 2005.
- Schnaiter, M., Büttner, S., Möhler, O., Skrotzki, J., Vragel, M., and Wagner, R.: Influence of particle size and shape on the backscattering linear depolarisation ratio of small ice crystals – cloud chamber measurements in the context of contrail and cirrus microphysics, *Atmos. Chem. Phys.*, 12, 10465-10484, 2012.
- 635 Segal-Rosenheimer, M., Dubowski, Y., and Linker, R.: Extraction of optical constants from mid-IR spectra of small aerosol particles, *J. Quant. Spectrosc. Radiat. Transfer*, 110, 415-426, 2009.
- 640 Segal-Rosenheimer, M., and Linker, R.: Impact of the non-measured infrared spectral range of the imaginary refractive index on the derivation of the real refractive index using the Kramers-Kronig transform, *J. Quant. Spectrosc. Radiat. Transfer*, 110, 1147-1161, 2009.
- Shen, Z. X., Kuok, M. H., Tang, S. H., and Sherman, W. F.: Infrared Study of Phases III, IV and V of NH_4NO_3 , *Spectrochim. Acta A*, 49, 21-29, 1993.
- 645 Shilling, J. E., Fortin, T. J., and Tolbert, M. A.: Depositional ice nucleation on crystalline organic and inorganic solids, *J. Geophys. Res. (Atmos.)*, 111, D12204, 10.1029/2005JD006664, 2006.
- 650 Signorell, R., and Luckhaus, D.: Aerosol spectroscopy of dihydroxyacetone: Gas phase and nanoparticles, *J. Phys. Chem. A*, 106, 4855-4867, 2002.
- Théorêt, A., and Sandorfy, C.: Infrared Spectra and Crystalline Phase Transitions of Ammonium Nitrate, *Can. J. Chem.*, 42, 57-62, 1964.

Thomason, L. W., and Vernier, J. P.: Improved SAGE II cloud/aerosol categorization and observations of the Asian tropopause aerosol layer: 1989-2005, *Atmos. Chem. Phys.*, 13, 4605-4616, 10.5194/acp-13-4605-2013, 2013.

Vernier, J. P., Thomason, L. W., and Kar, J.: CALIPSO detection of an Asian tropopause aerosol layer, *Geophys. Res. Lett.*,
660 38, L07804, 10.1029/2010GL046614, 2011.

Vernier, J. P., Fairlie, T. D., Natarajan, M., Wienhold, F. G., Bian, J., Martinsson, B. G., Crumeyrolle, S., Thomason, L. W., and Bedka, K. M.: Increase in upper tropospheric and lower stratospheric aerosol levels and its potential connection with Asian pollution, *J. Geophys. Res. (Atmos.)*, 120, 1608-1619, 2015.

Vernier, J. P., Fairlie, T. D., Deshler, T., Ratnam, M. V., Gadhavi, H., Kumar, B. S., Natarajan, M., Pandit, A. K., Raj, S. T. A., Kumar, A. H., Jayaraman, A., Singh, A. K., Rastogi, N., Sinha, P. R., Kumar, S., Tiwari, S., Wegner, T., Baker, N., Vignelles, D., Stenchikov, G., Shevchenko, I., Smith, J., Bedka, K., Kesarkar, A., Singh, V., Bhate, J., Ravikiran, V., Rao, M. D., Ravindrababu, S., Patel, A., Vernier, H., Wienhold, F. G., Liu, H., Knepppp, T. N., Thomason, L., Crawford, J., Ziembmba,
670 L., Moore, J., Crumeyrolle, S., Williamson, M., Berthet, G., Jegou, F., and Renard, J. B.: BATAL: The Balloon Measurement Campaigns of the Asian Tropopause Aerosol Layer, *Bull. Amer. Meteor. Soc.*, 99, 955-973, 2018.

Wagner, R., Benz, S., Möhler, O., Saathoff, H., and Schurath, U.: Probing ice clouds by broadband mid-infrared extinction spectroscopy: case studies from ice nucleation experiments in the AIDA aerosol and cloud chamber, *Atmos. Chem. Phys.*, 6,
675 4775-4800, 2006a.

Wagner, R., Bunz, H., Linke, C., Möhler, O., Naumann, K. H., Saathoff, H., Schnaiter, M., and Schurath, U.: Chamber Simulations of Cloud Chemistry: The AIDA Chamber, Proceedings of the NATO Advances Research Workshop on Environmental Simulation Chambers: Application to Atmospheric Chemical Processes, held in Zakopane, Poland, from 1 to
680 4 October 2004, 2006b.

Wagner, R., Linke, C., Naumann, K. H., Schnaiter, M., Vragel, M., Gangl, M., and Horvath, H.: A review of optical measurements at the aerosol and cloud chamber AIDA, *J. Quant. Spectrosc. Radiat. Transfer*, 110, 930-949, 2009.

685 Wagner, R., Höhler, K., Huang, W., Kiselev, A., Möhler, O., Mohr, C., Pajunoja, A., Saathoff, H., Schiebel, T., Shen, X. L., and Virtanen, A.: Heterogeneous ice nucleation of α -pinene SOA particles before and after ice cloud processing, *J. Geophys. Res. (Atmos.)*, 122, 4924-4943, 2017.

Wagner, R., Bertozzi, B., Höpfner, M., Höhler, K., Möhler, O., Saathoff, H., and Leisner, T.: Solid ammonium nitrate aerosols
690 as efficient ice nucleating particles at cirrus temperatures, *J. Geophys. Res. (Atmos.)*, 125, e2019JD032248,
10.1029/2019JD032248, 2020.

Wang, M., Kong, W., Marten, R., He, X.-C., Chen, D., Pfeifer, J., Heitto, A., Kontkanen, J., Dada, L., Kürten, A., Yli-Juuti,
T., Manninen, H. E., Amanatidis, S., Amorim, A., Baalbaki, R., Baccarini, A., Bell, D. M., Bertozzi, B., Bräkling, S., Brilke,
695 S., Murillo, L. C., Chiu, R., Chu, B., De Menezes, L.-P., Duplissy, J., Finkenzeller, H., Carracedo, L. G., Granzin, M., Guida,
R., Hansel, A., Hofbauer, V., Krechmer, J., Lehtipalo, K., Lamkaddam, H., Lampimäki, M., Lee, C. P., Makhmutov, V., Marie,
G., Mathot, S., Mauldin, R. L., Mentler, B., Müller, T., Onnela, A., Partoll, E., Petäjä, T., Philippov, M., Pospisilova, V.,
Ranjithkumar, A., Rissanen, M., Rörup, B., Scholz, W., Shen, J., Simon, M., Sipilä, M., Steiner, G., Stolzenburg, D., Tham,
Y. J., Tomé, A., Wagner, A. C., Wang, D. S., Wang, Y., Weber, S. K., Winkler, P. M., Wlasits, P. J., Wu, Y., Xiao, M., Ye,
700 Q., Zauner-Wieczorek, M., Zhou, X., Volkamer, R., Riipinen, I., Dommen, J., Curtius, J., Baltensperger, U., Kulmala, M.,
Worsnop, D. R., Kirkby, J., Seinfeld, J. H., El-Haddad, I., Flagan, R. C., and Donahue, N. M.: Rapid growth of new
atmospheric particles by nitric acid and ammonia condensation, *Nature*, 581, 184-189, 10.1038/s41586-020-2270-4, 2020.

Yu, P. F., Rosenlof, K. H., Liu, S., Telg, H., Thornberry, T. D., Rollins, A. W., Portmann, R. W., Bai, Z. X., Ray, E. A., Duan,
705 Y. J., Pan, L. L., Toon, O. B., Bian, J. C., and Gao, R. S.: Efficient transport of tropospheric aerosol into the stratosphere via
the Asian summer monsoon anticyclone, *Proc. Natl. Acad. Sci. U.S.A.*, 114, 6972-6977, 10.1073/pnas.1701170114, 2017.

Zasetsky, A. Y., Khalizov, A. F., Earle, M. E., and Sloan, J. J.: Frequency Dependent Complex Refractive Indices of
Supercooled Liquid Water and Ice Determined from Aerosol Extinction Spectra, *J. Phys. Chem. A*, 109, 2760-2764, 2005.

710

Zasetsky, A. Y., Earle, M. E., Cosic, B., Schiwon, R., Grishin, I. A., McPhail, R., Pancescu, R. G., Najera, J., Khalizov, A. F.,
Cook, K. B., and Sloan, J. J.: Retrieval of aerosol physical and chemical properties from mid-infrared extinction spectra, *J.*
Quant. Spectrosc. Radiat. Transfer, 107, 294-305, 2007.







Figure 1: Scheme of the instrumentation of the AIDA aerosol and cloud chamber facility employed in the present study. The abbreviations are explained in Sect. 2.1.




725 **Figure 2:** (a) Infrared extinction spectrum of crystalline AN particles recorded at 223 K. The notation of the AN vibrational modes is according to Fernandes et al. (1979). (b) Concomitant number size distribution of the almost pure AN particles crystallized from solution droplets with 99.4 mol% AN and 0.6 mol% AS. The data are shown as a function of the equal-volume sphere diameter, d_p , as obtained by converting the mobility and aerodynamic size spectra from the SMPS and APS measurements with $\rho(\text{AN}) = 1.72 \text{ g cm}^{-3}$ and $\chi = 1.1$. (c) Exemplary electron microscope images of filter-sampled, continuously cooled AN crystals from a crystallization experiment with 97.1 mol% AN and 2.9 mol% AS.




730 **Figure 3: Flowchart of the retrieval scheme to deduce the optical constants of the crystalline AN particles. All abbreviations, symbols, and steps in the procedure (numbered green circles) are explained in Sect. 2.2.**




735 **Figure 4:** Real (a) and imaginary (b) parts of the complex refractive for solid AN at 223 K derived from this work (black lines) in comparison with room-temperature data from Jarzembki et al. (2003) (red lines). The insert (c) shows an expanded view of the k spectrum in the regime of the $v_2(\text{NO}_3^-)$ mode.



740 **Figure 5:** Sensitivity of the retrieved $k(\tilde{\nu})$ spectrum on the parameters $n(\tilde{\nu}_x)$, χ , and ϕ in different wavenumber regimes. (a) – (c) Variation of $n(\tilde{\nu}_x)$ with $\chi = 1.1$ and $\phi = 1$. (d) – (f) Variation of χ with $n(\tilde{\nu}_x) = 1.56$ and $\phi = 1$. (g) and (h) Variation of ϕ with $n(\tilde{\nu}_x) = 1.56$ and $\chi = 1.1$.




745 **Figure 6:** (a) AN mass concentrations retrieved from GLORIA measurements during the StratoClim research flight on 31 July 2017 using the $n(\tilde{\nu}_x) = 1.56$, $\chi = 1.1$, and $\phi = 1$ refractive index data set (see Fig. 3b in Höpfner et al. (2019) as a comparison). Red line: aircraft altitude. (b) Difference of mean GLORIA spectra between 04:15 – 04:21 and 04:35 – 04:43 UTC at 16.5 – 16.75 km altitude (blue) and best spectral fit with the AN optical constants (orange) in the range of the $\nu_2(\text{NO}_3^-)$ mode.




750

Figure A1: Comparison between the measured (black lines) and the computed (red lines) infrared extinction spectrum of crystalline AN particles in five different wavenumber regimes after convergence of the minimization algorithm with $n(\tilde{\nu}_s) = 1.56$, $\chi = 1.1$, and $\phi = 1$. The notation of the AN vibrational modes is according to Fernandes et al. (1979).



755

Figure B1: Comparison between the measured infrared extinction spectrum of crystalline AN particles (a) and the retrieved optical constants (b) in the $1200 - 1600 \text{ cm}^{-1}$ wavenumber range. The retrieval was performed with the parameters $n(\tilde{\nu}_x) = 1.56$, $\chi = 1.1$, and $\phi = 1$. Vertical dashed lines indicate peak positions in the particle extinction spectrum (blue) and the $k(\tilde{\nu})$ spectrum (red).



760

Figure B2: Comparison between the measured infrared extinction spectrum of crystalline AN particles (a) and the retrieved optical constants (b) in the 800 – 860 cm^{-1} wavenumber range. The retrieval was performed with the parameters $n(\tilde{\nu}_*) = 1.56$, $\chi = 1.1$, and $\phi = 1$. Vertical dashed lines indicate the peak position in the particle extinction spectrum (blue) and the $k(\tilde{\nu})$ spectrum (red).

765