

Author Response to Luis Guanter (LG)

First, we want to thank Luis Guanter for the careful and concise review and the helpful comments. In the following, we will first repeat the review comment (LG-X), and then give our response.

LG-1: Comparison to Matched Filter (MF) and IMAP-DOAS methods: according to the authors, the motivation to propose a WFM-DOAS retrieval for AVIRIS-NG is to “fill a gap” between MF and IMAP-DOAS approaches. However, nothing is said about the retrieval performance of the proposed WFM-DOAS with respect to the other two methods. I believe that the study would greatly benefit from a quantitative comparison of the three retrievals. The MF XCH₄ data for the same ABOVE data set seem to be already available (p4, L20), and looking at the author list I understand that it wouldn’t be difficult to also have IMAP-DOAS retrievals for at least a data subset containing one of the identified plumes.

We agree that this comparison between all three methods would greatly enhance the paper. Consequently, we included one coal mine ventilation shaft plume observed in the Four Corners measurement campaign, for which also already IMAP-DOAS and MF results existed, and retrieved this plume with the WFM-DOAS method presented in this paper for comparison of all three methods. Additionally, we compared the MF and WFM-DOAS results for the plumes which were analyzed quantitatively in the main text in the new Section 4.3.

LG-2: Selection of the AVIRIS-NG data set used for this: the authors don’t provide any explanation of why they select this particular ABOVE AVIRIS-NG data set for the study, but I would say that there could be better AVIRIS / AVIRIS-NG data sets for a study focused on the presentation of a new retrieval method. For example, the data sets from the different campaigns that JPL has run over the last years, including e.g. the one over Four Corners (Frankenberg et al., 2016) and California (Duren et al., 2019), for which the L1B data are surely available, could offer better observation conditions, properly documented plumes and fluxes, and a wider range of emission intensities and types. So I would recommend the authors to extend the study to other AVIRIS-NG showing other acquisition conditions, and in particular larger emissions. On the other hand, if the authors were interested in the particularly challenging conditions of the Canada sites, this should be discussed in the text.

We chose the ABoVE data as it provided a wide range of surface types and shapes and observation geometries to test the retrieval under different, partly challenging circumstances. We added the explanation for the selection of this data set to Section 2.1. To include a source with higher enhancements of CH₄, we added a flight line over a coal mine ventilation shaft observed during the Four Corners measurement campaign.

LG-3: Abstract: it might benefit from shortening and removal of line breaks

We agree and shortened the abstract.

LG-4: p4 L25: “Copernicus”, nested brackets

Done

LG-5: Fig. 1 caption: “can not be fully resolved”

Done

LG-6: p7, L28: plus surface reflectance shows in general less variability at 1600 nm than at 2300 nm

The mean reflectance for different surface types varies similarly for 1600 nm and 2300 nm. Also some interfering absorbers have spectral features as large around 1650 nm as around 2300 nm (See for example Foote et al., 2020, Fig. 2). So we do not think that this is a major advantage of the 1600 nm spectral region.

LG-7: Section 3.3., sensitivity analysis: it could also be relevant to include uncertainties in the spectral calibration (position and shape of the spectral response function), if this is not optimized in the retrieval

The spectral calibration is optimized during the L0 → L1b processing (see Chapman et al., 2019). Nevertheless, we fitted additional wavelength shift and squeeze parameters in the retrieval. We added the explanation to Sect. 3.3.

LG-8: p15, L13 “determined by the following”

Done

LG-9: p17, L9 – the cross sectional method hasn’t been mentioned until now

We changed the sentence in the following way to address this criticism: “However, many plumes were faint or located near infrastructure, making unambiguous detection difficult or the plumes were very short. Therefore, we show plumes which extend over longer ranges and have unambiguous morphology in Fig. 8 and Fig. 9. The additional CH₄ plumes and enhancements can be found in the appendix in Sect. B.”

LG-10: p17, L17: “A wind speed higher by a factor 1.6 means...”? why is this? can you please explain this further?

Assuming a constant flux, if the wind speed is higher by a constant factor, the total column enhancements along the cross section have to be lower by that factor. However, a reduction in total column enhancements is less visible above the background and dilutes faster. This is only true if we assume a constant emitting source, which indeed might not be the case here. Due to the wind speed uncertainties however we can not dismiss this possibility.

LG-11: Section 4.2 and 4.3 could be merged. In general, a manuscript structure with separate Methods and Results sections would be better.

We decided against such a structure, as the selection and description of some methods relied on the results from previous parts. However, merging Section 4.2 and 4.3 is indeed more logical in the current structure.

LG-12: Figure captions: perhaps just a matter of taste, but I think they could be shorter and avoid the explanations already included in the main text (e.g. Fig. 8 and 9).

We agree and shortened excessive interpretation in the figures.

LG-13: Fig. 9, Plume P4: at least from visual inspection, it is not obvious to me that that is a real plume. It could also be the effect of background reflectance variations. Would it make sense to show a map of at-sensor radiance at 2200 nm and its slope to discard this possibility?

We produced this plot for comparison between the radiance at 2140 nm and the plume. The structures visible in the gray scale image of radiance at 2140 nm do not seem to match the faint plume structure. Consequently, we believe this is a real CH₄ plume.




Figure 0.1: Plume P4 (b) and Radiance at 2140 nm (a) to check for the possibility of surface effects being mistaken for a plume structure.

LG-14: Refs. At least Cusworth et al. (<https://doi.org/10.5194/amt-12-5655-2019>) and Foote et al. (<https://doi.org/10.1109/TGRS.2020.2976888>) should be added

We agree and added the citations.

Author Response to Reviewer #2 (R2)

We would like to thank Reviewer #2 for the helpful review. In the following, we will first repeat the review comment (R2-X), and then give our response.

R2-1: Please define the WFM-DOAS acronym early on in the text.

We added the definition of WFM-DOAS to the abstract and at its first mention in the introduction.

R2-2: The abstract is very long. Much of the content there seems better suited for the introduction. I would recommend shortening the abstract to one or two paragraphs so the reader can quickly get a sense of the paper's main findings/results.

We agree and shortened the abstract.

R2-3: P4, L11: Hamlin et al. (2011) cite 10 nm spectral sampling for AVIRIS and 5 nm spectral resolution for the next-generation instrument. I can't find a statement of 5 nm spectral sampling in their paper. If I'm not mistaken, can you add a reference for the 5 nm spectral sampling number?

We added Chapman et al. (2019) as a source for this.

R2-4: P5, L2: What kind of interpolation is used?

We used linear interpolation in time and space between the 4 nearest locations and the 2 nearest time steps. We added this explanation to Sect. 2.2.

R2-5: P5, L3: Any motivation for using the 50-m average? Is it just a convenient choice or is there an argument for this being the most relevant wind speed?

We originally used the 50 m average, as the sources P1 and P2 were estimated to release the plume at ~ 55 m height, and the other sources were assumed to be closer to the ground. As we are aware that the uncertainty in wind speed is one key error source for the emission estimate, we made an attempt to at least partly validate these nearsurface winds. For that we compared the 50 m height average calculated from ERA5 over weather station data near the plume locations (see new Sec. A). We found that the wind speed from ERA5 seemed to significantly underestimate the wind speed, with a mean difference of $+2.5 \text{ m s}^{-1}$ and single differences up to $+3.4 \text{ m s}^{-1}$. Additionally, for the source mapped in P5 the estimated emissions are $\gg 1000 \text{ kg hr}^{-1}$ according to other inversions and in situ data (e.g. Jongaramrungruang et al., 2019; Frankenberg et al., 2016). Also in this case the wind speeds from the ground station results in a considerable better match with the published flux estimates. Consequently, we changed our wind speed estimations consistently from the ERA5 50m height average to hourly wind data from ground stations

nearby and rewrote Sect. 2.2 and 4.3 accordingly, as well as the discussion and conclusions (Sect. 5 and 6). Additionally, we added the new Sect. A to the appendix for the comparison between ERA5 and weather station wind speeds.

R2-6: P5, L4: I couldn't find a description in the text of how the wind direction was assessed from the plume shape. Can you elaborate?

We visually inspected the plume direction for the best fitting line from the source along the plume. The cross sections were then calculated perpendicular to that. We complemented the according sentences in the manuscript.

R2-7: Table 1: Can you explain why the PSF-CH4-proxy uncertainty is (slightly) higher than the PSF-CH4 uncertainty? Error stacking from the uncorrelated errors in PSF-CH4 and PSF-CO2?

Yes, that is what we assumed, too. However, the standard deviation of the fraction is not exactly the root square sum of the single standard deviations and was calculated from the scene separately.

R2-8: I'm familiar with the CO2 proxy method being applied in the 1650 nm window where there is also strong CO2 absorption, but not in the 2300 nm window. Can you comment on the effectiveness of the method at 2300 nm compared to 1650 nm?

It works quite well using the strong absorption of CO₂ near 2000 nm to retrieve CO₂ scaling factors, as can be seen for example in Fig. 7. However, we did not compare the impact of the proxy on the two windows further than what is shown in Table 1.

R2-9: Can you explain more clearly the pros/cons of WFM-DOAS compared to IMAP-DOAS and Matched Filter? How much faster is WFM than IMAP? When/why is it better than Matched Filter? You say in the abstract that WFM "fills a gap" between the other two retrieval algorithms, and mention some of the differences between them in section 1 and section 5, but I didn't get a clear picture of the motivation behind WFM, except that it's computationally cheaper than IMAP. What are the trade-offs? A concise explanation in the introduction of when/why we should use WFM instead of IMAP or MF would be very helpful.

We added a quantitative comparison between IMAP-DOAS, MF and WFM-DOAS (see also answer to LG-1). At least from our study we conclude that WFM-DOAS overall produces less background noise. However, in its current form it is not fast enough for near real time analysis of AVIRIS-NG data, which the MF is. The IMAP-DOAS retrieval is more suited for observations deviating larger from the a priori, as through the iterative approach the non-linear behavior especially of the radiance dependent on gas concentration can be captured better.

R2-10: P18, L1-10: The double plume structure is intriguing. Could you briefly elaborate on the height of the source, solar zenith angle, and distance between the two plume images in this case? Also, how did you determine wind direction and do the cross-sectional method for this plume?

We do not know the stack height from direct height measurements or similar. However, from the solar zenith angle during overflight of $\sim 42^\circ$ and the length of the shadow of in the RGB image of ~ 50 m, we used trigonometry to calculate the stack height to be ~ 55 m. The apparent plume center distance is approximately 50 m. Accounting additionally for the viewing angle (which deviates slightly from nadir), we calculate a plume height of ~ 56 m shortly after the beginning of the plumes. Therefore we are confident that this is a real feature. We added the numbers and explained their origin in the manuscript.

For the wind direction, we applied the method described in answer R2-6 to the plume structure expanding from the top of the tower, not the top of the shadow. For the cross sectional flux method, one has to have in mind that for the double plume the light passes the plume only once, whereas it is assumed to pass the plume twice in the retrieval and conversion to columns. Summing over the total enhancements of a cross section through both plume structures yields the total enhancement of the plume.

R2-11: P18, L28: How do you define inside/outside the plume? I think previous AVIRIS studies used a threshold, did you do that too?

We did not use a threshold, but defined spatial boundaries which were definitely aside the plume. This was done to normalize to the background prior to the inversion (similar to e.g. Krautwurst et al., 2017, Fig. 6).

R2-12: P21, L1: Is the wind speed calculation referred to here the 50-m average from ERA5, or something else?

Yes, this was the 50-m average from ERA5. However, as described in our answer to R2-5, we changed to using weather station wind data, as the previously used ERA5 data likely strongly underestimated wind speeds near the surface.

R2-13 Section 4.3: Are these 1-sigma or 2-sigma errors? Or confidence intervals?

These are 1-sigma standard deviations. We added this to the text.

R2-14 P21, L26: I don't understand why the higher wind would cause you to underestimate the emission rate. Shouldn't the cross-sectional method be able to deal with this? Each plume transect gives you an emission estimate, and the higher wind speed will result in higher estimates. I can see how the precision would be lower for a weaker plume, since you might have fewer transects to analyze. But a negative bias would imply that you are missing part of each transect. In that case, couldn't you extend the

transects further from the plume axis? This goes back to my question about defining inside/outside the plume.

Indeed, it is partly compensated by the higher wind speed. On the other hand, a higher wind speed leads to less molecules in each air parcel. This might drop larger parts of the plume significantly below the background noise or the retrieval detection limit and therefore would not lead to a detectable increase above the normalized background anymore.

Bibliography

- Chapman, J. W., Thompson, D. R., Helmlinger, M. C., Bue, B. D., Green, R. O., Eastwood, M. L., Geier, S., Olson-Duvall, W., and Lundeen, S. R.: Spectral and Radiometric Calibration of the Next Generation Airborne Visible Infrared Spectrometer (AVIRIS-NG), *Remote Sensing*, 11, 2129, <https://doi.org/10.3390/rs11182129>, 2019.
- Foote, M. D., Dennison, P. E., Thorpe, A. K., Thompson, D. R., Jongaramrungruang, S., Frankenberg, C., and Joshi, S. C.: Fast and Accurate Retrieval of Methane Concentration From Imaging Spectrometer Data Using Sparsity Prior, *IEEE Transactions on Geoscience and Remote Sensing*, 58, 6480–6492, <https://doi.org/10.1109/tgrs.2020.2976888>, 2020.
- Frankenberg, C., Thorpe, A. K., Thompson, D. R., Hulley, G., Kort, E. A., Vance, N., Borchardt, J., Krings, T., Gerilowski, K., Sweeney, C., Conley, S., Bue, B. D., Aubrey, A. D., Hook, S., and Green, R. O.: Airborne methane remote measurements reveal heavy-tail flux distribution in Four Corners region, *Proceedings of the National Academy of Sciences*, <https://doi.org/10.1073/pnas.1605617113>, URL <http://www.pnas.org/content/early/2016/08/10/1605617113.abstract>, 2016.
- Jongaramrungruang, S., Frankenberg, C., Matheou, G., Thorpe, A. K., Thompson, D. R., Kuai, L., and Duren, R. M.: Towards accurate methane point-source quantification from high-resolution 2-D plume imagery, *Atmospheric Measurement Techniques*, 12, 6667–6681, <https://doi.org/10.5194/amt-12-6667-2019>, 2019.
- Krautwurst, S., Gerilowski, K., Jonsson, H. H., Thompson, D. R., Kolyer, R. W., Iraci, L. T., Thorpe, A. K., Horstjann, M., Eastwood, M., Leifer, I., Vigil, S. A., Krings, T., Borchardt, J., Buchwitz, M., Fladeland, M. M., Burrows, J. P., and Bovensmann, H.: Methane emissions from a Californian landfill, determined from airborne remote sensing and in situ measurements, *Atmospheric Measurement Techniques*, 10, 3429–3452, <https://doi.org/10.5194/amt-10-3429-2017>, 2017.

Detection and Quantification of CH₄ Plumes using the WFM-DOAS retrieval on AVIRIS-NG hyperspectral data

Jakob Borchardt¹, Konstantin Gerilowski¹, Sven Krautwurst¹, Heinrich Bovensmann¹, Andrew K. Thorpe², David R. Thompson², Christian Frankenberg^{3,2}, Charles E. Miller², Riley M. Duren^{4,2}, and John Philip Burrows¹

¹Institute of Environmental Physics (IUP), University of Bremen, Bremen, Germany

²Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, United States of America

³California Institute of Technology, Division of Geological and Planetary Sciences, Pasadena, CA, United States of America

⁴Institutes for Resilience, University of Arizona, Tuscon, AZ, United States of America

Correspondence: Jakob Borchardt (jakob.borchardt@iup.physik.uni-bremen.de)

Abstract. Methane is the second most important anthropogenic greenhouse gas in the Earth's atmosphere. ~~Reducing methane emissions is consequently an important element in limiting the global temperature increase below 2 °C compared to preindustrial times. Therefore~~ To effectively reduce these emissions, a good knowledge of source ~~strengths and source locations~~ locations and strengths is required. ~~Anthropogenic methane emissions often originate from point sources or small areal sources, such as fugitive emissions at oil and gas production sites or landfills.~~ Airborne remote sensing instruments such as the Airborne Visible InfraRed Imaging Spectrometer - Next Generation (AVIRIS-NG) with meter scale imaging capabilities are able to yield information about the locations and magnitudes of methane sources, ~~especially in areas with many potential emission sources.~~

~~To extract methane column enhancement information from spectra recorded with the.~~ In this study we successfully applied
10 the Weighting Function Modified Differential Optical Absorption Spectroscopy (WFM-DOAS) algorithm to AVIRIS-NG ~~instrument, different retrieval algorithms have been used, e. g. the data measured in Canada and the Four Corners region.~~
The WFM-DOAS retrieval is conceptually located between the statistical matched filter (MF) ~~or the and the optimal estimation~~
based Iterative Maximum A Posteriori DOAS (IMAP-DOAS) retrieval ~~. The WFM-DOAS algorithm, successfully applied~~
15 ~~algorithm, which already were applied successfully~~ to AVIRIS-NG data in this study, fills a gap between those retrieval
~~approaches by being a fast, non-iterative algorithm.~~ The WFM-DOAS algorithm is based on a first order Taylor series approx-
imation of the Lambert-Beer law ~~, which calculates the change in gas concentrations from deviations from one background~~
using only one precalculated radiative transfer calculation ~~using precalculated weighting functions specific to the state of the~~
atmosphere during the overflight. This allows the per scene. This yields fast quantitative processing of large data sets. ~~Although~~
20 ~~developed for high spectral resolution measurements from satellite instruments such as SCIAMACHY, TROPOMI and the~~
MAMAP airborne sensor, the algorithm can be applied well to lower spectral resolution ~~We detected several methane plumes~~
in the AVIRIS-NG ~~measurements, images recorded during the ABoVE Airborne Campaign and successfully retrieved a coal~~
mine ventilation shaft plume observed during the Four Corners measurement campaign. Comparison between IMAP-DOAS,
MF and WFM-DOAS showed good agreement for the coal mine ventilation shaft plume. Additional comparison between MF

and WFM-DOAS for a subset of plumes showed good agreement for one plume and some differences for the other. For five plumes the emissions were estimated using a simple cross sectional flux method. The retrieved fluxes originated from well pads, cold vents and a coal mine ventilation shaft and ranged between $(155 \pm 71) \text{ kg (CH}_4\text{) h}^{-1}$ and $(1220 \pm 450) \text{ kg (CH}_4\text{) h}^{-1}$. The wind velocity was a significant source of uncertainty for all plumes, followed by the single pixel retrieval noise and the uncertainty due to atmospheric variability. The ~~data-set examined here was recorded in Canada over different gas and coal extraction sites as part of the larger Arctic Boreal Vulnerability Experiment (ABoVE) Airborne Campaign in 2017.~~

The noise of the retrieved CH₄ imagery over bright surfaces ($> 1 \mu\text{W cm}^{-2} \text{ nm}^{-1} \text{ sr}^{-1}$ at 2140 nm) was typically $\pm 2.3\%$ of the background total column of CH₄ when fitting strong absorption lines around 2300 nm, but could reach over $\pm 5\%$ for darker surfaces ($< 0.3 \mu\text{W cm}^{-2} \text{ nm}^{-1} \text{ sr}^{-1}$ at 2140 nm). Additionally, a worst case large scale bias due to the assumptions made in the WFM-DOAS retrieval was estimated to be $\pm 5.4\%$. Radiance and fit quality filters were implemented to exclude the most uncertain results from further analysis, mostly due to either dark surfaces or surfaces, where the surface spectral reflection structures are similar to CH₄ absorption features at the spectral resolution of the AVIRIS-NG instrument.

~~We detected several methane plumes in the AVIRIS-NG images recorded during the ABoVE Airborne Campaign. For four of those plumes, the emissions were estimated using a simple cross sectional flux method. The retrieved fluxes originated from well pads and cold vents and ranged between $(89 \pm 46) \text{ kg (CH}_4\text{) h}^{-1}$ and $(141 \pm 87) \text{ kg (CH}_4\text{) h}^{-1}$. The wind uncertainty was a significant source of uncertainty for all plumes, followed by the single pixel retrieval noise and the uncertainty due to atmospheric variability. For one plume the wind was too low to estimate a trustworthy emission rate, although a plume was visible.~~

1 Introduction

Methane (CH₄) is an important greenhouse gas, with a global warming potential approximately 28 times larger than that of carbon dioxide (CO₂) on a timescale of 100 years (IPCC 2013, 2013). After a brief period of stable mixing ratios at the beginning of the 21st century, CH₄ concentrations have again begun rising (Dlugokencky et al., 2011; Dlugokencky, 2018); ~~though the~~. The origins of this stabilization and renewed increase are still debated (see for example Schaefer, 2019, and references therein). This uncertainty emphasizes the need to reduce anthropogenic CH₄ emissions to reach the goal of the Paris Agreement (Paris Agreement, 2015; Nisbet et al., 2019).

According to the Global Carbon Project (GCP, Saunio et al., 2016, 2019) between $\sim 50\%$ and $\sim 60\%$ of the global methane emissions are anthropogenic. Of those, roughly 55% result from agricultural practices and waste management, and nearly 35% from losses during fossil fuel extraction, delivery and use in energy production and transport, with a small contribution ($\sim 10\%$) coming from biomass and biofuel burning. Satellite instruments such as SCIAMACHY (spatial resolution $\sim 30 \times 60 \text{ km}^2$, Burrows et al., 1995; Bovensmann et al., 1999) and TROPOMI (spatial resolution $\sim 7 \times 7 \text{ km}^2$, Veefkind et al., 2012; Hu et al., 2016) have successfully been used to assess methane emissions from emission hot spots (Frankenberg et al., 2006; Schneising et al., 2009; Buchwitz et al., 2017; Hu et al., 2018; Schneising et al., 2019; Pandey et al., 2019). Additional efforts to make single CH₄ emitters visible from space by using measurements with high spatial resolution have been made, but so far only

strong single sources have been quantified (Thompson et al., 2016; Varon et al., 2019). Many anthropogenic CH₄ emissions occur over relatively large areas (e.g. rice paddies, animal herds, landfills), at previously unknown point sources (e.g. pipeline leaks, broken valves) or with highly varying emissions, making reliable detection and attribution of single sources from space challenging.

- 5 Airborne remote sensing campaigns can often gain a better knowledge of single emitter source strengths in emission hot spot regions due to their higher spatial resolution. In these campaigns, a defined area is sampled with stronger sensitivity to small localized CH₄ sources. For example, the Methane Airborne MAPper (MAMAP, Gerilowski et al., 2011), a non-imaging instrument with a nadir pointing field of view and a high spectral resolution of ~ 0.9 nm, successfully quantified emissions of known sources like coal mining shafts (Krings et al., 2013) or smaller areal sources like landfills (Krautwurst et al., 2017).
- 10 However, its viewing geometry requires a flight pattern orthogonal to the CH₄ plume for emission estimates and therefore limits the potential to pinpoint single unknown sources in a field of multiple potential sources. This problem is solved by using airborne imaging systems, which take multiple measurements across the flight track, thus creating an image of the area they pass over. However, to our knowledge, there is not yet an operational airborne imaging instrument specifically designed and optimized for the detection of CH₄~~available~~.
- 15 Nevertheless, data from multiple imaging instruments have been analyzed to map and/or quantify CH₄ emissions. For example, thermal imaging instruments such as SEBASS (Spatially-Enhanced Broadband Array Spectrograph System) could detect methane plumes as low as 0.4 kg h^{-1} of CH₄ flying 500 – 700 m above ground during a controlled release experiment (Scafutto et al., 2018). Also, the HyTES (Hyperspectral Thermal Emission Spectrometer) instrument has demonstrated detection of CH₄ leaks (Hulley et al., 2016). While these instruments have been able to detect very small sources at low flight altitude (500 m
- 20 above ground), performance may suffer at higher altitudes. For example, HyTES flying at 3 km had some difficulty consistently detecting a coal mine ventilation shaft plume with an estimated emission of ~~$\sim 1650 \text{ kg h}^{-1}$~~ $\sim 1200 \text{ kg h}^{-1}$ (Jongaramrungruang et al., 2019) due to the strongly varying sensitivity of the instrument to different atmospheric layers.

In the shortwave infrared (SWIR), the Airborne Visible-Infrared Imaging Spectrometer - Next Generation (AVIRIS-NG) was used for the detection and quantification of anthropogenic methane sources (Thompson et al., 2015; Frankenberg et al., 2016; Thorpe et al., 2016a, 2017; Duren et al., 2019; Cusworth et al., 2020; Thorpe et al., 2020). As the instrument was

25 not designed for the detection of atmospheric absorbers, it has a spectral resolution much coarser than SWIR instruments specifically designed to measure CO₂ and CH₄. However, AVIRIS-NG has a very high signal to noise ratio (SNR) and meter scale spatial resolution. The latter depends on flight altitude and flight speed, with typical values for the ground sampling distances for large scale methane surveys of $3 \times 3 \text{ m}^2$ to $5 \times 5 \text{ m}^2$. Successful algorithms for the retrieval of methane

30 comprised either a matched filter approach (Thompson et al., 2015), which uses a hypothesis test between presence and absence of additional CH₄ to infer CH₄ increases, or an adaption of the IMAP-DOAS retrieval (~~Frankenberg et al., 2005~~) (Frankenberg et al., 2005; Thorpe et al., 2013, 2017; Cusworth et al., 2019) to AVIRIS-NG airborne data (~~Thorpe et al., 2013, 2017~~), which is an iterative optimal estimation based algorithm. However, the latter is computationally very expensive, which makes it less suited to analyze large data sets acquired during longer measurement campaigns (Thorpe et al., 2017). Consequently, it

has only been applied to regions of special interest in the data. Recently, a new variant of the matched filter using a sparsity prior approach was successfully applied to AVIRIS-NG data (Foote et al., 2020).

In this study, we test and apply an adaption of the Weighting Function Modified Differential Optical Absorption Spectroscopy (WFM-DOAS) algorithm, used previously for the higher spectral resolution MAMAP measurements (spectral resolution ~ 0.9 nm Krings et al., 2011), to hyperspectral AVIRIS-NG data (spectral resolution ~ 6 nm). The data set was acquired during the Arctic Boreal Vulnerability Experiment Airborne Campaign (ABOVE, Miller et al., 2019b) in Canada and Alaska, which included overflights of multiple coal, oil and gas production sites. The WFM-DOAS approach uses assumptions on the background state of the atmosphere at the time and location of the overflight, including scattering. It performs a linear fit of atmospheric parameters deviating from this background state, making it a fast quantitative method compared to iterative retrievals. We identified multiple plumes in the retrieval results, and for ~~four of them~~, five of them the emissions were estimated by application of a cross sectional flux method.

This publication is organized as follows: Following this introduction, Sect. 2 gives an overview of the instrument and data sets. Section 2.1 describes the AVIRIS-NG instrument and radiance data, and Sect. 2.2 introduces the ~~ERA-5~~ used meteorological data briefly. In Sect. 3, we present the retrieval algorithm for CH₄ and the subsequent filtering. First, we describe the WFM-DOAS method used to infer methane enhancement maps from the spectra in Sect. 3.1. Section 3.2 justifies the fitting windows we use in the retrieval. Sect. 3.3 evaluates the sensitivity of the retrieval to assumptions in the forward model and in Sect. 3.4 we implement a filtering to remove certain error cases. We present experimental results in Sect. 4. First, the detection of plumes is described in Sect. 4.1. Second, ~~See~~ in Sect. 4.2 we compare the WFM-DOAS results with IMAP-DOAS and MF retrieval results. Finally, Sect. 4.3 illustrates a flux inversion using the cross sectional flux method ~~, and Sect. 4.4 finally and~~ shows the results and uncertainties of the emission estimate for ~~four~~ five plumes. The results are discussed in Sect. 5. Sect. 6 summarizes the findings of this study.

2 Instrument and data sets

2.1 The AVIRIS-NG instrument and measurements

AVIRIS-NG is a hyperspectral imaging spectrometer with a spectral sampling of ~ 5 nm and a spectral resolution of ~ 5 – 6 nm, depending on the wavelength (~~Hamlin et al., 2011~~) (Hamlin et al., 2011; Chapman et al., 2019). As a nadir looking instrument, it measures solar radiation reflected from the ground in the wavelength range from 380 to 2450 nm with a high signal-to-noise ratio of up to 800 at 2200 nm (Thorpe et al., 2016a). The instrument contains 600 spatial pixels, each having a 1 mrad field of view. This results in individual samples with 5 m spatial resolution and a 3 km swath from a typical flight altitude of 5 km above ground level. This allows it to scan large areas in short periods of time. The level-1 data distributed by the operations team contains orthorectified (and gridded) absolute radiances (Chapman et al., 2019), with additional data containing observation parameters such as flight altitude, both solar and instrument zenith and azimuth angles and surface elevation among others (see Miller et al., 2019a, for data description).

For this study, we analyzed a subset of the measurements collected during the ABoVE Airborne Campaign (Miller et al., 2019b) in 2017. The ABoVE campaign aimed to better understand the impacts of environmental changes in Alaska and western Canada. During the airborne campaign, several flight lines of the AVIRIS-NG instrument covered fossil fuel infrastructure in Canada, which contained multiple potential sources for CH₄ emission plumes.

- 5 The data analyzed in this paper had been preselected to cover a wide range of surface types (e.g. forest, mountainous regions, sand, grass) and at-sensor radiance levels, as well as different flight altitudes. Additionally, the tracks contained different emission sources detected using the matched filter (MF) algorithm (Thompson et al., 2015) to test the retrieval algorithm against known ~~plumes~~ plume locations over different terrain. The preselection contained 13 flight lines on 5 different days in August 2017, covering different types of sources and surface types. Additionally, to include another strong source under different
10 observation conditions, we included a coal mine ventilation shaft plume observed during the Four Corners measurement campaign in 2015 (Frankenberg et al., 2016).

For the coal mine ventilation shaft plume, already existent MF and IMAP-DOAS retrieval results were used for comparison with the WFM-DOAS results retrieved in the course of this study. Additionally, a subset of the MF results for the flight lines under consideration were utilized for quantitative comparison with WFM-DOAS.

15 2.2 ~~ERA-5 meteorological~~ Meteorological data from ERA5 and weather stations

The WFM-DOAS retrieval and the flux inversion require information about various atmospheric parameters in addition to the observed radiances. The following meteorological parameters were extracted from ERA-5 reanalysis data (Copernicus Climate Change Service (C3S) (2017), 2017): hourly data of temperature, pressure and water vapor profiles, as well as height resolved wind speeds and wind components.

- 20 For a given flight line, the atmospheric parameters of the nearest 4 spatial grid points and the nearest two time steps of the ERA-5 data set were linearly interpolated to the time and location of the flight line. For the wind speed ~~and the wind direction,~~
~~the wind speed components in East and North directions were averaged over the lowest 50 m. We used the averaged wind speed for the inversion, and included the averaged wind direction in the plots of the detected plumes,~~ we used hourly mean
25 surface wind speed data obtained from nearby weather stations (see also Sect. A for a brief, localized comparison of these data with ERA5 wind speed data). The wind direction for the inversion was estimated from the plume structure itself. For that, we visually inspected the plume direction for the best fitting line from the source along the plume.

3 Adaption of WFM-DOAS algorithm to AVIRIS-NG measurements

- The WFM-DOAS algorithm was first developed for SCIAMACHY measurements (Buchwitz et al., 2000; Schneising et al., 2008), where the absorption bands around 1580 nm and 1660 nm were used for the retrieval of CO₂ and CH₄. Recently, it
30 has been modified and applied to TROPOMI measurements by Schneising et al. (2019) for the simultaneous retrieval of CH₄ and CO. As TROPOMI was build without spectral bands around 1600 nm, the retrieval used the wavelength ranges from 2311 nm – 2315.5 nm for CO and from 2320 nm – 2338 nm for CH₄.

Additionally, the WFM-DOAS algorithm has been adapted and used since 2007 to retrieve local CH₄ and CO₂ enhancements from MAMAP aircraft measurements in the wavelength range between 1590 nm and 1690 nm (see e.g. Krings et al., 2011, 2013; Krautwurst et al., 2017; Krings et al., 2018).

3.1 Retrieval of total column increases with WFM-DOAS

5 The WFM-DOAS algorithm minimizes the difference between a measured and a modeled spectrum by scaling weighting functions for the different trace gas profiles such as CH₄ and CO₂, shifting the temperature profile, and fitting a low order polynomial for broad band absorption (e.g. at the surface) or scattering (e.g. by air molecules and aerosols). The weighting functions represent a linear relationship between the change in observed radiance and a change in the atmospheric parameters. A detailed mathematical description of the WFM-DOAS algorithm modified for aircraft measurements and an analysis of this
10 method for MAMAP measurements can be found in Krings et al. (2011).

Each flight track covered a different scene or different day. We calculated a modeled spectrum for each scene using the SCI-ATRAN radiative transfer model (Rozanov et al., 2017). There we used the solar zenith angle, viewing angle of the instrument, solar and instrument azimuth angle, surface elevation and flight altitude from the AVIRIS-NG level 1 orthorectified data set (see data set description in Miller et al., 2019a). For each flight track, we calculated the mean value of each parameter and
15 used the results as input for the radiative transfer calculation.

For the radiative transfer calculation with SCIATRAN, the state of the atmosphere for the location of the flight track during the time of overflight was equally important. Temperature, pressure and water vapor profiles were extracted from ECMWF ERA5 meteorological data (see 2.2). The background total columns of carbon dioxide (CO_{2,back}) were calculated using the Simple Empirical CO₂ Model (SECM) by Reuter et al. (2012) in the version SECM2018, which contains a recently updated
20 parameter set (see also Reuter et al., 2020). The background total columns of methane (CH_{4,back}) were calculated with the approach used by Schneising et al. (2019), where a climatology averaged over the years 2003 - 2005 was enhanced by the total increase in methane based on globally averaged marine NOAA surface data (Dlugokencky, 2018). The CH₄ and CO₂ profiles used in SCIATRAN ~~where were~~ then obtained by scaling a US Standard Atmosphere (Oceanic et al., 1976) so that the total column mixing ratio calculated from those profiles matched the a priori estimated local total column mixing ratio.
25 HITRAN 2016 (Gordon et al., 2017) was used as spectral line parameter data base in SCIATRAN for trace gas absorption. The SCIATRAN model predicted the radiance at the sensor for the background case and the height dependent weighting functions for CH₄, CO₂ and H₂O.

The WFM-DOAS retrieval produced profile scaling factors (PSF), which scaled the weighting functions of CH₄, CO₂, H₂O and temperature in a linear fit. An example fit of AVIRIS-NG data with the resulting residual structures is shown in Fig. 1.
30 The light passed the air mass above the aircraft once on the downward path to the Earth, but transected the air mass below the aircraft on both downward and upward paths. Consequently, the retrieval was more sensitive to atmospheric changes below the aircraft than above. This was captured by the averaging kernel, which represented the sensitivity of the instrument to changes in a specific altitude layer. In our case, strong local enhancements in atmospheric methane were confined below the aircraft,




Figure 1. Example fit result of the WFM-DOAS retrieval for AVIRIS-NG data. The left block ((a) - (d)) shows the CO₂ fitting window (2040 nm – 2100 nm), the right block ((e) - (h)) shows the CH₄ fitting window (2235 nm – 2380 nm). In sub figures (a) and (h) the fit result is shown as a solid blue line, while the actual measured intensities are shown as gray diamonds. The difference between the gray diamonds and the line is the residuum, which is shown in sub figures (b) and (e). In the second row, the scaled weighting functions for CO₂ and H₂O in the CO₂ window ((c) and (d)) and for CH₄ and H₂O in the CH₄ window ((g) and (h)) are shown as solid lines. The gray diamonds show the fit result plus the residuum, to assess, if the residual structures are larger than the actual fitted structures. In the CH₄ fit window the residuum shows some structure, which might indicate some residual correlation between water vapor and methane signature in the real measurements, which can not ~~fully~~ be ~~fully~~ resolved by the retrieval.

so we multiplied the total column enhancements by the inverse of the averaging kernel for the airmass underneath the aircraft (k_{AK}) to determine the true enhancement of CH₄ caused by an emission source near the ground.

We did not retrieve the pressure profile or scattering properties. This, and also other effects like surface elevation changes could alter the light path and therefore the absorption strength, which would be detected as an enhancement. We used the proxy method to correct the retrieved column enhancement for those effects (see also Frankenberg et al., 2005; Schneising et al., 2009; Krings et al., 2011). Specifically, we divided the derived scaling factor of CH₄ by the scaling factor of another well mixed gas, which was assumed to be constant for the region of interest and time of overflight. For this work, we used CO₂ as a proxy because of its spectral proximity to the CH₄ absorption band, resulting in the proxy

$$PSF_{CH_4, proxy} = PSF_{CH_4} / PSF_{CO_2}. \quad (1)$$

Finally, we corrected the enhancements in a detected plume for large scale effects by normalizing over the local background around the plume ($PSF_{\text{proxy, bg}}$). The local column enhancement of CH_4 below the aircraft in a plume ($\text{CH}_{4,\text{enh}}$) was then

$$\text{CH}_{4,\text{enh}} = \left(\frac{PSF_{\text{CH}_4}}{PSF_{\text{CO}_2}} \right) / (PSF_{\text{proxy, bg}} - 1) \cdot \text{CH}_{4,\text{back}} \cdot k_{\text{AK}}. \quad (2)$$

A discussion of the biases introduced by the assumptions made for the WFM-DOAS retrieval are studied in the sensitivity analysis in Sect. 3.3.

3.2 Comparison of major fitting windows in the SWIR spectral range

For the MAMAP instrument the fitting window was $\sim 1630\text{ nm} - 1675\text{ nm}$ for CH_4 and $\sim 1592\text{ nm} - 1617\text{ nm}$ for CO_2 at 0.9 nm spectral resolution due to the sensor design (see Krings et al., 2011; Gerilowski et al., 2011). AVIRIS-NG additionally offered the possibility to fit the CH_4 and CO_2 absorption lines between 2000 nm and 2400 nm for the retrieval of CH_4 , although at a coarser spectral resolution of $5.5\text{ nm} - 6.0\text{ nm}$. For example, the IMAP-DOAS retrieval successfully retrieved CH_4 concentrations from AVIRIS-NG data using the spectral regions of $2215\text{ nm} - 2410\text{ nm}$ for CH_4 and $1904\text{ nm} - 2099\text{ nm}$ for CO_2 (Thorpe et al., 2017).

For a first assessment of those absorption bands, we convolved a simulated high resolution spectrum and the corresponding weighting functions for CH_4 , CO_2 and H_2O with the AVIRIS-NG instrument spectral response function. We used a Gaussian spectral response, where the FWHM were distributed as part of the data set. Fig. 2 shows the results of convolution and resampling to the AVIRIS-NG wavelength grid.

Both fitting windows had their advantages and disadvantages, especially for the lower spectral resolution of AVIRIS-NG. Around 2300 nm , the absorption features of CH_4 were about a factor of 2 stronger and had a more pronounced structure, which could lead to a better detection of methane changes. Around 1650 nm , the at sensor radiance was nearly twice as high for the same albedo, which could mean a higher signal-to-noise ratio. Additionally, there was less overlap with water vapor absorption features near 1600 nm .

We used a two step approach to find the best fitting window: First, we created a spatially averaged spectrum over a homogeneous surface elevation and surface type to reduce the instrument noise and systematic influences. Then, we optimized the edges of both fit windows for fitting the gas features in each window. As a measure of fit quality, the root mean square error (RMSE) between measurement and fit result was used. For CH_4 , the best fitting windows were $1625\text{ nm} - 1700\text{ nm}$ and $2235\text{ nm} - 2380\text{ nm}$, and for CO_2 $1550\text{ nm} - 1620\text{ nm}$ and $2040\text{ nm} - 2100\text{ nm}$ (see also Fig. 2). For simplicity, the fitting windows between 1550 nm and 1700 nm will be called “weak windows”, and the fitting windows between 2040 nm and 2380 nm will be called “strong windows” in the following parts, according to the depth of the absorption features.

To assess the measurement precision in each window, we selected a homogeneous, flat, bright area which contained no potential sources. We then applied the retrieval to the whole flight line containing this test case for each of the fitting windows and gases. These initial results showed detector column dependent stripes (see Sect. B in the appendix). To correct this effect, we normalized the PSF_{CH_4} , PSF_{CO_2} and $PSF_{\text{CH}_4,\text{proxy}}$ for each pixel by the median PSF of its corresponding detector column. We selected the median for resilience against outliers, which could otherwise have a large impact on the correction

Spectrum and weighting functions




Figure 2. The high resolution simulated spectra (in (a) and (b) green) are convolved with the slit function of AVIRIS-NG and sampled to the AVIRIS-NG wavelength grid (solid black line in upper row). The lower row ((c) and (d)) shows the weighting functions, i.e. the change of intensity due to a change in atmospheric concentration for CH_4 (blue), CO_2 (black) and H_2O (red). The shaded areas denote the fitting windows for CO_2 (gray) and CH_4 (light orange). ~~These have been chosen to be large enough to have enough data points inside the fitting window, without interfering with other gases (Sect. 3.2).~~

After destriping, we compared the standard deviation in the test case region of the weak and strong window retrieval results of PSF_{CH_4} , PSF_{CO_2} and $PSF_{\text{CH}_4, \text{proxy}}$ (Table 1. The retrieved PSF_{CH_4} and $PSF_{\text{CH}_4, \text{proxy}}$ were noisier in the weak window by a factor of 3.3 and 2.9, respectively. The retrieved PSF_{CO_2} was noisier by a factor of 1.5 in the weak window. Therefore, we only used the strong windows in later analyses.

Table 1. Comparison of the standard deviation of PSF_{CH_4} , PSF_{CO_2} and $PSF_{CH_4,proxy}$ in the two fitting windows around 1645 nm and 2300 nm for the AVIRIS-NG FWHM (≈ 6 nm). The standard deviation was calculated over a homogeneous and flat area with no visible plume and possible source inside. The statistical uncertainties for PSF_{CH_4} and PSF_{CO_2} are therefore uncorrelated.

	Standard deviation PSF 1645 nm fitting window	Standard deviation PSF 2300 nm fitting window
PSF_{CH_4}	$\pm 6.4\%$	$\pm 1.9\%$
PSF_{CO_2}	$\pm 1.9\%$	$\pm 1.3\%$
$PSF_{CH_4,proxy}$	$\pm 6.6\%$	$\pm 2.3\%$

3.3 Sensitivity analysis

In addition to the noise in the spectra, uncertainties and variability in the assumed constant atmospheric background parameters could lead to errors in the retrieval results. To assess the magnitude and influence of these deviations, we performed multiple sensitivity analyses. We used a common set of geometric and atmospheric parameters to model the background spectrum. We then perturbed these atmospheric parameters to create synthetic AVIRIS-NG observations at instrument spectral resolution. Next, we applied the WFM-DOAS algorithm to these simulated measurements, and assessed the systematic offset from the expected PSF value for PSF_{CH_4} , PSF_{CO_2} and $PSF_{CH_4,proxy}$. To assess the influence of linearization on the retrieval results, we did not include instrument noise in this analysis. The background simulation was based on the parameters extracted for one flight line observed with a nadir viewing angle. The CH_4 enhancements and a plume from this flight line are shown in Fig. 8.

For the sensitivity analysis, we perturbed the following set of parameters (Table 2): the aircraft altitude, the surface elevation, the instrument viewing angle (i.e. the instrument zenith angle) and the surface albedo as geometric parameters; and the total columns of CH_4 , CO_2 and H_2O , and the pressure and temperature profiles as atmospheric parameters. Additionally, we used selected spectral reflectance spectra of different surfaces instead of a spectrally uniform albedo and examined two additional aerosol scenarios. We did not analyze the sensitivity to the solar zenith and azimuth angles, since these angles were effectively constant over the timespan of a flight line. In addition, we did not analyze the instrument azimuth angle dependency since the flight tracks were nearly straight and the azimuth angle therefore effectively constant for a flight line. Finally, we did not evaluate sensitivity to spectral response function and wavelength calibration, since these were adjusted during conversion of the raw digital numbers to radiometric data cubes (Chapman et al., 2019).

The viewing angle variations were chosen to represent the range of the AVIRIS-NG viewing angles. The surface elevation and aircraft altitude deviation were chosen to represent plausible deviations over one flight line. Temperature deviations were chosen to be relatively large, as the temperature profile at the time of overflight for the specific ground scene might deviate quite a lot from the ERA-5 reanalysis due to the spatial and temporal resolution of the model output. The pressure scaling was chosen to represent a possible range of deviations, erring in favor of a conservatively high deviation for the observed scales. The albedo deviations covered the range which was expected around 2100 nm – 2300 nm (Chen et al., 2006).

The scaling of the CO₂ column and the H₂O column spanned the natural deviations of CO₂ and water vapor from the assumed background to establish an upper bound on errors from these effects. The range for the total column of CH₄ covered the range, which might be observed directly over or near a strong source. However, in most ground scenes containing a plume signal, the enhancement was well below 20 %, and for smaller plumes normally even near the source well below 10 %.

- 5 The reflectance spectra in the sensitivity analysis included surfaces present in the survey region or associated with oil and gas infrastructure. The spectral reflectances were based on the ECOSTRESS Spectral library (Meerdink et al., 2019; Baldrige et al., 2009) and on the US Geological Survey Spectral Library, Version 7 (Kokaly et al., 2017). They contained spectra from a surface covered by a typical plant of the Canadian savanna, sandstone, sand, and rangeland surfaces, and anthropogenic structures such as aluminium, steel and paving substances. The reflectance spectra are shown in Fig. 3.
- 10 The background aerosol scenario was assumed to be an OPAC (Hess et al., 1998) urban aerosol scenario (same as used in Krings et al., 2011), as we were interested in emissions from anthropogenic infrastructure. To determine the magnitude of influence of the aerosol scenario on the retrieval, we used additionally simulated measurements with an OPAC background and desert aerosol scenario.

After retrieving the profile scaling factors of CH₄ and CO₂ for each simulation, we calculated their deviation from the ground truth defined in the simulations. We also calculated the deviations for the CH₄ proxy method described in Sect. 3.1, and plotted the errors as a function of the perturbation of each parameter in Fig. 4. While the observed uncertainties for the single profile scaling factors were quite high (orange and blue curves), for example up to 10 % for an elevation change of approximately 400 m, they were highly reduced by the proxy method (green curve). The influence of the surface spectral reflectance is shown in Table 3 and discussed at the end of the section.

- 20 For aircraft altitude, temperature shifts, surface pressure scaling and viewing angle of the instrument, the maximum deviation from the ground truth for $PSF_{CH_4, proxy}$ remained well below 0.5 %. Also for most albedo values the maximum deviation remained below 0.5 %. For a very low albedo of 0.01, results of the single retrievals as well as the proxy method both degraded considerably. We examined low radiance ground scenes further in Sect. 3.4. For large perturbations of the surface elevation of 400 m, the proxy method reduced the error only to around ± 3.6 %. The different aerosol scenarios did not introduce major errors either. For the OPAC background scenario, the error was well below 0.1 %, and even for the OPAC desert aerosol load, only an error of -0.2 % was introduced.

For perturbations of CH₄ and H₂O, errors between the true and retrieved PSF grewedgrew, the larger the perturbations got. WFM-DOAS assumes a linear relationship between gas enhancement and radiance, but this assumption does not hold for large deviations from the background. This will also occur for CO₂ when choosing larger deviations from the background.

- 30 When we perturbed CO₂, the application of the proxy method increased the error in methane. When only CO₂ was varied, the methane column alone was retrieved correctly in the standard retrieval. Similarly, the retrieval correctly estimated the total column of CO₂. However, in the proxy method the retrieved PSF_{CH_4} was divided by the retrieved PSF_{CO_2} , so that a decrease in CO₂ led to an apparent increase in CH₄ and vice versa. This meant, that CO₂ emission sources could mask CH₄ emissions, if the relative single column enhancement of CO₂ is similar or greater than that of CH₄. As the retrieval noise is similar for both gases (Table 1), this would be visible as a CO₂ point source in the PSF_{CO_2} map.




Figure 3. In (a), the weighting function for CH₄ is displayed. In (b), the reflectance spectra covered in the sensitivity analysis are shown. Especially for the paving concrete one can see a similar broad band shape compared to the weighting function of CH₄, which is caused by calcium carbonate (limestone).

For the scaling of CH₄, the proxy method did not reduce the deviation of the retrieved from the true enhancements, as expected. However, the large deviations for strong enhancements (11% underestimation for 100% increase) would nevertheless mean a clearly detectable signal in the retrieved CH₄ maps. Smaller deviations ($\pm 20\%$) from the background profile would induce only small ($< 1\%$) underestimations. Consequently, for inversions of large emitters, the emission might be underestimated near the source, where the large enhancements are located. In cases with large concentrations near the source, emission estimates should only be performed further down the plume.

To estimate the total systematic uncertainty, we combined all uncertainties in Table 4 aside from the extreme CH₄ case in quadrature. This led to maximum systematic uncertainties of $\pm 8.0\%$ for PSF_{CH_4} , $\pm 11.3\%$ for PSF_{CO_2} , and $\pm 5.4\%$ for $PSF_{\text{CH}_4, \text{proxy}}$ due to the simplification of the radiative transfer calculation to one single background spectrum and set of weighting

Table 2. Parameters studied in the sensitivity analysis and the range in which deviations were analyzed. The second column shows the background scenario used as “truth” in the sensitivity study. The third column notes the range of the perturbation of the parameters. Parameters not mentioned here were constant and estimated as described in Sect. 3.1 for the flight line ang20170811t192639.

Parameter	Standard value	Studied range
Aircraft altitude	5.33 km	4.93 km to 5.93 km
Surface elevation	0.39 km	0.0 km to 0.6 km
Instrument viewing angle	0.00 °	± 18 °
Surface albedo	0.1	0.01 to 0.5
xCH ₄	1.833 ppm	(0.5 to 2) · 1.833 ppm
xCO ₂	399.2 ppm	(0.97 to 1.03) · 399.2 ppm
H ₂ O	5.94 · 10 ²² molec cm ⁻²	(0.5 to 2) · 5.94 · 10 ²² molec cm ⁻²
Pressure profile	US standard scaled to 1015 hPa at sea level	(0.95 to 1.05) · 1015 hPa
Temperature profile	US standard shifted to 299 K at surface	± 10 K
Aerosol scenario	OPAC urban	OPAC background, OPAC desert
		Rangeland vegetation
		Barbed goatgrass
		Soil (Entisol)
Surface reflectance	Constant albedo 0.1	Grey sandstone
		Weathered aluminium
		Weathered steel
		Paving asphalt
		Paving concrete

functions. This uncertainty defined the large scale deviations possible in one flight track and should not be confused with the single pixel precision of the column enhancement, nor did that automatically limit detection. As parameters such as surface elevation in normal cases only vary smoothly, a plume signal on top of this bias is still detectable. However, a problem may occur, if large amounts of CO₂ are co-emitted with a weak methane source. In this case there could be a (partial) masking of the plume due to the negative bias introduced by CO₂.

In contrast to those biases, the different surface types induce widely varying biases (Table 3) at AVIRIS-NG spectral resolution. The proxy method reduced these errors for some surfaces, but not all. For rangeland vegetation, soil, gray sandstone and weathered aluminium the bias after application of the proxy was well below 1%. However, for weathered steel and paving asphalt, the bias increased to 1 – 2%, while for barbed goat grass and paving concrete (i.e. limestone) the bias due to the reflection properties was greater than 7%. This meant, that even after application of the proxy, some residual influence of the surface reflectance will remain. Paving concrete would be especially likely to cause a false positive, since it induced a large




Figure 4. Sensitivity analysis of WFM-DOAS to the examined input parameters of the SCIATRAN radiative transfer calculation. The absolute deviation of the retrieved from the expected PSF_{CH_4} (blue), PSF_{CO_2} (orange) and $PSF_{CH_4,proxy}$ (green) is plotted for each parameter. The largest deviations are present in surface elevation and scaling of the total column of CH₄. With the proxy method, the deviations are reduced for all parameters except for a scaling in the total columns. For enhancements deviations of CO₂, the proxy is worse than the single CH₄ retrieval due to division by the amount of CO₂ (see also Sect. 3.3).

Table 3. Uncertainty estimate of PSF_{CH_4} , PSF_{CO_2} and $PSF_{CH_4,proxy}$ due to the assumption of a constant albedo over different surfaces. As long as there is a constant surface type, a large value does not principally hinder source detection. However, especially paving or man made structures vary spatially with non man made structures, so false positive hits due to surface reflectance are possible.

Surface type	Uncertainty on PSF_{CH_4}	Uncertainty on PSF_{CH_2}	Uncertainty on $PSF_{CH_4,proxy}$
Rangeland vegetation	0.38 %	0.34 %	0.04 %
Barbed goat grass	−7.16 %	−0.06 %	−7.11 %
Soil (Entisol)	0.26 %	0.79 %	−0.53 %
Grey sandstone	0.25 %	0.62 %	−0.37 %
Weathered aluminium	1.30 %	1.05 %	0.25 %
Weathered steel	−0.94 %	0.18 %	−1.11 %
Paving asphalt	2.23 %	0.33 %	1.90 %
Paving concrete	11.89 %	0.77 %	11.04 %

Table 4. Uncertainty estimate resulting from the assumed constant atmospheric and geometric background parameters. For each parameter, the maximum deviations for PSF_{CH_4} and PSF_{CO_2} , as well as for $PSF_{CH_4,proxy}$, are listed. For albedo, the largest value was excluded from this table (see main text). For CH_4 , two different cases are regarded. The case CH_4 ($\pm 20\%$) is valid for most of the plumes and is relevant for detection of smaller sources. The extreme case (100 % increase) is only relevant near very strong sources and is excluded from the averaged systematic uncertainty. The absolutely correct retrieval of PSF_{CO_2} when changing CO_2 is due to the relatively small range of change in CO_2 investigated. However, this induces relatively large uncertainties for $PSF_{CH_4,proxy}$.

Parameter	Uncertainty on PSF_{CH_4}	Uncertainty on PSF_{CO_2}	Uncertainty on $PSF_{CH_4,proxy}$
Aircraft altitude	$\pm 2.3\%$	$\pm 2.1\%$	$\pm 0.2\%$
Surface elevation	$\pm 6.3\%$	$\pm 10.2\%$	$\pm 3.6\%$
Temperature shift	$\pm 3.8\%$	$\pm 3.7\%$	$\pm 0.2\%$
Surface pressure	$\pm 0.5\%$	$\pm 1.0\%$	$\pm 0.5\%$
Viewing angle	$+1.4\%$	$+1.8\%$	-0.4%
Albedo	$\pm 0.5\%$	$\pm 0.8\%$	$\pm 0.4\%$
Water vapor	$+0.9\%$	$+0.6\%$	$+0.2\%$
CH_4 ($\pm 20\%$)	-1.0%	$\pm 0.0\%$	-1.0%
CH_4 (extreme)	-10.0%	$\pm 0.0\%$	-10.0%
CO_2	$\pm 0.0\%$	$\pm 0.0\%$	$\pm 3.1\%$
Aerosol scenario	$\pm 0.1\%$	$\pm 0.3\%$	$\pm 0.2\%$
Systematic uncertainty	$\pm 8.0\%$	$\pm 11.3\%$	$\pm 5.4\%$

positive bias. However, this would be highly correlated to structures visible in the RGB images of the scene. Barbed goat grass,




Figure 5. Raw retrieval results of CH_4 (pure PSF_{CH_4} , not filtered and destriped, no proxy method applied) over a scene with large forest areas and a lake as an example of a dark scene. The noise in the profile scaling factors results from the retrieval not being able to distinguish between absorption features of CH_4 and surface reflectance due to the low signal and therefore low signal to produced large noise ratio. This effect is especially strong over water, but also significant over the forest areas surfaces with low spectral reflectance like forests or lakes.

on the other hand, led to a large underestimation of the total column. However, this surface type normally occurs over large patches of land, so that local enhancements on top of this bias should be detectable in most cases.

3.4 Filtering of poor fits

With the estimate of the influence of the background assumptions in place, we performed radiative transfer calculations for the different flight tracks and applied the retrieval to the whole data set. Examining the data it was obvious, that the retrieval sometimes failed to retrieve meaningful results. Especially over surfaces with low spectral reflectance, and therefore low signal on the detector, it produced mostly noise with profile scaling factors ranging from below 0 to largely over 2 and dramatic changes between neighboring ground scenes (see Fig. 5). This effect, due to the low SNR over dark surfaces, indicated the need to filter out low-radiance ground scenes. For IMAP-DOAS, Ayasse et al. (2018) concluded in a simulation study, that at sensor radiances below $0.1 \mu\text{W cm}^{-2} \text{ nm}^{-1} \text{ sr}^{-1}$ in the background signal led to significantly more inaccurate estimate of the methane column. In this study, we analyzed measured radiance spectra to estimate the radiance below which the retrieval results were not trustworthy.

The threshold was determined by the following procedure: For each ground scene the difference between the measured and the fitted spectrum was calculated for each spectral pixel after the retrieval. These values were added in quadrature to get

the root mean square difference between fit and measurement (RMS). This RMS value was then plotted over the radiance at 2140.0nm in box plots with $0.05 \mu\text{W cm}^{-2} \text{ nm}^{-1} \text{ sr}^{-1}$ bins on the horizontal axis (Fig. 6) for the whole data set. For low radiances this difference increased drastically, implying a strongly reduced fit quality. As a compromise between coverage and quality, we introduced a threshold of $0.1 \mu\text{W cm}^{-2} \text{ nm}^{-1} \text{ sr}^{-1}$. The filter rejected all retrieval results where the radiances at 2140.0nm were below this value. We also rejected measurements with an RMS over 2% to remove the worst outliers. Interestingly, for very bright surfaces the spread of the upper whisker, denoting the 75 to 95 percentile, is increased. This could have been from surfaces such as paving materials or other anthropogenic structures, for which the reflected spectrum already had interfering features similar to the absorption of CH_4 at the AVIRIS-NG spectral resolution (see also Sect. 3.3 and Table 3). This results agree with the findings of Ayasse et al. (2018).

4 Detection and inversion of plumes

4.1 Detection of plumes

For the detection of plumes, we filtered the retrieved CH_4 values (Sect. 3.4), removed striping (Sect. 3.2 and Sect. B in the appendix) and calculated the final $\text{CH}_{4,\text{enh}}$ according to Eq. 2. The proxy method proved necessary; otherwise diffuse CH_4 enhancements would be have been mistaken for true enhancements due to emissions of CH_4 . This can be seen in Fig. 7, where diffuse enhancements in the pure CH_4 results vanished completely after applying the proxy method.

The final $\text{CH}_{4,\text{enh}}$ were plotted as images and were manually inspected for methane plumes. For the reduced data set used in this study (13 tracks), this approach detected several plumes in 10 out of the 13 tracks. However, many plumes were faint or located near infrastructure, making unambiguous detection difficult ~~and/or preventing the application of the cross-sectional flux method (White et al., 1976, and Sect. 4.3) or the plumes were very short.~~ Therefore, we show plumes which are well-suited for the cross-sectional flux method extend over longer ranges and have unambiguous morphology in Fig. 8 and Fig. 9. The additional CH_4 plumes and enhancements ~~not inverted in the following sections~~ can be found in the appendix in Sect. C. Those comprise, among others, emissions most likely resulting from open cast coal mining (Fig. C1) or a well pad located in a forest (Fig. C2).

In Fig. 8, two overpasses of the same source on two different days are shown. On the first day (Fig. 8 (b)), the plume structure was recognizable over a relatively long distance, while on the second day (Fig. 8 (d)), the plume was only faintly visible in the vicinity of the source. This was most likely due to the wind speed, which was significantly higher on the second day (~~4.4ms⁻¹~~7.6ms⁻¹) compared to the first day (~~2.6ms⁻¹~~3.7ms⁻¹). Assuming a constant emission, a wind speed higher by a factor of ~~1.6~~roughly 2 means a decrease in the column enhancements by a factor of ~~1.6, which reduced roughly 2, which would significantly reduce~~ the visibility of the plume in the retrieval results by diluting larger parts of the plume faster below the background noise.

In Fig. 8(b), a new interesting feature was observable at the source. There, we observed a double plume structure that was especially prominent during the first overpass. Comparison with the RGB image revealed, that one part seemed to originate from the vent, while the other part seemed to originate from the top of the shadow of the vent. The vent released the plume

several meters above the surface. Because the plume was very narrow near the source, the sunlight only passed the plume either before or after hitting the ground. As those two light paths were attributed to different ground scenes, the absorption and therefore the apparent CH₄ enhancements were visible at two locations leading to the double plume structure. Further down the plume, atmospheric mixing took place and the plume widened. Then, the sunlight passed through the plume both before and after hitting the ground, and the double plume structure vanished. A simple geometric consideration of the distance between the two plume structures (~ 50 m), the solar and instrument zenith ~~angle and angles~~ ($\sim 42^\circ$ and $\sim 2.5^\circ$), and the vent height ~~estimated-calculated~~ from the shadow of the structure (~ 55 m) and the solar zenith angle supported this hypothesis.

In Fig. 9, two plumes originating from well pads are shown. Both extended linearly from their source and were visible over approximately 100 m. While the first plume (Fig. 9, b) originated from a cold vent (similar to Fig. 8), the emitting structure for the plume in Fig. 9 (d) could not be identified from the RGB images. It seemed, however, that the source was located near the surface. This would also explain the large deviation of the plume direction from the wind direction acquired from the ERA-5 model data, since the nearby forests could have significantly altered the wind direction near the surface.

Additionally, the plume detected over a coal mine ventilation shaft during the Four Corners measurement campaign in 2015 is shown in Fig. 10. The plume was nearly one kilometer long, with a straight profile except for a small diversion at the tip that suggested a change in wind direction.

4.2 Comparison of WFM-DOAS retrieval results with IMAP-DOAS and MF results

To assess the performance of the WFM-DOAS retrieval with respect to the IMAP-DOAS and MF retrieval results, a subset of the data was compared quantitatively. We focused here on the coal mine ventilation shaft plume for all three retrievals, and additionally compared the WFM-DOAS results to MF results over the plumes P1 - P4. As IMAP-DOAS and the MF retrieve CH₄ enhancements below the aircraft in ppm · m (CH_{4,enh,ppm m}), we converted these measurements to enhancements in molec cm⁻² (CH_{4,enh,MF}) via the following equation:

$$\text{CH}_{4,\text{enh,MF/IMAP}} = \frac{\text{CH}_{4,\text{enh,ppm m}}}{h_{\text{airc}}} \cdot \text{subcol}_{\text{tot}} \cdot 10^{-6}, \quad (3)$$

where h_{airc} is the distance between instrument and ground in nadir direction, and $\text{subcol}_{\text{tot}}$ is the number of dry molecules per square centimeter between aircraft and ground. The sub column below the aircraft was calculated on the basis of the profiles used for the WFM-DOAS retrieval.

We compared the retrieval results in two different ways, focusing on retrieval scatter and retrieved enhancements. First, for the retrieval scatter, we calculated the standard error (1-sigma) of retrieved enhancements over areas near the plumes P1 - P5. This is summarized in Table 5. For all plume regions except P5 the retrieval scatter is lower for WFM-DOAS than for the MF results, which indicate less retrieval noise. For P5, where all three retrieval results were available, the background noise is very similar for all retrieval methods.

Second, to evaluate the agreement of the retrieval results, we additionally produced scatter plots of the central plume area. We added the 1:1 line and a linear fit to the scatter plots. For the coal mine ventilation shaft plume all three retrievals agreed well

Table 5. Comparison of background noise of the retrievals based on the standard deviation of retrieval results near the plumes. As we only have IMAP-DOAS results for one plume, the other fields are left empty.

	P1 (molec cm ⁻²)	P2 (molec cm ⁻²)	P3 (molec cm ⁻²)	P4 (molec cm ⁻²)	P5 (molec cm ⁻²)
WFM-DOAS	$\pm 1.3 \cdot 10^{17}$	$\pm 8.3 \cdot 10^{17}$	$\pm 7.8 \cdot 10^{17}$	$\pm 1.8 \cdot 10^{18}$	$\pm 5.2 \cdot 10^{17}$
MF	$\pm 2.0 \cdot 10^{18}$	$\pm 2.1 \cdot 10^{18}$	$\pm 1.7 \cdot 10^{18}$	$\pm 2.4 \cdot 10^{18}$	$\pm 4.1 \cdot 10^{17}$
IMAP-DOAS					$\pm 5.5 \cdot 10^{17}$

in the retrieved enhancements, as can be seen in Fig. 11. Compared to the IMAP-DOAS retrieval, WFM-DOAS did retrieve slightly lower enhancements, while it did estimate slightly higher enhancements than MF.

Additionally, we compared the WFM-DOAS and MF results for the plumes P1 - P4 (see Fig. D1). For P3 especially the higher enhancements located inside the plume agree well. However, for the other plumes there is quite some mismatch between MF results and WFM-DOAS results. The cause for the disagreement is currently under investigation and might be related to the use of a standard gas target for the matched filter rather than calculating a target specifically parameterized for a given scene.

4.3 Cross Flux and uncertainty estimation based on cross sectional flux method

To estimate the fluxes for the selected sources, we applied the cross sectional flux method (White et al., 1976). As this method is computationally and conceptually simple, it could well be used for initial estimations of source strengths. It had successfully been utilized for estimating of emissions detected in remote sensing measurements (for example in Krings et al., 2011; Frankenberg et al., 2016; Krautwurst et al., 2017; Varon et al., 2019). In this method, one calculates the flux F in kilogram per hour through a transect orthogonal to the wind direction with length segments dx_i in meter based-on-using the total column enhancements $CH_{4,enh,i}$ in molecules per square centimeter at position i along the transect:

$$F = f \cdot \sum_i CH_{4,enh,i} \cdot u \cdot dx_i. \quad (4)$$

The wind speed u in meter per second is assumed to be constant in time and space for the time of overflight. For the detected AVIRIS-NG plumes, this assumption was valid, as these plumes had been sampled within seconds. The wind speed was extracted from the-ECMWF-ERA5-nearby weather station data (see Sect. 2.2 and Sect. A), while the wind direction was visually estimated from the observed plume directions by drawing the best fitting line from the source along the plume. The factor $f = 9.589e - 23 \text{ s kg h}^{-1} \text{ molec}^{-1}$ converted the flux to kilogram per hour.

We defined the local background for $CH_{4,enh}$ for each cross section as the region outside of the plume on each side. Then, we calculated the $PSF_{\text{proxy,bg}}$ for the normalization to the local background using a linear fit between both local background regions. This background fit reduced slight gradients present in the background concentration to accurately estimate the column enhancements originating from the source.

As is observable in Fig. 8 and Fig. 10, there were gaps and accumulations along the plume. These were caused by eddies and short gusts which disrupted the plume structure. To account for that atmospheric variability, we defined multiple cross sections along the plume, each one pixel apart. We then calculated the flux for each of the cross sections. The final flux estimate was the mean value of the single fluxes through all cross sections.

5 4.4 Flux and uncertainty estimation based on cross sectional flux method

For the plumes shown in Fig. 8 (in the following P1 and P2 for (b) and (d) respectively) and Fig. 9 (P3 and P4 for (b) and (d) respectively) and Fig. 10 (P5), the methane flux was calculated using the cross sectional flux method. We selected plumes P1 and P2 for two reasons: P1 was visible for approximately 200m before crossing the road, making it possible to define multiple cross sections through the plume and thus leading to a strong reduction in the uncertainties. This was the only source observed twice in an emitting state by AVIRIS-NG, which allowed comparison of the flux estimates for two overflight times. It originated from a vent in a bitumen extraction site. P3 and P4 shown in Fig. 9 (b) and (d) also showed a well shaped straight plume, which was favorable for the cross sectional flux method. In Fig. 8 (b) and Fig. 9 (b), a clear bias due to the underlying road surface was visible. In both cases, cross tracks that overlapped with this bias were excluded from the flux estimation. P5 showed the strongest enhancements over the longest range, which allowed to calculate the flux through over 100 different cross tracks through the plume, which greatly reduced the uncertainty due to atmospheric variability.

Application of the cross sectional flux method on all plumes yielded the following emission estimates: For P1, a mean flux of $(141 \pm 87) \text{ kg h}^{-1}$ $(196 \pm 94) \text{ kg h}^{-1}$ was calculated, while for P2, the mean flux was $(89 \pm 46) \text{ kg h}^{-1}$ $(155 \pm 71) \text{ kg h}^{-1}$. For P3 and P4, the mean fluxes were $(111 \pm 76) \text{ kg h}^{-1}$ and $(70 \pm 79) \text{ kg h}^{-1}$ $(157 \pm 109) \text{ kg h}^{-1}$ and $(204 \pm 116) \text{ kg h}^{-1}$, respectively. We calculated the For P5, the mean flux was $(1220 \pm 450) \text{ kg h}^{-1}$. The wind speeds for the different plumes to be 2.7 ms^{-1} for P1, 4.4 ms^{-1} for P2, 2.8 ms^{-1} for P3 and 1.5 ms^{-1} for P4 according to the weather stations were 3.7 ms^{-1} , 7.6 ms^{-1} , 3.9 ms^{-1} , 4.2 ms^{-1} and 4.1 ms^{-1} for P1 - P5, respectively.

We estimated the uncertainty of the flux via Gaussian error propagation from the uncertainties standard error of the single components of Eq. 4 and Eq. 2, with while additionally accounting for the atmospheric variability. The contribution and derivation of the main error sources for P1 is shown below. We used the same procedure to estimate the uncertainty for the other sources.

We estimated the single pixel precision for each plume region by analyzing the local background of the plume. We used the $1 - \sigma$ standard deviation of the local background $PSF_{\text{CH}_4, \text{proxy}}$ as measure of the retrieval noise and included the uncertainty due to small variations of CO_2 , which might had been coemitted from the flare or collocated to the plume but which were below the noise of the pure PSF_{CO_2} . This led to a single pixel precision of $\sim 3\%$ of the CH_4 background column, which translated to an uncertainty on the final flux of $\pm 26 \text{ kg h}^{-1}$ $\pm 36 \text{ kg h}^{-1}$.

The uncertainty due to atmospheric stability was calculated as the confidence interval of the fluxes through all cross sections. This amounted for an uncertainty on the final flux of $\pm 24 \text{ kg h}^{-1}$ $\pm 33 \text{ kg h}^{-1}$.

For the the-wind, we assumed an uncertainty of $\pm 1.5 \text{ ms}^{-1}$, according to the $1 - \sigma$ standard deviation of the hourly wind measurements of ERA5 compared to inland measurement stations (Minola et al., 2020). For most cases the wind direction

~~matched quite well~~, so we assumed that the wind speed was also reasonably well captured by the model to account for the partly larger distance between weather stations and plume locations as well as the usage of hourly mean data from the weather stations. As the wind speed directly influenced the flux, and the uncertainty could not be reduced by simply taking more cross sections into account, this directly propagated to an uncertainty of the final flux of $\pm 79 \text{ kg h}^{-1}$.

- 5 The estimate of the background column was dependent on a scaled climatology. We assumed a $\pm 5\%$ deviation as an upper limit of the uncertainty for the total column of CH_4 around the plume compared to this climatology. This led to an uncertainty of the final flux of $\pm 7 \text{ kg h}^{-1} \pm 10 \text{ kg h}^{-1}$, which was very small compared to the other uncertainties.

Assuming that those sources of uncertainty were uncorrelated, we combined them in quadrature. This resulted in an uncertainty for P1 of $\pm 87 \text{ kg h}^{-1} \pm 94 \text{ kg h}^{-1}$. An overview over the contribution of the single error sources for all sources is given
10 in Table 6. We emphasize that for all of these cases, the fluxes were calculated from snapshots and were only valid for the time of overflight.

Our estimates for P1 and P2 overlapped within their respective uncertainties, suggesting that the source may have been approximately constant over the two days. ~~This~~ The difference in the mean estimate could be explained by uncertainties in our assumptions of the wind speed, and the stronger dilution of the plume on the second day (Fig. 8). ~~Additionally, the ratio between the mean fluxes (P1/For P2) was nearly the same as the inverse ratio of the wind speeds ($4.4 \text{ ms}^{-1}/2.7 \text{ ms}^{-1}$), which further supported the theory, that dilution due to the higher wind speed played an important role for the lower flux on the second day.~~

~~The plume P4 was observed under unfavorable wind conditions with low winds of 1.5 ms^{-1} . This led to an uncertainty of more than 100%. For such low winds, however, the cross-sectional flux method is not well suited, as there horizontal transport due to diffusion is completely neglected. This however plays an increasing important role the lower the wind speed is (Sharan et al., 1996), additionally the wind speed uncertainty might be even higher, as in such strong wind situations also stronger wind gusts could momentarily dilute the plume.~~

5 Discussion

The WFM-DOAS retrieval provided an efficient and accurate way to handle AVIRIS-NG data quantitatively. In contrast to the
25 IMAP-DOAS retrieval (Frankenberg et al., 2005; Thorpe et al., 2014), the WFM-DOAS retrieval is a non-iterative retrieval with precalculated radiative transfer calculations. This reduced the computational time needed for the retrieval, while still delivering reliable local total column enhancements. On the other hand, the modeled background spectrum is adapted to the physical properties of the scene by scaling the trace gas columns and adapting the geometric parameters necessary to model the average light path over the scene, in contrast to the more statistical approach used in the matched filter (Thompson et al., 2015).
30 For most scenes, WFM-DOAS results showed less background noise than the MF results, apart from the coal mine ventilation shaft plume.

The WFM-DOAS retrieval of CH_4 in the two SWIR fit windows of AVIRIS-NG data produced much larger noise in the results for the weak window than for the strong window. The significantly less noisy WFM-DOAS results around 2300 nm

Table 6. Inversion results and uncertainty estimate ~~based on using~~ the cross sectional flux method for P1 and P2 (two overpasses of the same source on two consecutive days) ~~and~~ ~~P3~~ ~~and~~ ~~P4~~ ~~and~~ ~~P5~~. The single pixel precision was calculated from the $1-\sigma$ standard deviation of a background region for each plume. The atmospheric variability resulted from the confidence interval of the multiple cross sectional fluxes. The wind uncertainty and total column uncertainty was assumed to be $\pm 1.5 \text{ m s}^{-1}$ and 5 %, respectively (see also text for explanation). For all inversions, the wind speed uncertainty is a very large uncertainty. While for P2 additionally the atmospheric variability induces similar large errors, for P3 and P4 this is much less the case. However, there the different surfaces lead to a much larger single pixel uncertainty. ~~For~~ ~~P5 the error is dominated by the wind speed uncertainty.~~

	P1	P2	P3	P4	
Single pixel precision	$\pm 26 \text{ kg h}^{-1}$	$\pm 21 \text{ kg h}^{-1}$	$\pm 40 \text{ kg h}^{-1}$	$\pm 27 \text{ kg h}^{-1}$	
Atmospheric variability	$\pm 24 \text{ kg h}^{-1}$	$\pm 24 \text{ kg h}^{-1}$	$\pm 20 \text{ kg h}^{-1}$	$\pm 12 \text{ kg h}^{-1}$	
Wind uncertainty	$\pm 79 \text{ kg h}^{-1}$	$\pm 26 \text{ kg h}^{-1}$	$\pm 59 \text{ kg h}^{-1}$	$\pm 72 \text{ kg h}^{-1}$	
Total column uncertainty	$\pm 7 \text{ kg h}^{-1}$	$\pm 7 \text{ kg h}^{-1}$	$\pm 4 \text{ kg h}^{-1}$	$\pm 6 \text{ kg h}^{-1}$	
Total flux	$(141 \pm 87) \text{ kg h}^{-1}$	$(89 \pm 46) \text{ kg h}^{-1}$	$(111 \pm 76) \text{ kg h}^{-1}$	$(70 \pm 79) \text{ kg h}^{-1}$	
	P1	P2	P3	P4	<u>P5</u>
Single pixel precision	<u>$\pm 36 \text{ kg h}^{-1}$</u>	<u>$\pm 36 \text{ kg h}^{-1}$</u>	<u>$\pm 78 \text{ kg h}^{-1}$</u>	<u>$\pm 79 \text{ kg h}^{-1}$</u>	<u>$\pm 25 \text{ kg h}^{-1}$</u>
Atmospheric variability	<u>$\pm 33 \text{ kg h}^{-1}$</u>	<u>$\pm 42 \text{ kg h}^{-1}$</u>	<u>$\pm 37 \text{ kg h}^{-1}$</u>	<u>$\pm 35 \text{ kg h}^{-1}$</u>	<u>$\pm 43 \text{ kg h}^{-1}$</u>
Wind uncertainty	$\pm 79 \text{ kg h}^{-1}$	$\pm 26 \text{ kg h}^{-1}$	<u>$\pm 60 \text{ kg h}^{-1}$</u>	<u>$\pm 73 \text{ kg h}^{-1}$</u>	<u>$\pm 447 \text{ kg h}^{-1}$</u>
Total column uncertainty	<u>$\pm 10 \text{ kg h}^{-1}$</u>	$\pm 7 \text{ kg h}^{-1}$	<u>$\pm 8 \text{ kg h}^{-1}$</u>	<u>$\pm 10 \text{ kg h}^{-1}$</u>	<u>$\pm 61 \text{ kg h}^{-1}$</u>
Total flux	<u>$(196 \pm 94) \text{ kg h}^{-1}$</u>	<u>$(155 \pm 71) \text{ kg h}^{-1}$</u>	<u>$(157 \pm 109) \text{ kg h}^{-1}$</u>	<u>$(204 \pm 116) \text{ kg h}^{-1}$</u>	<u>$(1220 \pm 450) \text{ kg h}^{-1}$</u>

most likely originated from the higher number of spectral data points for the fit, and the stronger absorption features. Even though there was approximately half the amount of light reaching the detector at these wavelengths, which reduced the SNR on the detector, the SNR was still high enough for a good retrieval.

Problems however arose over very dark surfaces or surfaces with reflection properties similar to absorption features of CH_4 at the spectral resolution of AVIRIS-NG. This led to residual structures in the retrieved CH_4 maps. Especially paved roads or other anthropogenic structures were observable. Even though the CH_4/CO_2 proxy method reduced false positives in many cases, there was still a remaining dependency of the CH_4 results from the surface spectral reflectance for some surfaces like concrete or barbed goat grass. Additionally, the noise on the retrieval results varied over different surfaces, which is reflected in the uncertainties for the flux inversions for the 4 plumes.

Those effects could be mitigated by deploying and utilizing an imaging spectrometer specifically designed for the task of monitoring CH_4 and CO_2 concentrations, such as the proposed Airborne Methane Plume Spectrometer (AMPS Thorpe et al., 2016b), or the MAMAP 2D system currently being developed and built at the University of Bremen, Germany. Due to the higher spectral resolution, those instruments will have a higher sensitivity to smaller enhancements and should be less influenced by the surface reflectance properties, as is already the case for the MAMAP instrument (see e.g. Krings et al., 2011).

A large uncertainty in the flux inversion of emission sources, which can not be solved by advancing the imaging remote sensing instrument's characteristics, arises from the wind speed estimation. Additional measures have to be taken to reduce this uncertainty. For example, in situ wind measurements in the boundary layer at the plume location could additionally be made (see e.g. Krautwurst et al., 2017; Krings et al., 2018). This approach is especially useful when airborne in situ measurements are included in the campaign design. Another possibility includes the deployment of wind lidars (Wildmann et al., 2020), similar to the approach taken in the CO₂ and Methane mission (CoMet). However, for large surveys or transects, this is not feasible anymore, especially when the source locations of the plumes are not known prior to the flights. There, either advancing to local wind models with much higher spatial resolution such as the GRAL model (Berchet et al., 2017) or the MECO(n) model (Kerkweg and Jöckel, 2012) could lead to a significant uncertainty reduction, but those models have to be carefully validated. Also, methods such as the integrated mass estimation (IME, Jongaramrungruang et al., 2019), which use empirically derived correlations between surface wind, flux rates, plume shape and mass enhancement in the plume to estimate the wind speed and the flux, could help estimating the emissions independent from local wind models or wind measurements. Additionally, more sensitive remote sensing instruments could observe the plume over longer distances, where the plume is likely better mixed in the boundary layer and the horizontal ~~extend~~extent of the plume is less influenced by turbulence and gusts, so that the modeled wind speed ~~most~~in the boundary layer likely better matches the wind speed inside the plume.

6 Summary and Conclusions

We successfully adapted and applied the WFM-DOAS retrieval to AVIRIS-NG data and estimated the uncertainties of this method. In the data set, we were able to detect several point sources. An estimation of the methane emissions of a vent revealed emissions of ~~(141 ± 87) kg h⁻¹ and (89 ± 46) kg h⁻¹~~ (196 ± 94) kg h⁻¹ and (155 ± 71) kg h⁻¹ on two consecutive days, while two other sources related to gas extraction emitted ~~(111 ± 76) kg h⁻¹ and (70 ± 79) kg h⁻¹~~ (157 ± 109) kg h⁻¹ and (204 ± 116) kg h⁻¹, respectively. The emission of the coal mine ventilation shaft was estimated to (1220 ± 450) kg h⁻¹. These source strengths are quite common as indicated by the log normal distribution of sources in the Four Corners region (Frankenberg et al., 2016). A large source of uncertainty for the flux inversion was the wind speed estimate, as no collocated wind speed measurements near the surface were collected. Also the atmospheric variability played an important role for shorter (i.e. smaller) plumes. This influence is reduced for longer plumes, as more tracks are available for the flux estimation. The noise on the retrieval results varied with different surfaces, which contributed notably to the uncertainty of the flux estimate. For the high wind situation, parts of the plume might have been additionally missed due to the higher dilution.

The comparison between IMAP-DOAS/MF and WFM-DOAS resulted in good agreement for the coal mine ventilation shaft and P3. However, larger discrepancies were observed for the plumes P1, P2 and P4. The cause for this discrepancy is under investigation.

The dependency of the resulting total column CH₄ retrieval results from the parameter values assumed in the radiative transfer calculation have been examined. For most parameters, the induced bias was reduced to well below 1 % when using CO₂ as a proxy for light path correction. Large perturbations in elevation resulted in a residual bias, however, the elevation varied

mostly smooth (for example over hills) or only by smaller amounts over buildings or vegetation changes such as from grassland to forests. In addition, very strong CH₄ enhancements led to a systematic underestimation of CH₄, which in consequence could lead to an underestimation of very strong emitters. However, such strong enhancements only occur near to strong sources (not present in this study) and therefore inversion estimates could be performed further down the plume, where the enhancements are lower and therefore the bias is negligible. Deviations of CO₂ from the background on the other hand were retrieved correctly for typical variations of the total column. However, due to the use of CO₂ as proxy for light path correction, these deviations led to a bias in the proxy. Consequently, large amounts of CO₂ coemitted to CH₄ may mask a weak CH₄ plume. Additionally, the influence of some surface reflectance spectra on the retrieved $PSF_{CH_4, proxy}$ was examined. While for some surface types the bias on the retrieved $PSF_{CH_4, proxy}$ could be reduced to well below 1%, some surfaces introduced larger biases, reaching up to 11% for paving concrete.

As dark surfaces mostly produced noise in the retrieval results, ground scenes with at sensor radiances below $0.1 \mu W cm^{-2} nm^{-1} sr^{-1}$ were excluded from the analysis. In the future, this radiance filter described in Sect. 3.4 could be applied to the data before the application of the WFM-DOAS retrieval. This reduces the amount of data which has to be retrieved, without rejecting possible good retrieval results. Additional retrieval improvements could be achieved by fitting CH₄ and CO₂ in both the weak and the strong window simultaneously.

While this and previous studies have demonstrated detection and quantification of methane emission sources with AVIRIS-NG, the residual structures due to the relatively coarse spectral resolution make unambiguous detection and especially quantification of small sources difficult. To mitigate this problem, spectrometers dedicated to the detection and quantification of CH₄ are currently developed, such the proposed AMPS system (Thorpe et al., 2016b) or the MAMAP-2D system, which is being assembled.

Data availability. The AVIRIS-NG data set of the ABOVE campaign is available at https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1569 (Miller et al., 2019a).

Appendix A: Comparison of weather station wind data with ERA5 50 m height average data

We compared the average wind speed over the lowest 50 m height average of ERA5 data to the hourly mean wind speed data obtained from weather stations near the sources. For P1 and P2, we used wind speed data obtained at the Firebag weather station (Wood Buffalo Environmental Association, 2020) which was located slightly below 20 km away from the plumes. For P3 and P4, we used wind speed data from the weather stations “Sundre A” and “Patricia AGCM”¹, located 5 km and 17 km away from the source. For P5, we used wind speed data from the Four Corners Regional Airport weather station from the MesoWest network (Horel et al., 01 Feb. 2002) ~ 15 km away from the plume. A comparison between those ground stations and the average wind speed over the lowest 50 m of ERA5 data is shown in Fig. A1.

¹Data provided by Alberta Agriculture and Forestry, Alberta Climate Information Service (ACIS) <https://acis.alberta.ca> (retrieved in Dec. 2020)

As is clearly visible, the ERA5 data significantly deviated from the wind speeds present at the weather stations. The mean deviation was $+2.5\text{ m s}^{-1}$, with single differences up to $+3.4\text{ m s}^{-1}$. Consequently, we used the weather station wind data for further analysis in the flux inversion.

Appendix B: Striping Effect

- 5 In a push-broom imager such as AVIRIS-NG each detector column acts as one separate line detector looking at a different ground scene. Even with very good calibration and characterization (described in Chapman et al., 2019, for AVIRIS-NG), there will still be small differences in the spectra recorded by two different lines. In the retrieval results, this leads to a line (or column) dependent difference in the retrieval results called “striping” (see for example Fig. 5). To correct this effect, we normalize the PSF_{CH_4} , PSF_{CO_2} and $PSF_{\text{CH}_4,\text{proxy}}$ for each pixel by the median PSF of its corresponding detector column. We
10 select the median for resilience against outliers, which could otherwise have a large impact on the correction

Appendix C: Other detected CH_4 plumes

In this section, additional plumes are shown, which have been detected in the data, but were either too inconsistent or the conditions did not allow for the application of the cross sectional flux method.

- A very interesting observation are the plumes and CH_4 accumulations due to open cast coal mining, as can be seen in Fig. C1.
15 During coal mining, the enclosed methane is released to the environment. For underground mining, the methane concentration in the air in the mining shafts are kept well below the explosion limit by ventilating the shafts with fresh air (Özgen Karacan, 2008). The ventilation shafts then emit the total emissions from the whole mine. In an open coal mine the emissions may come out diffuse, however, according to Fig. C1, it seems as significant amounts of methane may be emitted from the brim, perhaps during cutting of the coal.
20 Additional plumes are shown in Fig. C2 and Fig. C3. Those plumes originate from oil and gas infrastructure such as a well pad in the forest and vents. Especially the vents in C3 are only very faintly visible.




Figure 6. Difference between fit and measurement (RMS) over radiance at 2140.0 nm. The box indicates the first to third quartile range, the whiskers denote the 5 to 95-percentile, and the small orange line inside the box the median RMS value of the according radiance bin. The small blue dots denote outliers outside the 95 percentile. For low radiances, the fitting quality decreases significantly, ~~as shown in the high values and wide spread of the RMS.~~ Therefore, all measurements over surfaces with radiances below $0.1 \mu W \cdot cm^{-2} \cdot nm^{-1} \cdot sr^{-1}$ are filtered out (dashed black vertical line), ~~as the results are unreliable.~~ Results with RMS higher than 2% are filtered out as additional quality flag (dashed orange line). ~~The increase in RMS spread especially at $1.5 \mu W \cdot cm^{-2} \cdot nm^{-1} \cdot sr^{-1}$ might be due to high reflecting structures, which have broad band reflection features similar to absorption features of CH_4 (see Ayasse et al., 2018, and Sect. 3.3).~~




Figure 7. Example of the effect/necessity of the proxy method from flight line ang20170823t180156. In (a), the destriped PSF_{CH_4} is shown. Simply analyzing this image, one could assume a diffuse enhancement due to perhaps coal mining activities, as this is part of an open cast coal mine. In figure (b) the destriped PSF_{CO_2} is shown. There, a similar enhancement is visible. In (c) the destriped proxy results are shown, where this diffuse enhancement vanishes.





Figure 8. Plume resulting from a cold vent. The black arrow denotes the source position, while the red arrow indicates the wind direction and wind speed according to ERA-5 data for comparison to the plume direction. In (b) the plume is much fainter, most likely due to the higher wind speed. Especially in (a) a double plume structure is visible. ~~This structure likely results from in the light passing through the plume only once before reflection on the ground for one first part of the double structure, and only once after reflection for the second plume.~~ As vertical and horizontal mixing takes place the further the emissions travel, the light passes through the plume twice, and the double plume structure vanishes. The roads are prominently visible in the retrieval results. It might be that this road is made of concrete, or otherwise contains limestone. ~~The roads were excluded from the flux analysis due to the high bias induced by it.~~



Flight line ang20150422t164506 - Source P5




Figure 10. Similar to Fig. 8, but for the exhaust plume of a coal mine ventilation shaft in the Four Corners region.

Comparison of retrieval results for ang20150422t164506




Figure 11. Scatter plot comparison between IMAP-DOAS and WFM-DOAS (a) and MF and IMAP-DOAS (b) respectively for the coal mine ventilation shaft plume. For the red line only enhancements above the background noise stated in Table 5 were analyzed, which should be largely dominated by the plume.




Figure A1. Comparison between wind speed averaged over the lowest 50 m above ground for ERA5 data and collocated wind station data. ERA5 data significantly underestimated the wind speed present at a given time.




Figure C1. ~~Methane~~ Same as Fig. 9, but for methane enhancements after application of the proxy resulting from bituminous coal extraction. On the left (a, c, e) the RGB-images and on the right (b, d, f) the according total column-enhancements of CH₄ are shown. The upper two measurements ~~where were~~ taken within ~ 10 minutes, and show the flight lines lay very close to each other, so that the brim is the same brim. The dashed line is plotted on the same location in both images. While in (b) a plume emanating from the brim is visible, in (d) strong accumulations near the brim are visible. In (f) the plume is more diffuse and the highest enhancements are ~~located~~ visible near a brim. However, it is not as clear as in (b) and (d), if it really originates from the brim or might be an artifact.




Figure C2. Similar to Fig. C1, but for methane enhancements from oil/gas infrastructure. In (a) and (b), the emissions originate from a well pad in a forest. Due to the low radiance over the trees, only the well pad itself passes the quality filters. The enhancements seem accumulated and no clear wind direction is visible. In (c) and (d), a facility located at a bitumen extraction site is shown and meanders through the facility.




Figure C3. Similar to Fig. C1, but for two overpasses over cold vents at a bitumen extraction site. The plumes are only faint, especially for the second overpass (d). In the results a small striping effect is still visible. The destriping reduces the effect, but is not able to totally eliminate it.

Appendix D: Additional comparison between MF and WFM-DOAS

In addition to the comparison of IMAP-DOAS and MF for the same plume, we compared MF and WFM-DOAS retrieval results for subregions zoomed in on the plumes P1 - P4. For the scene P3 the retrievals seem to agree quite well. Especially the higher enhancements correlated with the plume lay near the 1:1 line. However, for the plumes P1, P2 and P4, there is a bigger discrepancy between MF and WFM-DOAS, which can not be explained by the larger scatter alone. The cause for this is currently under investigation.

Comparison WFM-DOAS with MF




Figure D1. Similar to Fig. 11, but for the plume regions P1 - P4.

Author contributions. JB contributed to the study design, adapted the retrieval, analyzed the data and wrote the manuscript, KG and HB initialized the study, KG and SK contributed to the retrieval adaption, study design and paper draft, AKT, DRT and CF helped in data set selection and with the handling of the data, RD and CM designed the flight plans, organized and led the flights, HB and JPB supervised the study, contributing to the scientific objectives and interpretation of results. All authors contributed to the final manuscript.

5 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This work was funded in parts by the BMBF project AIRSPACE MAMAP2D (FKZ01LK1701B). We would also like to acknowledge the contributions of the AVIRIS flight and instrument teams. The AVIRIS-NG data were collected as part of the Arctic Boreal Vulnerability Experiment (ABoVE), a NASA Terrestrial Ecology program, and other methane studies supported by NASA's Earth Science Division. A portion of this work was performed at the Jet Propulsion Laboratory, California Institute of Technology, under contract

10 with NASA. [The hourly resolved wind data for P3 and P4 were provided provided by Alberta Agriculture and Forestry, Alberta Climate Information Service \(ACIS\), <https://acis.alberta.ca> for stations Patricia AGCM and Sundre A \(Data retrieved on Dec, 2020\)](#)

References

- Ayasse, A. K., Thorpe, A. K., Roberts, D. A., Funk, C. C., Dennison, P. E., Frankenberg, C., Steffke, A., and Aubrey, A. D.: Evaluating the effects of surface properties on methane retrievals using a synthetic airborne visible/infrared imaging spectrometer next generation (AVIRIS-NG) image, *Remote Sensing of Environment*, 215, 386–397, <https://doi.org/10.1016/j.rse.2018.06.018>, 2018.
- 5 Baldrige, A., Hook, S., Grove, C., and Rivera, G.: The ASTER spectral library version 2.0, *Remote Sensing of Environment*, 113, 711–715, <https://doi.org/10.1016/j.rse.2008.11.007>, 2009.
- Berchet, A., Zink, K., Muller, C., Oetl, D., Brunner, J., Emmenegger, L., and Brunner, D.: A cost-effective method for simulating city-wide air flow and pollutant dispersion at building resolving scale, *Atmospheric Environment*, 158, 181–196, <https://doi.org/10.1016/j.atmosenv.2017.03.030>, 2017.
- 10 Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission Objectives and Measurement Modes, *Journal of the Atmospheric Sciences*, 56, 127–150, [https://doi.org/10.1175/1520-0469\(1999\)056<0127:smoamm>2.0.co;2](https://doi.org/10.1175/1520-0469(1999)056<0127:smoamm>2.0.co;2), 1999.
- Buchwitz, M., Rozanov, V. V., and Burrows, J. P.: A near-infrared optimized DOAS method for the fast global retrieval of atmospheric CH₄, CO, CO₂, H₂O, and N₂O total column amounts from SCIAMACHY Envisat-1 nadir radiances, *Journal of Geophysical Research: Atmospheres*, 105, 15 231–15 245, <https://doi.org/10.1029/2000jd900191>, 2000.
- 15 Buchwitz, M., Schneising, O., Reuter, M., Heymann, J., Krautwurst, S., Bovensmann, H., Burrows, J. P., Boesch, H., Parker, R. J., Somkuti, P., Detmers, R. G., Hasekamp, O. P., Aben, I., Butz, A., Frankenberg, C., and Turner, A. J.: Satellite-derived methane hotspot emission estimates using a fast data-driven method, *Atmospheric Chemistry and Physics*, 17, 5751–5774, <https://doi.org/10.5194/acp-17-5751-2017>, 2017.
- 20 Burrows, J., Hölzle, E., Goede, A., Visser, H., and Fricke, W.: SCIAMACHY—scanning imaging absorption spectrometer for atmospheric cartography, *Acta Astronautica*, 35, 445–451, [https://doi.org/10.1016/0094-5765\(94\)00278-T](https://doi.org/10.1016/0094-5765(94)00278-T), 1995.
- Chapman, J. W., Thompson, D. R., Helmlinger, M. C., Bue, B. D., Green, R. O., Eastwood, M. L., Geier, S., Olson-Duvall, W., and Lundeen, S. R.: Spectral and Radiometric Calibration of the Next Generation Airborne Visible Infrared Spectrometer (AVIRIS-NG), *Remote Sensing*, 11, 2129, <https://doi.org/10.3390/rs11182129>, 2019.
- 25 Chen, Y., Sun-Mack, S., Arduini, R. F., Trepte, Q., and Minnis, P.: Clear -Sky and Surface Narrowband Albedo Datasets Derived from MODIS Data, <https://doi.org/10.1117/12.511180>, 2006.
- Copernicus Climate Change Service (C3S) (2017): ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate, Copernicus Climate Change Service Climate Data Store (CDS), <https://cds.climate.copernicus.eu/cdsapp#!/home>, 2017.
- Cusworth, D. H., Jacob, D. J., Varon, D. J., Miller, C. C., Liu, X., Chance, K., Thorpe, A. K., Duren, R. M., Miller, C. E., Thompson, D. R., 30 Frankenberg, C., Guanter, L., and Randles, C. A.: Potential of next-generation imaging spectrometers to detect and quantify methane point sources from space, *Atmospheric Measurement Techniques*, 12, 5655–5668, <https://doi.org/10.5194/amt-12-5655-2019>, 2019.
- Cusworth, D. H., Duren, R. M., Thorpe, A. K., Tseng, E., Thompson, D., Guha, A., Newman, S., Foster, K. T., and Miller, C. E.: Using remote sensing to detect, validate, and quantify methane emissions from California solid waste operations, *Environmental Research Letters*, 15, 054 012, <https://doi.org/10.1088/1748-9326/ab7b99>, 2020.
- 35 Dlugokencky, E.: Globally Averaged marine surface annual mean growth rate, www.esrl.noaa.gov/gmd/ccgg/trends_ch4/, 2018.

- Dlugokencky, E. J., Nisbet, E. G., Fisher, R., and Lowry, D.: Global atmospheric methane: budget, changes and dangers, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369, 2058–2072, <https://doi.org/10.1098/rsta.2010.0341>, 2011.
- Duren, R. M., Thorpe, A. K., Foster, K. T., Rafiq, T., Hopkins, F. M., Yadav, V., Bue, B. D., Thompson, D. R., Conley, S., Colombi, N. K.,
 5 Frankenberg, C., McCubbin, I. B., Eastwood, M. L., Falk, M., Herner, J. D., Croes, B. E., Green, R. O., and Miller, C. E.: California’s methane super-emitters, *Nature*, 575, 180–184, <https://doi.org/10.1038/s41586-019-1720-3>, 2019.
- Foote, M. D., Dennison, P. E., Thorpe, A. K., Thompson, D. R., Jongaramrungruang, S., Frankenberg, C., and Joshi, S. C.: Fast and Accurate Retrieval of Methane Concentration From Imaging Spectrometer Data Using Sparsity Prior, *IEEE Transactions on Geoscience and Remote Sensing*, 58, 6480–6492, <https://doi.org/10.1109/tgrs.2020.2976888>, 2020.
- 10 Frankenberg, C., Platt, U., and Wagner, T.: Iterative maximum a posteriori (IMAP)-DOAS for retrieval of strongly absorbing trace gases: Model studies for CH₄ and CO₂ retrieval from near infrared spectra of SCIAMACHY onboard ENVISAT, *Atmospheric Chemistry and Physics*, 5, 9–22, <https://doi.org/10.5194/acp-5-9-2005>, <http://www.atmos-chem-phys.net/5/9/2005/>, 2005.
- Frankenberg, C., Meirink, J. F., Bergamaschi, P., Goede, A. P. H., Heimann, M., Körner, S., Platt, U., van Weele, M., and Wagner, T.: Satellite chartography of atmospheric methane from SCIAMACHY on board ENVISAT: Analysis of the years 2003 and 2004, *Journal of Geophysical Research*, 111, <https://doi.org/10.1029/2005jd006235>, 2006.
 15
- Frankenberg, C., Thorpe, A. K., Thompson, D. R., Hulley, G., Kort, E. A., Vance, N., Borchardt, J., Krings, T., Gerilowski, K., Sweeney, C., Conley, S., Bue, B. D., Aubrey, A. D., Hook, S., and Green, R. O.: Airborne methane remote measurements reveal heavy-tail flux distribution in Four Corners region, *Proceedings of the National Academy of Sciences*, <https://doi.org/10.1073/pnas.1605617113>, <http://www.pnas.org/content/early/2016/08/10/1605617113.abstract>, 2016.
- 20 Gerilowski, K., Tretner, A., Krings, T., Buchwitz, M., Bertagnolio, P. P., Belemezov, F., Erzinger, J., Burrows, J. P., and Bovensmann, H.: MAMAP – a new spectrometer system for column-averaged methane and carbon dioxide observations from aircraft: instrument description and performance analysis, *Atmospheric Measurement Techniques*, 4, 215–243, <https://doi.org/10.5194/amt-4-215-2011>, <http://www.atmos-meas-tech.net/4/215/2011/>, 2011.
- Gordon, I., Rothman, L., Hill, C., Kochanov, R., Tan, Y., Bernath, P., Birk, M., Boudon, V., Campargue, A., Chance, K., Drouin, B., Flaud,
 25 J.-M., Gamache, R., Hodges, J., Jacquemart, D., Perevalov, V., Perrin, A., Shine, K., Smith, M.-A., Tennyson, J., Toon, G., Tran, H., Tyuterev, V., Barbe, A., Császár, A., Devi, V., Furtenbacher, T., Harrison, J., Hartmann, J.-M., Jolly, A., Johnson, T., Karman, T., Kleiner, I., Kyuberis, A., Loos, J., Lyulin, O., Massie, S., Mikhailenko, S., Moazzen-Ahmadi, N., Müller, H., Naumenko, O., Nikitin, A., Polyansky, O., Rey, M., Rotger, M., Sharpe, S., Sung, K., Starikova, E., Tashkun, S., Auwera, J. V., Wagner, G., Wilzewski, J., Wcisło, P., Yu, S., and Zak, E.: The HITRAN2016 molecular spectroscopic database, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 203, 3–69,
 30 <https://doi.org/10.1016/j.jqsrt.2017.06.038>, 2017.
- Hamlin, L., Green, R. O., Mouroulis, P., Eastwood, M., Wilson, D., Dudik, M., and Paine, C.: Imaging spectrometer science measurements for Terrestrial Ecology: AVIRIS and new developments, in: 2011 Aerospace Conference, IEEE, <https://doi.org/10.1109/aero.2011.5747395>, 2011.
- Hess, M., Koepke, P., and Schult, I.: Optical Properties of Aerosols and Clouds: The Software Package OPAC, *Bulletin of the American Meteorological Society*, 79, 831–844, [https://doi.org/10.1175/1520-0477\(1998\)079<0831:opoaac>2.0.co;2](https://doi.org/10.1175/1520-0477(1998)079<0831:opoaac>2.0.co;2), 1998.
 35
- Horel, J., Splitt, M., Dunn, L., Pechmann, J., White, B., Ciliberti, C., Lazarus, S., Slemmer, J., Zaff, D., and Burks, J.: MESOW-EST: COOPERATIVE MESONETS IN THE WESTERN UNITED STATES, *Bulletin of the American Meteorological Society*, 83,

- 211 – 226, [https://doi.org/10.1175/1520-0477\(2002\)083<0211:MCMITW>2.3.CO;2](https://doi.org/10.1175/1520-0477(2002)083<0211:MCMITW>2.3.CO;2), https://journals.ametsoc.org/view/journals/bams/83/2/1520-0477_2002_083_0211_mcmittw_2_3_co_2.xml, 01 Feb. 2002.
- Hu, H., Hasekamp, O., Butz, A., Galli, A., Landgraf, J., de Brugh, J. A., Borsdorff, T., Scheepmaker, R., and Aben, I.: The operational methane retrieval algorithm for TROPOMI, *Atmospheric Measurement Techniques*, 9, 5423–5440, [https://doi.org/10.5194/amt-9-5423-](https://doi.org/10.5194/amt-9-5423-2016)
5 2016, 2016.
- Hu, H., Landgraf, J., Detmers, R., Borsdorff, T., de Brugh, J. A., Aben, I., Butz, A., and Hasekamp, O.: Toward Global Mapping of Methane With TROPOMI: First Results and Intersatellite Comparison to GOSAT, *Geophysical Research Letters*, 45, 3682–3689, <https://doi.org/10.1002/2018gl077259>, 2018.
- Hulley, G. C., Duren, R. M., Hopkins, F. M., Hook, S. J., Vance, N., Guillevic, P., Johnson, W. R., Eng, B. T., Mihaly, J. M., Jovanovic, V. M., Chazanoff, S. L., Staniszewski, Z. K., Kuai, L., Worden, J., Frankenberg, C., Rivera, G., Aubrey, A. D., Miller, C. E., Malakar, N. K., Tomás, J. M. S., and Holmes, K. T.: High spatial resolution imaging of methane and other trace gases with the airborne Hyperspectral Thermal Emission Spectrometer (HyTES), *Atmospheric Measurement Techniques*, 9, 2393–2408, [https://doi.org/10.5194/amt-9-2393-](https://doi.org/10.5194/amt-9-2393-2016)
10 2016, 2016.
- IPCC 2013: climate change 2013: the physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change, [Stocker, TF and Qin, D and Plattner, GK and Tignor, M and Allen, SK and Boschung, J and Nauels, A and Xia, Y and Bex, B and Midgley, BM], Cambridge University Press, <https://doi.org/10.1017/CBO9781107415324>, www.cambridge.org/9781107661820, 2013.
- Jongaramrungruang, S., Frankenberg, C., Matheou, G., Thorpe, A. K., Thompson, D. R., Kuai, L., and Duren, R. M.: Towards accurate methane point-source quantification from high-resolution 2-D plume imagery, *Atmospheric Measurement Techniques*, 12, 6667–6681, <https://doi.org/10.5194/amt-12-6667-2019>, 2019.
20
- Kerkweg, A. and Jöckel, P.: The 1-way on-line coupled atmospheric chemistry model system MECO(n) – Part 1: Description of the limited-area atmospheric chemistry model COSMO/MESSy, *Geoscientific Model Development*, 5, 87–110, [https://doi.org/10.5194/gmd-5-87-](https://doi.org/10.5194/gmd-5-87-2012)
2012, 2012.
- Kokaly, R. F., Clark, R. N., Swayze, G. A., Livo, K. E., Hoefen, T. M., Pearson, N. C., Wise, R. A., Benzal, W. M., Lowers, H. A., Driscoll, R. L., and Klein, A. J.: USGS Spectral Library Version 7, <https://doi.org/10.3133/ds1035>, 2017.
25
- Krautwurst, S., Gerilowski, K., Jonsson, H. H., Thompson, D. R., Kolyer, R. W., Iraci, L. T., Thorpe, A. K., Horstjann, M., Eastwood, M., Leifer, I., Vigil, S. A., Krings, T., Borchardt, J., Buchwitz, M., Fladeland, M. M., Burrows, J. P., and Bovensmann, H.: Methane emissions from a Californian landfill, determined from airborne remote sensing and in situ measurements, *Atmospheric Measurement Techniques*, 10, 3429–3452, <https://doi.org/10.5194/amt-10-3429-2017>, 2017.
- 30 Krings, T., Gerilowski, K., Buchwitz, M., Reuter, M., Tretner, A., Erzinger, J., Heinze, D., Pflüger, U., Burrows, J. P., and Bovensmann, H.: MAMAP – a new spectrometer system for column-averaged methane and carbon dioxide observations from aircraft: retrieval algorithm and first inversions for point source emission rates, *Atmospheric Measurement Techniques*, 4, 1735–1758, [https://doi.org/10.5194/amt-4-](https://doi.org/10.5194/amt-4-1735-2011)
1735-2011, <http://www.atmos-meas-tech.net/4/1735/2011/>, 2011.
- Krings, T., Gerilowski, K., Buchwitz, M., Hartmann, J., Sachs, T., Erzinger, J., Burrows, J. P., and Bovensmann, H.: Quantification of methane emission rates from coal mine ventilation shafts using airborne remote sensing data, *Atmospheric Measurement Techniques*, 6, 151–166, <https://doi.org/10.5194/amt-6-151-2013>, <http://www.atmos-meas-tech.net/6/151/2013/>, 2013.
35

- Krings, T., Neininger, B., Gerilowski, K., Krautwurst, S., Buchwitz, M., Burrows, J. P., Lindemann, C., Ruhtz, T., Schüttemeyer, D., and Bovensmann, H.: Airborne remote sensing and in situ measurements of atmospheric CO₂ to quantify point source emissions, *Atmospheric Measurement Techniques*, 11, 721–739, <https://doi.org/10.5194/amt-11-721-2018>, 2018.
- Meerdink, S. K., Hook, S. J., Roberts, D. A., and Abbott, E. A.: The ECOSTRESS spectral library version 1.0, *Remote Sensing of Environment*, 230, 111 196, <https://doi.org/10.1016/j.rse.2019.05.015>, 2019.
- Miller, C., Green, R., Thompson, D., Thorpe, A., Eastwood, M., McCubbin, I., Olson-Duvall, W., Bernas, M., Sarture, C., Nolte, S., Rios, L., Hernandez, M., Bue, B., and Lundeenh, S.: ABoVE: Hyperspectral Imagery from AVIRIS-NG, Alaskan and Canadian Arctic, 2017-2018, <https://doi.org/10.3334/ornl daac/1569>, 2019a.
- Miller, C. E., Griffith, P., Goetz, S., Hoy, E., Pinto, N., McCubbin, I., Thorpe, A. K., Hofton, M. M., Hodkinson, D. J., Hansen, C., Woods, J., Larsen, E. K., Kasischke, E. S., and Margolis, H. A.: An overview of ABoVE airborne campaign data acquisitions and science opportunities, *Environmental Research Letters*, <https://doi.org/10.1088/1748-9326/ab0d44>, 2019b.
- Minola, L., Zhang, F., Azorin-Molina, C., Pirooz, A. A. S., Flay, R. G. J., Hersbach, H., and Chen, D.: Near-surface mean and gust wind speeds in ERA5 across Sweden: towards an improved gust parametrization, *Climate Dynamics*, <https://doi.org/10.1007/s00382-020-05302-6>, 2020.
- Nisbet, E. G., Manning, M. R., Dlugokencky, E. J., Fisher, R. E., Lowry, D., Michel, S. E., Myhre, C. L., Platt, S. M., Allen, G., Bousquet, P., Brownlow, R., Cain, M., France, J. L., Hermansen, O., Hossaini, R., Jones, A. E., Levin, I., Manning, A. C., Myhre, G., Pyle, J. A., Vaughn, B. H., Warwick, N. J., and White, J. W. C.: Very Strong Atmospheric Methane Growth in the 4 Years 2014–2017: Implications for the Paris Agreement, *Global Biogeochemical Cycles*, 33, 318–342, <https://doi.org/10.1029/2018gb006009>, 2019.
- Oceanic, U. S. N., Administration, A., and Force, U. S. A.: US standard atmosphere, 1976, vol. 76, National Oceanic and Atmospheric Administration, 1976.
- Pandey, S., Gautam, R., Houweling, S., van der Gon, H. D., Sadavarte, P., Borsdorff, T., Hasekamp, O., Landgraf, J., Tol, P., van Kempen, T., Hoogeveen, R., van Hees, R., Hamburg, S. P., Maasackers, J. D., and Aben, I.: Satellite observations reveal extreme methane leakage from a natural gas well blowout, *Proceedings of the National Academy of Sciences*, p. 201908712, <https://doi.org/10.1073/pnas.1908712116>, 2019.
- Paris Agreement: United Nations Framework Convention on Climate Change, Paris, France, 2015.
- Reuter, M., Buchwitz, M., Schneising, O., Hase, F., Heymann, J., Guerlet, S., Cogan, A. J., Bovensmann, H., and Burrows, J. P.: A simple empirical model estimating atmospheric CO₂ background concentrations, *Atmospheric Measurement Techniques*, 5, 1349–1357, <https://doi.org/10.5194/amt-5-1349-2012>, <http://www.atmos-meas-tech.net/5/1349/2012>, 2012.
- Reuter, M., Buchwitz, M., Schneising, O., Noël, S., Bovensmann, H., Burrows, J. P., Boesch, H., Noia, A. D., Anand, J., Parker, R. J., Somkuti, P., Wu, L., Hasekamp, O. P., Aben, I., Kuze, A., Suto, H., Shiomi, K., Yoshida, Y., Morino, I., Crisp, D., O’Dell, C. W., Notholt, J., Petri, C., Warneke, T., Velasco, V. A., Deutscher, N. M., Griffith, D. W. T., Kivi, R., Pollard, D. F., Hase, F., Sussmann, R., Té, Y. V., Strong, K., Roche, S., Sha, M. K., Mazière, M. D., Feist, D. G., Iraci, L. T., Roehl, C. M., Retscher, C., and Schepers, D.: Ensemble-based satellite-derived carbon dioxide and methane column-averaged dry-air mole fraction data sets (2003–2018) for carbon and climate applications, *Atmospheric Measurement Techniques*, 13, 789–819, <https://doi.org/10.5194/amt-13-789-2020>, 2020.
- Rozanov, V., Dinter, T., Rozanov, A., Wolanin, A., Bracher, A., and Burrows, J.: Radiative transfer modeling through terrestrial atmosphere and ocean accounting for inelastic processes: Software package SCIATRAN, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 194, 65–85, <https://doi.org/10.1016/j.jqsrt.2017.03.009>, 2017.

- Saunois, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken, D., Houweling, S., Janssens-Maenhout, G., Tubiello, F. N., Castaldi, S., Jackson, R. B., Alexe, M., Arora, V. K., Beerling, D. J., Bergamaschi, P., Blake, D. R., Brailsford, G., Brovkin, V., Bruhwiler, L., Crevoisier, C., Crill, P., Covey, K., Curry, C., Frankenberg, C., Gedney, N., Höglund-Isaksson, L., Ishizawa, M., Ito, A., Joos, F., Kim, H.-S., Kleinen, T., Krummel, P., Lamarque, J.-F., Langenfelds, R., Locatelli, R., Machida, T., Maksyutov, S., McDonald, K. C., Marshall, J., Melton, J. R., Morino, I., Naik, V., O'Doherty, S., Parmentier, F.-J. W., Patra, P. K., Peng, C., Peng, S., Peters, G. P., Pison, I., Prigent, C., Prinn, R., Ramonet, M., Riley, W. J., Saito, M., Santini, M., Schroeder, R., Simpson, I. J., Spahni, R., Steele, P., Takizawa, A., Thornton, B. F., Tian, H., Tohjima, Y., Viovy, N., Voulgarakis, A., van Weele, M., van der Werf, G. R., Weiss, R., Wiedinmyer, C., Wilton, D. J., Wiltshire, A., Worthy, D., Wunch, D., Xu, X., Yoshida, Y., Zhang, B., Zhang, Z., and Zhu, Q.: The global methane budget 2000–2012, *Earth System Science Data*, 8, 697–751, <https://doi.org/10.5194/essd-8-697-2016>, 2016.
- 10 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isakson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murgia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentretter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, P. L., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000-2017, *Earth System Science Data Discussions*, pp. 1–138, <https://doi.org/10.5194/essd-2019-128>, 2019.
- 20 Scafutto, R. D. M., de Souza Filho, C. R., Riley, D. N., and de Oliveira, W. J.: Evaluation of thermal infrared hyperspectral imagery for the detection of onshore methane plumes: Significance for hydrocarbon exploration and monitoring, *International Journal of Applied Earth Observation and Geoinformation*, 64, 311–325, <https://doi.org/10.1016/j.jag.2017.07.002>, 2018.
- Schaefer, H.: On the Causes and Consequences of Recent Trends in Atmospheric Methane, *Current Climate Change Reports*, <https://doi.org/10.1007/s40641-019-00140-z>, 2019.
- 25 Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Reuter, M., Notholt, J., Macatangay, R., and Warneke, T.: Three years of greenhouse gas column-averaged dry air mole fractions retrieved from satellite – Part 1: Carbon dioxide, *Atmospheric Chemistry and Physics*, 8, 3827–3853, <https://doi.org/10.5194/acp-8-3827-2008>, 2008.
- Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Bergamaschi, P., and Peters, W.: Three years of greenhouse gas column-averaged dry air mole fractions retrieved from satellite – Part 2: Methane, *Atmospheric Chemistry and Physics*, 9, 443–465, <https://doi.org/10.5194/acp-9-443-2009>, 2009.
- 30 Schneising, O., Buchwitz, M., Reuter, M., Bovensmann, H., Burrows, J. P., Borsdorff, T., Deutscher, N. M., Feist, D. G., Griffith, D. W. T., Hase, F., Hermans, C., Iraci, L. T., Kivi, R., Landgraf, J., Morino, I., Notholt, J., Petri, C., Pollard, D. F., Roche, S., Shiomi, K., Strong, K., Sussmann, R., Velasco, V. A., Warneke, T., and Wunch, D.: A scientific algorithm to simultaneously retrieve carbon monoxide and methane from TROPOMI onboard Sentinel-5 Precursor, *Atmospheric Measurement Techniques*, 12, 6771–6802, <https://doi.org/10.5194/amt-12-6771-2019>, 2019.
- 35 Sharan, M., Yadav, A. K., Singh, M., Agarwal, P., and Nigam, S.: A mathematical model for the dispersion of air pollutants in low wind conditions, *Atmospheric Environment*, 30, 1209–1220, [https://doi.org/10.1016/1352-2310\(95\)00442-4](https://doi.org/10.1016/1352-2310(95)00442-4), 1996.

- Thompson, D. R., Leifer, I., Bovensmann, H., Eastwood, M., Fladeland, M., Frankenberg, C., Gerilowski, K., Green, R. O., Kratwurst, S., Krings, T., Luna, B., and Thorpe, A. K.: Real-time remote detection and measurement for airborne imaging spectroscopy: a case study with methane, *Atmospheric Measurement Techniques*, 8, 4383–4397, <https://doi.org/10.5194/amt-8-4383-2015>, <http://www.atmos-meas-tech.net/8/4383/2015>, 2015.
- 5 Thompson, D. R., Thorpe, A. K., Frankenberg, C., Green, R. O., Duren, R., Guanter, L., Hollstein, A., Middleton, E., Ong, L., and Ungar, S.: Space-based remote imaging spectroscopy of the Aliso Canyon CH₄ superemitter, *Geophysical Research Letters*, 43, 6571–6578, <https://doi.org/10.1002/2016gl069079>, 2016.
- Thorpe, A., Frankenberg, C., Aubrey, A., Roberts, D., Nottrott, A., Rahn, T., Sauer, J., Dubey, M., Costigan, K., Arata, C., Steffke, A., Hills, S., Haselwimmer, C., Charlesworth, D., Funk, C., Green, R., Lundeen, S., Boardman, J., Eastwood, M., Sarture, C., Nolte,
10 S., McCubbin, I., Thompson, D., and McFadden, J.: Mapping methane concentrations from a controlled release experiment using the next generation airborne visible/infrared imaging spectrometer (AVIRIS-NG), *Remote Sensing of Environment*, 179, 104–115, <https://doi.org/10.1016/j.rse.2016.03.032>, 2016a.
- Thorpe, A. K., Roberts, D. A., Bradley, E. S., Funk, C. C., Dennison, P. E., and Leifer, I.: High resolution mapping of methane emissions from marine and terrestrial sources using a Cluster-Tuned Matched Filter technique and imaging spectrometry, *Remote Sensing of Environment*,
15 134, 305–318, <https://doi.org/10.1016/j.rse.2013.03.018>, 2013.
- Thorpe, A. K., Frankenberg, C., and Roberts, D. A.: Retrieval techniques for airborne imaging of methane concentrations using high spatial and moderate spectral resolution: application to AVIRIS, *Atmospheric Measurement Techniques*, 7, 491–506, <https://doi.org/10.5194/amt-7-491-2014>, <http://www.atmos-meas-tech.net/7/491/2014>, 2014.
- Thorpe, A. K., Frankenberg, C., Green, R. O., Thompson, D. R., Aubrey, A. D., Mouroulis, P., Eastwood, M. L., and Matheou, G.: The Air-
20 borne Methane Plume Spectrometer (AMPS): Quantitative imaging of methane plumes in real time, in: 2016 IEEE Aerospace Conference, IEEE, <https://doi.org/10.1109/aero.2016.7500756>, 2016b.
- Thorpe, A. K., Frankenberg, C., Thompson, D. R., Duren, R. M., Aubrey, A. D., Bue, B. D., Green, R. O., Gerilowski, K., Krings, T., Borchardt, J., Kort, E. A., Sweeney, C., Conley, S., Roberts, D. A., and Dennison, P. E.: Airborne DOAS retrievals of methane, carbon dioxide, and water vapor concentrations at high spatial resolution: application to AVIRIS-NG, *Atmospheric Measurement Techniques*, 10,
25 3833–3850, <https://doi.org/10.5194/amt-10-3833-2017>, 2017.
- Thorpe, A. K., Duren, R. M., Conley, S., Prasad, K. R., Bue, B. D., Yadav, V., Foster, K. T., Rafiq, T., Hopkins, F. M., Smith, M. L., Fischer, M. L., Thompson, D. R., Frankenberg, C., McCubbin, I. B., Eastwood, M. L., Green, R. O., and Miller, C. E.: Methane emissions from underground gas storage in California, *Environmental Research Letters*, 15, 045 005, <https://doi.org/10.1088/1748-9326/ab751d>, 2020.
- Varon, D. J., McKeever, J., Jervis, D., Maasakkers, J. D., Pandey, S., Houweling, S., Aben, I., Scarpelli, T., and Jacob, D. J.: Satellite
30 Discovery of Anomalously Large Methane Point Sources From Oil/Gas Production, *Geophysical Research Letters*, 46, 13 507–13 516, <https://doi.org/10.1029/2019gl083798>, 2019.
- Veefkind, J., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H., de Haan, J., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air
35 quality and ozone layer applications, *Remote Sensing of Environment*, 120, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
- White, W., Anderson, J., Blumenthal, D., Husar, R., Gillani, N., Husar, J., and Wilson, W.: Formation and transport of secondary air pollutants: ozone and aerosols in the St. Louis urban plume, *Science*, 194, 187–189, <https://doi.org/10.1126/science.959846>, 1976.

Wildmann, N., Päschke, E., Roiger, A., and Mallaun, C.: Towards improved turbulence estimation with Doppler wind lidar VAD scans, <https://doi.org/10.5194/amt-2020-8>, 2020.

Wood Buffalo Environmental Association: Ambient Air Monitoring Station Site Documentation - AMS 19 Firebag, available under <https://wbea.org/stations/firebag/>, 2020.

- 5 Özgen Karacan, C.: Modeling and prediction of ventilation methane emissions of U.S. longwall mines using supervised artificial neural networks, *International Journal of Coal Geology*, 73, 371–387, <https://doi.org/10.1016/j.coal.2007.09.003>, 2008.