



TROPOMI/S5P formaldehyde validation using an extensive network of ground-based FTIR stations

Corinne Vigouroux¹, Bavo Langerock¹, Carlos Augusto Bauer Aquino², Thomas Blumenstock³, Martine De Mazière¹, Isabelle De Smedt¹, Michel Grutter⁴, James Hannigan⁵, Nicholas Jones⁶, Rigel Kivi⁷, Erik Lutsch⁸, Emmanuel Mahieu⁹, Maria Makarova¹⁰, Jean-Marc Metzger¹¹, Isamu Morino¹², Isao Murata¹³, Tomoo Nagahama¹⁴, Justus Notholt¹⁵, Ivan Ortega⁵, Mathias Palm¹⁵, Gaia Pinardi¹, Amelie Röhlings³, Dan Smale¹⁶, Wolfgang Stremme⁴, Kim Strong⁸, Ralf Sussmann¹⁷, Yao Té¹⁸, Michel van Roozendael¹, Pucai Wang¹⁹, and Holger Winkler¹⁴

¹Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

²Instituto Federal de Educação, Ciência e Tecnologia de Rondônia (IFRO), Porto Velho, Brazil

³Karlsruhe Institute of Technology (KIT), Institute for Meteorology and Climate Research (IMK-ASF), Karlsruhe, Germany

⁴Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México (UNAM), Mexico City, México

⁵Atmospheric Chemistry, Observations & Modeling, National Center for Atmospheric Research (NCAR), Boulder, CO, USA

⁶Centre for Atmospheric Chemistry, University of Wollongong, Wollongong, Australia

⁷Finnish Meteorological Institute (FMI), Sodankylä, Finland

⁸Department of Physics, University of Toronto, Toronto, Canada

⁹Institut d'Astrophysique et de Géophysique, Université de Liège, Liège, Belgium

¹⁰Saint Petersburg State University, Atmospheric Physics Department, St. Petersburg, Russia

¹¹Observatoire des Sciences de l'Univers Réunion (OSU-R), UMS 3365, Université de la Réunion, Saint-Denis, France

¹²National Institute for Environmental Studies (NIES), Tsukuba, Ibaraki 305-8506, Japan

¹³Graduate School of Environment Studies, Tohoku University, Sendai 980-8578, Japan

¹⁴Institute for Space-Earth Environmental Research (ISEE), Nagoya University, Nagoya, Japan

¹⁵Institute of Environmental Physics, University of Bremen, Bremen, Germany

¹⁶National Institute of Water and Atmospheric Research Ltd (NIWA), Lauder, New Zealand

¹⁷Karlsruhe Institute of Technology, IMK-IFU, Garmisch-Partenkirchen, Germany

¹⁸LERMA-IPSL, Sorbonne Université, CNRS, Observatoire de Paris, PSL Université, 75005 Paris, France

¹⁹Institute of Atmospheric Physics, Chinese Academy of Sciences (CAS), Beijing, China

Correspondence to: C. Vigouroux
(corinne.vigouroux@aeronomie.be)

Abstract. TROPOMI (the TROPOspheric Monitoring Instrument), on-board the Sentinel-5 Precursor satellite, has been monitoring the Earth's atmosphere since October 2017, with an unprecedented horizontal resolution (initially $7 \times 3.5 \text{ km}^2$, upgraded to $5.5 \times 3.5 \text{ km}^2$ since August 2019). Monitoring air quality is one of the main objectives of TROPOMI, with the measurements of important pollutants such as nitrogen dioxide, carbon monoxide, and formaldehyde (HCHO). In this paper we assess the quality of the latest HCHO TROPOMI products (version 1.1.[5-7]), using ground-based solar-absorption FTIR (Fourier Transform Infrared) measurements of HCHO from twenty-five stations around the world, including high, mid, and low latitude sites. Most of these stations are part of the Network for the Detection of Atmospheric Composition Change (NDACC), and they provide a wide range of observation conditions from very clean remote sites to those with high HCHO levels from anthropogenic



or biogenic emissions. The ground-based HCHO retrieval settings have been optimized and harmonized at all the stations, ensuring a consistent validation among the sites.

In this validation work, we first assess the accuracy of TROPOMI HCHO tropospheric columns, using the median of the relative differences between TROPOMI and FTIR ground-based data (BIAS). We observe that, at all sites, the TROPOMI accuracy is below the upper limit of the pre-launch requirements of 80%, and below the lower limit of 40% for 20 of the 25 stations. The provided TROPOMI systematic uncertainties are well in agreement with the observed biases at most of the stations, except for the highest HCHO levels site where it is found to be underestimated. We find that, while the BIAS has no latitudinal dependence, it is dependent on the HCHO concentration levels: an overestimation ($+26\pm 5\%$) of TROPOMI is observed for very small HCHO levels ($<2.5\times 10^{15}$ molec/cm²), while an underestimation ($-30.8\%\pm 1.4\%$) is found for high HCHO levels ($>8.0\times 10^{15}$ molec/cm²). This demonstrates the great value of such a harmonized network covering a wide range of concentration levels, the sites with high HCHO concentrations being crucial for the determination of the satellite bias at the regions of emissions, and the clean sites allowing a small TROPOMI offset to be determined. The wide range of sampled HCHO levels within the network allows the robust determination of the significant constant and proportional TROPOMI HCHO biases ($\text{TROPOMI} = +1.10 (\pm 0.05) \times 10^{15} + 0.64 (\pm 0.03) \times \text{FTIR}$, in molec/cm²).

Second, the precision of TROPOMI HCHO data is estimated by the median absolute deviation (MAD) of the relative differences between TROPOMI and FTIR ground-based data. The clean sites are especially useful to minimize a possible additional collocation error. The precision requirement of 1.2×10^{16} molec/cm² for a single pixel is reached at most of the clean sites, where it is found that the TROPOMI precision can even be twice better ($0.5\text{--}0.8 \times 10^{15}$ molec/cm² for a single pixel). However, we find that the provided TROPOMI random uncertainties may be underestimated by a factor of 1.6 (for clean sites) to 2.3 (for high HCHO levels). The correlation is very good between TROPOMI and FTIR data ($R=0.88$ for 3 hours-mean coincidences; $R=0.91$ for monthly means coincidences). Using about 17 months of data (from May 2018 to September 2019), we show that the TROPOMI seasonal variability is in very good agreement at all of the FTIR sites.

The FTIR network demonstrates the very good quality of the TROPOMI HCHO products which is well within the pre-launch requirements for both accuracy and precision. This paper advises for a refinement of the TROPOMI random uncertainty budget and of the TROPOMI quality assurance values for a better filtering of the remaining outliers.

1 Introduction

TROPOMI (the TROPOspheric Monitoring Instrument), on-board the Sentinel-5 Precursor (S5P) satellite, has been monitoring the column amounts of atmospheric constituents since October 2017, at the unprecedented horizontal resolution of 7×3.5 km², upgraded to 5.5×3.5 km² since August 2019. This huge amount of data, delivered to the public and the scientific community, represents a big step to improve our knowledge of chemical and dynamical processes in the atmosphere. It is crucial to validate the quality of these new satellite data to trust and benefit their scientific exploitation. This paper focuses on the first quality assessment of the latest publicly available TROPOMI HCHO data products (v.1.1.[5-7]).



In the past, the HCHO satellite products have been validated at a few locations only, mainly using aircraft (Martin et al., 2004; Zhu et al., 2016), MAX-DOAS (Multi-AXis Differential Optical Absorption Spectroscopy) technique (Wittrock et al., 2006; De Smedt et al., 2015) and FTIR (Fourier Transform Infra-Red) technique (Jones et al., 2009; Vigouroux et al., 2009; De Smedt et al., 2015). However, given the high spatial heterogeneity of HCHO concentrations due to its short lifetime (a few hours), there is a crucial need for a more extended world coverage to assess unambiguously the satellites' achieved accuracy and precision. Furthermore, increasing the number of ground-based locations is not sufficient: it is also important to harmonize the reference data obtained at all the stations, in order to facilitate the interpretation of the satellite validation by minimizing the site-to-site biases. In this view, and in particular in the framework of the TROPOMI Cal/Val activities, we have developed HCHO retrieval settings that are suitable for any ground-based FTIR site, which have been consistently applied in Vigouroux et al. (2018) at twenty-one FTIR stations, most of them affiliated with NDACC (Network for the Detection of Atmospheric Composition Change). Vigouroux et al. (2018) described in detail the retrieval settings and the HCHO harmonized time-series obtained at these stations which cover a large range of HCHO concentrations, from very clean Arctic and oceanic sites to high HCHO levels sites, such as polluted cities (e.g. Paris or Mexico City) and sites close to large biogenic emissions like the Amazon basin (Porto Velho).

This paper presents the validation of the TROPOMI HCHO product (v.1.1.[5-7]) using an updated network of twenty-five ground-based FTIR stations. In the first section, the TROPOMI HCHO data are introduced with their uncertainty budget and their quality flag criteria. The second section describes the ground-based FTIR HCHO network and the characterization of these reference data (uncertainties and averaging kernels). Then, the validation procedure (collocation criteria, smoothing technique, definition of the quantities to be used in the quality assessment) is explained in Sect. 4. Finally, Sect. 5 shows the validation results using comparisons between TROPOMI and FTIR ground-based network data, leading to an assessment of the TROPOMI HCHO accuracy and precision, and the observed TROPOMI bias.

2 TROPOMI HCHO data

TROPOMI, on the S5P platform, is in a low-Earth afternoon polar orbit with a swath of 2600 km resulting in a daily global coverage (Veefkind et al., 2012). Operational Level 2 (L2) products include vertical columns of O₃, SO₂, NO₂, HCHO, CO and CH₄, as well as O₃ profile, aerosol layer height, cloud information and aerosol index. The spatial resolution, originally of 3.5x7 km² has been increased to 3.5x5.5 km² on 6 August 2019.

The prototype algorithm of the formaldehyde product is being developed at the Royal Belgian Institute for Space Aeronomy (BIRA-IASB) and operationally implemented by the Remote Sensing Technology Institute (IMF) of the German Aerospace Center (DLR) in the Payload Ground Data Segment (PDGS). The product has been declared operational and released to the public at the end of 2018. At the time of writing this paper, the latest product versions 1.1.[5-7] provide a consistent time series of Reprocessed+Offline (RPRO+OFFL) data, covering the period between May 2018 up to (at least) December 2019 (last access). The Near-Real-Time (NRTI) product, for the same versions 1.1.[5-7], started in December 2018. Details are



found in the Readme file ¹. The S5P HCHO retrieval algorithm is based on the DOAS method, and is directly inherited from the OMI QA4ECV product retrieval algorithm (<https://doi.org/10.18758/71021031>). It consists in a 3-step method (slant column retrieval, air mass factor calculation, and conversion to tropospheric column), fully described in De Smedt et al. (2018). The retrieval of the slant columns (N_s) is performed in the UV part of the spectra (in TROPOMI channel 3), in a fitting interval of 328.5–359 nm. The HCHO cross-section is from Meller and Moortgat (2000). All cross-sections have been pre-convolved for every row separately with an instrumental slit function adjusted just after launch. The DOAS reference spectrum is updated daily with an average of Earth radiances selected in the Equatorial Pacific region on the previous day. The result of the fit is therefore a differential slant column, showing increases over continental sources compared to the remote background. The conversion from slant to tropospheric columns (N_v) is performed using a look up table of vertically resolved air mass factors (M) calculated at 340 nm with the radiative transfer model VLIDORT v2.6 (Spurr, 2008). Parameters for each ground pixel are the observation geometry, the surface elevation and reflectivity, including the clouds (that are treated as reflecting surfaces), and a priori tropospheric profiles. The surface albedo is taken from the monthly OMI albedo climatology (minimum Lambertian equivalent reflectivity, Kleippl et al. (2008)) at the spatial resolution of $1^\circ \times 1^\circ$. A priori vertical profiles are specified using the TM5-MP daily forecast, at the same spatial resolution (Williams et al., 2017). Cloud properties are provided by the S5P operational product in its CRB mode (Cloud as Reflecting Boundary, Loyola et al. (2018)). A cloud correction based on the independent pixel approximation (Boersma et al., 2004) is applied for cloud fractions larger than 0.1. In order to correct for any remaining global offset and stripes, a background correction is applied based on HCHO slant columns from the 5 previous days in the Pacific Ocean ($N_{(s,0)}$), as described in De Smedt et al. (2018). Finally, the background vertical column of HCHO, due to methane oxidation, is taken from the TM5 model in the reference region ($N_{(v,0)}^{CTM}$). The equation of the tropospheric HCHO vertical column can be written as follows: $N_v = \frac{(N_s - N_{(s,0)})}{M} + \frac{M_0}{M} \cdot N_{(v,0)}^{CTM}$. Intermediate quantities and auxiliary data are all provided in the L2 files (http://www.tropomi.eu/sites/default/files/files/Sentinel-5P-Level-2-Product-User-Manual-Formaldehyde_v1.01.01_20180716.pdf).

Several diagnostic variables are provided together with the measurements. Quality assurance values (QA) are defined to perform a quick selection of the observations. QA > 0.5 filters out most observations presenting an error flag or a solar zenith angle (SZA) larger than 70° , a cloud radiance fraction at 340 nm larger than 0.6 or an air mass factor smaller than 0.1. The product Readme file reports that in the current version, the QA values are not always correctly set over snow/ice regions or above 75° of SZA. They also need to be further checked over cloudy scenes. In the forthcoming S5P version 2, QA values will be refined, and will exclude data with surface albedo larger than 0.2 and snow/ice warning, and remaining SZA larger than 75° .

The tropospheric column uncertainty is divided into random (precision) and systematic components (accuracy), and is provided per pixel. It varies with the observation conditions. Over remote regions at moderate solar zenith angle, the precision of an individual observation is about 5×10^{15} molec/cm². This value agrees with the standard deviation of the columns in the same region for a particular day. The random uncertainty is dominated by the random error on the slant columns. The tropospheric column accuracy is the combined systematic uncertainty resulting from the slant column, the air mass factor and the back-

¹<http://www.tropomi.eu/sites/default/files/files/publicSentinel-5P-Formaldehyde-Readme.pdf>; doi: 10.5270/S5P-tjlxfd2




Figure 1. Network of ground-based FTIR stations providing HCHO total columns data. The background is the September 2018 monthly mean of TROPOMI HCHO tropospheric columns.

ground correction errors. It varies between 30 and 60% of the columns. The column averaging kernel and the a priori profiles are provided for every observation.

3 Ground-based FTIR HCHO data

We show in Fig. 1 a map of the ground-based FTIR stations used in this TROPOMI validation. The background image represents the global TROPOMI monthly mean tropospheric columns for September 2018, illustrating the different HCHO levels sampled by the ground-based network: from clean Arctic and oceanic sites to very high-concentrations sites such as Porto Velho, in the Amazon basin.

Table 1 lists the ground-based FTIR stations, their coordinates and altitude, the spectrometer type, the retrieval code, and the team involved in the measurements and/or the retrievals of HCHO. For more details on the monitoring of FTIR solar absorption spectra at these stations, we refer to Vigouroux et al. (2018) and references therein, and, for the FTIR retrieval principles, to e.g. Vigouroux et al. (2009).

The same retrieval settings are used at all the stations to avoid introducing possible bias in the HCHO total columns between the stations and inconsistent comparisons with the satellite. Details are given in Vigouroux et al. (2018). The main settings that might be responsible for internal biases within the network are the spectroscopic database and the fitted spectral windows, the spectroscopic parameters being the main source of the FTIR HCHO systematic uncertainties. The HCHO spectral signatures



Table 1. FTIR stations that are contributing to the present work: location and altitude (in km a.s.l.), instrument type, retrieval code, team.

Station	Latitude	Longitude	Altitude	Instrument	Code	Team
Eureka	80.05° N	86.42° W	0.61	Bruker 125 HR	SFIT4	U. of Toronto
Ny-Ålesund	78.92° N	11.92° E	0.02	Bruker 120 HR	SFIT4	U. of Bremen
Thule	76.52° N	68.77° W	0.22	Bruker 120 M	SFIT4	NCAR
Kiruna	67.84° N	20.40° E	0.42	Bruker 120/5 HR	PROFFIT	KIT-ASF ; IRF Kiruna
Sodankylä	67.37° N	26.63° E	0.19	Bruker 125 HR	SFIT4	FMI ; BIRA
St-Petersburg	59.88° N	29.83° E	0.02	Bruker 125 HR	SFIT4	SPbU
Bremen	53.10° N	8.85° E	0.03	Bruker 125 HR	SFIT4	U. of Bremen
Paris	48.97° N	2.37° E	0.06	Bruker 125 HR	PROFFIT	Sorbonne U.
Zugspitze	47.42° N	10.98° E	2.96	Bruker 120/5 HR	PROFFIT	KIT-IFU
Jungfraujoch	46.55° N	7.98° E	3.58	Bruker 120 HR	SFIT4	U. of Liège
Toronto	43.60° N	79.36° W	0.17	Bomem DA8	SFIT4	U. of Toronto
Rikubetsu	43.46° N	143.77° E	0.38	Bruker 120/5 HR	SFIT4	Nagoya U. ; NIES
Boulder	40.04° N	105.24° W	1.61	Bruker 120 HR	SFIT4	NCAR
Xianghe	39.75° N	116.96° E	0.05	Bruker 125 HR	SFIT4	CAS ; BIRA
Tsukuba	36.05° N	140.12° E	0.03	Bruker 125 HR	SFIT4	NIES ; Tohoku U.
Izaña	28.30° N	16.48° W	2.37	Bruker 120/5 HR	PROFFIT	AEMET ; KIT-ASF
Mauna Loa	19.54° N	155.57° W	3.40	Bruker 120/5 M	SFIT4	NCAR
Mexico City	19.33° N	99.18° W	2.26	Bruker Vertex 80	PROFFIT	UNAM
Altzomoni	19.12° N	98.66° W	3.98	Bruker 120/5 HR	PROFFIT	UNAM
Palau	7.34° N	134.47° E	0.03	Bruker 120/5 M	SFIT4	U. of Bremen
Paramaribo	5.81° N	55.21° W	0.03	Bruker 120/5 M	SFIT4	U. of Bremen
Porto Velho	8.77° S	63.87° W	0.09	Bruker 125 M	SFIT4	BIRA
Maïdo	21.08° S	55.38° E	2.16	Bruker 125 HR	SFIT4	BIRA
Wollongong	34.41° S	150.88° E	0.03	Bruker 125 HR	SFIT4	U. of Wollongong
Lauder	45.04° S	169.68° E	0.37	Bruker 120 HR	SFIT4	NIWA

lie in the 3.6 μm region and belong to the ν_1 and ν_5 bands (fitted windows are: 2763.42 - 2764.17; 2765.65 - 2766.01 ; 2778.15 - 2779.1 ; 2780.65 - 2782.0, in cm^{-1}). The spectroscopic database used is the atm16 linelist by G. Toon (JPL), which can be found at <http://mark4sun.jpl.nasa.gov/toon/linelist/linelist.html>. This linelist is optimized for the main absorbing gases in the fitted windows (HDO, CH_4 , O_3 , N_2O , CO_2) and is based on HITRAN 2012 (Rothman et al., 2013) for HCHO, which used the work of Jacquemart et al. (2010).

The retrieval codes used in the FTIR NDACC community are PROFITT9 (Hase et al., 2006) and SFIT4.0.9.4 (updated from SFIT2 (Pougatchev et al., 1995)), which are both based on the optimal estimation method (Rodgers, 2000). Based on a priori




Figure 2. Left panels: typical total column averaging kernel (AK) from FTIR (blue) and TROPOMI (red) measurements at the Maïdo station (altitude: 2.2 km). Right panels: a priori profile used in the FTIR retrievals for the whole time series at Maïdo (blue), and an example of TROPOMI a priori profile at Maïdo (the 18th January 2019).

profile information (from the model WACCM, Garcia et al. (2007)), and a L1 Tikhonov regularization matrix (Tikhonov, 1963), low vertical resolution profiles can be retrieved in principle, as well as total columns. However, as described in Vigouroux et al. (2018), the degrees of freedom for signal are very low for HCHO (median value of 1.1 for all FTIR sites), meaning that we essentially have one piece of information. The FTIR total column averaging kernel shows a decrease of the sensitivity at the surface, which is quite similar to the TROPOMI sensitivity. This can be seen in Fig. 2, as an example for the Maïdo station. We also show in Fig. 2 the FTIR a priori profile at Maïdo, which is based on a climatology (1980-2020) from the WACCM model calculated at Maïdo. A single profile is used for the whole time series at a specific station (Vigouroux et al., 2018), while TROPOMI uses daily a priori profiles from TM5 (Sect. 2). An example is shown in Fig. 2 for the 18th January 2019.

The FTIR uncertainty budget is calculated following the formalism of Rodgers (2000) and is described in Vigouroux et al. (2018). It is separated into random and systematic components. The random uncertainty is dominated at all sites by the measurement noise uncertainty, which can vary from site to site depending on the spectrometer. The uncertainty on the retrieved FTIR total columns for individual sites is given in Vigouroux et al. (2018) for the 21 sites involved at that time. We obtain a median random uncertainty of 2.3×10^{14} molec/cm², with a large value of 11.1×10^{14} molec/cm² only at Mexico City where a lower resolution instrument is used (Vertex 80). The smoothing uncertainty on the total column has a non negligible random component (median value of 1.2×10^{14} molec/cm²). With the inclusion of the smoothing error in the uncertainty budget, the median total random uncertainty is 2.9×10^{14} molec/cm², which is very close to our empirical uncertainty estimation of 2.8×10^{14} molec/cm², based on the standard deviation of the differences between two FTIR individual subsequent measurements (within a maximum of 30 minutes interval), confirming our theoretical uncertainty calculation. Since the Vigouroux et al. (2018) paper, five more sites have joined the HCHO harmonized network. The mean random errors on individual FTIR measurements are (in molec/cm²): 1.4×10^{14} , 2.7×10^{14} , 2.2×10^{14} , 5.2×10^{14} , and 5.6×10^{14} , for Jungfraujoch, Rikubetsu, Tsukuba, Palau, and Xianghe, respectively.

The forward model parameters median systematic uncertainty on the HCHO FTIR total columns is 13% in the network of Vigouroux et al. (2018). As already mentioned, the dominating systematic uncertainty sources are the spectroscopic parameters: the line intensities and the pressure broadening coefficients of the fitted HCHO absorption lines. We use 10% for the three parameters: the line intensity, and the air- and self- broadening coefficients. The systematic uncertainty can be larger



at the stations (up to 21-26%) using the PROFFIT9 retrieval code, due to an uncertainty on the channeling that is not taken into account yet in the SFIT4 code. The median smoothing systematic uncertainty is 3.4%. For the 5 added sites, the median systematic uncertainty is 13% (Jungfraujoch, Tsukuba, Palau), or 14% (Rikubetsu, Xiangue).

4 Validation method

5 4.1 Collocation criteria

The precision of a single pixel TROPOMI HCHO measurement is expected to be below 1.2×10^{16} molec/cm² (pre-launch requirements) or even better, as 5×10^{15} molec/cm² for remote areas (after launch uncertainty analysis, see Sect. 2). These values are quite large compared to the measured levels of HCHO (around 1.5×10^{15} molec/cm² for very clean sites to e.g. around 9×10^{15} molec/cm² for a city such Paris). It is therefore necessary to average several pixels in order to reduce the random uncertainty of the TROPOMI mean HCHO data, improve the detection level and increase the TROPOMI sensitivity to day-to-day variability. For this reason, we choose to average the TROPOMI pixels located within 20 km from the FTIR station. Once we filter out the TROPOMI pixels that do not reach the recommended quality criteria (QA flag > 0.5; see Sect. 2), we obtain a median value of 34 pixels to average. In cloudy conditions, this number can be smaller. A collocation pair is kept when at least 10 pixels can be averaged. Higher number of pixels can be averaged for Arctic stations (around 45-60), which is useful due to the very low HCHO levels to be detected there. At sub-tropical / tropical stations, the median number of pixels is around 20-29. The higher number of pixels in Arctic is due to the fact that each FTIR measurement is co-located to all S5P pixels that match the co-location criteria, even if these pixels originate from different orbits, with different overpass times.

The time coincidence criterion is set to ± 3 hours. This choice is a compromise to obtain significant number of coincidences between TROPOMI and FTIR data, noting that the median FTIR measurement frequency is 5 per day (with a range of 3 to 10 depending on the station). A shorter time coincidence criterion decreases significantly the numbers of sampled collocated days and sometimes months, which is a limitation for checking the TROPOMI seasonality (sampled months: 267 for 1 h, and 305 for 3 h criteria). Note that a 6 h criterion would provide 20 additional sampled months: the critical stations are Mauna Loa, Alzomoni and Paramaribo, for which we would have coincidences back to May 2018. The standard deviations of the TROPOMI / FTIR comparisons are usually smaller with a longer time coincidence criterion, but this can be explained by the increased number of pixels (so improved TROPOMI precision on the mean) in the 6 h collocation, mainly at Arctic sites with increased number of multiple orbits. Despite the smaller standard deviations usually obtained within a 6 h criterion, we finally choose 3 h to reduce the possible impact of some passing plumes and of the HCHO diurnal cycle on the comparisons. The diurnal cycle at most of the FTIR stations can be found in Vigouroux et al. (2018) and its Supplement. At many stations no significant diurnal cycle was observed but, in some cases, mainly polluted sites, we obtained a maximum around noon-1 p.m., close to the TROPOMI overpass time. At the Mexico City station, where the diurnal cycle amplitude is the greatest, the effect of collocation time (6 h vs 3 h) on the statistical bias is 4%.



4.2 Building inter-comparable products

Some manipulation of the original data products is needed before looking at the differences between TROPOMI and FTIR data. Both measurements provide total columns (for FTIR) or tropospheric columns (for TROPOMI) that have a lower sensitivity near the ground (see Fig. 2), and their retrievals use a priori profile information that have been chosen differently (TROPOMI: 5 daily a priori profiles from TM5; FTIR: single a priori profile from climatology of WACCM). To correct for this, for each S5P individual pixel collocated with each FTIR measurement, we use the comparison method described in Rodgers and Connor (2003). First, the a priori substitution is applied, using the S5P a priori profile as the common a priori profile. For this, the S5P a priori profile is regridded to the FTIR retrieval grid using a mass conservation algorithm (Langerock et al., 2015). In the rare situation where the satellite pixel elevation is above the FTIR site, the S5P a priori profile is extended to the FTIR 10 instrument's altitude. The regridded S5P a priori $\mathbf{x}_{S,a}$ is then substituted following Rodgers and Connor (2003), and we finally use the corrected FTIR retrieved profile \mathbf{x}'_F in the comparisons:

$$\mathbf{x}'_F = \mathbf{x}_F + (\mathbf{A}_F - \mathbf{I})(\mathbf{x}_{F,a} - \mathbf{x}_{S,a}), \quad (1)$$

where \mathbf{x}_F is the original FTIR retrieved profile, \mathbf{A}_F is the FTIR averaging kernel matrix, \mathbf{I} is the unit matrix, and $\mathbf{x}_{F,a}$ is the FTIR a priori profile.

15 The next step, following Rodgers and Connor (2003), is to smooth the corrected FTIR profile with the S5P column averaging kernel \mathbf{a}_S . For that purpose we regrid the corrected FTIR profile \mathbf{x}'_F to the S5P column averaging kernel grid and apply the smoothing equation:

$$c_F^{smoo} = c_{S,a} + \mathbf{a}_S(\mathbf{x}'_F - \mathbf{x}_{S,a}) \quad (2)$$

with $c_{S,a}$ the S5P a priori column derived from the S5P a priori profile. We obtain a smoothed FTIR column c_F^{smoo} associated 20 with a collocated TROPOMI pixel. In the case of mountain sites where the pixel altitude is below the instrument's height, the regridding of the FTIR profile \mathbf{x}'_F is done such that the FTIR profile is extended with the S5P a priori profile (such an extension is invariant under the latter smoothing equation). Note that this FTIR regridding to the satellite grid has also the advantage that only the FTIR profile up to the altitude of the satellite product (which is only a tropospheric column) remains in the regridded column: we therefore finally compare tropospheric columns in both products.

25 Next, we need to take into account that, for mountain stations, the difference between satellite columns and the original ground-based columns can be significant. To bring both satellite and smoothed FTIR column c_F^{smoo} (which is calculated as a column valid at the satellite's pixel surface) values to the scale of the original FTIR columns, we apply a scaling factor f representative for the fraction of the partial column between the satellite pixel altitude and the FTIR station altitude. This scaling factor is derived from the satellite a priori profile and is defined as:

$$30 \quad f = 1 - \frac{c_{S,a}^{\Delta z}}{c_{S,a}}, \quad (3)$$



where $c_{S,a}^{\Delta z}$ denotes the partial column derived from the S5P a priori profile between the pixel surface and the FTIR station. The TROPOMI column c_S and its random and systematic uncertainties are also scaled with the same factor, so that finally the collocated products are all expressed at the altitude of the FTIR site (and not of the pixel surface). For mountain stations, the scaling factor f , calculated for each satellite's pixel, can reach a minimum of 0.5 for stations located at about 2 km altitude
 5 from the satellite's pixel surface (Maïdo, Izaña, or Altzomoni), or even 0.3 at the higher sites Jungfraujoch and Zugspitze, while at sea-level sites it is of course close to 1.0. In the rare cases where the satellite pixel is above the FTIR station, we apply the conversion factor $f = 1 + c_{S,a}^{\Delta z}/c_{S,a}$, where the satellite a priori profile is extrapolated to the station surface in order to calculate the partial column of the a priori between both altitudes.

The final step is to average the individual smoothed and scaled FTIR columns $c_F^{smoo} \times f$ that are taken within 3 h, and the
 10 TROPOMI $c_S \times f$ individual pixel columns that are available within 20 km (which can belong to different orbits), to form the collocated pair FTIR_{*i*} and TROP_{*i*} used in the next section.

4.3 Estimation of the TROPOMI accuracy and precision

In Sect. 5.1, we assess whether the TROPOMI accuracy is compliant with pre-launch requirements (40-80%, as reported in the ESA official document S5P-RS-ESA-SY-164, 2014, Table 3, p. 19). The accuracy of the TROPOMI HCHO measurements
 15 will be estimated by deriving the median of the relative differences (BIAS) between the collocated TROP_{*i*} and the reference FTIR_{*i*} data at each station:

$$\text{BIAS} = \text{med}\left(\frac{\text{TROP}_i - \text{FTIR}_i}{\text{FTIR}_i}\right). \quad (4)$$

We can note that the applied scaling factor f (see previous section) does not affect the BIAS estimation, even at high mountains stations, because it cancels in the division.

20 For robust statistics, the median is preferred to the mean due to the presence of outliers (a few remaining TROPOMI outliers after the QA filter, and some very small FTIR values that give very large relative difference after the division in Eq. 4). The presence of TROPOMI outliers is minimized by using the median, but they should be ideally removed by the QA filter. An improvement of the QA value is foreseen in the next product version, which should improve, e.g., the filtering at Arctic sites ($\text{SZA} > 75^\circ$).

25 In the next section, we also compare the obtained BIAS with the systematic uncertainty on the difference σ_{sys} , to evaluate the TROPOMI uncertainty budget:

$$\sigma_{\text{sys}}^2 = (\sigma_{S,\text{sys}})^2 + \mathbf{a}_S^T \mathbf{S}_{F,\text{sys}} \mathbf{a}_S + \mathbf{a}_S^T (\mathbf{I} - \mathbf{A}_F) \mathbf{S}_{\text{var},\text{sys}} (\mathbf{I} - \mathbf{A}_F)^T \mathbf{a}_S, \quad (5)$$

where $\sigma_{S,\text{sys}}$ is the systematic uncertainty of TROPOMI columns, as provided in the public release database (but scaled for altitude, see Sect. 4.2), \mathbf{a}_S is the TROPOMI total column averaging kernel, and $\mathbf{S}_{F,\text{sys}}$ is the FTIR systematic covariance matrix
 30 provided in vmr^2 in the standardized GEOMS format converted in partial columns units. The last term is the impact of different low vertical resolution profile measurements (the smoothing error) on the comparisons (see Eq. 27 in Rodgers and Connor (2003)), where for the systematic uncertainty part, we account for possible bias on $x_{S,a}$ by following von Clarmann (2014):



$$\mathbf{S}_{\text{var,syst}} = (\mathbf{x}_{S,a} - \langle \mathbf{x} \rangle)(\mathbf{x}_{S,a} - \langle \mathbf{x} \rangle)^T.$$

The $\mathbf{x}_{S,a} - \langle \mathbf{x} \rangle$ is not known and we follow Vigouroux et al. (2018), with $\mathbf{x}_{S,a} - \langle \mathbf{x} \rangle = -50\%$, -20% , -10% , $+10\%$, $+8\%$, $+5\%$ for the ground-4 km; 4-8 km; 8-13 km; 13-25 km; 25-40 km; 40-120 km layers, respectively (expressed in molec/cm²).

The last term of Eq. 5 is found to be of the order of a few percent, therefore negligible in σ_{syst} . In practice, the systematic uncertainty on the difference σ_{syst} is dominated by the TROPOMI systematic uncertainty of about 40%, FTIR having a median systematic uncertainty of only 13% with a maximum of 26% (See Sect. 3).

Similarly, the precision of the TROPOMI HCHO products is estimated in Sect. 5.2, not with the usual standard deviation which is not robust in case of outliers, but with the median absolute deviation (MAD, see Huber (1981)) of the differences ($\text{DIFF}_i = \text{TROP}_i - \text{FTIR}_i$):

$$\text{MAD} = k \times \text{med}(\text{abs}(\text{DIFF}_i - \text{med}(\text{DIFF}_i))), \quad (6)$$

where $k = 1.4826$ for a correspondence with the 1- σ standard deviation for normal distribution without outliers.

In Sect. 5.2, we compare the obtained MAD to the random uncertainty on the differences σ_{rand} , which is calculated following Rodgers and Connor (2003):

$$\sigma_{\text{rand}}^2 = (\sigma_{S,\text{rand}})^2 + \mathbf{a}_S^T \mathbf{S}_{S,\text{rand}} \mathbf{a}_S + \mathbf{a}_S^T (\mathbf{I} - \mathbf{A}_F) \mathbf{S}_{\text{var,rand}} (\mathbf{I} - \mathbf{A}_F)^T \mathbf{a}_S, \quad (7)$$

where $\mathbf{S}_{\text{var,rand}}$, to take into account the impact of low vertical resolution in the random part of the uncertainty, is the natural variability matrix chosen to be 50%, 50%, 40%, 35%, 30%, 30%, 10% for the ground-4 km; 4-8 km; 8-13 km; 13-25 km; 25-40 km; 40-120 km layers, respectively (expressed in molec/cm²). As for the systematic uncertainty part, the random uncertainty on the difference is dominated by the TROPOMI random uncertainty (median of about 1.1×10^{15} molec/cm² for TROP_{*i*} within 20 km), while FTIR_{*i*} has a median random uncertainty of 2.0×10^{14} molec/cm². The last term of Eq. 7 is comparable to the FTIR one (median value of 2.4×10^{14} molec/cm²).

We can use MAD as an upper limit of the TROPOMI precision, since collocation in space and time of the sounded air-masses are never perfect. It is compared in the next section to the pre-launch precision requirement. The MAD estimation is influenced by the scaling factor f , which is important only for high altitude sites (Sect. 4.2). It should be interpreted as an estimation of the precision of a TROPOMI column that would be measured at the altitude of the FTIR site. The random uncertainty on the differences are also expressed at the altitude of the FTIR site, so that the comparison between MAD and σ_{rand} is always valid.

The observed BIAS between TROPOMI and the reference FTIR data is statistically significant if it exceeds its statistical error: $\text{ERR}_B = 2 \times \text{MAD} / \sqrt{n}$ (with n the number of coincidences).

5 Validation results

In this section, we provide a table and plots for the offline (RPRO+OFFL) HCHO TROPOMI product. We do not show detailed results for the near real time (NRTI) product (version 1.1.[5-7]) because they are very similar to the offline version. Numbers for the main conclusions will be given in the text for this NRTI product.



5.1 TROPOMI observed BIAS and accuracy

In Table 2, we provide, at each individual FTIR station, the mean of the FTIR HCHO total columns (mean FTIR), the obtained median of the relative differences BIAS (in % to compare with the pre-launch TROPOMI accuracy requirements of 40-80%, Eq. 4), the error on the BIAS (ERR_B), and the number of collocated pairs n . The systematic uncertainty on a single difference is also given (in %, calculated from Eq. 5 where each term has been expressed in %, dividing by each instrument individual HCHO column).

We have ordered the stations, not in decreasing latitudes as in Table 1, but in increasing mean HCHO FTIR columns. The reason is that we observe a tendency of the BIAS between TROPOMI and FTIR: while the BIAS is always (with the exception of Eureka) positive or not significant (if $BIAS < ERR_B$) for very clean to clean sites with HCHO mean levels lower than 6.5×10^{15} molec/cm², it is negative for higher levels and very consistent for the stations from 8.7 to 28.6×10^{15} molec/cm² (-29 to -36 %) with small error on the bias (2 to 6 %). Note that the BIAS is also consistent at Paramaribo (-26%) but with larger error (14%), due to small number of collocations. This dependence of the TROPOMI bias on the HCHO concentration levels can be visualized in Fig. 3, where the BIAS at each station is plotted as a function of the mean FTIR columns. It is therefore not appropriate to use the median bias obtained using the data from all stations together (-10%), if one wants to correct the

TROPOMI HCHO data in model inversion studies. If we calculate the median of the differences for HCHO FTIR columns $> 8.0 \times 10^{15}$ molec/cm², we obtain a significant negative bias of $-30.8 \pm 1.4\%$. The detection of this bias is especially important for modeling studies that use satellite data to optimize the volatile organic compound emissions sources, as done in e.g. Fortems-Cheiney et al. (2012); Stavrakou et al. (2015) with OMI and GOME-2. The bias for clean HCHO levels ($< 2.5 \times 10^{15}$ molec/cm²) is significantly positive ($+26 \pm 5\%$).

The validation results for the NRTI TROPOMI products give very similar results: a negative BIAS ($-31.7 \pm 1.8\%$) for the high HCHO levels ($> 8.0 \times 10^{15}$ molec/cm²) and a positive one ($+22 \pm 7\%$) for low HCHO levels ($< 2.5 \times 10^{15}$ molec/cm²). The small differences are mainly due to the different sampling of the comparisons (NRTI data are since December 2018, while the OFFL data are since May 2018).

The different TROPOMI BIAS at different HCHO levels is pointing to the presence of two kinds of bias: a constant one and a proportional one. They can be obtained by using the scatter plot of the two instruments, shown in Fig. 4: the constant bias is the intercept of the linear relationship between TROPOMI and FTIR, while the proportional bias is given by its slope. But this has to be done carefully: a usual linear regression by ordinary least squares (OLS) is not statistically robust and can give spurious results in the presence of outliers and/or heteroscedasticity. We are confronted to both problems in our scatter plot: we do have outliers and the uncertainty is increasing with HCHO levels. Therefore, we use the robust Theil-Sen estimator (Sen, 1968) where the slope s of the scatter plot is the median of the slopes of the lines through all pairs of data points ($TROP_j - TROP_i$)/($FTIR_j - FTIR_i$), with $FTIR_j \neq FTIR_i$. The intercept b is then the median of ($TROP_i - s \times FTIR_i$). Using this robust estimator, we obtain the relation: $TROP = 0.64 \times FTIR + 1.10 \times 10^{15}$ molec/cm². We have calculated the uncertainties in s and b using $2 \times MAD / \sqrt{n}$, with MAD the median absolute deviation of the slopes and intercepts of the pairs of data points, and n



Table 2. Validation of TROPOMI RPRO+OFFL. Please note that the ordering of the sites is by increasing mean HCHO column. For each station: mean of the HCHO FTIR total columns (in molec/cm²), median of the relative differences $BIAS = med((TROP_i - FTIR_i)/FTIR_i)$ and its error ERR_B (in %, see text), number of collocated pairs n , systematic uncertainty on a single difference σ_{syst} (in %, Eq. 5), median absolute deviation (MAD, in molec/cm², Eq. 6), random uncertainty on a single difference σ_{rand} (in molec/cm², Eq. 7), and pre-launch TROPOMI precision requirements associated to the choice of 20 km around the station $Requ = 1.2 \times 10^{16} / \sqrt{n_{pix}}$ molec/cm², with n_{pix} the mean number of pixels averaged in the collocated TROPOMI data. The Pearson correlation coefficient R is given for individual coincidences (± 3 h) and for monthly means of coincident data.

Station	mean FTIR molec/cm ²	BIAS $\pm ERR_B$ %	n	σ_{syst} %	MAD molec/cm ²	σ_{rand} molec/cm ²	Requ. molec/cm ²	n_{pix}	R indiv.	R monthly
Jungfraujoch	1.24E+15	19 ± 15	87	58	9.0E14	5.6E14	2.5E15	24	0.61	0.70
Zugspitze	1.36E+15	52 ± 10	184	59	7.8E14	5.0E14	2.1E15	33	0.71	0.86
Mauna Loa	1.60E+15	52 ± 22	52	54	1.3E15	8.8E14	2.5E15	23	-0.09	-0.05
Eureka	1.65E+15	-40 ± 11	114	97	1.1E15	5.3E14	1.8E15	45	0.22	0.43
Maïdo	1.86E+15	5 ± 9	155	43	1.0E15	7.1E14	3.0E15	16	0.45	0.53
Ny-Ålesund	1.90E+15	43 ± 20	47	41	1.2E15	4.9E14	1.7E15	52	0.35	0.38
Thule	2.06E+15	-3 ± 5	346	57	9.6E14	4.8E14	1.6E15	60	0.56	0.82
Izaña	2.07E+15	13 ± 10	97	83	8.9E14	6.4E14	2.5E15	24	0.47	0.79
Altzomoni	2.44E+15	66 ± 18	67	42	1.6E15	8.9E14	2.6E15	22	0.50	0.86
Kiruna	2.44E+15	50 ± 12	146	67	1.5E15	8.8E14	1.7E15	60	0.64	0.72
Lauder	2.54E+15	-11 ± 14	225	78	2.6E15	1.3E15	2.1E15	33	0.38	0.65
Rikubetsu	3.16E+15	26 ± 40	16	50	2.8E15	1.0E15	1.9E15	41	0.45	0.60
Palau	3.80E+15	0 ± 15	10	36	9.8E14	8.2E14	2.7E15	20	0.15	0.33
Sodankyla	4.15E+15	8 ± 7	307	51	2.5E15	1.2E15	1.7E15	48	0.51	0.69
Boulder	5.91E+15	-1 ± 9	103	50	2.2E15	1.3E15	2.2E15	31	0.79	0.90
St-Petersburg	6.21E+15	-4 ± 8	158	44	3.0E15	1.2E15	1.9E15	42	0.68	0.78
Wollongong	6.36E+15	9 ± 8	322	54	3.3E15	1.9E15	2.3E15	27	0.78	0.94
Tsukuba	7.05E+15	-23 ± 12	34	44	3.1E15	1.2E15	2.2E15	31	0.68	0.51
Bremen	7.77E+15	-5 ± 12	46	39	3.2E15	1.4E15	1.8E15	43	0.63	0.68
Paramaribo	8.43E+15	-26 ± 14	15	36	3.3E15	1.3E15	2.5E15	23	0.12	0.14
Paris	8.72E+15	-29 ± 6	128	44	3.1E15	1.2E15	1.9E15	41	0.76	0.79
Toronto	1.06E+16	-34 ± 4	251	38	3.7E15	1.6E15	2.3E15	29	0.73	0.95
Xianghe	1.43E+16	-33 ± 2	384	38	5.5E15	2.1E15	2.0E15	36	0.86	0.97
Mexico City	1.92E+16	-27 ± 4	154	32	5.9E15	2.6E15	2.5E15	25	0.34	0.27
Porto Velho	2.86E+16	-36 ± 3	81	31	8.3E15	3.6E15	2.2E15	29	0.81	1.00
All stations BIAS	6.60E+15	-10 ± 2	3529	48	2.4E15	1.2E15	2.1E15	34	0.87	0.91
Low HCHO BIAS FTIR < 2.5 × 10 ¹⁵	1.64E+15	+26 ± 5	1321	52	1.3E15	7.7E14	2.2E15	31	0.40	
High HCHO BIAS FTIR > 8.0 × 10 ¹⁵	1.57E+16	-30.8 ± 1.4	952	46	4.8E15	2.1E15	2.1E15	33	0.88	



the numbers of pairs. We obtain an uncertainty of 0.03 and 0.05×10^{15} molec/cm² for s and b respectively. Therefore, both the constant ($1.10 \pm 0.05 \times 10^{15}$ molec/cm²) and proportional ($0.64 \pm 0.03\%$) biases are significant.

Using the scatter plot to derive the constant and proportional biases is very sensitive to the range of observed values. As an example, if one would only use HCHO FTIR data $> 8.5 \times 10^{15}$ molec/cm², one would obtain a slope of 0.51 and an intercept of 3.2×10^{15} molec/cm², which would point to a strong overestimation and underestimation of the constant and proportional biases, respectively. This would lead to erroneous prediction of TROPOMI overall bias outside the range of observed values. This demonstrates why such a ground-based network, covering very clean sites to high HCHO level sites, is crucial to provide a good estimate of both constant and proportional biases of TROPOMI.




Figure 3. BIAS at each station (in %) as a function of the mean FTIR total columns (molec/cm²). The gray bars are the systematic uncertainty on the differences σ_{sys} , and the colored error bars are the $2\text{-}\sigma$ error on the bias ERR_B . Black markers are for mountains stations, blue for Arctic stations, cyan for Oceania/Australia/New Zealand, magenta for China/Japan, red for mid-latitude Europe/North America, and green for Central/South America.

The BIAS given in Table 2 are a combination of the constant and proportional biases, and can be used to statistically assess the TROPOMI HCHO overall accuracy. We can easily see from Table 2 that all BIAS values are within the upper limit of the pre-launch requirement of 80%, and they are within the 40% requirement lower limit for 20 of the 25 stations. The five stations exceeding a 40% BIAS are clean (Arctic or mountains) sites, with mean HCHO columns below 2.5×10^{15} molec/cm². But these are sites where the systematic uncertainty on the differences (see Table 2 and Eq. 5) are usually also the largest, leading to a good correspondence between observed higher BIAS and higher calculated uncertainty for 3 of these 5 stations (Zugspitze, Mauna Loa, and Kiruna).

Therefore, we can conclude that the TROPOMI HCHO accuracy satisfies the pre-launch requirements and that the system-




Figure 4. Scatter plots of TROPOMI versus FTIR data, for individual collocated pairs (± 3 h; left panel), and for the monthly means of collocated data (right panel). The non robust OLS fit of the data is given in the legend and plotted as a black line, while the slope and intercept obtained with robust Theil-Sen estimator is given by the red line and text.

atic uncertainty budget is in very good agreement with observed bias, except at a very few stations (Ny-Ålesund $43 > 41\%$, Altzomoni $71 > 42\%$, and Porto Velho $36 > 31\%$). At most of the other stations, the reported systematic uncertainty tends to be larger than the BIAS. We find the same conclusions on TROPOMI accuracy when making comparisons with the NRTI products.

5 5.2 Observed TROPOMI precision

For discussing the observed TROPOMI precision, we provide in Table 2, the MAD for each station (in absolute value to compare with the pre-launch precision requirement of 1.2×10^{16} molec/cm² for a single pixel, Eq. 6). The precision pre-launch requirement is provided at each site taking into account the mean number of pixels n_{pix} involved in the collocated TROPOMI data (Requ. = 1.2×10^{16} molec/cm² / $\sqrt{n_{pix}}$). We see that for all the cleanest sites ($< 2.5 \times 10^{15}$ molec/cm²), where an additional collocation uncertainty is expected to be small, the MAD is well within the pre-launch requirements. The MAD for these cleanest sites has a median of 1.3×10^{15} molec/cm², and a minimum of 0.9×10^{15} molec/cm². This is a good estimate of the precision that TROPOMI can reach in remote conditions. For a single pixel, the TROPOMI best precision at remote conditions is therefore $5-8 \times 10^{15}$ molec/cm².



It must be noted that the pre-launch HCHO precision requirements were chosen based on pre-launch requirements for the instrument signal to noise ratio (equivalent to OMI). The actual signal to noise of the measurements appears to be better than the requirements, especially in the HCHO wavelength fitting range. Furthermore, the good quality of the recorded spectra allowed to increase the size of the TROPOMI HCHO fitting spectral interval just after launch, further improving the precision of the slant columns. Indeed, as seen in Table 2, only at the three highest HCHO levels sites (Xianghe, Mexico City, and Porto Velho) the provided random uncertainties are as high as the pre-launch requirements. The actual provided random uncertainty are smaller, and we can see that, even for clean sites, the observed MAD is larger than the random uncertainty on the differences by a factor of 1.6. This factor increases up to 1.8 if we take into account all the stations, but this is expected due to a collocation uncertainty that should have more impact at high-levels sites (the factor rises up to 2.3 for high HCHO levels $>8.0 \times 10^{15}$ molec/cm²). Our comparisons suggest that the TROPOMI random uncertainty is underestimated by at least a factor of 1.6 and up to maximum of 2.3 (if one would assume the collocation uncertainty to be smaller than the TROPOMI uncertainty). This underestimation could be due to the fact that currently the uncertainties associated to the air mass factor calculation and to the background correction step are assumed to be fully systematic. The discrimination between random and systematic part of the uncertainties might be refined in the future, based on such validation results.

5.3 Observed TROPOMI monthly variability

The Pearson correlation coefficient is very good for the collocated monthly means of TROPOMI and FTIR data (0.91, see Table 2 and Fig. 4), and is usually good for individual sites. However, Pearson correlation is not robust and can give a wrong conclusion when only few data are coincident, especially when outliers are present. We have 17 months of coincident TROPOMI and FTIR measurements in the best cases, while only 4 for the newest stations Palau and Porto Velho. We therefore verify that the TROPOMI precision allows the seasonal variability to be well captured, even at very clean sites which can be at the limit of the satellite detection, by plotting the individual monthly mean time-series in Fig. 5.

The seasonal variability, with a maximum in July-August, is well observed at all the Arctic sites (Eureka, Ny-Ålesund, Thule, Kiruna, and Sodankylä). The monthly mean correlation is better than 0.69, except at Eureka and Ny-Ålesund. It can be seen in Fig. 5 that Sept. 2019 is very high in TROPOMI data at Ny-Ålesund, and only 1 coincidence is found for this month. Removing this last outlier gives a 0.76 correlation coefficient at this station. The northern mid-latitude clean sites (mountains: Jungfrauoch, Zugspitze, Izaña) also display a seasonal variability in very good agreement, with correlation coefficients higher than 0.70. The Japanese clean site Rikubetsu shows poorer correlation (0.60) but only few data are in coincidence. The stations where we find the poorer correlations are the oceanic sites. The poorest one is Mauna Loa, but this is expected due to the very small seasonal variability there, associated to a small number of coincidences. A similar situation is observed at the other recent oceanic site Palau, where only 4 months of data are available. At the oceanic site Maïdo, we find a good agreement in most of the months but not in October-December, which are the predominant biomass burning months in the region so the collocation of the plumes might play a role there. Finally at Lauder, TROPOMI shows many negative values in the beginning of the period (May-Sept. 2018), which is responsible for a lower correlation (0.65) and for the negative bias there (although not significant), while other clean sites show usually positive ones (see Table 2).



The higher HCHO level sites show a TROPOMI seasonal variability in very good agreement with FTIR, with correlation larger than 0.90 for Boulder, Wollongong, Toronto, Xianghe, and Porto Velho. At Tsukuba, removing the outlier of 1 coincidence in November 2018 increases the correlation to 0.93. The poorest correlation (0.14) is found at the coastal site Paramaribo, where usually only one coincidence per month is found. Looking at the highest HCHO level sites, these monthly mean time-series also confirm that TROPOMI has more difficulty to reproduce the months with the highest enhancements, which is responsible for the significant negative bias (-31%) found in the previous section for high HCHO levels ($>8.0 \times 10^{15}$ molec/cm²).




Figure 5. Monthly means time-series of FTIR raw data (black), FTIR data smoothed with the TROPOMI column averaging kernel (blue), and TROPOMI (red) at each site. Only data in coincidences are included in the monthly mean to avoid sampling bias. When the number of coincidences within one month is smaller than 10, it is written below the monthly mean.




Figure 5. Continued.

6 Conclusions

We have used a network of twenty-five FTIR stations, most of them affiliated to NDACC, to validate the latest TROPOMI HCHO tropospheric columns (v.1.1.[5-7]). This network covers a wide range of concentrations, from very clean Arctic, oceanic and mountain sites, with columns that can be lower than 10^{14} molec/cm², to high HCHO level sites such Mexico city or Porto Velho, near the Amazonian forest, where columns up to 7×10^{16} molec/cm² have been observed.

We found an overestimation ($+26 \pm 5\%$) of TROPOMI OFFL products for very small HCHO columns ($< 2.5 \times 10^{15}$ molec/cm²) and an underestimation of TROPOMI of about -30.8% ($\pm 1.4\%$) for high HCHO columns ($> 8.0 \times 10^{15}$ molec/cm²), which can be used, e.g., to correct TROPOMI data near emissions sources. The results are very similar for NRTI products ($+22 \pm 7\%$ and $-31.7 \pm 1.8\%$ for small and high columns, respectively), and the differences are mainly due to the different period of available TROPOMI v.1.1.[5-7] products. Our wide range of HCHO levels and the use of the Theil-Sen method allow us to derive robust and significant constant (intercept) and proportional (slope) biases of TROPOMI ($\text{TROP} = +1.10 (\pm 0.05) \times 10^{15} + 0.64 (\pm 0.03) \times \text{FTIR}$, in molec/cm²). Such different BIAS for low/high target species concentration levels, due to the presence of both constant and proportional biases, was also recently observed (although with less FTIR sites involved) in another nadir satellite product, the formic acid observed by IASI (Supporting Information in Franco et al. (2020)). The NDACC FTIR network, which covers a large number of atmospheric species at wide ranges of concentrations, is a powerful source of reference



data to detect such nadir satellites' biases.

Although significant, the observed overestimation and underestimation of TROPOMI are within the lower limits of the pre-launch requirements ($\pm 40\%$), as are the biases at individual sites for 20 of the 25 stations. The TROPOMI systematic uncertainty budget is in very good agreement with the observed bias, larger uncertainties being reported at stations where the bias
5 exceeds the 40% requirements.

The precision of TROPOMI OFFL products is estimated by the median absolute deviation (MAD) at the clean sites, where the collocation effect is expected to be small. For FTIR HCHO levels lower than 2.5×10^{15} molec/cm², the MAD is 1.3×10^{15} molec/cm², corresponding to a single pixel precision of 7×10^{15} molec/cm² (5 to 8×10^{15} at individual sites), which is well below the pre-launch precision requirements of 1.2×10^{16} molec/cm². However, the provided TROPOMI random uncertainties
10 (after launch) were indeed found to be better than the pre-launch requirements, but they are too small by a factor of 1.6 compared to the MAD at the clean sites. There is a factor of 2.3 difference between MAD and the random uncertainty on the comparisons (dominated by TROPOMI random uncertainty) at the high-level sites, where an additional effect of collocation might take a role as well. The underestimation of the TROPOMI random uncertainty could be due to a random effect of the uncertainty associated to the air mass factor calculation that is not currently included in the budget. This would also explain
15 a larger underestimation of random error at high-levels sites (factor 2.3 vs 1.6 at clean sites). Also, a systematic uncertainty component on a short-term (so not included in the TROPOMI random uncertainty) can have a random effect on our longer-term comparisons.

We have shown that the TROPOMI data capture very well the HCHO seasonal variability, even at very clean sites. The Pearson correlation coefficient for monthly mean coincident data is 0.91. Although we have found room for a refinement of
20 the TROPOMI random uncertainty estimation and for an improvement of the QA values for a better filtering of the remaining few outliers and negative columns (exceeding the expected statistical distribution), this validation work has demonstrated the very good quality of the TROPOMI HCHO product, which is well within the pre-launch requirements for both accuracy and precision. This work has also shown the high value of the FTIR HCHO network, providing harmonized and well-characterized data covering a wide range of HCHO columns. These ground-based FTIR data are continuously extended by new measurements
25 and will be used in the coming years for the routine S5P validation within the ESA dedicated validation server (<https://mpc-vdaf-server.tropomi.eu/>). The FTIR network will also be used in the near future for the validation of previous satellite missions such as OMI or GOME-2. New FTIR measurements are continuously performed and can be used in the coming years for the validation of new satellite generation, such as TEMPO, GEMS, Sentinel 5P, or Sentinel 4.

An extension of this TROPOMI HCHO validation with ground-based MAX-DOAS and Pandora instruments, especially
30 at sites where both FTIR and UV-Visible techniques are available (e.g. Xianghe, Maïdo, Lauder,...) or at uncovered regions (Africa) would bring additional knowledge. However, there is first a need for a data product harmonization within the MAX-DOAS network, as was done with the FTIR network used here. This work is ongoing as part of the ESA FRM4DOAS and Pandonia projects.

Data availability:



The TROPOMI HCHO data are publicly available at <https://scihub.copernicus.eu>. The access and use of any Copernicus Sentinel data available through the Copernicus Sentinel Data Hub is governed by the Legal Notice on the use of Copernicus Sentinel Data and Service Information which is given here:

https://sentinels.copernicus.eu/documents/247904/690755/Sentinel_Data_Legal_Notice.

5 The FTIR data sets can be provided in the public NDACC repository (<ftp://ftp.cpc.ncep.noaa.gov/ndacc/station/>, last access: January 2020) depending on each PI decision. Please pay attention to the NDACC data policy. The whole data set used in this publication can be provided upon request to Corinne Vigouroux (corinne.vigouroux@aeronomie.be) and data per station can be requested from the individual PIs.

10 *Author contribution:* Corinne Vigouroux and Bavo Langerock performed the validation using HCHO TROPOMI and FTIR data at all sites. They are also involved in the FTIR measurements at Maïdo and Porto Velho. Corinne Vigouroux also analyzed the Sodankylä and Xiangue data. Isabelle De Smedt and Michel Van Roozendael are responsible for the TROPOMI HCHO data. Gaia Pinardi was involved in the validation method section through her expertise in validation using UV-Visible techniques, which is part of the projects TROVA and TROVA-2 that funded this work. All other co-authors provided the FTIR HCHO data for the station(s) they are responsible for.

15 *Competing interests:* The authors declare that they have no conflict of interest.

Acknowledgements. This study has been supported by the ESA PRODEX projects TROVA and TROVA-E2 funded by the Belgian Science Policy Office (Belspo). The measurements at Reunion Island have been also supported by the Université de La Réunion and CNRS (LACY-UMR8105 and UMS3365), and at Porto Velho by the BRAIN-pioneer project IKARE, funded by Belspo, with the collaboration of the Instituto Federal de Educação, Ciência e Tecnologia de Rondônia (IFRO). SPbU FTIR team has been supported by the Russian Foundation for Basic Research Project #18-05-00011. St. Petersburg FTIR measurements were carried out by the instrumentation of the Geomodel resource center of SPbU. The NDACC stations Bremen, Izaña, Ny-Ålesund and Paramaribo have been supported by the German Bundesministerium für Wirtschaft und Energie (BMWi) via DLR under grants 50EE1711A, B and D. We thank the EU-project STRATOCLIM for financial support to U. of Bremen. Measurements made at Lauder, by NIWA, are funded by New Zealand's Ministry of Business, Innovation and Employment through the Strategic Science Investment Fund. We thank the AWI Bremerhaven for logistical support and the station personnel in Ny Alesund. We acknowledge the Meteorologische Dienst van Suriname for logistical and on-site support in Paramaribo, Suriname. We acknowledge the AWI Potsdam and the Coral Reef Foundation for logistical and on-site support in Koror, Palau. ULiège has received support from the F.R.S. - FNRS, from the Fédération Wallonie-Bruxelles and from the GAW-CH programme of MeteoSwiss. We further acknowledge the International Foundation High Altitude Research Stations Jungfraujoch and Gornergrat (HFSJG, Bern) for supporting the facilities needed to perform the Jungfraujoch observations. E. Mahieu is Research Associate with the F.R.S. - FNRS. The National Center for Atmospheric Research is sponsored by the National Science Foundation. The NCAR FTS observation programs at Thule, GR, Boulder, CO, and Mauna Loa, HI are supported under contract by the National Aeronautics and Space Administration (NASA). The Thule work is also supported by the NSF Office of Polar Programs (OPP). We wish to thank the Danish Meteorological Institute for support at the Thule site and NOAA for support at the Mauna Loa. Financial support from grants by DGAPA-UNAM (07417 & 111418) and CONACYT (290589) are acknowledged as well as the University Network of Atmospheric Observatories (RUOA) for the maintenance and operation of the Mexican stations. The Paris TCCON site has received fundings from Sorbonne Université, the French research center CNRS, the French space agency



CNES and Région Île-de-France. Eureka measurements were made at the Polar Environment Atmospheric Research Laboratory (PEARL) under the CANDAC and PAHA projects led by James R. Drummond, and in part by the Canadian Arctic ACE/OSIRIS Validation Campaigns, led by Kaley A. Walker. Funding was provided by AIF/NSRIT, CFI, CFCAS, CSA, ECCC, GOC-IPY, NSERC, NSTP, OIT, PCSP, and ORF. Logistical and operational support was provided by PEARL Site Manager Pierre Fogal, the CANDAC operators, and the ECCC Weather
5 Station. Toronto measurements were made at the University of Toronto Atmospheric Observatory, supported by CFCAS, ABB Bomem, CFI, CSA, ECCC, NSERC, ORDCF, PREA, and the University of Toronto.

The authors would like to thanks all the people responsible for the FTIR measurements and/or data analysis at the different sites: C. Hermans, N. Kumps, M. Zhou, BIRA-IASB; L. Gatti, INPE; Uwe Raffalski, IRF Kiruna; Omaira García and Eliezer Sepulveda, AEMet; C. Becker, Paramaribo; J. Robinson, Lauder; A.V. Poberovskii, H.H. Imkhasin, S.I. Osipov, SPbU; A. Bezanilla and C. Guarín, UNAM; P.
10 Jeseck, Sorbonne Université; M. Rettinger, IMK-IFU; H. Nakajima, NIES.



References

- Boersma, K. F., Eskes, H. J., and Brinkma, E. J.: Error analysis for tropospheric NO₂ retrieval from space, *J. Geophys. Res.*, 109, doi:10.1029/2003JD003962, 2004.
- De Smedt, I., Stavrou, T., Hendrick, F., Danckaert, T., Vlemmix, T., Pinardi, G., Theys, N., Lerot, C., Gielen, C., Vigouroux, C., Hermans, C., Fayt, C., Veefkind, P., Müller, J.-F., and Van Roozendaal, M.: Diurnal, seasonal and long-term variations of global formaldehyde columns inferred from combined OMI and GOME-2 observations, *Atmos. Chem. Phys.*, 15, 12519-12545, doi:10.5194/acp-15-12519-2015, 2015.
- De Smedt, I., Theys, N., Yu, H., Danckaert, T., Lerot, C., Compernelle, S., Van Roozendaal, M., Richter, A., Hilboll, A., Peters, E., Pedernana, M., Loyola, D., Beirle, S., Wagner, T., Eskes, H., van Geffen, J., Boersma, K. F., and Veefkind, P.: Algorithm theoretical baseline for formaldehyde retrievals from S5P TROPOMI and from the QA4ECV project, *Atmos. Meas. Tech.*, 11, 2395-2426, https://doi.org/10.5194/amt-11-2395-2018, 2018.
- Fortems-Cheiney, A., Chevallier, F., Pison, I., Bousquet, P., Saunio, M., Szopa, S., Cressot, C., Kurosu, T. P., Chance, K., and Fried, A.: The formaldehyde budget as seen by a global-scale multi-constraint and multi-species inversion system, *Atmos. Chem. Phys.*, 12, 6699-6721, doi:10.5194/acp-12-6699-2012, 2012.
- Franco, B., Clarisse, L., Stavrou, T., Müller, J.-F., Taraborrelli, D., Hadji-Lazaro, J., Hannigan, J.W., Hase, F., Hurtmans, D., Jones, N., Lutsch, E., Mahieu, E., Ortega, I., Schneider, M., Strong, K., Vigouroux, C., Clerbaux, C., Coheur, P.-F.: Spaceborne Measurements of Formic and Acetic Acid: A Global View of the Regional Sources, *Geophysical Research Letters*, 47, e60094., https://doi.org/10.1029/2019GL086239, 2020.
- Garcia, R. R., Marsh, D. R., Kinnison, D. E., Boville, B. A., and Sassi, F.: Simulation of secular trends in the middle atmosphere, 1950-2003, *J. Geophys. Res.*, 112, D09301, doi:10.1029/2006JD007485, 2007.
- Hase, F., Demoulin, P., Sauval, A. J., Toon, G. C., Bernath, P. F., Goldman, A., Hannigan, J. W., and Rinsland, C. P.: An empirical line-by-line model for the infrared solar transmittance spectrum from 700 to 5000 cm⁻¹, *J. Quant. Spectros. Radiat. Transfer*, 102, 450-463, 2006.
- Huber, P.J., *Robust Statistics*, Wiley, New York, 1981.
- Jacquemart, D., Laraia, A., Kwabia Tchana, F., Gamache, R. R., Perrin, A., and Lacombe, N.: Formaldehyde around 3.5 and 5.7 μm: measurement and calculation of broadening coefficients, *J. Quant. Spectrosc. Radiat. Transfer*, 111, 1209-1222, 2010.
- Jones, N. B., Riedel, K., Allan, W., Wood, S., Palmer, P. I., Chance, K., and Notholt, J.: Long-term tropospheric formaldehyde concentrations deduced from ground-based fourier transform solar infrared measurements, *Atmos. Chem. Phys.*, 9, 7131-7142, doi:10.5194/acp-9-7131-2009, 2009.
- Kleipool, Q. L., Dobber, M. R., de Haan, J. F., and Levelt, P. F.: Earth surface reflectance climatology from 3 years of OMI data, *J. Geophys. Res.*, 113, D18308, doi:10.1029/2008JD010290, 2008.
- Langerock, B., De Mazière, M., Hendrick, F., Vigouroux, C., Desmet, F., Dils, B., and Niemeijer, S.: Description of algorithms for co-locating and comparing gridded model data with remote-sensing observations, *Geosci. Model Dev.*, 8, 911-921, https://doi.org/10.5194/gmd-8-911-2015, 2015.
- Loyola, D. G., Gimeno García, S., Lutz, R., Argyrouli, A., Romahn, F., Spurr, R. J. D., Pedernana, M., Doicu, A., Molina García, V., and Schüssler, O.: The operational cloud retrieval algorithms from TROPOMI on board Sentinel-5 Precursor, *Atmos. Meas. Tech.*, 11, 409-427, https://doi.org/10.5194/amt-11-409-2018, 2018.



- Martin, R. V., Parrish, D. D., Ryerson, T. B., Nicks, D. K. Jr., Chance, K., Kurosu, T. P., Jacob, D. J., Sturges, E. D., Fried, A., and Wert, B. P.: Evaluation of GOME satellite measurements of tropospheric NO₂ and HCHO using regional data from aircraft campaigns in the southeastern United States, *J. Geophys. Res.*, 109, D24307, doi:10.1029/2004JD004869, 2004.
- Meller, R. and Moortgat, G. K.: Temperature dependence of the absorption cross section of HCHO between 223 and 323 K in the wavelength range 225–375 nm, *J. Geophys. Res.*, 105, 7089–7102, doi: 10.1029/1999JD901074, 2000.
- Pougatchev, N. S., Connor, B. J., and Rinsland, C. P.: Infrared measurements of the ozone vertical distribution above Kitt Peak, *J. Geophys. Res.*, 100, 16689–16697, 1995.
- Rodgers, C. D.: Inverse methods for atmospheric sounding: Theory and Practice, Series on Atmospheric, Oceanic and Planetary Physics - Vol. 2, World Scientific Publishing Co., Singapore, 2000.
- Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *J. Geophys. Res.* 108, 4116–4129, 2003.
- Rothman, L. S., Gordon, I. E., Babikov, Y., Barbe, A., Benner, D. C., Bernath, P. F., Birk, M., Bizzocchi, L., Boudon, V., Brown, L. R., Campargue, A., Chance, K., Cohen, E. A., Coudert, L. H., Devi, V. M., Drouin, B. J., Fayt, A., Flaud, J.-M., Gamache, R. R., Harrison, J. J., Hartmann, J.-M., Hill, C., Hodges, J. T., Jacquemart, D., Jolly, A., Lamouroux, J., Roy, R. J. L., Li, G., Long, D. A., Lyulin, O. M., Mackie, C. J., Massie, S. T., Mikhailenko, S., Müller, H. S. P., Naumenko, O. V., and Nikitin, A. V.: The HITRAN2012 molecular spectroscopic database, *J. Quant. Spectrosc. Ra.*, 130, 4–50, doi:10.1016/j.jqsrt.2013.07.002, 2013.
- Sen, Pranab Kumar: Estimates of the regression coefficient based on Kendall's tau, *Journal of the American Statistical Association*, 63 (324): 1379–1389, doi:10.2307/2285891, 1968.
- Spurr, R. J. D.: LIDORT and VLIDORT: Linearized pseudo-spherical scalar and vector discrete ordinate radiative transfer models for use in remote sensing retrieval problems, in *Light Scattering Reviews*, edited by: Kokhanovsky, A., 229–271, Berlin, 2008.
- Stavrou, T., Müller, J.-F., Bauwens, M., De Smedt, I., Van Roozendael, M., De Mazière, M., Vigouroux, C., Hendrick, F., George, M., Clerbaux, C., Coheur, P.-F., and Guenther, A.: How consistent are top-down hydrocarbon emissions based on formaldehyde observations from GOME-2 and OMI?, *Atmos. Chem. Phys.*, 15, 11861–11884, doi:10.5194/acp-15-11861-2015, 2015.
- Tikhonov, A.: On the solution of incorrectly stated problems and a method of regularization, *Dokl. Acad. Nauk SSSR*, 151, 501–504, 1963.
- Veefkind, J.P., Aben, I., McMullan, K., Förster, H., De Vries, J., Otter, G., Claas, J., Eskes, H.J., De Haan, J.F., Kleipool, Q., Van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P.F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sensing of Environment*, 120, 70–83, doi:10.1016/j.rse.2011.09.027, 2012.
- Vigouroux, C., Hendrick, F., Stavrou, T., Dils, B., De Smedt, I., Hermans, C., Merlaud, A., Scolas, F., Senten, C., Vanhaelewyn, G., Fally, S., Carleer, M., Metzger, J.-M., Müller, J.-F., Van Roozendael, M., and De Mazière, M.: Ground-based FTIR and MAX-DOAS observations of formaldehyde at Reunion Island and comparisons with satellite and model data, *Atmos. Chem. Phys.*, 9, 9523–9544, 2009.
- Vigouroux, C., Bauer Aquino, C. A., Bauwens, M., Becker, C., Blumenstock, T., De Mazière, M., García, O., Grutter, M., Guarín, C., Hannigan, J., Hase, F., Jones, N., Kivi, R., Koshelev, D., Langerock, B., Lutsch, E., Makarova, M., Metzger, J.-M., Müller, J.-F., Notholt, J., Ortega, I., Palm, M., Paton-Walsh, C., Poberovskii, A., Rettinger, M., Robinson, J., Smale, D., Stavrou, T., Stremme, W., Strong, K., Sussmann, R., Té, Y., and Toon, G.: NDACC harmonized formaldehyde time series from 21 FTIR stations covering a wide range of column abundances, *Atmos. Meas. Tech.*, 11, 5049–5073, https://doi.org/10.5194/amt-11-5049-2018, 2018.
- von Clarmann, T.: Smoothing error pitfalls, *Atmos. Meas. Tech.*, 7, 3023–3034, https://doi.org/10.5194/amt-7-3023-2014, 2014.



- Williams, J. E., Boersma, K. F., Le Sager, P., and Verstraeten, W. W.: The high-resolution version of TM5-MP for optimized satellite retrievals: description and validation, *Geosci. Model Dev.*, 10, 721-750, doi:10.5194/gmd-10-721-2017, 2017.
- Wittrock, F., Richter, A., Oetjen, H., Burrows, J. P., Kanakidou, M., Myriokefalitakis, S., Volkamer, R., Beirle, S., Platt, U., and Wagner, T.: Simultaneous global observations of glyoxal and formaldehyde from space, *Geophys. Res. Lett.*, 33, L16804, doi:10.1029/2006GL026310, 5 2006.
- Zhu, L., Jacob, D. J., Kim, P. S., Fisher, J. A., Yu, K., Travis, K. R., Mickley, L. J., Yantosca, R. M., Sulprizio, M. P., De Smedt, I., González Abad, G., Chance, K., Li, C., Ferrare, R., Fried, A., Hair, J. W., Hanisco, T. F., Richter, D., Jo Scarino, A., Walega, J., Weibring, P., and Wolfe, G. M.: Observing atmospheric formaldehyde (HCHO) from space: validation and intercomparison of six retrievals from four satellites (OMI, GOME2A, GOME2B, OMPS) with SEAC⁴RS aircraft observations over the southeast US, *Atmos. Chem. Phys.*, 16, 10 13477-13490, doi:10.5194/acp-16-13477-2016, 2016.