



Application of chemical derivatization techniques combined with chemical ionization mass spectrometry to detect stabilized Criegee intermediates and peroxy radicals in the gas phase

Alexander Zaytsev¹, Martin Breitenlechner¹, Anna Novelli², Hendrik Fuchs², Daniel A. Knopf³, Jesse H. Kroll⁴, and Frank N. Keutsch^{1,5,6}

¹John A. Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138, USA

²Institute of Energy and Climate Research – Troposphere (IEK-8), Forschungszentrum Jülich GmbH, 52428 Jülich, Germany

³School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11794, USA

⁴Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

⁵Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA

⁶Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA 02138, USA

Correspondence to: Alexander Zaytsev (zaytsev@g.harvard.edu) and Frank N. Keutsch (keutsch@seas.harvard.edu)

Abstract. Short-lived highly reactive atmospheric species, such as organic peroxy radicals (RO₂) and stabilized Criegee intermediates (SCIs), play an important role in controlling the oxidative removal and transformation of many natural and anthropogenic trace gases in the atmosphere. Direct speciated measurements of these components are extremely helpful for understanding their atmospheric fate and impact. We describe the development of an online method for measurements of SCIs and RO₂ in laboratory experiments using chemical derivatization and spin trapping techniques combined with H₃O⁺ and NH₄⁺ chemical ionization mass spectrometry (CIMS). Using chemical derivatization agents with low proton affinity, such as electron-poor carbonyls, we scavenge all SCIs produced from a wide range of alkenes without depleting CIMS reagent ions. Comparison between our measurements and results from numeric modelling, using a modified version of the Master Chemical Mechanism, shows that the method can be used for quantification of SCIs in laboratory experiments with detection limit of 1.4×10^7 molecule cm⁻³ for 30 s integration time with the instrumentation used in this study. We show that spin traps are highly reactive towards atmospheric radicals and form stable adducts with them by studying the gas-phase kinetics of their reaction with hydroxyl radical (OH). We also demonstrate that spin trap adducts with SCIs and RO₂ can be simultaneously probed and quantified under laboratory conditions with detection limit of 1.6×10^8 molecule cm⁻³ for 30 s integration time for RO₂ species with the instrumentation used in this study. Spin trapping prevents radical secondary reactions and cycling, which ensures that measurements are not biased by chemical interferences, and can be implemented for detecting RO₂ species in the ambient atmosphere.

1 Introduction

Earth's atmosphere is an oxidizing environment. The initial oxidation step of volatile organic compounds (VOCs) involves reaction of a parent hydrocarbon with an oxidant. The hydroxyl radical (OH) is the most important oxidant in the atmosphere,



although oxidation can be also initiated by O_3 , NO_3 and Cl- or Br-atoms. Generally, reaction of VOCs with OH, NO_3 and Cl-atoms occurs via H-abstraction or via addition to unsaturated carbon double bonds leading to the formation of alkyl radicals. This reaction is quickly followed by O_2 addition resulting in the production of organic peroxy radicals (RO_2). In an NO-rich environment, RO_2 radicals predominantly react with NO, while at lower NO concentrations reactions with the hydroperoxy radical (HO_2), potentially other RO_2 , and unimolecular reactions become more important. The common tendency is the formation of closed-shell, more oxidized VOCs (OVOCs). OVOCs may have lower volatilities than the parent hydrocarbons and may partition to the particle phase, thereby contributing to secondary organic aerosol (SOA) formation. OH, HO_2 and RO_2 radicals can form a catalytic reaction cycle, which can lead to production of tropospheric ozone as a consequence of the shift in the NO/ NO_2 ratio to favor formation of NO_2 . This cycle is terminated by the formation of organic hydroperoxides and nitrates, which can be viewed as reservoirs of the corresponding radicals. Overall, atmospheric radicals, especially their cycling, play an important role in the formation of SOA and tropospheric ozone, as well as in controlling atmospheric oxidation capacity.

Organic peroxy radicals can also be formed via ozonolysis of unsaturated organic compounds. Ozonolysis of alkenes results in the formation of primary ozonides that promptly decompose to a stable carbonyl and a vibrationally excited carbonyl oxide, also known as a Criegee intermediate (CI), some of which are thermally stabilized (SCI). SCI primarily decompose or react with water vapor (Vereecken et al., 2017) but are also believed to play a role in oxidation of SO_2 to form H_2SO_4 in the tropical regions (Khan et al., 2018). *syn*-SCI can undergo a unimolecular reaction and form a vinyl hydroperoxide, which rapidly decomposes to an OH radical and a vinyl radical. This radical is in resonance with an acetonyl-type radical, which can combine with molecular oxygen to form an RO_2 species (Johnson and Marston, 2008).

Measurements of atmospheric radicals and reactive intermediates, such as RO_2 and SCIs, are challenging because of their high reactivity towards trace gases and surfaces and rapid cycling, which may lead to potential interferences. Highly sensitive detection systems are required to determine the minute concentrations of these species, which are typically on the order of 10^8 molecule cm^{-3} for organic peroxy radicals (Fuchs et al., 2008) and are expected to be less than 10^5 molecule cm^{-3} for SCIs (Novelli et al., 2017). With respect to RO_2 species, there are several field-deployable measurement techniques available for non-speciated measurements of the sum of RO_2 . Matrix Isolation Electron Spin Resonance Spectroscopy (MIESR) is an established, but rarely used, method for field measurements (Mihelcic et al., 1985). MIESR is an offline technique with a low time resolution (~ 30 min), however, its main advantage is that it does not require instrument calibration. Besides MIESR, chemical amplification and conversion systems represent another class of instruments for field studies (Edwards et al., 2003; Hornbrook et al., 2011; Cantrell et al., 1984; Wood and Charest, 2014). In these systems peroxy radicals are not measured directly but are rather converted to other radicals or closed-shell molecules (e.g., NO_2 or H_2SO_4). A detection limit of 10^7 molecule cm^{-3} can be achieved at a temporal resolution of 15 s, however, discrimination of different RO_2 species is not possible (Edwards et al., 2003). In addition, secondary chemistry, i.e., additional sources of radical production and destruction, has to be considered, and care needs to be taken to ensure that measurements are not biased by any chemical interferences (Reiner et al., 1997). Finally, laser-induced fluorescence (LIF) was also applied for ambient measurements of RO_2 radicals (Fuchs et al.,



2008). This technique is characterized by an excellent detection limit of $(2 - 7) \times 10^7$ molecule cm^{-3} for an integration time of 30 s. Similarly to chemical amplifier systems, LIF does not allow for differentiation of various RO_2 species, however, although it is indirect and converts RO_2 to OH it does not have an amplification chain. Recently, novel mass spectrometric techniques using different ionization schemes to directly detect individual RO_2 species were developed (Hansel et al., 2018; Berndt et al., 2018; Berndt et al., 2019; Nozière and Vereecken, 2019).

As for SCIs, indirect measurement techniques have been widely used. In these techniques SCIs are chemically converted to other species (e.g., H_2SO_4 or hydroxymethyl hydroperoxide, HMHP) (Berndt et al., 2014; Sipilä et al., 2014; Neeb et al., 1997). In 2008, the simplest SCI, CH_2OO , was directly detected for the first time (Taatjes et al., 2008). Later, synchrotron photoionization mass spectrometry was combined with the CI generation technique using diiodoalkane photolysis (Welz et al., 2012), which spurred several studies to examine kinetics of bimolecular and unimolecular SCI reactions (Taatjes et al., 2012; Lewis et al., 2015; Chhantyal-Pun et al., 2016). Recently two new techniques for direct measurements of SCIs using Fourier transform microwave spectroscopy and chemical ionization mass spectrometry (CIMS) were introduced (Womack et al., 2015; Berndt et al., 2017).

Despite the abundance of different analytical methods used for detection of atmospheric radicals and reactive intermediates, there is still a need for an online, direct, field-deployable technique for measuring these short-lived highly reactive compounds in a speciated way. Free radicals have been conventionally detected by chemical derivatization (CD) techniques including spin trapping in condensed-phase biological and chemical systems (Hawkins and Davies, 2014; Nosaka and Nosaka, 2017). Non-radical spin traps (e.g., nitron spin traps) are known to react with free radicals to form stable radical adducts that can be detected with electron paramagnetic resonance spectroscopy (Roberts et al., 2016). In addition, radical spin traps (e.g., nitroxide radicals) are also highly reactive towards radical species such as C-centered radicals and form closed-shell adducts with them (Bagryanskaya and Marque, 2014). However, there are only few studies in which these techniques were applied for probing atmospheric radicals and intermediates. Watanabe et al. (1982) presented an offline method to quantify hydroxyl radicals using the spin trap α -4-pyridyl-*N*-*tert*-butylnitron α -1-oxide (4-POBN) where condensed-phase stable adducts were detected by electron spin resonance. Recently, Giorio et al. (2017) used the spin trap 5,5-Dimethyl-1-pyrroline *N*-oxide (DMPO) to characterize SCIs by detecting gas-phase spin trap adducts with online mass spectrometry.

Here we explore three types of CD agents, including two spin trapping agents, and show how they can be used for detection and quantification of various atmospheric radicals and reactive intermediates (Fig. 1). First, we implement the CD agent hexafluoroacetone (HFA) to characterize a wide range of gas-phase SCIs. HFA is selectively reactive towards SCIs (i.e., it is unreactive towards OH, HO_2 and RO_2), forms stable secondary ozonides with them, and has high vapor pressure and low proton affinity (Fig. 1). Next, we use the radical spin trap (2,2,6,6-Tetramethylpiperidin-1-yl)oxyl (TEMPO) to demonstrate that spin traps are highly reactive towards radicals in the gas phase, by studying kinetics of TEMPO+OH reaction, and therefore can effectively scavenge atmospheric radicals. Finally, we utilize the non-radical spin trap DMPO to simultaneously detect atmospheric gas-phase radicals and intermediates, including SCIs and RO_2 species (Fig. 1). Spin trap adducts and secondary ozonides with CD agents are observed and quantified using H_3O^+ and NH_4^+ CIMS, which allows for speciated online



100 measurements of stabilized Criegee intermediates and speciated RO₂ radicals formed via ozonolysis of a wide range of parent hydrocarbons. The analytical methods presented here can be used for quantification of speciated SCIs and RO₂ formed in laboratory experiments as well as for field measurements.

2 Methods

2.1 Ozonolysis experiments with chemical derivatization agent HFA

105 The ozonolysis experiments of multiple hydrocarbons including tetramethylethylene (TME), isoprene, pentene, hexene, α -pinene and limonene were conducted in a flow tube reactor at ambient pressure and temperature (~290 K) and dry conditions (relative humidity < 2%). The experimental setup consisted of a flow reactor system with a residence time of ~10 s. The parent hydrocarbon was mixed with ozone and the chemical derivatization agent HFA (C₃F₆O) in the flow reactor leading to the formation of SCIs and their scavenging as SCI·HFA adducts. SCIs are known to be highly reactive towards ketones, especially
110 electron poor ones such as HFA (Horie et al., 1999; Drozd et al., 2011; Drozd and Donahue, 2011; Taatjes et al., 2012). The other advantage of employing this chemical derivatization agent is its relatively low proton affinity (PA 670.4 kJ/mol; Hunter and Lias, 1998). Since the PA of HFA is lower than that of water, HFA cannot be protonated in H₃O⁺ CIMS. Hence, one can introduce significant amount of HFA to the system to make sure that all SCIs are scavenged very rapidly without any concern that H₃O⁺ reagent ions would be depleted. The parent hydrocarbon was vaporized from a flask filled with pure substance by
115 passing a constant flow of zero air regulated via a 0.1-10 cm³ min⁻¹ mass flow controller (Bronkhorst). HFA flow was regulated by another mass flow controller (Bronkhorst). Ozone was produced by passing zero air through an ozone generator using a low-pressure mercury ultraviolet lamp. Ozone concentration was measured using an ozone monitor (2B Technologies). A proton-transfer-reaction mass spectrometer (PTR-8000, IONICON Analytik) was used to observe formed SCI·HFA adducts as well as parent hydrocarbons and their oxidation products. This instrument was operated using H₃O⁺ reagent ions (H₃O⁺
120 CIMS) and was directly calibrated to 10 VOCs with different functional groups (Isaacman-VanWertz et al., 2017; Isaacman-VanWertz et al., 2018).

2.2 Experiments with spin traps

2.2.1 Kinetics experiments with spin trap TEMPO

Highly reactive spin traps are needed for effective derivatization of radicals and reactive intermediates in the gas phase. A set
125 of experiments, in which the reaction rate coefficient between the spin trap TEMPO (C₉H₁₈NO) and OH was measured, was conducted in a flow tube experimental setup at Forschungszentrum Jülich. TEMPO is commonly used to detect carbon-centered radicals in chemical and biological systems (Bagryanskaya and Marque, 2014) and is known to be highly reactive towards OH in the aqueous phase (Samuni et al., 2002). TEMPO was introduced in the flow tube setup using a liquid calibration unit (LCU, IONICON Analytik). The LCU quantitatively evaporates aqueous standards into the gas stream. TEMPO standard



130 was prepared gravimetrically with aqueous volume mixing ratio of 485 parts per million (ppm). A known amount (up to 10 $\mu\text{L min}^{-1}$) of this solution was then evaporated into a humidified gas stream of synthetic air (31 SLPM), resulting in the gas-phase TEMPO concentration of up to 4.5×10^{11} molecule cm^{-3} . One part of the setup outflow was drawn to a laser photolysis – laser-induced fluorescence (LP-LIF) instrument (Lou et al., 2010), with which OH reactivity of TEMPO was measured. Laser flash photolysis of ozone was used to produce OH in the experimental setup, while LIF was applied to monitor the time
135 dependent OH decay. Another part of the outflow was drawn to a CIMS instrument PTR3 (IONICON Analytik) to monitor concentrations of TEMPO and its oxidation products. This instrument was operated in two ionization modes: using $\text{H}_3\text{O}^+(\text{H}_2\text{O})_n$, $n = 0-1$ (as H_3O^+ CIMS; Breitenlechner et al., 2017) and $\text{NH}_4^+(\text{H}_2\text{O})_n$, $n = 0-2$ (as NH_4^+ CIMS; Zaytsev et al., 2019) reagent ions. The PTR3 is designed to minimize inlet losses of sampled compounds. It was directly calibrated to 10 VOCs with different functional groups using LCU. Collision-dissociation methods were used to constrain sensitivities of NH_4^+
140 CIMS to compounds that cannot be calibrated directly (Zaytsev et al., 2019). Sensitivities were calculated in normalized duty-cycle-corrected counts per second per part per billion by volume (ndcps ppb^{-1} ; the duty-cycle correction was done to the reference $m/z = 100$; ion signals were normalized to the primary ion signal of 10^6 dcps).

2.2.2 Ozonolysis experiments with spin trap DMPO

An additional set of ozonolysis experiments of several hydrocarbons including TME and α -pinene were conducted in a double
145 flow reactor setup (Fig. 2). The goal of these experiments was to examine how spin traps can be used for simultaneous detection of stabilized Criegee intermediates and peroxy radicals. The parent hydrocarbon was mixed with ozone in the first flow tube, while the spin trap DMPO ($\text{C}_6\text{H}_{11}\text{NO}$) was introduced in the second flow tube using an LCU. A known amount (up to $10 \mu\text{L min}^{-1}$) of the DMPO solution was evaporated into a humidified gas stream of synthetic air (5.4-7 SLPM), resulting in the gas-phase DMPO concentration of up to 1.1×10^{13} molecule cm^{-3} . DMPO represents a class of non-radical spin traps and is
150 widely used to detect oxygen-centered radicals, such as OH, HO_2 and RO_2 , in chemical and biological systems (Roberts et al., 2016; Van Der Zee et al., 1996). Recently, DMPO was also employed to detect SCIs in the gas phase (Giorio et al., 2017). Similar to the previous ozonolysis experiments described in Sect. 2.1, the parent hydrocarbon was vaporized from a flask filled with pure substance by passing zero air regulated by a mass flow controller, and ozone was generated using a low-pressure mercury ultraviolet lamp. The PTR3 was implemented to detect spin trap adducts with SCIs and RO_2 species, while ozone
155 levels were observed using an ozone monitor (2B Technologies).

2.3 Kinetic model and quantum-chemical calculations

The Framework for 0-D Atmospheric Modelling v3.1 (F0AM; Wolfe et al., 2016) containing reactions from the Master Chemical Mechanism (MCM v3.3.1) (Jenkin et al., 1997; Saunders et al., 2003) was used to simulate photooxidation of studied alkenes in the flow reactor system and to compare the modeled concentrations of the products with the measurements. Model
160 calculations were constrained using physical parameters of the experimental setup (pressure and temperature) as well as to observed concentrations of the parent hydrocarbon, ozone and the chemical derivatization agent.



In order to estimate proton affinities of SCI·HFA adducts, we performed geometry optimization and proton affinity calculations with the Gaussian 09 package (Frisch et al., 2009) using the B3LYP functional (Stephens et al., 1994) and TZVP basis sets.

3 Results and discussion

165 3.1 Detection of speciated stabilized Criegee intermediates using chemical derivatization techniques

The primary goal of the first set of experiments was detection of speciated stabilized Criegee intermediates as adducts with the chemical derivatization agent HFA to prevent secondary reactions within the experimental setup. Starting with (CH₃)₂COO, an SCI produced via ozonolysis of TME, we tested the formation of SCI·HFA adducts under different experimental conditions (Fig. 3). (CH₃)₂COO·HFA (C₆H₆O₃F₆·H⁺, *m/z* 241.03) can be easily identified in the mass-spectrum due to its unique mass defect associated with six F-atoms (Fig. S2). SCI·HFA adducts were observed when TME, ozone, and HFA were present in
170 the experimental setup. Ozonolysis of TME also results in the formation of acetone (C₃H₆O·H⁺, *m/z* 59.05), which was detected in the presence of TME and ozone and was not affected by HFA addition (Fig. 3). Since the reaction rate constant of SCI·HFA with H₃O⁺ ions is unknown, we assumed that all SCI·HFA adducts were ionized via proton transfer from hydronium ions and therefore used the sensitivity we obtained from acetone calibration to quantify detected SCI·HFA species. In addition, we did
175 not take into account possible fragmentation of SCI·HFA adducts which may impede their detection, although a first bond cleavage would likely only break the ozonide ring structure without loss of mass. These assumptions may lead to measurement uncertainties as discussed later in this section.

We measured the (CH₃)₂COO·HFA adduct signal as a function of different reactant conditions: initial TME concentration were in the range of (1.48 – 1.85) × 10¹¹ molecule cm⁻³, ozone, (6.77 – 108.2) × 10¹² molecule cm⁻³, HFA (1.17 – 6.13) ×
180 10¹⁵ molecule cm⁻³. The measurements are compared to the predictions of the kinetic model in Fig. 4. Concentrations of (CH₃)₂COO species were calculated using the MCM with updated kinetics data from the literature (Newland et al., 2015; Chhantyal-Pun et al., 2016; Long et al., 2018). For more details see the Supplement.

In the presence of HFA, SCI can react with HFA and form stable adducts:



185 The reaction rate coefficient *k*₁ was not measured experimentally, and we used the *k*-value for CH₂OO + HFA reaction: *k*₁ = 3 × 10⁻¹¹ cm³ molecule⁻¹ s⁻¹ (Taatjes et al., 2012). It has been suggested that the reaction between HFA and acetone oxide may be slower compared to the CH₂OO one (Murray et al., 1965; Taatjes et al., 2012). However, the concentration of HFA was two orders of magnitude higher than concentrations of other chemical compounds, so even at lower *k*-values reaction with HFA remains the major chemical loss pathway for (CH₃)₂COO (Fig. S3).

190 Observed concentrations of (CH₃)₂COO·HFA agree to within a factor of 3 with concentrations predicted by the kinetic model (Fig. 4). This discrepancy can be explained by a combination of the following factors: (1) wall losses of (CH₃)₂COO·HFA in the experimental setup and the PTR 8000 instrument; (2) uncertainty in the sensitivity at which the SCI·HFA adducts were



195 detected; (3) potential ion fragmentation of protonated SCI·HFA adducts; and (4) uncertainty in the SCI yield, and unimolecular and bimolecular reaction rate coefficients used in the kinetic model. The detection limit for (CH₃)₂COO·HFA adducts was 1.4×10^7 molecule cm⁻³ and was calculated for 30 s integration time as 3 standard deviations of measured background divided by derived sensitivity.

Besides TME, we also observed formation of SCI·HFA for a series of precursors including isoprene, pentene, and hexene. (Figs. S4-S6). Proton affinities (PAs) of different CI·HFA adducts were calculated using DFT methods (Table 1). A variety of these adducts can be detected using H₃O⁺ CIMS since their PAs are significantly higher than that of water which is in agreement with experimental data (Figs. S4-S6). CH₂OO·HFA cannot be detected because of its low PA (Table 1). We also did not observe SCI·HFA adducts for larger C₁₀ SCIs produced via ozonolysis of α-pinene and limonene. This can be explained by the lower reactivity of larger SCIs with HFA, potential instability of these secondary ozonides in the gas phase, or their gas-wall partitioning in tubing and inside the PTR-8000 instrument.

3.2 Reactivity of spin traps with OH

205 Spin traps have been shown to be highly reactive towards free radicals and efficiently form adducts with them in the aqueous phase. However, their reactivity with atmospheric radicals and stability of formed adducts in the gas phase remain largely unknown. In order to address these questions, we conducted a set of experiments to estimate the reaction rate between the spin trap TEMPO and the hydroxyl radical by measuring its OH reactivity.

OH reactivity of a specific reactant can be calculated as a product of the reactant concentration and its reaction rate with OH (Fuchs et al., 2017):

$$k_{\text{OH}} = k_{\text{OH}+\text{TEMPO}} \cdot [\text{TEMPO}] \quad (1)$$

k_{OH} was measured as a function of TEMPO concentration by varying the amount of TEMPO introduced in the experimental setup using the LCU (Fig. 5). The slope of the fitted line in Fig. 5 determines the reaction rate coefficient $k_{\text{OH}+\text{TEMPO}} = (9.3 \pm 0.9) \times 10^{-11}$ cm³ molecule⁻¹ s⁻¹. This rate constant is close to the collisional limit of typical radical-molecule reactions in the atmosphere and is one order of magnitude greater than the rate constant for the same reaction in the aqueous phase ($k_{\text{aqueous}} = 7.5 \times 10^{-12}$ cm³ molecule⁻¹ s⁻¹; Samuni et al., 2002). This demonstrates that TEMPO is highly reactive towards OH in the gas phase, emphasizing the applicability of spin trapping for atmospheric measurements. Furthermore, TEMPO + OH reaction leads to the formation of stable TEMPO·OH adducts that can be detected by H₃O⁺ CIMS (C₉H₁₈NO·H⁺, *m/z* 174.149) and therefore could be used for quantification of hydroxyl radicals in the atmosphere (Fig. S7). Further tests are needed to compare the measurement capability of this method (e.g., sensitivity, wall losses, and potential interferences) with that of a well-established technique, such as LIF.



3.3 Simultaneous detection of SCIs and RO₂ species from ozonolysis of alkenes using spin trapping techniques

Next, we implemented spin trapping for detection of speciated SCIs and RO₂ species formed via ozonolysis of alkenes, starting with TME. Decomposition of the TME primary ozonide leads to formation of acetone oxide (CH₃)₂COO. This SCI can further
225 undergo a unimolecular reaction followed by O₂ addition to form a peroxy radical CH₃C(=O)CH₂OO· and OH (Fig. 1). In order to detect SCIs and RO₂ species produced via ozonolysis of TME, we used a measurement method based on stabilization of these species using the spin trap DMPO followed by detection by NH₄⁺ and H₃O⁺ CIMS. DMPO was shown to form stable secondary ozonides with SCIs in the gas phase (Giorio et al., 2017):



230 DMPO is shown to be highly reactive towards SCIs ($k_{CI+DMPO} \geq 6 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$; see the Supplement for more details).

In addition, DMPO is known to be highly reactive towards oxygen-centered radicals, such as RO₂, and form stable radical adducts with them (Fig. 1):



235 We observed the formation of SCI·DMPO and RO₂·DMPO adducts both in NH₄⁺ CIMS (e.g., C₉H₁₇NO₃·NH₄⁺, *m/z* 205.155 and C₉H₁₆NO₄·NH₄⁺, *m/z* 220.142) and H₃O⁺ CIMS (e.g., C₉H₁₇NO₃·H⁺, *m/z* 188.128 and C₉H₁₆NO₄·H⁺, *m/z* 203.116) under different experimental conditions (Figs. 6 and S8). SCI·DMPO and RO₂·DMPO were only detected when TME, ozone, and DMPO were present in the experimental setup. Acetone, also formed via ozonolysis of TME, was observed in the presence of TME and ozone and was not affected by addition of DMPO (Figs. 6 and S8). One of the benefits of NH₄⁺ CIMS is the
240 possibility of quantifying compounds for which authentic standards are not available, using a voltage scanning procedure based on collision-induced dissociation (Zaytsev et al., 2019). Based on this method, DMPO adducts with SCIs and RO₂ were detected at high sensitivities: 2,400 ndcps ppbv⁻¹ for SCI·DMPO and 2,000 ndcps ppbv⁻¹ for RO₂·DMPO (Table S1). Sensitivities were experimentally determined in each ozonolysis experiment and depend on the operational conditions of the PTR3 instrument. Detection limits for SCI·DMPO and RO₂·DMPO adducts were 3.4×10^7 and $1.6 \times 10^8 \text{ molecule cm}^{-3}$,
245 respectively. These limits of detection are calculated for 30 s integration time as 3 standard deviations of measured background divided by derived sensitivity.

In addition, we compare measured concentrations of RO₂·DMPO adducts with the concentrations predicted by the kinetic model (Fig. 7). The observed values are an order of magnitude lower than the modeled ones. Several factors can contribute to this discrepancy: (1) gas-wall partitioning of RO₂ species and RO₂·DMPO adducts in the flow tube setup and inside the PTR3
250 instrument; (2) uncertainty in sensitivity at which RO₂·DMPO adducts were detected; (3) potential fragmentation of RO₂·DMPO; and (4) uncertainties in the reaction rate coefficient k_{RO_2+DMPO} . In our model we assumed that the major fraction of RO₂ species was scavenged by DMPO. This assumption is valid if k_{RO_2+DMPO} is larger than $1 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. Otherwise, other loss channels for peroxy radicals, especially the RO₂+RO₂ reaction, become more important (Fig. S9).



Additional experiments under different conditions and intercomparison with established methods (i.e., LIF) are needed to
255 further estimate the measurement capability of the proposed analytical method.

Finally, we employed spin trapping for detection of SCIs and organic peroxy radicals formed via ozonolysis of larger cyclic
alkenes, such as α -pinene. Decomposition of the α -pinene primary ozonide yields four different C_{10} -SCIs, all of which have
the same molecular formula $C_{10}H_{16}O_3$ (Newland et al., 2018). These SCIs can further isomerize to form primary peroxy radicals
 $C_{10}H_{15}O_4$ and OH. Autoxidation of $C_{10}H_{15}O_4$ - RO_2 species can in turn result in formation of several more oxygenated peroxy
260 radicals $C_{10}H_{15}O_x$, $x = 5-9$ (Zhao et al., 2018). Signals of $SCI \cdot DMPO$ ($C_{10}H_{16}O_3 \cdot DMPO$) and $RO_2 \cdot DMPO$ ($C_{10}H_{15}O_x \cdot DMPO$,
 $x = 4-9$) adducts were observed both in NH_4^+ and H_3O^+ CIMS (Figs. 8, S10, S11). This demonstrates that this analytical method
allows for simultaneous detection of a wide range of atmospheric radicals, including the ones with high oxygen content (an
O:C ration of up to 0.9) that are formed via autoxidation pathway, and can be used to study kinetics of these species in the
laboratory.

265 4 Conclusions

In summary, we experimentally demonstrated the measurement of speciated, short-lived highly reactive atmospheric
compounds, such as Criegee intermediates, organic peroxy radicals and hydroxyl radicals, formed via ozonolysis of alkenes.
The analysis was carried out using chemical derivatization techniques, including spin trapping, while the detection of formed
radical adducts and closed-shell secondary ozonides was performed by the means of H_3O^+ and NH_4^+ CIMS. Detected adducts
270 and secondary ozonides have unique mass defects and can therefore be clearly separated from other observed compounds in
the mass spectrum. Implementation of chemical derivatization agents with lower proton affinity allows for the full scavenging
and quantification of stabilized Criegee intermediates without depleting CIMS reagent ions. We show that spin traps can be
used to effectively scavenge atmospheric radicals and reactive intermediates by demonstrating their high reactivity with
radicals in the gas phase using the TEMPO+OH reaction as an example. Using the spin trap DMPO, SCIs and RO_2 species
275 can be simultaneously detected while quantification of observed adducts can be done without their direct calibration. The
detection limits of spin trap and chemical derivatization agent adducts of 1.4×10^7 molecule cm^{-3} for SCIs and 1.6×10^8
molecule cm^{-3} for RO_2 for 30 s integration time were estimated for the instrumentation used here and show promise that these
techniques would also work when sampling ambient air. In particular, this method fundamentally enables any CIMS instrument
to detect radicals and SCIs. Since spin traps, such as DMPO and TEMPO, are reactive towards a plethora of atmospheric
280 radicals and reactive intermediates, including RO_2 , SCIs, and OH, implementation of such spin traps results in the effective
suppression of the radical secondary chemistry and, thus, elimination of potential chemical interferences. The direct method
for speciated SCIs and RO_2 measurements provides a means to study the atmospheric chemistry of these compounds. We stress
that the quantification of RO_2 species was done under well-defined laboratory conditions using the CID technique such that
the estimated sensitivities are likely unique to the electric fields, pressures and flows of the NH_4^+ CIMS instrument. Further



285 validation of the proposed analytical methods in more complex environments closer to the ambient conditions and
intercomparison with established methods (i.e., LIF) are needed.

For the future application of the method in field and laboratory experiments, various modifications of the experimental setup
can be implemented to improve its measurement capability. We plan to synthesize and test new chemical derivatization agents
optimized for the gas-phase measurements with respect to their vapor pressure, selective reactivity and by labelling with atomic
290 isotopes to simplify mass spectrometric detection and improve detection limits. With labeled spin traps, the identification of
reactive intermediates may be greatly simplified and detection limits may be further improved, as the spin trap can provide a
unique signature in the complex mass-spectrum and move the observed m/z to a region with very low background.

Acknowledgements. This work has received funding from the Harvard Global Institute. Martin Breitenlechner acknowledges
295 support from the Austrian science fund (FWF; grant J-3900). Hendrik Fuchs and Anna Novelli acknowledge support from the
European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (SARLEP
grant agreement No. 681529). Daniel Knopf acknowledges support from the U.S. National Science Foundation (grant no.
AGS-1446286). Jesse Kroll acknowledges support from the U.S. National Science Foundation (grant no. AGS-1638672).



References

- 300 Bagryanskaya, E.G. and Marque, S.R.: Scavenging of organic C-centered radicals by nitroxides. *Chem. Rev.*, 114(9), 5011-5056, DOI: 10.1021/cr4000946, 2014.
- Berndt, T., Jokinen, T., Sipilä, M., Mauldin III, R.L., Herrmann, H., Stratmann, F., Junninen, H. and Kulmala, M.: H₂SO₄ formation from the gas-phase reaction of stabilized Criegee Intermediates with SO₂: Influence of water vapour content and temperature. *Atmospheric Environment*, 89, 603-612, DOI: 10.1016/j.atmosenv.2014.02.062, 2014.
- 305 Berndt, T., Herrmann, H., and Kurten, T.: Direct Probing of Criegee Intermediates from Gas-Phase Ozonolysis Using Chemical ionization Mass Spectrometry, *J. Am. Chem. Soc.*, 139, 13387-13392, DOI: 10.1021/jacs.7b05849, 2017.
- Berndt, T., Mentler, B., Scholz, W., Fischer, L., Herrmann, H., Kulmala, M. and Hansel, A.: Accretion product formation from ozonolysis and OH radical reaction of α -pinene: mechanistic insight and the influence of isoprene and ethylene. *Environmental science & technology*, 52(19), 11069-11077, DOI: 10.1021/acs.est.8b02210, 2018.
- 310 Berndt, T., Hyttinen, N., Herrmann, H., and Hansel, A.: First oxidation products from the reaction of hydroxyl radicals with isoprene for pristine environmental conditions. *Communications Chemistry*, 2(1), 1-10, DOI: 10.1038/s42004-019-0120-9, 2019.
- Breitenlechner, M., Fischer, M., Hainer, M., Heinritzi, M., Curtius, M., and Hansel, A.: PTR3: An instrument for Studying the Lifecycle of Reactive Organic Carbon in the Atmosphere, *Anal. Chem.*, 89, 5824–5831, DOI: 10.1021/acs.analchem.6b05110, 315 2017.
- Cantrell, C.A., Stedman, D.H., and Wendel, G.J.: Measurement of atmospheric peroxy radicals by chemical amplification, *Anal. Chem.*, 56, 1496–1502. DOI: 10.1021/ac00272a065, 1984.
- Chhantyal-Pun, R., Welz, O., Savee, J.D., Eskola, A.J., Lee, E.P.F., Blacker, L., Hill, H.R., Ashcroft, M., Khan, M.A.H., Lloyd-Jones, G.C., Evans, L., Rotavera, B., Huang, H., Osborn, D.L., Mok, D.K.W., Dyke, J.M., Shallcross, D.E., Percival, C.J., 320 Orr-Ewing, A.J., and Taatjes, C.A.: Direct Measurements of Unimolecular and Bimolecular Reaction Kinetics of the Criegee Intermediate (CH₃)₂COO, *J. Phys. Chem. A*, 121, 4–15, DOI: 10.1021/acs.jpca.6b07810, 2016.
- Drozd, G.T., Kroll, J.H., and Donahue, N.M.: 2,3-Dimethyl-2-butene (TME) Ozonolysis: Pressure Dependence of Stabilized Criegee Intermediates and Evidence of Stabilized Vinyl Hydroperoxides, *J. Phys. Chem. A*, DOI: 10.1021/jp108773d, 2011.
- Drozd, G.T. and Donahue, N.M.: Pressure Dependence of Stabilized Criegee Intermediate Formation from a Sequence of 325 Alkenes, *J. Phys. Chem. A*, DOI: 10.1021/jp2001089, 2011.
- Edwards, G.D., Cantrell, C.A., Stephens, S., Hill, B., Goyea, O., Shetter, R.E., Mauldin III, R.L., Kosciuch, E., Tanner, D.J., and Eisele, F.L.: Chemical Ionization Mass Spectrometer Instrument for the Measurement of Tropospheric HO₂ and RO₂, *Anal. Chem.*, 75, 5317-5327, DOI: 10.1021/ac034402b, 2003.
- Frisch, M. J., Trucks, G. W., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Scalmani, G., Barone, V., 330 Mennucci, B., Petersson, G. A., Nakatsuji, H., Caricato, M., Li, X., Hratchian, H. P., Izmaylov, A. F., Bloino, J., Zheng, G., Sonnenberg, J. L., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O.,



- Nakai, H., Vreven, T., Montgomery, J. A., Jr., Peralta, J. E., Ogliaro, F., Bearpark, M., Heyd, J. J., Brothers, E., Kudin, K. N., Staroverov, V. N., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A., Burant, J. C., Iyengar, S. S., Tomasi, J., Cossi, M., Rega, N., Millam, J. M., Klene, M., Knox, J. E., Cross, J. B., Bakken, V., Adamo, C., Jaramillo, J., Gomperts, R.,
335 Stratmann, R. E., Yazyev, O., Austin, A. J., Cammi, R., Pomelli, C., Ochterski, J. W., Martin, R. L., Morokuma, K., Zakrzewski, V. G., Voth, G. A., Salvador, P., Dannenberg, J. J., Dapprich, S., Daniels, A. D., Farkas, Ö., Foresman, J. B., Ortiz, J. V., Cioslowski, J., and Fox, D. J.: Gaussian 09, Revision D.01, Gaussian, Inc.: Wallingford, CT, 2009.
- Fuchs, H., Holland, F., and Hofzumahaus, A.: Measurement of tropospheric RO₂ and HO₂ radicals by a laser-induced fluorescence instrument, *Rev. Sci. Instrum.*, 79, 084104, DOI: 10.1063/1.2968712, 2008.
- 340 Fuchs, H., Novelli, A., Rolletter, M., Hofzumahaus, A., Pfannerstill, E. Y., Kessel, S., Edtbauer, A., Williams, J., Michoud, V., Dusanter, S., Locoge, N., Zannoni, N., Gros, V., Truong, F., Sarda-Esteve, R., Cryer, D. R., Brumby, C. A., Whalley, L. K., Stone, D., Seakins, P. W., Heard, D. E., Schoemaeker, C., Blocquet, M., Coudert, S., Batut, S., Fittschen, C., Thames, A. B., Brune, W. H., Ernest, C., Harder, H., Muller, J. B. A., Elste, T., Kubistin, D., Andres, S., Bohn, B., Hohaus, T., Holland, F., Li, X., Rohrer, F., Kiendler-Scharr, A., Tillmann, R., Wegener, R., Yu, Z., Zou, Q., and Wahner, A.: Comparison of OH
345 reactivity measurements in the atmospheric simulation chamber SAPHIR, *Atmos. Meas. Tech.*, 10, 4023–4053, DOI: 10.5194/amt-10-4023-2017, 2017.
- Giorio, C., Campbell, S.J., Bruschi, M., Tampieri, F., Barbon, A., Toffoletti, A., Tapparo, A., Paijens, C., Wedlake, A.J., Grice, P., Howe, D.J., and Kalberer, M.: Online Quantification of Criegee Intermediates of α -Pinene Ozonolysis by Stabilization with Spin Traps and Proton-Transfer Reaction Mass Spectrometry Detection, *J. Am. Chem. Soc.*, 139, 3999–4008, DOI:
350 10.1021/jacs.6b10981, 2017.
- Hansel, A., Scholz, W., Mentler, B., Fischer, L. and Berndt, T.: Detection of RO₂ radicals and other products from cyclohexene ozonolysis with NH₄⁺ and acetate chemical ionization mass spectrometry. *Atmospheric Environment*, 186, 248-255, DOI: 10.1016/j.atmosenv.2018.04.023, 2018.
- Hawkins, C.L. and Davies, M.J.: Detection and characterisation of radicals in biological materials using EPR methodology,
355 *Biochimica et Biophysica Acta (BBA)-General Subjects*, 1840(2), 708-721, DOI: 10.1016/j.bbagen.2013.03.034, 2014.
- Horie, O., Schäfer, C. and Moortgat, G.K.: High reactivity of hexafluoro acetone toward criegee intermediates in the gas-phase ozonolysis of simple alkenes. *International journal of chemical kinetics*, 31(4), 261-269, DOI: 10.1002/(SICI)1097-4601(1999)31:4<261::AID-KIN3>3.0.CO;2-Z, 1999.
- Hornbrook, R.S., Crawford, J.H., Edwards, G.D., Goyea, O., Mauldin Iii, R.L., Olson, J.S., Cantrell, C.A: Measurements of
360 tropospheric HO₂ and RO₂ by oxygen dilution modulation and chemical ionization mass spectrometry. *Atmos Meas Tech* 4, 735–756. DOI: 10.5194/amt-4-735-2011, 2011.
- Hunter, E.P. and Lias, S.G.: Evaluated Gas Phase Basicities and Proton Affinities of Molecules: An Update, *J. Phys. Chem. Ref. Data*, 27, 3, 413-656, DOI: 10.1063/1.556018, 1998.
- Isaacman-VanWertz, G., Massoli, P., O'Brien, R.E., Nowak, J.B., Canagaratna, M.R., Jayne, J.T., Worsnop, D.R., Su, L.,
365 Knopf, D.A., Misztal, P.K., Arata, C., Goldstein, A.H. and Kroll, J.H.: Using advanced mass spectrometry techniques to fully



- characterize atmospheric organic carbon: current capabilities and remaining gaps. *Faraday Discussions*, 200, 579-598, DOI: 10.1039/c7fd00021a, 2017.
- Isaacman-VanWertz, G., Massoli, P., O'Brien, R., Lim, C., Franklin, J.P., Moss, J.A., Hunter, J.F., Nowak, J.B., Canagaratna, M.R., Misztal, P.K., Arata, C., Roscioli, J.R., Herndon, S.T., Onasch, T.B., Lambe, A.T., Jayne, J.T., Su, L., Knopf, D.A.,
370 Goldstein, A.H., Worsnop, D.R. and Kroll, J.H.: Chemical evolution of atmospheric organic carbon over multiple generations of oxidation. *Nature Chem.*, 10, 462-468, DOI: 10.1038/s41557-018-0002-2, 2018.
- Jenkin, M.E., Saunders, S.M., Pilling, M.J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31, 81-104, DOI: 10.1016/S1352-2310(96)00105-7, 1997.
- Johnson, D. and Marston, G.: The gas-phase ozonolysis of unsaturated volatile organic compounds in the
375 troposphere. *Chemical Society Reviews*, 37(4), 699-716, DOI: 10.1039/b704260b, 2008.
- Khan, M.A.H., Percival, C.J., Caravan, R.L., Taatjes, C.A. and Shallcross, D.E.: Criegee intermediates and their impacts on the troposphere, *Environ. Sci.: Processes Impacts*, 20, 437, DOI: 10.1039/C7EM00585G, 2018.
- Lewis, T.R., Blitz, M.A., Heard, D.E. and Seakins, P.W.: Direct evidence for a substantive reaction between the Criegee intermediate, CH₂OO, and the water vapour dimer. *Physical Chemistry Chemical Physics*, 17(7), 4859-4863, DOI:
380 10.1039/C4CP04750H, 2015.
- Long, B., Bao, J.L., and Truhlar, D.G. Unimolecular reaction of acetone oxide and its reaction with water in the atmosphere, *PNAS*, 115, 6135-6140, DOI: 10.1073/pnas.1804453115, 2018.
- Lou, S., Holland, F., Rohrer, F., Lu, K., Bohn, B., Brauers, T., Chang, C. C., Fuchs, H., Häsel, R., Kita, K., Kondo, Y., Li, X., Shao, M., Zeng, L., Wahner, A., Zhang, Y., Wang, W., and Hofzumahaus, A.: Atmospheric OH reactivities in the Pearl
385 River Delta – China in summer 2006: measurement and model results, *Atmos. Chem. Phys.*, 10, 11243–11260, <https://doi.org/10.5194/acp-10-11243-2010>, 2010.
- Mihelcic, D., Müsgen, P., and Ehhalt, D.H.: An improved method of measuring tropospheric NO₂ and RO₂ by matrix isolation and electron spin resonance, *J. Atmos. Chem.*, 3, 341-361, DOI: 10.1007/BF00122523, 1985.
- Murray, R.W., Story, P.R., and Loan, L.D.: Ozonides from Aldehydic Zwitterions and Acetone, *J. Am. Chem. Soc.* 87, 13,
390 3025-3026, DOI: 10.1021/ja01091a054, 1965.
- Neeb, P., Sauer, F., Horie, O., and Moortgat, G. K.: Formation of hydroxymethyl hydroperoxide and formic acid in alkene ozonolysis in the presence of water vapour, *Atmos. Environ.*, 31, 1417– 1423, DOI: 10.1016/S1352-2310(96)00322-6, 1997.
- Newland, M.J., Rickard, A.R., Alam, M.S., Vereecken, L., Muñoz, A., Rodenas, M. and Bloss, W.J.: Kinetics of stabilised Criegee intermediates derived from alkene ozonolysis: reactions with SO₂, H₂O and decomposition under boundary layer
395 conditions. *Phys. Chem. Chem. Phys.*, 17, 4076-4088, DOI: 10.1039/C4CP04186K, 2015.
- Newland, M. J., Rickard, A. R., Sherwen, T., Evans, M. J., Vereecken, L., Muñoz, A., Ródenas, M., and Bloss, W. J.: The atmospheric impacts of monoterpene ozonolysis on global stabilised Criegee intermediate budgets and SO₂ oxidation: experiment, theory and modelling, *Atmos. Chem. Phys.*, 18, 6095–6120, DOI: 10.5194/acp-18-6095-2018, 2018.



- Nosaka, Y. and Nosaka, A.Y.: Generation and detection of reactive oxygen species in photocatalysis, *Chem. Rev.*, 117(17), 11302–11336, DOI: 10.1021/acs.chemrev.7b00161, 2017.
- Novelli, A., Hens, K., Tatum Ernest, C., Martinez, M., Nölscher, A. C., Sinha, V., Paasonen, P., Petäjä, T., Sipilä, M., Elste, T., Plass-Dülmer, C., Phillips, G. J., Kubistin, D., Williams, J., Vereecken, L., Lelieveld, J., and Harder, H.: Estimating the atmospheric concentration of Criegee intermediates and their possible interference in a FAGE-LIF instrument, *Atmos. Chem. Phys.*, 17, 7807–7826, DOI: 10.5194/acp-17-7807-2017, 2017.
- Nozière, B. and Vereecken, L.: Direct Observation of Aliphatic Peroxy Radical Autoxidation and Water Effects: An Experimental and Theoretical Study. *Angew. Chem.*, 58(39), 13976–13982, DOI: 10.1002/anie.201907981, 2019.
- Orlando, J.J., and Tyndall, G.S.: Laboratory studies of organic peroxy radical chemistry: an overview with emphasis on recent issues of atmospheric significance, *Chem. Soc. Rev.*, 41, 6294–6317, DOI: 10.1039/c2cs35166h, 2012.
- Pattison, D.I., Lam, M., Shinde, S.S., Anderson, R.F. and Davies, M.J.: The nitroxide TEMPO is an efficient scavenger of protein radicals: Cellular and kinetic studies. *Free Radical Biology and Medicine*, 53, 1664–1674, DOI: 10.1016/j.freeradbiomed.2012.08.578, 2012.
- Reiner, T., Hanke, M. and Arnold, F.: Atmospheric peroxy radical measurements by ion molecule reaction-mass spectrometry: A novel analytical method using amplifying chemical conversion to sulfuric acid. *Journal of Geophysical Research: Atmospheres*, 102(D1), 1311–1326, DOI: 10.1029/96JD02963, 1997.
- Roberts, J.G., Voinov, M.A., Schmidt, A.C., Smirnova, T.I. and Sombers, L.A.: The hydroxyl radical is a critical intermediate in the voltammetric detection of hydrogen peroxide. *J. Am. Chem. Soc.*, 138(8), 2516–2519, DOI: 10.1021/jacs.5b13376, 2016.
- Samuni, A., Goldstein, S., Russo, A., Mitchell, J.B., Krishna, M.C. and Neta, P.: Kinetics and mechanism of hydroxyl radical and OH-adduct radical reactions with nitroxides and with their hydroxylamines. *J. Am. Chem. Soc.*, 124(29), 8719–8724, DOI: 10.1021/ja017587h, 2002.
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, DOI: 10.5194/acp-3-161-2003, 2003.
- Sipilä, M., Jokinen, T., Berndt, T., Richters, S., Makkonen, R., Donahue, N. M., Mauldin III, R. L., Kurtén, T., Paasonen, P., Sarnela, N., Ehn, M., Junninen, H., Rissanen, M. P., Thornton, J., Stratmann, F., Herrmann, H., Worsnop, D. R., Kulmala, M., Kerminen, V.-M., and Petäjä, T.: Reactivity of stabilized Criegee intermediates (sCIs) from isoprene and monoterpene ozonolysis toward SO₂ and organic acids, *Atmos. Chem. Phys.*, 14, 12143–12153, DOI: 10.5194/acp-14-12143-2014, 2014.
- Stephens, P. J., Devlin, F. J., Chabalowski, C. F. N., and Frisch, M. J.: Ab initio calculation of vibrational absorption and circular dichroism spectra using density functional force fields. *The Journal of physical chemistry*, 98, 11623–11627, DOI: 10.1021/j100096a001, 1994.
- Taatjes, C.A., Meloni, G., Selby, T.M., Trevitt, A.J., Osborn, D.L., Percival, C.J. and Shallcross, D.E.: Direct observation of the gas-phase Criegee intermediate (CH₂OO). *J. Am. Chem. Soc.*, 130(36), 11883–11885, DOI: 10.1021/ja804165q, 2008.




- Taatjes, C.A., Welz, O., Eskola, A.J., Savee, J.D., Osborn, D.L., Lee, E.P.F., Dyke, J.M., Mok, D.W.K., Shallcross D.E., and Percival, C.J.: Direct measurement of Criegee intermediate (CH_2OO) reactions with acetone, acetaldehyde, and hexafluoroacetone, *Phys. Chem. Chem. Phys.*, 14, 10391–10400, DOI: 10.1039/C2CP40294G, 2012.
- 435 Van Der Zee, J., Barr, D.P. and Mason, R.P.: ESR spin trapping investigation of radical formation from the reaction between hematin and tert-butyl hydroperoxide. *Free Radical Biology and Medicine*, 20(2), 199–206, DOI: 10.1016/0891-5849(95)02031-4, 1996.
- Vereecken, L., Novelli, A. and Taraborrelli, D.: Unimolecular decay strongly limits the atmospheric impact of Criegee intermediates. *Phys. Chem. Chem. Phys.*, 19(47), 31599–31612, DOI: 10.1039/C7CP05541B, 2017.
- 440 Watanabe, T., Yoshida, M., Fujiwara, S., Abe, K., Onoe, A., Hirota, M. and Igarashi, S.: Spin trapping of hydroxyl radical in the troposphere for determination by electron spin resonance and gas chromatography/mass spectrometry. *Anal. Chem.*, 54(14), 2470–2474, DOI: 10.1021/ac00251a015, 1982.
- Welz, O., Savee, J.D., Osborn, D.L., Vasu, S.S., Percival, C.J., Shallcross, D.E., and Taatjes, C.A.: Direct kinetic measurements of Criegee intermediate (CH_2OO) formed by reaction of CH_2I with O_2 , *Science*, 335, 204–207, DOI: 10.1126/science.1213229, 2012.
- 445 Wolfe, G. M., Marvin, M. R., Roberts, S. J., Travis, K. R., and Liao, J.: The Framework for 0-D Atmospheric Modeling (F0AM) v3.1, *Geosci. Model Dev.*, 9, 3309–3319, DOI: 10.5194/gmd-9-3309-2016, 2016.
- Womack, C.C., Martin-Drumel, M.A., Brown, G.G., Field, R.W. and McCarthy, M.C.: Observation of the simplest Criegee intermediate CH_2OO in the gas-phase ozonolysis of ethylene. *Sci. Adv.*, 1(2), DOI: 10.1126/sciadv.1400105, 2015.
- 450 Wood, E.C. and Charest, J.R.: Chemical amplification - cavity attenuated phase shift spectroscopy measurements of atmospheric peroxy radicals. *Anal. Chem.* 86, 10266–10273. DOI: 10.1021/ac502451m, 2014.
- Wright, P.J. and English, A.M.: Scavenging with TEMPO• to identify peptide-and protein-based radicals by mass spectrometry: Advantages of spin scavenging over spin trapping. *Journal of the American Chemical Society*, 125(28), 8655–8665, DOI: 10.1021/ja0291888, 2003.
- 455 Zaytsev, A., Breitenlechner, M., Koss, A. R., Lim, C. Y., Rowe, J. C., Kroll, J. H., and Keutsch, F. N.: Using collision-induced dissociation to constrain sensitivity of ammonia chemical ionization mass spectrometry (NH_4^+ CIMS) to oxygenated volatile organic compounds, *Atmos. Meas. Tech.*, 12, 1861–1870, DOI: 10.5194/amt-12-1861-2019, 2019.
- Zhao, Y., Thornton, J.A., and Pye, H.O.: Quantitative constraints on autoxidation and dimer formation from direct probing of monoterpene-derived peroxy radical chemistry. *Proceedings of the National Academy of Sciences*, 115, 12142–12147, DOI: 10.1073/pnas.1812147115, 2018.
- 460




Table 1: Proton affinities (PAs) of HFA, water and secondary ozonides produced in reactions of SCIs with HFA. Species with PAs higher than that of water can be detected in H_3O^+ CIMS.

Species	PA, kcal/mol	Reference
$\text{CH}_2\text{OO}\cdot\text{HFA}$	662.9	This work
HFA	670.4	Hunter and Lias (1998)
H_2O	691	Hunter and Lias (1998)
$\text{CH}_3\text{CH}_2\text{CHOO}\cdot\text{HFA}$	720.7	This work
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CHOO}\cdot\text{HFA}$	730.8	This work
$(\text{CH}_3)_2\text{COO}\cdot\text{HFA}$	747.2	This work
$(\text{CH}_2=\text{C}(\text{CH}_3))\text{CHOO}\cdot\text{HFA}$	779.6	This work




465 **Figure 1: Mechanism of tetramethylethylene (TME) ozonolysis. Stabilized Crigee intermediate (shown in red) can be scavenged by the chemical derivatization agent HFA or the spin trap DMPO, or decompose to peroxy radical (shown in yellow) and OH. RO_2 and OH species can in turn react with spin traps. Reactions involving SCI are from MCM v3.3.1 (Jenkin et al., 1997) and Giorio et al. (2017).**




470

Figure 2: Schematic of experimental setup used to detect SCIs and RO₂ with the spin trap DMPO. DMPO was introduced in the experimental setup using a liquid calibration unit (LCU, IONICON Analytik).




475

Figure 3: Ion tracers observed in a TME ozonolysis experiment as a function of different reactant conditions. Reactant concentrations are [TME] = 1.85×10^{12} ; [O₃] = 1.67×10^{13} ; [HFA] = 6.09×10^{15} molecule cm⁻³. (CH₃)₂COO·HFA·H⁺ ion (red tracer, *m/z* 241.03) is observed when TME, HFA, and O₃ are present in the system.



480 **Figure 4: Correlation plot comparing measured and calculated concentrations of $(\text{CH}_3)_2\text{COO}\cdot\text{HFA}$. The adducts were detected using H_3O^+ CIMS as $(\text{CH}_3)_2\text{COO}\cdot\text{HFA}\cdot\text{H}^+$ (m/z 241.03). The slope is calculated using weighted least squares (WLS). A 95% confidence interval is estimated via a Monte Carlo simulation ($N=5000$) and shown using red shading.**



485 **Figure 5: OH reactivity measured as a function of TEMPO concentration. The slope determining the reaction rate coefficient $k_{\text{TEMPO}+\text{OH}} = (9.3 \pm 0.9) \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ is calculated using weighted least squares (WLS). A 95% confidence interval is estimated via a Monte Carlo simulation ($N=5000$) and shown using blue shading. The intercept $(5.1 \pm 2.4) \text{ s}^{-1}$ can be explained by other OH reactants such as O_3 , NO , NO_2 , and CO .**






Figure 6: Ion tracers observed by NH_4^+ CIMS in a TME ozonolysis experiment as a function of different reactant conditions. Reactant concentrations are $[\text{TME}] = 3.69 \times 10^{11}$; $[\text{O}_3] = 7.87 \times 10^{12}$; $[\text{DMPO}] = 2.01 \times 10^{12}$ molecule cm^{-3} .



490 Figure 7: Correlation plot comparing measured and calculated concentrations of $\text{CH}_3\text{C}(=\text{O})\text{CH}_2\text{OO}\cdot\text{DMPO}$. The adducts were detected using NH_4^+ CIMS as $\text{CH}_3\text{C}(=\text{O})\text{CH}_2\text{OO}\cdot\text{DMPO}\cdot\text{NH}_4^+$ (m/z 220.142). The slope is calculated using weighted least squares (WLS). A 95% confidence interval is estimated via a Monte Carlo simulation ($N=5000$) and shown using yellow shading.



495 **Figure 8: Mass spectra of SCI-DMPO (red) and RO₂-DMPO (yellow) adducts in α -pinene ozonolysis experiments observed using (a) NH_4^+ CIMS and (b) H_3O^+ CIMS. Primary RO₂ species ($\text{C}_{10}\text{H}_{15}\text{O}_4$) are formed via CI isomerization and can in turn undergo various autoxidation reactions resulting in formation of several organic peroxy radicals ($\text{C}_{10}\text{H}_{15}\text{O}_x$, x=5-9), which were detected as adducts with the spin trap DMPO by the means of NH_4^+ and H_3O^+ CIMS.**