

Overview:

Wang et al have submitted a manuscript pertaining to ground based infrared-FTIR measurements of ammonia (NH₃) at Hefei, China. Two years (Dec 2016 to Nov 2018) of spectral measurements and retrieval processes are detailed, followed with a comparison of the retrieved NH₃ total columns to that of co-located NH₃ column measurement from the satellite-based IASI instruments. Thereafter the ground-based NH₃ columns are used in conjunction with other datasets (in situ carbon monoxide, meteorological data, traffic density) and a back-trajectory modelling study to help attribute NH₃ emissions to specific source types. The authors conclude that agricultural NH₃ emissions are the main source in spring, autumn and winter, and urban anthropogenic emission (vehicles) dominate in summer. Over the two-years of measurements, an increase of 22.14% in NH₃ is reported.

The novelty of this study is that this is the first-time ground based FTIR NH₃ measurements at Hefei are reported and compared to satellite data. The high-quality ground based NH₃ dataset will be a welcome and important addition to international scientific databases. The manuscript is overall logically structured and well referenced. The writing style is fluent and easily understood. Information on data availability is given and there are no registered conflicts of interest.

Improvements can be made to the manuscript. In its current form I do not recommend publication without major revision to address issues concerning the methodology, trend analysis and the source emission attribution (especially sections 3.4 and 3.5).

The manuscript can be seen as comprised of two parts, that describing the ground based NH₃ measurements and satellite comparison, and that of NH₃ source attribution using the NH₃ column measurements, meteorological & air quality datasets, and back trajectory modelling. The first part is within scope of the AMT journal whereas the NH₃ source emission analysis and hypothesis testing in the second part is investigative.

The source emission work is not a simple investigation to support novel measurement validity. To this end, is outside (or bordering on) the scope of AMT. Due to the high level of source attribution investigation and critical issues raised surrounding the analysis of source emission attribution I recommend that this manuscript be either:

- 1) split into two manuscripts, along the lines of the two parts mentioned previously. The first part submitted to a technical atmospheric journal and the second part submitted to an atmospheric sciences/chemistry journal, or
- 2) submitted to an atmospheric sciences journal with the focus on the source attribution work.

It is my opinion that the work presented in part 1 requires only minor/moderate changes and technical clarification whereas part 2 requires substantial revision to support the interpretation of the findings or a possible reinterpretation of findings.

Commentary and questions are listed below, and referenced to the documents: amt-2020-39.pdf and amt-2020-39-supplement.pdf

Specific comments:**S1/ Trend analysis.**

It states in the manuscript that there is a 22.14%/yr annual trend in NH₃ (L32, L261, L282, L395, L402-L403). This is calculated from the relative difference between two annual means. I think it is a bit premature to calculate an annual trend based on only 2 data points. The difference could be a step function instead of a trend. I recommend that an annual trend is not reported, but the authors may

wish to state the annual means of the two years. It is too speculative at this stage to report a trend, wait for a longer time series.

S2/ Comparison of ground-based and IASI measurements.

L294: "Since the IASI-NH3 retrieval does not provide averaging kernels and vertical profiles (Van Damme, et al 2014a), this method for comparison is not applied. Here we therefore compare the IASI satellite and FTIR data directly, without considering the effect of different a priori profiles and averaging kernels."

Such an issue was also confronted by Dammers, et al. 2016., see section 2.3.2. The authors could use the same approach on a subset of the Hefei data, and comment if this alters comparison results. This does assume proxy profiles that are scaled, again see Dammers et al. 2016 for details.

Dammers, E., et al., An evaluation of IASI-NH3 with ground-based Fourier transform infrared spectroscopy measurements, Atmos. Chem. Phys., 16, 10351–10368, 525 <https://doi.org/10.5194/acp-16-10351-2016>, 2016.

S3/ The descriptive language of correlation coefficients.

There are inconsistencies in the use of descriptive language used in commentary of correlation coefficients. In the two sentences below, correlations of 0.47 and 0.48 are described as both 'weak' and 'high'.

L345: "NH3 columns show weak correlation (R=0.47) with PM2.5 concentrations"

L359: "Further, the scatter plot of NH3 columns with air temperature in spring and autumn season displays relatively high correlation, with correlation coefficients of 0.53 and 0.48".

Furthermore, at L417: "there was a clear correlation between NH3 columns and air temperature in spring and autumn over Hefei, with correlation coefficients of 0.53 and 0.48, respectively."

I recommend that a standard way of describing the correlations be adopted to achieve consistency. For instance, below is a table from Schober, et al.

Table. Example of a Conventional Approach to Interpreting a Correlation Coefficient	
Absolute Magnitude of the Observed Correlation Coefficient	Interpretation
0.00–0.10	Negligible correlation
0.10–0.39	Weak correlation
0.40–0.69	Moderate correlation
0.70–0.89	Strong correlation
0.90–1.00	Very strong correlation

Schober, Patrick, Christa Boer, and Lothar A. Schwarte. "Correlation coefficients: appropriate use and interpretation." Anesthesia & Analgesia 126.5 (2018): 1763-1768.

Using this example, all the correlations mentioned above would be described as moderate.


S4/ Calculation of correlation coefficients.

Could the authors also state the method used to calculate the correlation coefficients, most importantly did the method use uncertainty estimates in both the abscissa and ordinate data points, as both can have large uncertainties.

S5/ NH3 Air temperature correlation.

L352: “High correlation of NH3 columns with air temperature is obvious from their diurnal variation during the observation period, as seen in Figure 8.”

From figure 8, I could not see a ‘high’ correlation. The calculated correlation coefficient is also not given. I read data values from figure 8 (and assuming no uncertainty in the read values), then made a scatter plot and calculated the correlation coefficient. See the figure below. I calculated $R^2 \approx 0.08$. This dataset is imprecise and just an indicator. Could the authors please provide the correlation coefficient from the proper dataset and remark if they think the correlation is high. If not, how does this affect the conclusions the authors have made. In my opinion, fig 8 shows a negligible correlation between NH3 and temperature over the analysed averaged time period. There are two spikes in NH3, in the mid-morning and mid-afternoon. Does this correlate with traffic volumes or traffic patterns?



Lastly figure 9 should also include the winter and summer correlation plots, like that for spring and autumn.

S6/ Section 3.4 and 3.5: NH3 and CO industrial emissions.

Industrial ammonia:

Throughout the manuscript there is a lack of detail and discussion on possible industrial (point source) emissions of NH3 (such as in Wang, et al, 2015). In section 3.4 analysis only concerns NH3 from vehicle sources. Section 3.5 talks about agricultural emissions (transported into the Hefei region) with a passing mention of “urban anthropogenic emissions” (L388). The only specific mention of point sources is in figure 1. Two large NH3 point sources are identified based on Table A1 of Clarisse et al. 2019.


Clarisse, L., Van Damme, M., Clerbaux, C., and Coheur, P.-F.: Tracking down global NH3 point sources with wind-adjusted superresolution, Atmos. Meas. Tech., 12, 5457–5473, <https://doi.org/10.5194/amt-12-5457-2019>, 2019.

Wang, S., Nan, J., Shi, C., Fu, Q., Gao, S., Wang, D., Cui, H., Saiz-Lopez, A., and Zhou, B.: Atmospheric ammonia and its impacts on regional air quality over the megacity of Shanghai, China, Sci. Rep., 5, 15842, <https://doi.org/10.1038/srep15842>, 2015.

Industrial CO emissions:

L329 “To examine the contribution of traffic to NH3 columns, we analyze the relationship of NH3 columns with CO surface concentrations” and L342 “Atmospheric CO is regarded as a primary

pollutant mainly emitted from vehicles in urban areas, and there is no significant biomass burning source around the Hefei site, thus the elevated NH₃ columns are likely partly caused by urban emissions from vehicles.”. Again, this is a lack of detail on CO industrial (point source) emissions and the effect this would have on any NH₃-CO correlations. For example, there is a 700MW coal-fired power station ([https://www.gem.wiki/Hefei-2 power station](https://www.gem.wiki/Hefei-2_power_station)) to the east of Hefei (~8km), the red flag in the image below, and two power stations in the Hefei region identified in Tan, et al. 2019



Without detailed analysis that includes probable/possible industrial sources of NH₃ and CO, conclusions drawn in sections 3.4 and 3.5 are highly speculative and qualitative (using vague phrases such as “likely partly” (L344)).

Tan, W., Zhao, S., Liu, C., Chan, K., L., Xie, Z., Zhu, Y., Su, W., Zhang, C., Liu, H., Xing, C., and Liu, J., Estimation of winter time NO_x emissions in Hefei, a typical inland city of China, using mobile MAX-DOAS observations, Atmos. Environ., 200, 228-242, <https://doi.org/10.1016/j.atmosenv.2018.12.009>, 2019.

S7/ Traffic pollutant correlations.

L336: “The results show that there exists no correlation between NH₃ columns and concentrations of NO₂, SO₂, PM₁₀ and the value of AQI over the four seasons (not shown).”

Mathematically, there is always a correlation coefficient, if the correlation is less than 0.1, maybe better to state correlation is negligible. Even better, would be to list the correlation coefficient value after each species, i.e. NO₂ (XX), SO₂ (XX) etc...

It is interesting to read that there is no (negligible) correlation with NO₂. NO_x (NO₂ + NO) is a primary pollutant from vehicle emissions (Zhou et al., 2014, Phillips et al, 2019). What does a NH₃-CO moderate correlation with a negligible NH₃-NO₂ correlation indicate? Could the authors please comment on the reported NH₃-NO₂ correlation.

Zhou, Rui, et al. "Study on the traffic air pollution inside and outside a road tunnel in Shanghai, China." PloS one 9.11 (2014).

Phillips, F. A., et al.: Vehicle ammonia emissions measured in an urban environment in Sydney, Australia, using open path Fourier Transform Infra-Red Spectroscopy. Atmosphere, 10, 208, <https://doi.org/10.3390/atmos10040208>, 645 2019.

There is also no quantitative measurand of traffic volumes. If NH₃ emissions were primarily (or even partly) from vehicle emissions, then more in-depth quantitative analysis is required.

S8/ NH₃ and PM_{2.5}.

L345: “Meanwhile, NH₃ columns show weak correlation ($R=0.47$) with PM_{2.5} concentrations (Fig. 7(b)), meaning that NH₃ contributed to the formation of fine particulates significantly in summer.”.

This qualitative statement needs expanding. More in-depth analysis of the Hefei dataset is needed. A more quantitative investigation is required into PM_{2.5} and NH₃ correlation and/or causation. It is well known NH₃ contributes to PM_{2.5}.

S9/ Column to point correlations

The NH₃ measurements are partial columns extending from ~0-4km in altitude. The meteorological and air quality datasets are from point source ground level measurements, located a distance from the NH₃ measurements. These two datasets will have quite different footprints. The in-situ ground-based measurements will have a more localised footprint, whereas the column measurements will have a larger regional footprint. Could the authors comment on this fact and how it affects correlation interpretation.

S10/ Agricultural NH₃ emissions

L362 – 367: “In Hefei, the cropland in spring is characterized by the growing season of wheat and early rice, and synthetic fertilizer is applied during this period. The fertilizer application also tends to occur in autumn, when winter wheat is sowed, and a second rice crop is growing. The agricultural practice may explain the correlation between NH₃ columns and temperature over Hefei over these two seasons, which suggests that agriculture was the main source of ammonia in spring and autumn.”

This a good starting hypothesis to investigate, but it is left as a conjecture. Too vague language is used, e.g. “May explain” and “suggests”. These are good starting points, but quantitative substantial evidence is required. This is lacking. Seasonal (hence temperature) correlations is not a strong causal link without additional collaborating evidence. The authors need to add such evidence to prove the statement in the manuscript.

S11/ Windrose

In such wind rose polar plots the wind strength is usually plotted as the radial component and the NH₃ abundance as coloured points. I would strongly recommend replotting with wind speed as the radial axis of the polar plot and NH₃ concentrations as coloured dots.

Looking at the current plots, the greatest wind speed is less than 4.5ms^{-1} . This is classified as a gentle breeze using the Beaufort scale. This indicates that the majority of measured NH₃ is from local sources (assuming hourly- low daily NH₃ lifetimes). The highest concentrations of NH₃ is measured during easterly winds, the direction of the main Hefei urban area and the lowest concentrations from the west, a less populated and dense area. At L368 the authors comment on the wind rose data, but do not use it to help attribute the urban emission sources (traffic vs industrial). High NH₃ concentration points are in the NW quadrant, the direction of an NH₃ point source shown in figure 1. There is no commentary on this fact.

S12/ NH₃ lifetime.

There is only passing mention of NH₃ lifetime (L55). Is there an expected difference in NH₃ lifetime depending on emission source (i.e. rural vs industrial) and would this influence NH₃ source attribution?

S13/ Back-trajectory calculations using HYSPLIT.

At L380 the authors introduce a new method to help assess NH₃ source emissions. Back trajectory modelling is introduced, but I cannot understand why the bulk of the findings are relegated to the supplement. I recommend that the supplement content and the main manuscript from L380 -390 be concatenated into a new section, e.g. section 3.6.

In addition, a much more detailed explanation of the model set up (meteorological fields, resolution, cluster analysis details, etc) and analysis (source attribution) needs to be undertaken. Example: the PSCF's in figure S2 are hard to see in detail and require further inquiry. The interpretation of the results at L386-390, and the supplement, need more rigorous assessment in a scale more regional to Hefei (due to low wind speeds). There is no mention of carbon monoxide transport either, and the effect this would have on NH₃-CO correlations and source attribution.

S14/ Concluding statements in section 4.

This section requires major revision based upon the issues raised in this review.

Currently there is not enough collaborating evidence in the manuscript to explicitly state with certainty the origin and abundance of NH₃ emissions measured at Hefei.

Minor/Technical:

M1/ L30: "emission sources and potential sources". What is the difference between emission and potential sources? Do you mean known emission sources and unknown emission sources? Can you please make this clearer and correct through-out the manuscript?

M2/ L34: "The results demonstrate the IASI data are in broad agreement with our FTIR data.". I think it best to rephrase as "The results demonstrate that the IASI and FTIR data, over Hefei, are in broad agreement.".

M3/ L64: "This is mainly caused by lack of representative measurements of atmospheric NH₃". Could a reference to this statement be added?

M4/ NH₄⁺: There is no mention of NH₄⁺ with respect to NH₃ partition equilibrium in NH_x, and how this will affect the measured NH₃ abundances. Could the authors please comment on this aspect. Lachatre, et al, 2019 would be a good starting point.

M5/ L68: There are ~12 references to a single statement. Can this be cut down to only the essential references needed. Passive sampling is not the focus of this manuscript.

M6/ L71: Pedantic but...could 'Sticky' be replaced with a more scientific descriptive term, or place in quotation i.e. 'sticky'.

M7/ L78: Replace 'ground-based' with 'ground level'

M8/ L120: "Furthermore, ground based observations of NH₃ have been sparse throughout China (Liu et al., 2011; Xu et al., 2015)." Ground based or ground level in situ?

M9/ L132: Replace "reduces" with "reductions of"

M10/ L145: I think "Materials" is the incorrect word, maybe "Measurements" would be better.

M11/ L164: "liquid-nitrogen-cooled MCT/InSb detector". Just to clarify, is this a single detector or dual HgCT and InSb detectors?

M12/ L165: Replace 'suit' with 'suite'

M13/ L171: "At the same time, the indoor pressure, temperature and relative humidity are logged continuously.". Are these important parameters for the FTIR measurements?

M14/ L173: "similar to the NH₃ retrieval strategies in Dammers et al. (2015)". What are the differences? Are the Microwindows the same?

M15/ L181: "A priori profiles of NH₃ and interfering gases are taken from the Whole Atmosphere Community Climate Model (WACCM, v.6_120_99) in combination with initial measurement values."

There is no NH₃ modelling output from WACCMv6. Could the authors please correctly reference the NH₃ apriori source used.

Is a static apriori used? Or does it change throughout the retrieval time series?

M16/ L181: "...in combination with initial measurement values.". Sorry, I do not understand this. Can you please explain what initial measurements are, and how they are combined with (so-called) WACCMv6 profiles to obtain an apriori?

M17/ L182: "WACCM, v.6_120_99". I think this can be referenced better. '_120_99' is not part of the WACCM simulation identify code. Maybe better to reference it this way: (WACCMv6, Garcia et al. 2007).

Garcia, R. R., Marsh, D. R., Kinnison, D. E., Boville, B. A., and Sassi, F.: Simulation of secular trends in the middle atmosphere, 1950-2003, 20 J. Geophys. Res., 112, D09301, doi:10.1029/2006JD007485, 2007.

M18/ L183: "for ammonia was constructed to be diagonal, with standard deviations of 100% for all layers." What was the basis for assuming an uncertainty of 100% and is there any inter-layer covariance/correlation constraint?

M19/ I would also encourage the authors to include another sentence (or two) after the description of the apriori profile and uncertainties describing other state vector retrieval parameters such as phase fitting, instrument line shape, continuum fit, zero level offset etc.

M20/ Also...could be authors also state the SNR (or S_e) of the measurements, and either if the SNR is held static or varies on a spectrum by spectrum basis.

M21/ How is the H₂O apriori set? For instance, is it from NCEP daily profiles, or from WACCM, or from a previous H₂O retrieval?

M22/ L190: "about 0.498 % and 0.505%", could these values be rounded to the correct number of significant figures.

M23/ Fig 2 and L188: "The measured spectrum is shown in blue, the fitted spectrum in red and the residual in black". There are two blue lines, 'observed' and H₂O. Either state as 'dark blue' or switch colour.

M24/ I recommend that section 3.1.2 lines L238 to 246 be moved to the end of section 2.2. This is because it details retrieval information content and logically comes after spectral fitting.

M25/ L239: "It is evident that the retrieval is most sensitive to the troposphere, where the concentration of ammonia peaks." Add in 'lower': "It is evident that the retrieval is most sensitive in the lower troposphere, where the concentration of ammonia peaks"

M26/ L243: "The degrees of freedom for signal (DOFS) value is 1.10 given by this measurement". Can the authors indicate at what height, from the surface up, does the DOFS reach 1.0, essentially showing the partial column range in which there is a signal?

M27/ Fig 4. The individual layer averaging kernels are all the same colour, very confusing. Maybe only plot for layers 0-10km. There are also no units for the VMR averaging kernels (I assume VMR/VMR)? Also, for the total column averaging kernel, what are the units?

M28/ Section 3.1.1 could be transferred to be section 2.3. (IASI becomes section 2.4). It makes more sense that the entire retrieval process from retrieval set up to retrieval output then error analysis is all together in one section.

M29/ L215: "The error calculation is based on attributing uncertainties to all parameters used in the profile retrieval". Maybe better stated as "The error calculation is based on attributing uncertainties to all parameters in the retrieval state vector".

M30/ L221: "Table 1 lists the uncertainties of the parameters assumed in the retrievals." Maybe better stated as: "Table 1 lists the uncertainties of the retrieval state vector parameters."

M31/ L222: "The total errors are about 11.42 % based on the combination of random and systematic errors". Is error analysis applied to each individual retrieval? If so, the mean total error and 1-sigma standard deviation of the time series could be given. This would be a better indication of the overall timeseries uncertainty than a single retrieval.

M32/ Table 1:

Could the authors please state where or how the temperature profile uncertainties were set/calculated.

Did the authors calculate the uncertainty associated with inferring species spectral uncertainties? i.e. the line intensity (and pressure and temperature broadening) uncertainties in CO₂, H₂O etc...

Could 'NH₃' be put in front of Line Intensity, and P, T broadening (broadening, not broadening). Could the authors also reference the origin of the 10% spectral uncertainties.

M33/ Section 3.1.2 and Fig 3: With Section 3.1.1 moved to section 2, along with the second half of section 3.1.2, this leaves a single paragraph (L230-237) in section 3.1.2. This paragraph could be moved into section 3.2. There is very little vertical information due to the low DOFs, basically profile scaling.

Figure 3 offers little information due to the low DOFs. The variation within a season is not presented so it is hard to conclude if the seasonal differences in profile are significant. Maybe reconstruct figure 3 to have 4 plots (2 by 2), one for each season and for each season include the spread (1-sigma SD?) at each level.

M34/ L235: "The seasonal averaged surface level of NH₃ decreased from 10.82 ppb in summer to 2.92 ppb in winter during 2017 and 2018, and the corresponding values are about 5.48 and 6.04 ppb in spring and autumn, respectively.". It would be advantageous if for each season the mean and spread (1-sigma SD) was given.

M35/ L250: “Many spectra ranging from 700 to 1350 cm^{-1} are saturated in summer (due to high humidity), causing the retrieved NH_3 data to be sparsely sampled relative to those in other seasons”. MW1 ($\sim 929.6\text{-}931.2\text{cm}^{-1}$) (fig 2a) shows small H_2O absorption features, as opposed to MW2. In summertime, could plausible retrievals be performed with only using MW1 without altering the DOFS too much? This may help increase the number of observations.

M36/ L276: “It follows therefore that seasonal variation of NH_3 columns in the Hefei area accords with that in other areas in China, with the main emission source being agriculture.” This sentence is not needed as a near repeat of the statement at L263.

M37/ L287: Maybe rewritten as “Here we present a comparison of the ground-based NH_3 column measurements with the IASI satellite measurements”.

M38/ L290. How was the co-location criteria of 0.5 degrees and 90 minutes decided upon?

M39/ L303: “The Relative differences larger than 100% were considered as outliers from the data”. On what basis, and why the value of 100%? Was the ground based or satellite measurements anomalous?

M40/ L305: “with standard deviation of 44.44% and 41.00%, respectively.”. Maybe 44% and 41%.

M41/ L308: “Additionally, the distributions of the relative difference of the two dataset show that, the relative bias mainly range from -60% to 80% for IASI A data, from -60% to 60% for IASI B data, and the bias from -20% to 0% as well as -40% to -20% has the highest frequency in both respective bins (Figure 6 (c) and (d)).”

Bias refers to the overall difference between the two datasets, so I think it is better worded as:

“Additionally, the distributions of the relative difference of the two datasets show that the relative differences of a measurement pair, mainly range from -60% to 80% for IASI A data, from -60% to 60% for IASI B data as seen in figure 6 (c) and (d).”

M42/ L313L First time the acronym NDACC is used. For the first instance can the full title please be used. Also maybe better stated as: “nine NDACC FTIR stations”.

M43/ L317: “The correlation coefficient (R) between the CrIS NH_3 total columns and FTIR data was 0.77, and the relative difference is 0-5 % with a standard deviation of 25–50 % for comparison of high levels of NH_3 .” Sorry I do not understand the ‘0-5%’ statement, can you please explain it more clearly, how can you have multiple mean differences?

M44/ L320: “results from other NDACC site data”. Is Hefei a NDACC affiliated FTIR site? If not, then the current wording can be read that it is. Maybe replace with “comparison results in the aforementioned literature”

M45/ L330: “The Dongpu Reservoir air quality monitoring station (31.91°N , 117.16°E) is very close to our site, part of a National Ambient Air Quality Monitoring Network, which monitors and routinely publishes the concentrations of main gaseous pollutants, including CO , NO_2 , $\text{PM}_{2.5}$, PM_{10} , SO_2 , O_3 and Air Quality Index (AQI) etc”. Could the distance from the air quality station to the FTIR site please be stated and a reference added pertaining to the National Ambient Air Quality Monitoring Network, (esp. for the Hefei site if possible). Could the Dongpu site please be marked in figure 1b.

M46/ L376: “Overall, the results indicate that air temperature, wind direction and wind speed are the main factors that influence gaseous NH_3 concentrations in Hefei.”. What about source emissions?

M47/ L393: the adjective 'great' can be omitted.

M48/ L415: replace "meteorological parameters" with "meteorological conditions".

M49/ L432: "This is the first time that ground-based FTIR remote sensing of NH₃ columns and comparison with satellite data are reported in China". A valid and important finding and should be reported at the start of the conclusion section.

M50/ L405: "To validate the satellite observations...". In this case since the satellite measurements have been compared against other measurements favourably, the satellite measurements are used to confirm/validate the Hefei site ground base measurements.

M51/ Fig 1: Could longitude and latitude units please be put on both axis. The area displayed could also be reduced to focus on the eastern and middle region of China. In map B, could the Dongpu AQ station also be identified. Could the location identifier 'dots' also be bigger in both maps. Could a distance scale also be put on the map?

Caption: "The regional distributions of NH₃ columns (molec cm⁻²) from 2008-2018 IASI-A and 2013-2018 IASI-B morning overpasses of ANNI-NH₃-v3R data". Are these annual mean columns? Good idea to specify the temporal period.

M52/ Fig 2a and b: the title consists of RMS. If possible, add in the DOFS and CHI² fit variables as well for completeness.

M53/ Fig 5: Could 1-sigma standard deviation spread/error bars be put on the daily averages.

M54/ Fig 6: A lot of white space in plots A & B, could the X-axis range be shortened, MAX X \sim 8e16.

M55/ Fig 7: Could the uncertainty bars be added for each point, like that in fig 6 a and b.

Caption: "Scatter plot of NH₃ columns (molec cm⁻²) with CO (mg m⁻³, a) and PM_{2.5} (μ g m⁻³, b) concentrations measured at the Dongpu Reservoir air quality monitoring site in summer." Are these daily averages? Please add in temporal period.

M56/ Fig 9: Caption: "Scatter plot of NH₃ column (molec cm⁻²) with air temperature ($^{\circ}$ C) in spring (a) and autumn (b)." Again, please add temporal period (hourly, daily, monthly...).

M57/ Fig S1 and S2, it is hard to ascertain information from both plots. As a first step, could the map lat/lon boundaries be reduced to concentrate on areas of interest better. I.e. zoomed in a bit.

M58/

Franco, B. Clarisse, L. , Stavrakou, T., Müller, J.-F, Van Damme, M. ,Whitburn, S., Hadji-Lazaro, J. , Hurtmans, D., Taraborrelli, D. , Clerbaux, C. , Coheur, P.-F.: A General Framework for Global Retrievals of Trace Gases From IASI: Application to Methanol, Formic Acid, and PAN, J. Geophys. Res.-Atmos., 123, 963-984, <https://doi.org/10.1029/2018JD029633>, 2018.

Is this reference relevant?