Long-term column-averaged greenhouse gas observations using a COCCON spectrometer at the high surface albedo site Gobabeb, Namibia

Matthias M. Frey^{1,a}, Frank Hase¹, Thomas Blumenstock¹, Darko Dubravica¹, Jochen Groß¹, Frank Göttsche¹, Martin Handjaba², Petrus Amadhila², Roland Mushi², Isamu Morino³, Kei Shiomi⁴, Mahesh Kumar Sha⁵, Martine de Mazière⁵, and David F. Pollard⁶

¹Karlsruhe Institute of Technology (KIT), Institute for Meteorology and Climate Research (IMK-ASF), Karlsruhe, Germany
 ²Gobabeb Namib Research Institute, Gobabeb, Namibia
 ³National institute for Environmental Studies (NIES), Tsukuba, Japan
 ⁴Japan Aerospace Exploration Agency (JAXA), Tsukuba, Japan
 ⁵Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

⁶National Institute of Water and Atmospheric Research (NIWA), Lauder, New Zealand

^anow at: National institute for Environmental Studies (NIES), Tsukuba, Japan

Correspondence: M. Frey (frey.matthias.max@nies.go.jp)

Abstract. In this study we present column-averaged dry-air mole fractions of CO_2 (XCO₂), CH_4 (XCH₄) and CO (XCO) from a recently established measurement site in Gobabeb, Namibia. Gobabeb is a hyperarid desert site at the sharp transition zone between the sand desert and the gravel plains, offering unique characteristics with respect to surface albedo properties. Measurements started in January 2015 and are performed utilizing a ground-based Fourier transform infrared (FTIR)

- 5 EM27/SUN spectrometer of the COllaborative Carbon Column Observing Network (COCCON). Gobabeb is the first measurement site observing XCO₂ and XCH₄ on the African mainland and improves the global coverage of ground-based remotesensing sites. In order to achieve the high level of precision and accuracy necessary for meaningful greenhouse gas observations, we performed calibration measurements for eight days between November 2015 and March 2016 with the COCCON reference EM27/SUN spectrometer operated at the Karlsruhe Institute of Technology. We derived scaling factors for XCO₂, XCH₄ and
- 10 XCO with respect to the reference instrument that are close to 1.0. We compare the results obtained in Gobabeb to measurements at Reunion Island and Lauder from the Total Carbon Column Observing Network (TCCON) sites at Reunion Island and Lauder. We choose these TCCON sites because, while 4000 km apart, the instruments at Gobabeb and Reunion Island operate at roughly the same latitude. The Lauder station is the southernmost TCCON station and functions as a background site without a pronounced XCO₂ seasonal cycle. We find a good agreement for the absolute Xgas valuesand representative diurnal
- ¹⁵ variability-, apart from an expected XCH_4 offset between Gobabeb and Lauder due to significantly different tropopause height, as well as representative intraday variability between TCCON and COCCON. Together with the absence of long term drifts, this highlights the quality of the COCCON measurements. In the Southern hemispheric summer we observe lower XCO_2 values at Gobabeb compared to the TCCON stations, likely due to the influence of the African biosphere. We performed coincident measurements with the Greenhouse Gases Observing Satellite (GOSAT), where GOSAT observed three nearby specific
- 20 observation points, over the sand desert south of the station, directly over Gobabeb and over the gravel plains to the north.

GOSAT H-gain XCO_2 and XCH_4 agree with the EM27/SUN measurements within the 1 σ uncertainty limit. The number of coincidence-coincident soundings is limited, but we confirm a bias of 1.2 - 2.6 ppm between GOSAT M-gain and H-gain XCO_2 soundings-retrievals depending on the target point. This is in agreement with results reported by a previous study and the GOSAT validation team. We also report a bias of 5.9 - 9.8 ppb between GOSAT M-gain and H-gain XCH_4 measure-

- 25 ments which is within the range given by the GOSAT validation team. Finally we use the COCCON measurements to evaluate inversion-optimized CAMS model data. For XCO_2 we find high biases of 0.9 ± 0.5 ppm for the OCO-2 assimilated product and 1.1 ± 0.6 ppm for the in situ-driven product with $R^2 > 0.9$ in both cases. These biases are comparable to reported offsets between the model and TCCON data. The OCO-2 assimilated model product is able to reproduce the drawdown of XCO_2 observed by the COCCON instrument at the beginning of 2017, as opposed to the in situ-optimized product. Also, for XCH_4 ,
- 30 the observed biases are in line with prior model comparisons with TCCON.

1 Introduction

In 2019, the The concentrations of the most important anthropogenic greenhouse gases (GHGs), carbon dioxide (CO_2) and methane (CH_4), have risen to unprecedented values values unprecedented since the beginning of high-frequency observational records (Dlugokencky et al., 2019a, b). Additionally, it was stated recently that recently fossil CO_2 emissions exceeded

- 35 10 GtCyr⁻¹ for the first time in history (Friedlingstein et al., 2019). Precise and accurate global observations of GHGs are therefore important for the estimation of emission strengths, flux changes (Olsen and Randerson, 2004) and model evaluation. Furthermore, these measurements can be directly-used for the verification of climate mitigation actions as demanded by international treaties, e.g. the Paris COP21 agreement (https://unfccc.int/files/essential_background/convention/application/pdf/ english_paris_agreement.pdf, last access: 15 October 2020).
- 40 Satellites like the Greenhouse Gases Observing Satellite (GOSAT) (Kuze et al., 2009; Morino et al., 2011; Yoshida et al., 2013), Orbiting Carbon Observatory-2 (OCO-2) (Frankenberg et al., 2015; Crisp et al., 2017; Eldering et al., 2017), Orbiting Carbon Observatory-3 (OCO-3) (Eldering et al., 2019), SENTINEL5-Precursor (S5P) (Veefkind et al., 2012) or Greenhouse Gases Observing Satellite-2 (GOSAT-2) (Suto et al., 2020) are well suited candidates for this task as they retrieve total column abundances of atmospheric GHGs on a global scale. However, current satellites, while offering quasi-global spatial coverage,
- 45 have coarse temporal resolution. The OCO-2 repeat cycle is 16 days, the GOSAT-2 repeat cycle is 6 days. S5P offers daily global coverage of CH_4 and CO. However; however, the measurements are mostly around local noon time. Future geostationary satellites will likely help to overcome this shortcoming (Moore III et al., 2018; Nivitanont et al., 2019). Due to the fact that satellites measure backscattered sunlight from the surface of the earth and its atmosphere, retrievals of GHGs are complicated and biases can easily occur, which need to be recognized and - if possible - corrected. Therefore satellite measure-
- 50 ments are commonly validated against ground-based remote-sensing instruments, as these measurements are not influenced by surface albedo effects and only minimally affected by aerosols (Dils et al., 2014; Wunch et al., 2017). The Total Carbon Column Observing Network (TCCON) is a ground-based network retrieving total columns of GHGs with reference precision and accuracy utilizing high-resolution solar-viewing Fourier transform infrared (FTIR) spectrometers (Wunch et al., 2011;

Washenfelder et al., 2006). TCCON is the reference instrument network and primary validation source for current satellites

55 (Inoue et al., 2016; Wu et al., 2018; Borsdorff et al., 2018)(Inoue et al., 2016; Wu et al., 2018; Borsdorff et al., 2018; Sha et al., 2021)

Recently, in an effort to further improve the global coverage of ground-based observations, the COllaborative Carbon Column Observing Network (COCCON) was established (Frey et al., 2019). This network employs compact, portable FTIR spectrometers. The spectrometers used have been developed by KIT in cooperation with Bruker (Gisi et al., 2012; Hase et al., 2016)

- 60 and are commercially available since 2014 (type designation EM27/SUN spectrometer). While lately-Lately a COCCON spectrometer was used in combination with two TCCON instruments to validate OCO-2 (Jacobs et al., 2020), to study boreal forests (Tu et al., 2020) and recently-Velazco et al. (2019) performed a campaign to validate GOSAT in central Australia. Apart from these studies, until now the major activity of the emerging network was to create the capability of permanent COCCON measurements at remote sites as a supplement of to the existing TCCON stations by developing the. To this end, procedures
- 65 for ensuring proper calibration and by providing the required were developed and evidence of the long-term stability of the EM27/SUN spectrometer was provided (Frey et al., 2015, 2019; Sha et al., 2020). Tasks which that can be accomplished by performing differential measurements using several spectrometers which can be calibrated side-by-side in the framework of campaigns are easier to achieve. Many successful campaigns for quantifying GHG emission strengths from regions of interest, as-like cities, coal mines, large dairy farms, etc., by arranging several spectrometers around the source have been performed
- 70 successfully using EM27/SUN spectrometers in the recent past (Hase et al., 2015; Vogel et al., 2019; Makarova et al., 2020; Viatte et al., 2017; Kille et al., 2019; Butz et al., 2017; Luther et al., 2019). In this work we introduce a COCCON station in Gobabeb, Namibia, where measurements are being have been conducted since January 2015. We demonstrate the excellent long-term stability of the COCCON instrument and its usefulness for satellite and model validation studies. The remainder of this paper is structured as follows. In section 2 we describe the measurement site, used instrumentation and data analysis,
- focusing on the COCCON EM27/SUN spectrometer. In section 3 we present the measurement results obtained over the last four years. In section 4, a comparison with respect to TCCON stations at Reunion Island and Lauder is conducted to illustrate the feasibility of our results. Additionally, the COCCON instrument is used to validate specific target mode observations from GOSAT, confirming a previously reported bias between GOSAT M-gain and H-gain soundings for XCO_2 (Velazco et al., 2019), and for the first time also reporting a bias in XCH_4 for the different gain settings. We also compare our measurements to
- 80 CAMS inversion-optimized model data. In section 5 we investigate the influence of the African biosphere on the observations performed in Gobabeb. In section 6 the results are discussed and an outlook for further studies is given.

2 Gobabeb site description, instrumentation and data analysis

2.1 Gobabeb site description

In 2015, we installed an EM27/SUN spectrometer of the COCCON network at the Gobabeb Namib Research Institute in 85 Namibia (23.561°S, 15.042°E, 410 m a.s.l.), see inset of Fig. 1. Gobabeb is located at the center of the hyperarid Namib desert. Moreover, Gobabeb is positioned next to the Kuiseb river, which marks the sharp transition zone between the gravel plains to the north and the sand desert to the south of the station, see Fig. 1. Gobabeb is situated 60 km east of the Atlantic ocean and the site is approximately 80 km southeast of the closest town, Walvis Bay, with a population of about 50000. The site is uninfluenced by nearby local emission sources of GHGs. Southwesterly winds prevail during austral summer, whereas in

90 winter easterly winds are predominant. The maximum temperature in summer can exceed 40 °C. Gobabeb is a high albedo station, together with the with a surface albedo derived from GOSAT retrievals at 1.6 μ m of 0.4 for the sand desert and 0.45 for the gravel plains. Together with the changing terrain this results in unique site characteristics desirable especially for satellite validation studies. In Fig. 2 we show the COCCON Gobabeb station in a broader context on a global map together with the TCCON Reunion Island and Lauder stations used in this study.

95 2.2 Description and history of the COCCON spectrometer operated at Gobabeb

The EM27/SUN spectrometer as used by COCCON has been described in great detail in the works of Gisi et al. (2012), Frey et al. (2015) and Hase et al. (2016). As a concise summary, the EM27/SUN is a solar-viewing Fourier transform infrared (FTIR) spectrometer measuring in the near infrared spectral range (5500 - 11000 cm⁻¹) with a spectral resolution of 0.5 cm⁻¹. One measurement takes 58 s and consists of 10 individual double-sided scans. This allows the retrieval of total column abundances,
VC_{gas}, of the target gases O₂, CO₂, CH₄ and H₂O. In 2018 the spectrometer used in this study was upgraded in Karlsruhe and a second, extended room temperature (RT) InGaAs detector (4000 - 5500 cm⁻¹) was added, allowing the detection of CO. During this service at KIT, the gold coating of the tracker mirrors was found to be degraded and therefore was removed (the mirror substrate is aluminium, so the operation was continued with aluminium mirrors since then). Finally, the mechanical parts of the solar tracker attached to the spectrometer was were serviced, as the very fine wind-blown dust particles tend to enter the motor stages during longer operation in the desert.

- The retrieved total column abundances of the trace gases are converted into column-averaged dry air mole fractions (DMFs), where the DMF of a gas is denoted $X_{gas} = \frac{VC_{gas}}{VC_{O_2}} \times 0.2095$. Here, both the column amounts of the target gas and O₂ are derived from the same spectroscopic observation reducing several potential error sources (Wunch et al., 2010). Furthermore, the dependence on the ground pressure is reduced improving comparability between different sites. A sensitive measure of the stability of a spectrometer is the column averaged amount of dry air (X_{air}) because for X_{air} there is no compensation of
- 110 the stability of a spectrometer is the column averaged amount of dry air (X_{air}) because for X_{air} there is no compensation of possible instrumental problems, in contrast to X_{gas} , where errors can partially cancel out. X_{air} compares the measured oxygen column (VC_{O2}) with surface pressure measurements (P_S):

$$X_{air} = \frac{g}{P_S} \cdot \left(\frac{V C_{O_2} \cdot \overline{\mu}}{0.2095} + V C_{H_2O} \cdot \mu_{H_2O} \right) \tag{1}$$

Here $\overline{\mu}$ and μ_{H_2O} denote the molecular masses of dry air and water vapour, respectively, *g* is the column averaged gravi-115 tational acceleration and VC_{H_2O} is the total column of water vapour. The correction with VC_{H_2O} is necessary as the surface pressure instruments measure the pressure of the total air column, including water vapour. Sudden changes in X_{air} indicate instrumental problems, e.g. errors with the surface pressure, pointing errors, timing errors or changes in the optical alignment of the instrument. Frey et al. (2019) present a comprehensive characterization for EM27/SUN spectrometers used by the COCCON network,

- 120 which included the instrument serial number 51 deployed in Gobabeb. In short, the instrumental line shape (ILS) of the EM27/SUN was optimized and characterized using open-path measurements as described in Frey et al. (2015), using version 14.5 of the LINEFIT retrieval software (Hase et al., 1999). Other detrimental effects, for example double-passing or channeling, were corrected if found. For more details see section 4.2 of Frey (2018). After this initial check in December 2014 side-by-side measurements with the reference EM27/SUN and the nearby TCCON instrument were performed on the observation platform
- 125 of the Institute for Meteorology and Climate Research (IMK-ASF) at the Karlsruhe Institute of Technology (KIT), Campus North (CN) near Karlsruhe (49.100° N, 8.439° E, 133 m a.s.l.). These measurements took place from November 2015 to March 2016 and once more in 2018 and 2019 in order to trace the results to TCCON (and thereby the WMO scale). This rigorous calibration routine is necessary in order to fulfill the high precision and accuracy requirements for GHG measurements. After the initial alignment check, no realignment was performed during the whole observation period.
- 130 The data analysis is performed differently from Frey et al. (2019). Spectra are generated from the raw interferograms (IFGs) using a FORTRAN 2003 preprocessing tool developed in the framework of the COCCON-PROCEEDS project and extensions (http://www.imk-asf.kit.edu/english/COCCON.php) of the European Space Agency (ESA). The IFGs are read from the OPUS file, the solar position is calculated, a correction for direct current (DC) fluctuations following Keppel-Aleks et al. (2007) is performed, the IFGs are truncated to the nominal resolution of 0.5 cm⁻¹, a numerical apodization function is applied and a fast
- 135 Fourier transformation including a phase correction routine and resampling scheme is implemented. Several quality filters are applied, for example requiring a minimum DC level , and of 5 % of the maximum detector signal level, restricting the tolerable DC variation in the IFG or to 10 % of the measured level, checking the centerburst location in the IFG and the centerburst amplitudes of forward and backward scans and the relative amplitude of out-of-band artifacts. All quality filters are summarized in Table 1.
- 140 For the retrieval of the EM27/SUN spectra we do not use the PROFFIT 9.6 retrieval algorithm (Schneider and Hase, 2009; Kiel et al., 2016; Chen et al., 2016). Here we use the recently developed non-linear least-squares PROFFAST retrieval algorithm, which fits atmospheric spectra by scaling a priori trace gas profiles. PROFFAST is a new efficient line-by-line forward model and retrieval code dedicated for COCCON data analysis. It is a source-open code accessible without restrictions and is designed to be numerically efficient and simple to use.source-open code for quantitative trace gas analysis, mainly intended for the use
- 145 with low-resolution FTIR spectrometers. Particular attention has been paid to achieve high processing speed without compromising the high level of accuracy required in the analysis of column-averaged greenhouse gas abundances. For achieving this goal, several measures are taken: (1) PROFFAST uses daily precalculated and tabulated molecular cross-sections derived from line-by-line calculations. (2) Instead of storing the cross sections per discrete layers, the cross-sections are expanded as function of solar zenith angle (SZA), which allows downsizing of the lookup tables by a factor of about five and accelerating
- 150 the subsequent calculation of atmospheric spectral transmission as function of SZA. (3) The process of convolution of the monochromatic spectrum with the instrumental line shape (ILS) is formulated as a two-step procedure, the first step thins the spectral grid before the convolution is performed. (4) The state vector of the previous solution is maintained for fitting the next spectrum, as typically the atmospheric variations from spectrum to spectrum are rather small. This strategy allows reducing

the number of required iterations to typically two. (5) PROFFAST provides averaging kernels not for each measurement, but

- 155 as function of a set of SZA values for each measurement day. Evaluation of data quality achieved with a COCCON spectrometer operated in Finland including using the PROFFAST data analysis chain has been investigated in the framework of ESA's FRM4GHG project and results are reported by Sha et al. (2019)Sha et al. (2020). The analysis of 4 years of Gobabeb data consisting of around 120000 spectra took about 40 h, which is approximately 30 times faster than the previously used PROFFIT 9.6 retrieval algorithm. In order to be consistent with TCCON, the GGG2014 generated a priori files (Wunch et al., 1000)
- 160 2015) are usedas a priori profiles, for trace gases as well as for temperature and pressure. The ground pressure was recorded using a MHB-382SD data logger with a pressure accuracy of 3 hPa (> 1000 hPa) or 2 hPa (< 1000 hPa) and a precision of 0.07 hPa. In order to increase the level of accuracy, we calibrate our sensor against a co-located CS 100 pressure sensor from the Southern African Science Service Centre for Climate Change and Adaptive Land Management (SASSCAL) network with a long-term stability better than 0.1 hPa per year and a stated accuracy of 0.5 hPa at 20 °C and 1.0 hPa for temperatures between</p>
- 165 0 °C and 40 °C. We do not directly use the data from the CS 100 pressure sensor for our analysis as only hourly data are available (http://www.sasscalweathernet.org/weatherstat_infosheet_we.php?loggerid_crit=8893, last access: 07 May 2021). We use the spectroscopic line lists and retrieval windows as described in Frey et al. (2019). The resulting XCO₂ and XCH₄ products are bias-corrected with respect to TCCON based on long-term comparisons between COCCON data products analysed with PROFFAST and official TCCON data products from Karlsruhe (2014 ongoing) and Sodankyla (2017 2019). In
- 170 the future it is planned to incorporate comparisons from additional stations to improve the basis of the bias-correction. For X_{air} a scaling factor of 0.9737 is derived from the long-term observations performed in Karlsruhe and Sodankylä centering the X_{air} data around 1.

2.3 TCCON Reunion Island and Lauder

- Measurement procedures and data analysis at both sites follow TCCON protocol (Wunch et al., 2011) using the GGG2014 software package (Wunch et al., 2015). As required by TCCON, the instrumentation consists of a high-resolution FTIR spectrometer, model BRUKER IFS 125HR, which offers a maximum is routinely operated at a spectral resolution of 0.0035-0.02 cm⁻¹. The instrument is housed inside a temperature-controlled building. The TCCON station in Reunion Island, France (20.901°S, 55.485°E, 87 m a.s.l.) is located on the university campus of the Université de La Réunion in St. Denis, approximately 2000 km east of the African mainland. The data are available via De Mazière et al. (2017). The TCCON station at
- 180 Lauder, New Zealand (45.038°S, 169.684°E, 370 m a.s.l.) is situated in a sparsely populated environment on the South Island of New Zealand (Pollard et al., 2017). The data are available via Sherlock et al. (2014); Pollard et al. (2019). In October 2018 a new TCCON instrument was installed at Lauder (Pollard et al., 2021). For this study we combine the data sets datasets of both spectrometers and for the overlap period (October 2018) we use the data from the old TCCON instrument.

2.4 GOSAT specific target observations

185 A detailed description of the GOSAT instrumental features and data analysis is given in Kuze et al. (2009) and Yoshida et al. (2013). GOSAT detects shortwave-infrared radiation in three narrow bands (0.76, 1.6 and 2.0 μ m) with a resolution of 0.2

 cm^{-1} . Additionally it is equipped with a sensor measuring in the thermal infrared range. The TANSO-FTS footprint diameter is about 10.5 km at sea level. The nominal single-scan acquisition time is 4 s. For this study the GOSAT FTS Short Wave InfraRed (SWIR) Level 2 data version V02.81 from NIES is used. The satellite is flying at an altitude of 666 km with a repeat

- 190 cycle of 3 days. Starting-From May 2016, GOSAT performed specific target mode observations over Gobabeb by performing observations at three distinct points , see-(Fig. 1. Directly-), directly at the Gobabeb COCCON site, approximately 10 km north east over the gravel plains and around 10 km south west over the sand desert. These points were chosen because of their different surface reflectance in order to study the sensitivity of the GOSAT retrieval with respect to the surface albedo. The satellite performed measurements with different gain settings, M-gain and H-gain. M-gain soundings are generally performed
- 195 over surfaces that are bright in the near infrared. For M-gain observations other validation sites with ground-based FTIR measurements are sparse(Yoshida et al., 2013; Velazeo et al., 2019). Yoshida et al. (2013) found no suitable TCCON site for the validation of GOSAT M-gain observations. In recent years, GOSAT M-gain soundings are mainly compared to the Edwards TCCON station, which was established 2013. More recently, Velazco et al. (2019) performed a campaign to validate GOSAT in central Australia using an EM27/SUN.

200 2.5 CAMS global CO₂ and CH₄ atmospheric inversion products

The CAMS model has been described previously in great detail, e.g. (Agustí-Panareda et al., 2014; Massart et al., 2016; Inness et al., 2019) Agustí-Panareda et al. (2014); Massart et al. (2016); Inness et al. (2019). Here we utilize the CAMS global inversion-optimized column-averaged dry air mole fractions DMFs for CO_2 and CH_4 . For CO_2 , we use an inversion product FT19r1 (Chevallier, 2020a) assimilating OCO-2 satellite observations (O'Dell et al., 2018; Kiel et al., 2019) as well as an in situ driven inversion product v18r3 (Chevallier, 2019). More details can be found in Chevallier et al. (2019); Chevallier (2020b). For CH_4 , an inversion product v18r1s assimilating a combination of surface and GOSAT satellite observations (Detmers and Hasekamp, 2016) as well as one product v18r1s using only surface observations are analyzed (Segers and Houweling, 2020a). A description of the inversion procedure together with comparisons against independent observational data sets-datasets is given in Segers and Houweling (2020b).

210 3 Measurement results

205

3.1 Side-by-side measurements at Karlsruhe

ILS measurements were carried out seven times since December 2014. This is depicted in Fig. 3. The modulation efficiency (ME) at maximum optical path difference (MOPD) ranges between 0.979 and 0.986 with a mean value of 0.983 and a standard deviation of 0.002. The mean phase error is 0.0019 ± 0.0003 . No drift is apparent and the ILS is stable. The spread in the ME

215 is in good agreement with the error budget of 0.003 given in Frey et al. (2019). This high instrumental stability is remarkable and not self-evident. Between the considering that between measurements the EM27/SUN was shipped from Karlsruhe to Gobabeb, including airlift and transport by car on bumpy gravel roads. Between November 2015 and March 2016 side-by-side comparison measurements were conducted on eight days together with the reference EM27/SUN were conducted on eight days to derive calibration factors for the different trace gases for this

- 220 spectrometer and thereby removing possible instrument-dependent biases. Some data had to be filtered out due to for different reasons. Because most measurements were performed during winter, the solar elevation was low, which sometimes led to a partially obstructed view due to railings and a metal frame on the terrace where the observations took place. In the morning the first measurements were omitted due to unusually high scatter caused by the quickly changing temperature of the not frequency-stabilized HeNe laser, which is not frequency-stabilized as already reported by Gisi et al. (2012). In rare cases, the
- tracking software failed, resulting in corrupted spectra, that were also filtered out. For this analysis only observations from the two instruments performed within one minute and solar zenith angles (SZAs) below 85° are taken into account, resulting in 1209 coincident measurements. The results are shown in Fig. 4. The derived instrument-specific calibration factors are 1.0002 ± 0.0003 for XCO₂, 1.0005 ± 0.0004 for XCH₄, 1.0011 ± 0.0029 for XH₂O and 0.9995 ± 0.0005 X_{air} between the reference instrument and the instrument deployed in Namibia. Although the scaling factors are close to nominal for all species,
- 230 to avoid biases due to instrumental differences these calibration factors are taken into account in the analysis of the Namibia data setdataset.

Additional side-by-side measurements were performed in February and March 2018 after the instrument came back from Namibia as well as between November 2018 and February 2019 after the dual channel upgrade and mirror exchange. The combined results are shown in Appendix A. A slight variation in the calibration factors is detectable, for XCH_4 and X_{air} the

- change is significant at the 1 σ level. The numeric values for the scaling factors are 1.0004 \pm 0.0004 for XCO₂, 0.9989 \pm 0.0004 for XCH₄, 0.9988 \pm 0.0016 for XH₂O and 1.0031 \pm 0.0007 for X_{*air*}. For the period between November 2018 and February 2019 we also derive a calibration factor of 0.9940 \pm 0.0050 for XCO. As the bias between the calibration factors obtained during the two side-by-side measurement periods is within 0.1 ppm for XCO₂, 3 ppb for XCH₄ and 3 ppm for XH₂O, for the analysis of the Namibia data we will only use the mean calibration factors derived from these observation periods.
- 240 In order to investigate if the difference in the calibration factors is linked to the upgrade of the EM27/SUN in 2018, we average the ME at MOPD obtained from the ILS measurements before (0.982) and after (0.985) the upgrade. The ME increased by 0.3 %, which is within the uncertainty budget of 0.3 % using this method. Therefore, we conclude that the changes in the instrumental line shape due to the upgrade of the COCCON instrument might contribute to the slightly different scaling factors, but they are not the main reason for the changes.

245 3.2 Gobabeb Xgas time series

For the subsequent analysis only observations with SZAs not exceeding 80° are taken into account, resulting in 113049 individual measurements on 319 days between 2015 and 2019. In Fig. 5 we present the XCO₂, XCH₄, XCO, XH₂O and X_{air} retrieval results from the COCCON Gobabeb observations. For better visibility, daily mean values are shown. Error bars denote the 1 σ standard deviation of the daily mean values. Between February and May 2015, no measurements could be performed

due to software problems. In October 2015, the spectrometer was shipped back to Karlsruhe due to customs requirements.Observations in Gobabeb were continued from April 2016. In February 2018 the spectrometer was shipped to Karlsruhe for

the dual channel upgrade. COCCON measurements were restarted in March 2019.

For XCO₂, the underlying trend of about 2 ppm / year can be seen. Correspondingly, a daily minimum value was observed at the beginning of the measurements on 24 January 2015 with 394.3 ± 0.2 ppm and the maximum daily value was observed on

- 15 October 2019 (410.6 \pm 0.2 ppm). A seasonal cycle is also detectable, with a peak-to-peak amplitude of 5.3 ppm in 2017. Here it is calculated as the difference between the maximum monthly mean of 404.0 \pm 1.1 ppm in September and the minimum monthly mean of 398.0 \pm 0.5 in March. This amplitude is higher than observed in other southern hemisphere TCCON stations in Australia and New Zealand (Deutscher et al., 2014), owing to a rather sharp drawdown of XCO₂ in February and March 2017. However, this is probably a real signal as the impact of the biosphere in Africa might lead to a larger seasonal cycle in
- Gobabeb. Also Olsen and Randerson predict a rather prominent XCO_2 seasonal cycle on the order of 5 ppm in southern Africa, see Figure 5 of Olsen and Randerson (2004). For XCH_4 , daily mean values range between 1759 \pm 1 ppb (2 June 2015) and 1828 ± 1 ppb (25 June 2019). The trend is roughly 0.01 ppm / year. The XCH_4 seasonal cycle has lowest values in southern hemispheric summer (January 2017: 1783 ± 5 ppb) and highest values throughout winter and early spring (September 2017: 1808 ± 5 ppb) resulting in a peak-to-peak amplitude of 25 ppb. Regarding XCO, the time series is limited to 2019 due to the
- fact that the dual channel upgrade was only performed in 2018. At this point, it can already be seen that this site observes highly variable amounts of carbon monoxide, ranging from very clean background conditions with daily mean XCO values as low as 49 ± 1 ppb (16 April 2019) to elevated results of up to 131 ± 9 ppb (4 September 2019). XH₂O is very low during large parts of the year, as expected for a desert site. The lowest value was reached on 29 June 2015 (357 ± 10 ppm). During late southern hemispheric summer and early spring, XH₂O can reach up to several thousand ppm. As mentioned in section 2.2, X_{air} is an
- 270 important parameter to monitor the instrumental stability. For the whole time series, daily X_{air} results are stable within 1 %. No apparent drift of X_{air} is visible during the four years of measurements performed at the COCCON Gobabeb station.

4 Gobabeb data comparisons

4.1 TCCON Reunion Island and Lauder

background site without a pronounced XCO₂ seasonal cycle.

In this section we compare the results obtained in Gobabeb with results from the TCCON stations at Reunion Island and Lauder. Although this is not a side-by-side comparison, Reunion Island as the second closest TCCON station is approximately 4000 km east of Gobabeb, this comparison will give us a measure of the feasibility of our results. The observations should be comparable qualitatively as the variation of XCO₂ is relatively low in the southern hemisphere compared to the northern hemisphere (Olsen and Randerson, 2004). Moreover, Gobabeb (24°S) and Reunion Island (21°S) are roughly at at roughly the same latitude. The TCCON station Ascension Island is slightly closer to Gobabeb with a distance of approximately 3600 km, but the latitudional difference is larger. Due to the latitudional gradient in XCH₄, we therefore chose to compare our COCCON measurements to Reunion Island rather than Ascension Island. Lauder is the southernmost TCCON station and functions as a

Daily mean XCO₂, XCH₄, XCO and XH₂O results are shown in Fig. 6 from COCCON Gobabeb (blue dots), TCCON Reunion Island (black dots) and TCCON Lauder (red dots) stations. Error bars denote the 1 σ standard deviation of the daily

- 285 mean values. Difference timeseries between the stations are shown in separate panels. For XCO_2 we see a good agreement between the sites, given the fact that they are spatially far apart. The annual increase of XCO_2 is similar for all stations. For Reunion Island and Lauder, no pronounced seasonal cycle is visible. Most The most prominent difference is the sharp decrease of XCO_2 at Gobabeb beginning of 2017, most pronounced in March. This is not seen for the TCCON data at the two other sites. There is no conspicuous change in X_{air} during this period for the COCCON instrument, and the other gases do not deviate.
- 290 We therefore do not think that instrumental issues are the cause of the difference. As discussed in the previous section, this difference is probably due to the impact of the African biosphere to on the measurements in Gobabeb. To a smaller extent this difference can also be seen at the beginning of 2018. Despite the similarities, it can also be seen that the Reunion Island values at the beginning of 2018 and then at the end of 2019 it can also be seen that the Reunion Island values somewhat diverge from the Gobabeb and Lauder values. XCH_4 at the Gobabeb and Reunion sites is similar, with lower absolute values at Lauder. The
- annual increase as well as and the seasonal variability are similar at all sites. Opposed to Unlike XCO_2 , there is no conspicuous difference between the data sets datasets at the beginning of 2017. For XCO, the sites do not have a long observation overlap, it seems that the variability is slightly larger in the COCCON data. Regarding XH_2O , the seasonality is similar between the sites, with highest values at Reunion Island throughout the year.

In order to affirm that the drawdown of Xat the beginning of 2017 at the Gobabeb station is due to the influence of the

- 300 African biosphere, we show 10-day backward trajectory ensemble simulations from the National Oceanic and Atmospheric Administration (NOAA) HYSPLIT model (Stein et al., 2016) for 16 February 2017, the day with the lowest Xvalues in 2017. Initial 3-hourly meteorological input data is provided by the NCEP Global Data Assimilation System (GDAS) model on a 1 degree latitude-longitude grid. The end point of the trajectory analysis is chosen at a height of 5000 m above ground level. We choose this height because in section 4.3 a comparison between COCCON data with CAMS model data shows that the CAMS
- 305 model version assimilating total column data reproduces the Xdrawdown, in contrast to the version assimilating in situ data. Therefore we think that the drawdown is driven by low concentrations of in the higher layers in the atmosphere rather than in the atmospheric boundary layer. Backward trajectories for Gobabeb are depicted in Fig. 18. All trajectories exhibit a long dwell time over the African continent, corroborating the conjecture that the low Xvalues at Gobabeb are due to the influence of the African biosphere. In contrast, the backward trajectories for Reunion Island shown in Fig. 19 dwell almost exclusively
- 310 over the ocean.In a next step, we show correlation plots for the COCCON site with respect to the TCCON sites for XCO_2 and XCH_4 in Fig. 7 and 8. Error bars denote the 1 σ standard deviation (STD) of the daily mean values. The colorbar denotes the measurement date. Focussing first on the comparison between Gobabeb and Reunion Island in Fig. 7, we find an agreement within one standard deviation of the averaged daily mean values for both gases. For XCO_2 a scaling factor of 1.0027 ± 0.0028 and a correlation coefficient R² of 0.911 are derived. For XCH_4 the scaling factor is 1.0028 ± 0.0045 and R² of 0.670. Bias
- and STD in absolute values are given in Table 2. Despite this good agreement, especially for XCO_2 , there is some divergence between the data before and after 2018, corresponding to larger scatter in the TCCON Reunion Island data setdataset, as can be seen by the larger error bars for the 2018 and 2019 data. For 2018 the reason for the increased scatter was continued mirror degradation as a result of sea salt deposition from the ocean. In Fig. 8 we see an excellent agreement between the COCCON Gobabeb and TCCON Lauder data for XCO_2 with a scaling factor of 0.9990 \pm 0.0027 and a correlation coefficient R² of

320 0.906. The only discernible anomaly are the lower COCCON values beginning of 2017, which is also seen in the time series in Fig. 6. Otherwise, no temporal drift between the two data sets datasets is apparent. For XCH₄ a scaling factor of 0.9800 \pm 0.0060 with R² = 0.556 is found. The large bias is to be expected due to the latitudinal gradient in atmospheric methane concentrations (Saeki et al., 2013).

Next, we examine several diurnal cycles for five days between Gobabeb and Reunion Island, one day each year, where data

- 325 is available for both sites. The results for Xand Xare shown in Fig. A2, COCCON measurements are shown as blue dots, TCCON measurements as black dots. In contrast to other graphs, here we show local time data, for better comparability of the diurnal cycles. For X, the diurnal curvature for both COCCON and TCCON is relatively flat, however a slight parabola shape is discernible. For southern hemispheric summer, compared to TCCON Reunion COCCON Gobabeb values are slightly lower as was already seen in the time series analysis. Xdiurnal variations are similar for both sites, also the absolute values
- 330 are in perfect agreement. A common feature for both data sets is the apparent parabola shape on most days. This is probably the result of a combination of non-perfect a priori profiles, residual airmass dependency and intraday changes of atmospheric temperature. In the next version of the TCCON trace gas retrieval algorithm, updated a priori profiles will be used that will help to further reduce these unwanted effects. For 13 July 2015 it seems that this effect is slightly more pronounced for the COCCON instrument. For the other days this is hard to assess as the scatter of the TCCON Reunion Island data continuously
- 335 increases with time due to degrading mirror quality. This finding is true for both Xand X.

4.2 GOSAT validation

In this section we validate specific target mode observations from the GOSAT satellite around Gobabeb at three distinct points with different surface albedo properties against COCCON Gobabeb observations. Target mode measurements started in 2016 and are ongoing. The time series of the GOSAT observations is shown in Fig. 9. Measurements over the gravel plains are

- displayed in red, observations directly at Gobabeb in black and measurements over the sand desert are presented in gold, with 59, 78 and 85 successful observations, respectively. In general, the agreement between GOSAT observations and COCCON measurements is reasonable, GOSAT data seem to be slightly biased high both in XCO_2 and XCH_4 . An interesting anomaly is observed in the GOSAT data, there seems to be a small decrease both in XCO_2 and XCH_4 during southern hemispheric winter, which is not observed by the COCCON instrument. For-However, for a rigorous assessment the data is too sparsehowever.
- An additional difference is that the drawdown of XCO_2 values at the beginning of 2017 is more pronounced for COCCON compared to the satellite data.

For a quantitative analysis, we analyze coincident observations between GOSAT and COCCON. To make the data sets datasets comparable, we correct for the influence of the different a priori profiles following Rodgers and Connor (2003). We adjust the GOSAT values to the ensemble profile, which we assume to be the GGG2014 generated a priori profile. In Fig. 10 we present

350 the XCO_2 and XCH_4 COCCON and GOSAT averaging kernels for different SZAs. Although the COCCON averaging kernels are shown for SZAs in the range of 0° and 85°, for all coincident overpasses the SZA was between 10° and 50°. Due to the similarities of the averaging kernels, we neglect the smoothing error in the following analysis.

The number of coincident measurements with COCCON observations are 13, 18 and 20 for the three specific observation

points and the chosen coincidence criteria is that COCCON observations were performed within thirty minutes of the satellite

overpass. Of these coincident measurements, the vast majority occurred in 2016. The correlation graphs for these three target points are presented in Fig. 11, 12 and 13. GOSAT M-gain observations are color-coded red, while H-gain observations are shown in blue. Error bars denote the 1 σ standard deviation of the hourly mean values for COCCON measurements and the measurement error for the GOSAT soundings.

For the GOSAT observations over the gravel plains, only GOSAT M-gain soundings were performed. The spread of the data

- 360 set dataset is relatively large, GOSAT is biased high and we derive a scaling factor with respect to the COCCON observations of 1.0062 ± 0.0026 and 1.0044 ± 0.0039 for XCO₂ and XCH₄, where the difference is statistically significant at the 1 σ level. This corresponds to a high bias of 2.5 ± 1.1 ppm for XCO₂ and 7.9 ± 7.1 ppb for XCH₄. In Table 3 and Table 4-3 the absolute values of the GOSAT - COCCON comparison are summarized. Directly over Gobabeb GOSAT M-gain as well as H-gain soundings were performed. Between COCCON and GOSAT M-gain data we derive a scaling factor of $1.0026 \pm$
- 365 0.0027 for XCO₂ and 1.0018 ± 0.0033 for XCH₄, corresponding to a high bias of 1.0 ± 1.1 ppm for XCO₂ and 3.1 ± 6.0 ppb for XCH₄. For H-Gain observations we derive a scaling factor of 0.9996 ± 0.0020 for XCO₂ and 0.9984 ± 0.0016 for XCH₄, corresponding to a low bias of 0.2 ± 0.8 ppm for XCO₂ and 2.8 ± 2.9 ppb for XCH₄. The differences between GOSAT and COCCON are not statistically different at the 1 σ level. Over the sand desert, the GOSAT M-gain data are biased high with respect to the COCCON data with a scaling factor of 1.0068 ± 0.0026 for XCO₂ and 1.0070 ± 0.0045 for XCH₄,
- corresponding to a high bias of 2.7 ± 1.1 ppm for XCO₂ and 12.5 ± 8.1 ppb for XCH₄. The H-gain data are in very good agreement with the COCCON observations with a scaling factor of 1.0003 ± 0.0008 for XCO₂ and 1.0015 ± 0.0028 for XCH₄, corresponding to a high bias of 0.1 ± 0.3 ppm for XCO₂ and a high bias of 2.7 ± 5.1 ppb for XCH₄.

Although not always statistically significant at the 1 σ level, clear differences are discernible between the different GOSAT gain settings. This is in agreement with results reported by Velazco et al. (2019) and the GOSAT validation team. For the H-gain soundings we report a good agreement with the COCCON observations within the 1 σ level for XCO₂ as well as XCH₄ with high correlation coefficients (R² > 0.9).

4.3 CAMS evaluation

COCCON and CAMS XCO_2 and XCH_4 timeseries are provided in Fig. 14. Hourly mean COCCON data are shown as blue dots, CAMS data as red (satellite data assimilation) and black dots (in situ data assimilation) with 3-hourly model output for

380 XCO_2 and 6-hourly model output for XCH_4 . For XCO_2 , note that the OCO-2 assimilated data is available until 2019 and the in situ assimilated data is available until 2018. For XCH_4 , both CAMS datasets are available until 2018.

As was shown in section 4.1, the COCCON measurements exhibit a small but discernible parabola shape during the day. For better comparability, we therefore only compare COCCON measurements around local noon with the CAMS model data. Although using all COCCON data instead of only noon data results in only a small bias of 0.2 ppm for XCO₂ and 2 ppb for

385 XCH_4 relative to the noon-only data, we feel that this is the more consistent comparison. The resulting correlation plots for XCO_2 and XCH_4 are presented in Fig. 15 and 16. For X, note that the OCO-2 assimilated data is available until 2019 and the in situ assimilated data is available until 2018. The left panel of Fig. 15 shows the OCO-2 assimilated model data. We see an

excellent agreement between the two data sets datasets with a bias of 0.9 ± 0.5 ppm and a correlation coefficient R² of 0.983. This offset agrees well with the bias between CAMS model and TCCON data presented in Chevallier (2020a). We do not

- 390 observe an increased bias at the beginning of 2017. This means that the OCO-2 assimilated model reproduces the drawdown of XCO_2 seen in the COCCON time series in Fig. 5 during this time. In contrast, we see an increased bias during the beginning of 2017 in the in situ assimilated data in the right panel. Apart from this anomaly, the agreement between the two data sets datasets is good. The CAMS model has a high bias of 1.1 ± 0.6 ppm and $R^2 = 0.927$. The absolute values of biases between the CAMS model evaluation are depicted simulation and the COCCON Gobabeb measurements are presented in Table 4.
- For XCH₄, both the combined GOSAT and in situ assimilated data as well as and the in situ assimilated data are available until 2018. The GOSAT and in situ assimilated CAMS data exhibit a low bias of 2.4 ± 8.0 ppb, R² = 0.455. From At the end of 2016 to through to the beginning of 2017 an anomaly is discernible with higher CAMS values. This is not seen in the comparison with the in situ assimilated dataset. The anomaly corresponds to a period of increased scatter in the GOSAT and in situ assimilated CAMS timeseries itself. see Fig. 14. Therefore we attribute this anomaly to the influence of the GOSAT
- 400 observations. For the in situ assimilated data we find a low bias of 5.8 ± 4.8 ppb and $R^2 = 0.645$. This is consistent with the low bias of CAMS with respect to TCCON measurements in the latitude band between 20°S and 30°S of around 10 ppb, as shown in figure 17 of Segers and Houweling (2020a).

5 Influence of the African biosphere on the Gobabeb observations

From the end of 2016 until the beginning of 2017, the XCO₂ values at the COCCON station at Gobabeb were significantly
lower compared to the TCCON stations Reunion Island and Lauder, see section 4.1. We rule out instrumental problems as the reason, as X_{air} is stable and the other observed gases do not show abnormal variations during this period. In order to investigate whether the drawdown of XCO₂ at the beginning of 2017 at the Gobabeb station is linked to the African biosphere, in Fig. 17 we present global OCO-2 assimilated CAMS a posteriori surface carbon fluxes for 16 February 2017 12 UTC, the day with the lowest XCO₂ values in 2017. We find that in the direct vicinity of Gobabeb, no strong negative carbon fluxes are apparent. From this, we deduce that air parcels with low CO₂ concentrations are transported to Gobabeb from other re-

- gions of the African mainland with negative surface fluxes. We therefore expect that the drawdown of XCO_2 is driven by low CO_2 concentrations in higher layers of the atmosphere that are representative for medium- or long-range transport. This is in agreement with the results of section 4.3, where a comparison between COCCON data with CAMS model data shows that the CAMS model version assimilating total column data reproduces the XCO_2 drawdown, in contrast to the version assimilating
- 415 in situ data only. We grant the possibility that the discrepancy between the different CAMS products could also stem from imperfections of the CAMS model.

In Fig. 18, we show 10-day backward trajectory ensemble simulations from the National Oceanic and Atmospheric Administration (NOAA) HYSPLIT model (Stein et al., 2016) for 16 February 2017. Initial 3-hourly meteorological input data is provided by the NCEP Global Data Assimilation System (GDAS) model on a 1 degree latitude-longitude grid. The end point of the

420 trajectory analysis is chosen at a height of 5000 m above ground level. All trajectories exhibit a long dwell time over the

African continent in regions with strong negative carbon surface fluxes, see Fig. 17. This corroborates the conjecture that the low XCO_2 values at Gobabeb are due to the influence of the African biosphere. Most of the trajectories that arrive at 5000 m height at Gobabeb originate from significantly lower levels of the atmosphere, close to the surface, and are then uplifted, as can be seen in the lower panel of Figure 18.

425 In contrast, the backward trajectories for Reunion Island shown in Fig. 19 dwell almost exclusively over the ocean. In Fig. 20 we additionally provide backward trajectories for Gobabeb ending at 1000 m above ground level. In contrast to the trajectories at 5000 m, these originate from the ocean.

6 Conclusions and Outlook

- 430 We present measurements from a new ground-based remote-sensing COCCON station in Namibia, the first FTIR site measuring GHGs on the African continent. We performed a thorough calibration scheme carried out in Karlsruhe in order to make the results traceable to TCCON (and thereby the WMO scale), including ILS measurements and side-by-side comparisons with a reference COCCON spectrometer. The results from Namibia show a typical global annual increase rates rate for both XCO₂ as well as XCH₄. In contrast to comparable FTIR measurements in the southern hemisphere, we observe a pronounced seasonal variability for XCO₂ with a peak-to-peak amplitude of 5.3 ppm in 2017, in agreement with OCO-2 assimilated CAMS model
- data and global transport model predictions (Olsen and Randerson, 2004). As expected for a desert site, we observe very low values of XH_2O , with a minimum value of 357 ppm. For the whole time series, daily X_{air} results are stable within 1 %. No apparent drift of X_{air} is visible during the four years of measurements performed at the COCCON Gobabeb station. To put our results in the broader geophysical context, we compare the COCCON Namibia results to measurements from the
- TCCON stations Reunion Island and Lauder. Given the fact that the stations are spatially far apart, the results are in good agreement. For XCO₂ both TCCON Lauder (-0.4 ± 1.1 ppm) and Reunion Island (1.1 ± 1.1 ppm) show biases compared to COCCON Gobabeb within the 1 σ uncertainty range and correlation coefficients R² > 0.9. For XCH₄ TCCON Reunion Island and COCCON Gobabeb data agree within the 1 σ uncertainty range (5.1 ± 8.1 ppb) while a large bias (-35.9 ± 10.6 ppb) is observed with respect to the Lauder data. This is a direct result of the strong latitudinal gradient in total column averaged methane concentrations. We further investigate the diurnal variations from TCCON Reunion Island and COCCON Gobabeb
- for XCO₂ and XCH₄. Both share a small but systematic downward parabola shape, probably the result of a combination of non-perfect a priori profiles, residual airmass dependency and intraday changes of atmospheric temperature. From a comparison of the two data sets datasets we also deduce that the Reunion Island data set dataset shows increased scatter during some time periods due to the degrading mirror quality as a result of sea salt deposition from the ocean. Compared to the TCCON results, the COCCON observations are of comparable quality.
- We show the usefulness of our station for satellite validation by comparing the COCCON results to GOSAT specific target mode observations at three points with different surface albedos close to or directly at the sitewith different surface albedos. The satellite performed measurements with different gain settings. Ground-based validation of the different gain settings is

difficult as very few sites worldwide have the necessary surface characteristics, further supporting the importance of this new

station. We find a good agreement between GOSAT H-gain and COCCON observations within the 1 σ uncertainty range with low biases of -0.2 ± 0.8 ppm for XCO₂ and -2.8 ± 2.9 ppb for XCH₄ at Gobabeb and high biases of 0.1 ± 0.3 ppm for XCO₂ and 2.7 ± 5.1 ppb for XCH₄ over the sand desert approximately 15 kilometers south-east of the station. For M-gain soundings, GOSAT measurements are always biased high with respect to the COCCON measurements, the differences over the gravel plains and the sand desert are statistically significant at the 1 σ level. Thereby we show the capability of this site to validate satellite measurements for different high albedo surfaces.

Then we evaluate the performance of the inversion-optimized CAMS model data against our ground-based COCCON data. For XCO_2 we find high biases of 0.9 ± 0.5 ppm for the OCO-2 assimilated product and 1.1 ± 0.6 ppm for the in situ-driven product with $R^2 > 0.9$ in both cases. These biases are comparable to offsets between the model and TCCON data. The OCO-2 assimilated model product is able to reproduce the drawdown of XCO_2 beginning of 2017, as opposed to the in situ-optimized product. Also for XCH_4 the biases found are in line with prior model comparisons with TCCON.

- With this work we show the potential of the COCCON network for satellite validation and atmospheric transport model validation. We expect that the availability of additional COCCON sites in the near future will be a great asset for future satellite and model studies as they are easy to deploy. In the course of the ESA funded COCCON PROCEEDS project COCCON data from several sites will be made available via a web portal. We conclude that instruments from the COCCON network offer stable
- 470 long-term records of GHGs in remote environments and can be used to close gaps in the global distribution of ground-based remote-sensing sites.

Data availability. COCCON data will be made available in the near future through a web portal hosted at the Karlsruhe Institute of Technology. TCCON Reunion Island and Lauder data can be obtained via: https://tccondata.org, last access: 20 October 2020. The GOSAT TANSO-FTS SWIR L2 data are available from the GOSAT Data Archive Service (GDAS) at https://data2.gosat.nies.go.jp/ (GDAS, last access: 20 October 2020).

475

Appendix A: Calibration measurements Karlsruhe 2018 and 2019

In Fig. A1 we present the results from the calibration measurements performed between February 2018 and 2019.

Appendix B: Intraday XCO₂ and XCH₄ variability at Gobabeb and Reunion Island

480 We examine several measurement days between Gobabeb and Reunion Island, one day each year, where data is available for both sites. The results for XCO_2 and XCH_4 are shown in Fig. A2, COCCON measurements are shown as blue dots, TCCON measurements as black dots. In contrast to other graphs, here we show local time data, for better comparability of the measurement days. For XCO_2 , the curvature for both COCCON and TCCON is relatively flat, however a slight parabola shape is discernible. For southern hemispheric summer, compared to TCCON Reunion, COCCON Gobabeb values are slightly

- lower as was already seen in the time series analysis. XCH₄ variations are similar for both sites, also the absolute values are in perfect agreement. A common feature for both datasets is the apparent parabola shape on most days. This is probably the result of a combination of non-perfect a priori profiles, residual airmass dependency and intraday changes of atmospheric temperature. In the next version of the TCCON trace gas retrieval algorithm, updated a priori profiles will be used that will help to further reduce these unwanted effects. For 13 July 2015 it seems that this effect is slightly more pronounced for the COCCON instrument. For the other days this is hard to assess as the scatter of the TCCON Reunion Island data continuously
 - increases with time due to degrading mirror quality. This finding is true for both XCO₂ and XCH₄.

Author contributions. MF, TB and FH planned the study. MF and TB installed the COCCON station in Gobabeb with help from FH, FG and JG. RM, MH and PA performed the day to day COCCON measurements. MKS and MDM provided the TCCON Reunion Island data. DFP provided the TCCON Lauder data. DD provided the COCCON Gobabeb data. IM provided the GOSAT data. KS organized the GOSAT specific target mode observations over Gobabeb. MF performed the data analysis and wrote the paper. All authors reviewed, edited and approved the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. CAMS data were generated using Copernicus Atmosphere Monitoring Service Information 2020.

The authors acknowledge support by the ACROSS research infrastructure of the Helmholtz Association of German Research Centres (HGF). 500 The authors acknowledge support by the MOSES research infrastructure of the HGF.

The authors acknowledge support by ESA in the framework of FRM4GHG, COCCON-PROCEEDS, COCCON-PROCEEDSII and QA4EO projects.

The Réunion Island station is operated by the Royal Belgian Institute for Space Aeronomy with financial support since 2014 by the EU project ICOS-Inwire and the ministerial decree for ICOS (FR/35/IC1 to FR/35/IC5) and local activities supported by LACy/UMR8105 –

505 Université de La Réunion.

495

The authors gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model used in this publication.

References

525

535

545

- 510 Agustí-Panareda, A., Massart, S., Chevallier, F., Boussetta, S., Balsamo, G., Beljaars, A., Ciais, P., Deutscher, N. M., Engelen, R., Jones, L., Kivi, R., Paris, J.-D., Peuch, V.-H., Sherlock, V., Vermeulen, A. T., Wennberg, P. O., and Wunch, D.: Forecasting global atmospheric CO₂, Atmospheric Chemistry and Physics, 14, 11959–11983, https://doi.org/10.5194/acp-14-11959-2014, https://www.atmos-chem-phys.net/ 14/11959/2014/, 2014.
 - Borsdorff, T., aan de Brugh, J., Hu, H., Hasekamp, O., Sussmann, R., Rettinger, M., Hase, F., Gross, J., Schneider, M., Garcia, O., Stremme,
- W., Grutter, M., Feist, D. G., Arnold, S. G., De Mazière, M., Kumar Sha, M., Pollard, D. F., Kiel, M., Roehl, C., Wennberg, P. O., Toon, G. C., and Landgraf, J.: Mapping carbon monoxide pollution from space down to city scales with daily global coverage, Atmospheric Measurement Techniques, 11, 5507–5518, https://doi.org/10.5194/amt-11-5507-2018, https://www.atmos-meas-tech.net/11/5507/2018/, 2018.
 - Butz, A., Dinger, A. S., Bobrowski, N., Kostinek, J., Fieber, L., Fischerkeller, C., Giuffrida, G. B., Hase, F., Klappenbach, F., Kuhn, J.,
- 520 Lübcke, P., Tirpitz, L., and Tu, Q.: Remote sensing of volcanic CO₂, HF, HCl, SO₂, and BrO in the downwind plume of Mt. Etna, Atmospheric Measurement Techniques, 10, 1–14, https://doi.org/10.5194/amt-10-1-2017, https://www.atmos-meas-tech.net/10/1/2017/, 2017.
 - Chen, J., Viatte, C., Hedelius, J. K., Jones, T., Franklin, J. E., Parker, H., Gottlieb, E. W., Wennberg, P. O., Dubey, M. K., and Wofsy, S. C.: Differential column measurements using compact solar-tracking spectrometers, Atmospheric Chemistry and Physics, 16, 8479–8498, https://doi.org/10.5194/acp-16-8479-2016, https://www.atmos-chem-phys.net/16/8479/2016/, 2016.
 - Chevallier, F.: Validation report for the CO₂ fluxes estimated by atmospheric inversion, v18r3. CAMS deliverable CAMS73_2018SC1_D73.1.4.1-2018-v2_201911., https://ads.atmosphere.copernicus.eu/, 2019.
- Chevallier, F.: Evaluation and Quality control document for the OCO-2-driven CO₂ inversion FT19r1. CAMS deliverable CAMS73_2018SC2_D73.4.3.1-2020-v1_202004_v1_Validation FT inversion CO2 fluxes_v1, https://ads.atmosphere.copernicus.eu/, 2020a.
 - Chevallier, F.: Description of the CO₂ inversion production chain 2020. CAMS deliverable CAMS73_2018SC2_D5.2.1-2020_202004_CO2 inversion production chain_v1, https://ads.atmosphere.copernicus.eu/, 2020b.
 - Chevallier, F., Remaud, M., O'Dell, C. W., Baker, D., Peylin, P., and Cozic, A.: Objective evaluation of surface- and satellite-driven carbon dioxide atmospheric inversions, Atmospheric Chemistry and Physics, 19, 14 233–14 251, https://doi.org/10.5194/acp-19-14233-2019, https://www.atmos-chem-phys.net/19/14233/2019/, 2019.
 - Crisp, D., Pollock, H. R., Rosenberg, R., Chapsky, L., Lee, R. A. M., Oyafuso, F. A., Frankenberg, C., O'Dell, C. W., Bruegge, C. J., Doran, G. B., Eldering, A., Fisher, B. M., Fu, D., Gunson, M. R., Mandrake, L., Osterman, G. B., Schwandner, F. M., Sun, K., Taylor, T. E., Wennberg, P. O., and Wunch, D.: The on-orbit performance of the Orbiting Carbon Observatory-2 (OCO-2) instrument and its radiometrically calibrated products, Atmospheric Measurement Techniques, 10, 59–81, https://doi.org/10.5194/amt-10-59-2017, https://doi.org/10.5194/amt-10-59-
- 540 //www.atmos-meas-tech.net/10/59/2017/, 2017.
 - De Mazière, M., Sha, M. K., Desmet, F., Hermans, C., Scolas, F., Kumps, N., Metzger, J.-M., Duflot, V., and Cammas, J.-P.: TCCON data from Réunion Island (RE), Release GGG2014.R1, https://doi.org/10.14291/tccon.ggg2014.reunion01.r1, https://data.caltech.edu/records/ 322, 2017.

Detmers, R. and Hasekamp, O.: Product User Guide (PUG) for the RemoTeC XCH₄ PROXY GOSAT Data Product v2.3.8, ftp://ftp.sron.nl/ pub/pub/RemoTeC/ECV2/CRDP4/DOCS_GOSAT/PUGv4p0{_}CH4{_}GOS{_}SRPR.pdf, 2016.

17

- Deutscher, N. M., Sherlock, V., Mikaloff Fletcher, S. E., Griffith, D. W. T., Notholt, J., Macatangay, R., Connor, B. J., Robinson, J., Shiona, H., Velazco, V. A., Wang, Y., Wennberg, P. O., and Wunch, D.: Drivers of column-average CO₂ variability at Southern Hemispheric Total Carbon Column Observing Network sites, Atmospheric Chemistry and Physics, 14, 9883–9901, https://doi.org/10.5194/acp-14-9883-2014, https://www.atmos-chem-phys.net/14/9883/2014/, 2014.
- Dils, B., Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Parker, R., Guerlet, S., Aben, I., Blumenstock, T., Burrows, J. P., Butz, A., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., De Mazière, M., Notholt, J., Sussmann, R., Warneke, T., Griffith, D., Sherlock, V., and Wunch, D.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON, Atmospheric Measurement Techniques, 7, 1723–1744, https://doi.org/10.5194/amt-7-1723-2014, https://www.atmos-meas-tech.
- 555 net/7/1723/2014/, 2014.

570

- Dlugokencky, E. J., M., C. A., Mund, J. W., J., C. M., and W., T. K.: Atmospheric Methane Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network 1983-2018, Version: 2019-07, https://doi.org/https://doi.org/10.15138/VNCZ-M766, 2019a.
- Dlugokencky, E. J., Mund, J. W., M., C. A., J., C. M., and W., T. K.: Atmospheric Carbon Dioxide Dry Air Mole
- 560 Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network 1968-2018, Version: 2019-07, https://doi.org/https://doi.org/10.15138/wkgj-f215, 2019b.
 - Eldering, A., O'Dell, C. W., Wennberg, P. O., Crisp, D., Gunson, M. R., Viatte, C., Avis, C., Braverman, A., Castano, R., Chang, A., Chapsky, L., Cheng, C., Connor, B., Dang, L., Doran, G., Fisher, B., Frankenberg, C., Fu, D., Granat, R., Hobbs, J., Lee, R. A. M., Mandrake, L., McDuffie, J., Miller, C. E., Myers, V., Natraj, V., O'Brien, D., Osterman, G. B., Oyafuso, F., Payne, V. H., Pollock,
- H. R., Polonsky, I., Roehl, C. M., Rosenberg, R., Schwandner, F., Smyth, M., Tang, V., Taylor, T. E., To, C., Wunch, D., and Yoshimizu,
 J.: The Orbiting Carbon Observatory-2: first 18 months of science data products, Atmospheric Measurement Techniques, 10, 549–563, https://doi.org/10.5194/amt-10-549-2017, https://www.atmos-meas-tech.net/10/549/2017/, 2017.
 - Eldering, A., Taylor, T. E., O'Dell, C. W., and Pavlick, R.: The OCO-3 mission: measurement objectives and expected performance based on 1 year of simulated data, Atmospheric Measurement Techniques, 12, 2341–2370, https://doi.org/10.5194/amt-12-2341-2019, https: //www.atmos-meas-tech.net/12/2341/2019/, 2019.
 - Frankenberg, C., Pollock, R., Lee, R. A. M., Rosenberg, R., Blavier, J.-F., Crisp, D., O'Dell, C. W., Osterman, G. B., Roehl, C., Wennberg, P. O., and Wunch, D.: The Orbiting Carbon Observatory (OCO-2): spectrometer performance evaluation using pre-launch direct sun measurements, Atmospheric Measurement Techniques, 8, 301–313, https://doi.org/10.5194/amt-8-301-2015, http://www.atmos-meas-tech. net/8/301/2015/, 2015.
- 575 Frey, M.: Characterisation and application of portable solar absorption spectrometers for the detection of greenhouse gas emissions from regional anthropogenic sources, Ph.D. thesis, Karlsruhe Institute of Technology, Karlsruhe, https://doi.org/10.5445/IR/1000088312, http://dx.doi.org/10.5445/IR/1000088312;http://nbn-resolving.org/urn:nbn:de:swb:90-883128;http://d-nb.info/1174252006/34;https: //publikationen.bibliothek.kit.edu/1000088312, 2018.
 - Frey, M., Hase, F., Blumenstock, T., Groß, J., Kiel, M., Mengistu Tsidu, G., Schäfer, K., Sha, M. K., and Orphal, J.: Calibration and instrumen-
- 580 tal line shape characterization of a set of portable FTIR spectrometers for detecting greenhouse gas emissions, Atmospheric Measurement Techniques, 8, 3047–3057, https://doi.org/10.5194/amt-8-3047-2015, https://www.atmos-meas-tech.net/8/3047/2015/, 2015.
 - Frey, M., Sha, M. K., Hase, F., Kiel, M., Blumenstock, T., Harig, R., Surawicz, G., Deutscher, N. M., Shiomi, K., Franklin, J. E., Bösch, H., Chen, J., Grutter, M., Ohyama, H., Sun, Y., Butz, A., Mengistu Tsidu, G., Ene, D., Wunch, D., Cao, Z., Garcia, O., Ra-

monet, M., Vogel, F., and Orphal, J.: Building the COllaborative Carbon Column Observing Network (COCCON): long-term stability

- 585 and ensemble performance of the EM27/SUN Fourier transform spectrometer, Atmospheric Measurement Techniques, 12, 1513–1530, https://doi.org/10.5194/amt-12-1513-2019, https://www.atmos-meas-tech.net/12/1513/2019/, 2019.
 - Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B., Anthoni, P., Barbero, L., Bastos, A., Bastrikov, V., Becker, M., Bopp, L., Buitenhuis, E., Chandra, N., Chevallier, F., Chini, L. P., Currie, K. I., Feely, R. A., Gehlen, M., Gilfillan, D., Gkritzalis, T., Goll, D. S.,
- Gruber, N., Gutekunst, S., Harris, I., Haverd, V., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Joetzjer, E., Kaplan, J. O., Kato, E., Klein Goldewijk, K., Korsbakken, J. I., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Marland, G., McGuire, P. C., Melton, J. R., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Neill, C., Omar, A. M., Ono, T., Peregon, A., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Séférian, R., Schwinger, J., Smith, N., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Werf, G. R., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2019, Earth System Science Data, 11, 1783–1838, https://doi.org/10.5194/essd-11-1783-2019, https://www.earth-syst-sci-data.net/11/1783/2019/, 2019.
- Gisi, M., Hase, F., Dohe, S., Blumenstock, T., Simon, A., and Keens, A.: XCO₂-measurements with a tabletop FTS using solar absorption spectroscopy, Atmospheric Measurement Techniques, 5, 2969–2980, https://doi.org/10.5194/amt-5-2969-2012, http://www. atmos-meas-tech.net/5/2969/2012/, 2012.

Hase, F., Blumenstock, T., and Paton-Walsh, C.: Analysis of the instrumental line shape of high-resolution Fourier transform IR spectrometers

- 600 with gas cell measurements and new retrieval software, Appl. Opt., 38, 3417–3422, https://doi.org/10.1364/AO.38.003417, http://ao.osa. org/abstract.cfm?URI=ao-38-15-3417, 1999.
 - Hase, F., Frey, M., Blumenstock, T., Groß, J., Kiel, M., Kohlhepp, R., Mengistu Tsidu, G., Schäfer, K., Sha, M. K., and Orphal, J.: Application of portable FTIR spectrometers for detecting greenhouse gas emissions of the major city Berlin, Atmospheric Measurement Techniques, 8, 3059–3068, https://doi.org/10.5194/amt-8-3059-2015, https://www.atmos-meas-tech.net/8/3059/2015/, 2015.
- 605 Hase, F., Frey, M., Kiel, M., Blumenstock, T., Harig, R., Keens, A., and Orphal, J.: Addition of a channel for XCO observations to a portable FTIR spectrometer for greenhouse gas measurements, Atmospheric Measurement Techniques, 9, 2303–2313, https://doi.org/10.5194/amt-9-2303-2016, https://www.atmos-meas-tech.net/9/2303/2016/, 2016.
 - Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., and Suttie, M.:
- 610 The CAMS reanalysis of atmospheric composition, Atmospheric Chemistry and Physics, 19, 3515–3556, https://doi.org/10.5194/acp-19-3515-2019, https://www.atmos-chem-phys.net/19/3515/2019/, 2019.
 - Inoue, M., Morino, I., Uchino, O., Nakatsuru, T., Yoshida, Y., Yokota, T., Wunch, D., Wennberg, P. O., Roehl, C. M., Griffith, D. W. T., Velazco, V. A., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Hase, F., Blumenstock, T., Rettinger, M., Sussmann, R., Kyrö, E., Kivi, R., Shiomi, K., Kawakami, S., De Mazière, M., Arnold, S. G., Feist, D. G., Barrow, E. A., Barney, J., Dubey, M.,
- 615 Schneider, M., Iraci, L. T., Podolske, J. R., Hillyard, P. W., Machida, T., Sawa, Y., Tsuboi, K., Matsueda, H., Sweeney, C., Tans, P. P., Andrews, A. E., Biraud, S. C., Fukuyama, Y., Pittman, J. V., Kort, E. A., and Tanaka, T.: Bias corrections of GOSAT SWIR XCO₂ and XCH₄ with TCCON data and their evaluation using aircraft measurement data, Atmospheric Measurement Techniques, 9, 3491–3512, https://doi.org/10.5194/amt-9-3491-2016, https://www.atmos-meas-tech.net/9/3491/2016/, 2016.
- Jacobs, N., Simpson, W. R., Wunch, D., O'Dell, C. W., Osterman, G. B., Hase, F., Blumenstock, T., Tu, Q., Frey, M., Dubey, M. K., Parker,
 H. A., Kivi, R., and Heikkinen, P.: Quality controls, bias, and seasonality of CO₂ columns in the Boreal Forest with OCO-2, TCCON,

and EM27/SUN measurements, Atmospheric Measurement Techniques Discussions, 2020, 1–42, https://doi.org/10.5194/amt-2019-505, https://www.atmos-meas-tech-discuss.net/amt-2019-505/, 2020.

- Keppel-Aleks, G., Toon, G. C., Wennberg, P. O., and Deutscher, N. M.: Reducing the impact of source brightness fluctuations on spectra obtained by Fourier-transform spectrometry, Appl. Opt., 46, 4774–4779, https://doi.org/10.1364/AO.46.004774, http://ao.osa.org/abstract. cfm?URI=ao-46-21-4774, 2007.
- Kiel, M., Hase, F., Blumenstock, T., and Kirner, O.: Comparison of XCO abundances from the Total Carbon Column Observing Network and the Network for the Detection of Atmospheric Composition Change measured in Karlsruhe, Atmospheric Measurement Techniques, 9, 2223–2239, https://doi.org/10.5194/amt-9-2223-2016, https://www.atmos-meas-tech.net/9/2223/2016/, 2016.

625

- Kiel, M., O'Dell, C. W., Fisher, B., Eldering, A., Nassar, R., MacDonald, C. G., and Wennberg, P. O.: How bias correction goes
 wrong: measurement of X_{CO2} affected by erroneous surface pressure estimates, Atmospheric Measurement Techniques, 12, 2241–2259, https://doi.org/10.5194/amt-12-2241-2019, https://www.atmos-meas-tech.net/12/2241/2019/, 2019.
 - Kille, N., Chiu, R., Frey, M., Hase, F., Sha, M. K., Blumenstock, T., Hannigan, J. W., Orphal, J., Bon, D., and Volkamer, R.: Separation of Methane Emissions From Agricultural and Natural Gas Sources in the Colorado Front Range, Geophysical Research Letters, 46, 3990–3998, https://doi.org/10.1029/2019GL082132, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019GL082132, 2019.
- 635 Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the Greenhouse Gases Observing Satellite for greenhouse gases monitoring, Appl. Opt., 48, 6716–6733, https://doi.org/10.1364/AO.48.006716, http://ao.osa.org/abstract.cfm?URI=ao-48-35-6716, 2009.
 - Luther, A., Kleinschek, R., Scheidweiler, L., Defratyka, S., Stanisavljevic, M., Forstmaier, A., Dandocsi, A., Wolff, S., Dubravica, D., Wildmann, N., Kostinek, J., Jöckel, P., Nickl, A.-L., Klausner, T., Hase, F., Frey, M., Chen, J., Dietrich, F., Necki, J., Swolkień, J., Fix, A.,
- 640 Roiger, A., and Butz, A.: Quantifying CH₄ emissions from hard coal mines using mobile sun-viewing Fourier transform spectrometry, Atmospheric Measurement Techniques, 12, 5217–5230, https://doi.org/10.5194/amt-12-5217-2019, https://www.atmos-meas-tech.net/12/ 5217/2019/, 2019.
 - Makarova, M. V., Alberti, C., Ionov, D. V., Hase, F., Foka, S. C., Blumenstock, T., Warneke, T., Virolainen, Y., Kostsov, V., Frey, M., Poberovskii, A. V., Timofeyev, Y. M., Paramonova, N., Volkova, K. A., Zaitsev, N. A., Biryukov, E. Y., Osipov, S. I., Makarov,
- 645 B. K., Polyakov, A. V., Ivakhov, V. M., Imhasin, H. K., and Mikhailov, E. F.: Emission Monitoring Mobile Experiment (EMME): an overview and first results of the St. Petersburg megacity campaign-2019, Atmospheric Measurement Techniques Discussions, 2020, 1–45, https://doi.org/10.5194/amt-2020-87, https://amt.copernicus.org/preprints/amt-2020-87/, 2020.
 - Massart, S., Agustí-Panareda, A., Heymann, J., Buchwitz, M., Chevallier, F., Reuter, M., Hilker, M., Burrows, J. P., Deutscher, N. M., Feist, D. G., Hase, F., Sussmann, R., Desmet, F., Dubey, M. K., Griffith, D. W. T., Kivi, R., Petri, C., Schneider, M., and Velazco, V. A.:
- Ability of the 4-D-Var analysis of the GOSAT BESD XCO₂ retrievals to characterize atmospheric CO₂ at large and synoptic scales, Atmospheric Chemistry and Physics, 16, 1653–1671, https://doi.org/10.5194/acp-16-1653-2016, https://www.atmos-chem-phys.net/16/ 1653/2016/, 2016.
 - Moore III, B., Crowell, S. M. R., Rayner, P. J., Kumer, J., O'Dell, C. W., O'Brien, D., Utembe, S., Polonsky, I., Schimel, D., and Lemen, J.: The Potential of the Geostationary Carbon Cycle Observatory (GeoCarb) to Provide Multi-scale Constraints on the Carbon Cycle in
- 655 the Americas, Frontiers in Environmental Science, 6, 109, https://doi.org/10.3389/fenvs.2018.00109, https://www.frontiersin.org/article/ 10.3389/fenvs.2018.00109, 2018.
 - Morino, I., Uchino, O., Inoue, M., Yoshida, Y., Yokota, T., Wennberg, P. O., Toon, G. C., Wunch, D., Roehl, C. M., Notholt, J., Warneke, T., Messerschmidt, J., Griffith, D. W. T., Deutscher, N. M., Sherlock, V., Connor, B., Robinson, J., Sussmann, R., and Ret-

tinger, M.: Preliminary validation of column-averaged volume mixing ratios of carbon dioxide and methane retrieved from GOSAT

- 660 short-wavelength infrared spectra, Atmospheric Measurement Techniques, 4, 1061–1076, https://doi.org/10.5194/amt-4-1061-2011, http://www.atmos-meas-tech.net/4/1061/2011/, 2011.
 - Nivitanont, J., Crowell, S. M. R., and Moore III, B.: A scanning strategy optimized for signal-to-noise ratio for the Geostationary Carbon Cycle Observatory (GeoCarb) instrument, Atmospheric Measurement Techniques, 12, 3317–3334, https://doi.org/10.5194/amt-12-3317-2019, https://www.atmos-meas-tech.net/12/3317/2019/, 2019.
- O'Dell, C. W., Eldering, A., Wennberg, P. O., Crisp, D., Gunson, M. R., Fisher, B., Frankenberg, C., Kiel, M., Lindqvist, H., Mandrake, L., Merrelli, A., Natraj, V., Nelson, R. R., Osterman, G. B., Payne, V. H., Taylor, T. E., Wunch, D., Drouin, B. J., Oyafuso, F., Chang, A., McDuffie, J., Smyth, M., Baker, D. F., Basu, S., Chevallier, F., Crowell, S. M. R., Feng, L., Palmer, P. I., Dubey, M., García, O. E., Griffith, D. W. T., Hase, F., Iraci, L. T., Kivi, R., Morino, I., Notholt, J., Ohyama, H., Petri, C., Roehl, C. M., Sha, M. K., Strong, K., Sussmann, R., Te, Y., Uchino, O., and Velazco, V. A.: Improved retrievals of carbon dioxide from Orbiting Carbon Observatory-2
- 670 with the version 8 ACOS algorithm, Atmospheric Measurement Techniques, 11, 6539–6576, https://doi.org/10.5194/amt-11-6539-2018, https://amt.copernicus.org/articles/11/6539/2018/, 2018.
 - Olsen, S. C. and Randerson, J. T.: Differences between surface and column atmospheric CO₂ and implications for carbon cycle research, Journal of Geophysical Research: Atmospheres, 109, https://doi.org/10.1029/2003JD003968, http://dx.doi.org/10.1029/2003JD003968, 2004.
- 675 Pollard, D. F., Sherlock, V., Robinson, J., Deutscher, N. M., Connor, B., and Shiona, H.: The Total Carbon Column Observing Network site description for Lauder, New Zealand, Earth System Science Data, 9, 977–992, https://doi.org/10.5194/essd-9-977-2017, https://www. earth-syst-sci-data.net/9/977/2017/, 2017.
 - Pollard, D. F., Robinson, J., and Shiona, H.: TCCON data from Lauder (NZ), Release GGG2014.R0, https://doi.org/10.14291/TCCON.GGG2014.LAUDER03.R0, https://data.caltech.edu/records/1220, 2019.
- 680 Pollard, D. F., Robinson, J., Shiona, H., and Smale, D.: Intercomparison of Total Carbon Column Observing Network (TCCON) data from two Fourier transform spectrometers at Lauder, New Zealand, Atmospheric Measurement Techniques, 14, 1501–1510, https://doi.org/10.5194/amt-14-1501-2021, https://amt.copernicus.org/articles/14/1501/2021/, 2021.
 - Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, Journal of Geophysical Research: Atmospheres, 108, https://doi.org/10.1029/2002JD002299, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2002JD002299, 2003.
- 685 Saeki, T., Saito, R., Belikov, D., and Maksyutov, S.: Global high-resolution simulations of CO₂ and CH₄ using a NIES transport model to produce a priori concentrations for use in satellite data retrievals, Geoscientific Model Development, 6, 81–100, https://doi.org/10.5194/gmd-6-81-2013, https://www.geosci-model-dev.net/6/81/2013/, 2013.
 - Schneider, M. and Hase, F.: Ground-based FTIR water vapour profile analyses, Atmospheric Measurement Techniques, 2, 609–619, https://doi.org/10.5194/amt-2-609-2009, http://www.atmos-meas-tech.net/2/609/2009/, 2009.
- 690 Segers, A. and Houweling, S.: Validation of the CH₄ surface flux inversion reanalysis 1990-2018. CAMS deliverable CAMS73_2018SC1_D73.2.4.1-2019_202021_validation_CH4_1990-2018_v1., https://ads.atmosphere.copernicus.eu/, 2020a.
 - Segers, A. and Houweling, S.: Description of the CH₄ inversion production chain. CAMS deliverable CAMS73_2018SC1_D73.5.2.2--2019_202001_production_chain_v1, https://ads.atmosphere.copernicus.eu/, 2020b.
- 695 Sha, M. K., De Mazière, M., Notholt, J., Blumenstock, T., Chen, H., Dehn, A., Griffith, D. W. T., Hase, F., Heikkinen, P., Hermans, C., Hoffmann, A., Huebner, M., Jones, N., Kivi, R., Langerock, B., Petri, C., Scolas, F., Tu, Q., and Weidmann, D.: Intercomparison of low

and high resolution-low- and high-resolution infrared spectrometers for ground-based solar remote sensing measurements of total column concentrations of CO₂, CH₄-CO₂, CH₄, and CO, Atmospheric Measurement Techniques Discussions, 2019, 1–67, 13, 4791–4839, https://doi.org/10.5194/amt-13-4791-2020, 2019. https://amt.copernicus.org/articles/13/4791/2020/, 2020.

700

- Sha, M. K., De Mazière, M., NotholtLangerock, B., Blavier, J.-F. L., Blumenstock, T., Chen, H., Borsdorff, T., Buschmann, M., Dehn, A., De Mazière, M., Deutscher, N. M., Feist, D. G., García, O. E., Griffith, D. W. T., Grutter, M., Hannigan, J. W., Hase, F., Heikkinen, P., Hermans, C., Hoffmann, A., Huebner, MIraci, L. T., Jeseck, P., Jones, N., Kivi, R., Langerock, B., Kumps, N., Landgraf, J., Lorente, A., Mahieu, E., Makarova, M. V., Mellqvist, J., Metzger, J.-M., Morino, I., Nagahama, T., Notholt, J., Ohyama, H., Ortega, I., Palm, M., Petri,
- 705 C., Seolas, Pollard, D. F., Tu, Q., and Weidmann, D.: Intercomparison of low- and high-resolution infrared spectrometers for ground-based solar remote sensing measurements of total column concentrations of Rettinger, M., Robinson, J., Roche, S., Roehl, C. M., Röhling, A. N., Rousogenous, C., Schneider, M., Shiomi, K., Smale, D., Stremme, W., Strong, K., Sussmann, R., Té, Y., Uchino, O., Velazco, V. A., Vrekoussis, M., Wang, P., Warneke, T., Wizenberg, T., Wunch, D., Yamanouchi, S., Yang, Y., and Zhou, M.: Validation of Methane and Carbon Monoxide from Sentinel-5 Precursor using TCCON and NDACC-IRWG stations, - and CO, Atmospheric Measurement
- 710 Techniques Discussions, 2021, 1–84, https://doi.org/10.5194/amt-2021-36, 13, 4791–4839, , , 2020. https://amt.copernicus.org/preprints/ amt-2021-36/, 2021.
 - Sherlock, V., Connor, B., Robinson, J., Shiona, H., Smale, D., and Pollard, D. F.: TCCON data from Lauder (NZ), 125HR, Release GGG2014.R0, https://doi.org/10.14291/TCCON.GGG2014.LAUDER02.R0/1149298, https://data.caltech.edu/records/281, 2014.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and
 Dispersion Modeling System, Bulletin of the American Meteorological Society, 96, 2059–2077, https://doi.org/10.1175/BAMS-D-14-
 - 00110.1, https://doi.org/10.1175/BAMS-D-14-00110.1, 2016.
 Suto, H., Kataoka, F., Kikuchi, N., Knuteson, R. O., Butz, A., Haun, M., Buijs, H., Shiomi, K., Imai, H., and Kuze, A.: Thermal and near-infrared sensor for carbon observation Fourier-transform spectrometer-2 (TANSO-FTS-2) on the Greenhouse Gases Observing Satellite-2 (GOSAT-2) during its first year on orbit, Atmospheric Measurement Techniques Discussions, 2020, 1–51, https://doi.org/10.5194/amt-

720 2020-360, https://amt.copernicus.org/preprints/amt-2020-360/, 2020.

- Tu, Q., Hase, F., Blumenstock, T., Kivi, R., Heikkinen, P., Sha, M. K., Raffalski, U., Landgraf, J., Lorente, A., Borsdorff, T., Chen, H., Dietrich, F., and Chen, J.: Intercomparison of atmospheric CO₂ and CH₄ abundances on regional scales in boreal areas using Copernicus Atmosphere Monitoring Service (CAMS) analysis, COllaborative Carbon Column Observing Network (COCCON) spectrometers, and Sentinel-5 Precursor satellite observations, Atmospheric Measurement Techniques, 13, 4751–4771, https://doi.org/10.5194/amt-13-4751-2020, https://amt.copernicus.org/articles/13/4751/2020/, 2020.
- Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, rse, 120, 70–83, https://doi.org/10.1016/j.rse.2011.09.027, 2012.
- Velazco, V. A., Deutscher, N. M., Morino, I., Uchino, O., Bukosa, B., Ajiro, M., Kamei, A., Jones, N. B., Paton-Walsh, C., and Griffith, D. W. T.: Satellite and ground-based measurements of XCO₂ in a remote semiarid region of Australia, Earth System Science Data, 11, 935–946, https://doi.org/10.5194/essd-11-935-2019, https://www.earth-syst-sci-data.net/11/935/2019/, 2019.
 - Viatte, C., Lauvaux, T., Hedelius, J. K., Parker, H., Chen, J., Jones, T., Franklin, J. E., Deng, A. J., Gaudet, B., Verhulst, K., Duren, R., Wunch,D., Roehl, C., Dubey, M. K., Wofsy, S., and Wennberg, P. O.: Methane emissions from dairies in the Los Angeles Basin, Atmospheric

- 735 Chemistry and Physics, 17, 7509–7528, https://doi.org/10.5194/acp-17-7509-2017, https://www.atmos-chem-phys.net/17/7509/2017/, 2017.
 - Vogel, F. R., Frey, M., Staufer, J., Hase, F., Broquet, G., Xueref-Remy, I., Chevallier, F., Ciais, P., Sha, M. K., Chelin, P., Jeseck, P., Janssen, C., Té, Y., Groß, J., Blumenstock, T., Tu, Q., and Orphal, J.: XCO₂ in an emission hot-spot region: the COCCON Paris campaign 2015, Atmospheric Chemistry and Physics, 19, 3271–3285, https://doi.org/10.5194/acp-19-3271-2019, https://www.atmos-chem-phys.net/19/3271/2019/, 2019.
 - Washenfelder, R. A., Toon, G. C., Blavier, J.-F., Yang, Z., Allen, N. T., Wennberg, P. O., Vay, S. A., Matross, D. M., and Daube, B. C.: Carbon dioxide column abundances at the Wisconsin Tall Tower site, Journal of Geophysical Research: Atmospheres, 111, https://doi.org/10.1029/2006JD007154, http://dx.doi.org/10.1029/2006JD007154, 2006.
 - Wu, L., Hasekamp, O., Hu, H., Landgraf, J., Butz, A., aan de Brugh, J., Aben, I., Pollard, D. F., Griffith, D. W. T., Feist, D. G., Koshelev,
- D., Hase, F., Toon, G. C., Ohyama, H., Morino, I., Notholt, J., Shiomi, K., Iraci, L., Schneider, M., de Mazière, M., Sussmann, R., Kivi, R., Warneke, T., Goo, T.-Y., and Té, Y.: Carbon dioxide retrieval from OCO-2 satellite observations using the RemoTeC algorithm and validation with TCCON measurements, Atmospheric Measurement Techniques, 11, 3111–3130, https://doi.org/10.5194/amt-11-3111-2018, https://www.atmos-meas-tech.net/11/3111/2018/, 2018.
 - Wunch, D., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Stephens, B. B., Fischer, M. L., Uchino, O., Abshire, J. B., Bernath, P., Biraud, S. C.,
- Blavier, J.-F. L., Boone, C., Bowman, K. P., Browell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher, N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst, D. F., Jiménez, R., Keppel-Aleks, G., Kort, E. A., Macatangay, R., Machida, T., Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl, C. M., Sawa, Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. A.: Calibration of the Total Carbon Column Observing Network using aircraft profile data, Atmospheric Measurement Techniques, 3, 1351–1362, https://doi.org/10.5194/amt-3-1351-2010, http://www.atmos-meas-tech.net/3/1351/2010/, 2010.
- 755 Wunch, D., C Toon, G., L Blavier, J.-F., A Washenfelder, R., Notholt, J., J Connor, B., W T Griffith, D., Sherlock, V., and Wennberg, P.: The Total Carbon Column Observing Network, 369, 2087–112, 2011.
 - Wunch, D., Toon, G. C., Sherlock, V., Deutscher, N. M., Liu, X., Feist, D. G., and Wennberg, P. O.: The Total Carbon Column Observing Network's GGG2014 Data Version, https://doi.org/10.14291/TCCON.GGG2014.DOCUMENTATION.R0/1221662, 2015.

Wunch, D., Wennberg, P. O., Osterman, G., Fisher, B., Naylor, B., Roehl, C. M., O'Dell, C., Mandrake, L., Viatte, C., Kiel, M., Griffith,

D. W. T., Deutscher, N. M., Velazco, V. A., Notholt, J., Warneke, T., Petri, C., De Maziere, M., Sha, M. K., Sussmann, R., Rettinger, M., Pollard, D., Robinson, J., Morino, I., Uchino, O., Hase, F., Blumenstock, T., Feist, D. G., Arnold, S. G., Strong, K., Mendonca, J., Kivi, R., Heikkinen, P., Iraci, L., Podolske, J., Hillyard, P. W., Kawakami, S., Dubey, M. K., Parker, H. A., Sepulveda, E., García, O. E., Te, Y., Jeseck, P., Gunson, M. R., Crisp, D., and Eldering, A.: Comparisons of the Orbiting Carbon Observatory-2 (OCO-2) X_{CO2} measurements with TCCON, Atmospheric Measurement Techniques, 10, 2209–2238, https://doi.org/10.5194/amt-10-2209-2017,

765 https://www.atmos-meas-tech.net/10/2209/2017/, 2017.

740

- Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data, Atmospheric Measurement Techniques,
- 770 6, 1533–1547, https://doi.org/10.5194/amt-6-1533-2013, https://www.atmos-meas-tech.net/6/1533/2013/, 2013.

Table 1. This table summarizes the quality filters applied in the FORTRAN based preprocessing tool.

Q1	Check of DC level as fraction of ADC range, require 0.05
Q2	Check maximum variability of DC level (max. 10 % relative variation in interferogram resulting from 10 coadded scans)
Q3	Check FWD / BWD centerburst amplitudes (should agree within 5 %)
Q4	Check centerburst location in interferogram record
Q5	Check relative amplitude of out-of-band artifacts
Q6	Check slope, curvature, and change of curvature of phase spectrum
Q7	Check spectral calibration based on cross-correlation of spectral structure wrt a wavenumber-calibrated reference spectrum
Q8	Compare spectra derived from forward and backward scans

Table 2. This table presents the results of the comparison between the COCCON station in Gobabeb and the TCCON stations in Lauder and Reunion Island. Bias-Difference and STD are given as the mean difference and one standard deviation between the coincident daily TCCON and COCCON XCO_2 and XCH_4 values.

Station	XCO_2 Bias Difference \pm STD [ppm]	XCH_4 Bias-Difference \pm STD [ppb]	Number of coincidences
Reunion Island	1.1 ± 1.1	5.1 ± 8.1	155
Lauder	-0.4 ± 1.1	-35.9 ± 10.6	241

Table 3. This table presents the results of the comparison between the COCCON station in Namibia and the GOSAT M-gain and H-gain specific target observations. Bias and STD are given as the mean difference and one standard deviation between the coincident GOSAT and COCCON observations.

GOSAT target point	M-gain XCO ₂ Bias \pm STD [ppm]	M-gain XCH ₄ Bias \pm STD [ppb]	Number of coincidences
Gravel plains	2.5 ± 1.1	7.9 ± 7.1	13
Gobabeb	1.0 ± 1.1	3.1 ± 6.0	13
Sand desert	2.7 ± 1.1	12.5 ± 8.1	12
GOSAT target point	H-gain XCO ₂ Bias \pm STD [ppm]	H-gain XCH ₄ Bias \pm STD [ppb]	Number of coincidences
	1	1	1
Gravel plains	-	-	0
Gravel plains Gobabeb	-0.2 ± 0.8	-2.8 ± 2.9	0 5

Table 4. This table presents the results of the comparison between the COCCON station in Namibia and the assimilated CAMS model data. Bias and STD are given as the mean difference and one standard deviation between the coincident hourly-pooled local noon COCCON and CAMS XCO_2 and XCH_4 values.

Assimilation data	XCO_2 Bias \pm STD [ppm]	$XCH_4 Bias \pm STD [ppb]$	Number of coincidences
OCO-2 data	0.9 ± 0.5	-	263
In situ data	1.1 ± 0.6	-	187
In situ and GOSAT data	-	-2.4 ± 8.0	187
In situ data	-	-5.8 ± 4.8	187



Figure 1. Google earth image (Map data: Google, Maxar technologies) of the measurement site at Gobabeb, Namibia. The blue pin denotes the position of the COCCON instrument. The yellow points show the positions of the GOSAT target observation points. A black circle with a radius of 10 km has been drawn around the COCCON site for visual reference. The inset in the upper right corner shows the EM27/SUN spectrometer at Gobabeb.



Figure 2. Global map showing the COCCON Gobabeb, TCCON Reunion Island and Lauder sites used in this study.



Figure 3. Timeseries of the modulation efficiency at MOPD of the EM27/SUN used in this study. ILS measurements were performed during periods when the instrument was in Karlsruhe for maintenance or detector upgrade. Yellow areas denote measurement periods in Gobabeb. The black bar denotes the time of the detector upgrade.



Figure 4. Side-by-side measurements between the reference EM27/SUN and the instrument deployed in Namibia performed between November 2015 and March 2016 in Karlsruhe. From left to right, the panels show correlation plots for XCO_2 , XCH_4 , XH_2O and X_{air} . The coincident criteria is that measurements for both instruments occured within one minute. The colorbar denotes the solar zenith angle. For the analysis, only measurements with zenith angles below 85° are considered.



Figure 5. Column-averaged dry air mole fraction time series for XCO_2 , XCH_4 , XCO, XH_2O and X_{air} measured at the COCCON site in Gobabeb, Namibia from January 2015 until November 2019. Daily mean values are shown for better visibility. Error bars denote the 1 σ standard deviation of the daily mean values. In 2018 the instrument was upgraded with a second channel. Therefore XCO observations only started in 2019.



Figure 6. Column-averaged dry air mole fraction daily mean time series for XCO_2 , XCH_4 , XCO and XH_2O measured at the COCCON site in Gobabeb, Namibia (blue dots) and at the TCCON sites Reunion Island (black dots) and Lauder (red dots). Error bars denote the 1 σ standard deviation of the daily mean values. Additionally, the difference XCO_2 , XCH_4 , XCO and XH_2O timeseries between Gobabeb and Reunion Island (black dots) and between Gobabeb and Lauder (red dots) are shown in separate panels.



Figure 7. Correlation plots between the COCCON Gobabeb and TCCON Reunion Island stations for XCO_2 and XCH_4 from 2015 to 2019. Shown are daily mean values, errorbars denote the 1 σ standard deviation. The colorbar denotes the date of the measurement.



Figure 8. Correlation plots between the COCCON Gobabeb and TCCON Lauder stations for XCO_2 and XCH_4 from 2015 to 2019. Shown are daily mean values, errorbars denote the 1 σ standard deviation. The colorbar denotes the date of the measurement.



Figure 9. Column-averaged dry air mole fraction daily mean time series for XCO_2 and XCH_4 measured at the COCCON site in Gobabeb (blue dots) and GOSAT observations from the three specific target observation points with different surface albedos close to Gobabeb are shown (red dots: gravel plains, black dots: COCCON site, golden dots: sand desert). Error bars denote the 1 σ standard deviation of the daily mean values for COCCON measurements and the measurement error for the GOSAT soundings.



Figure 10. XCO_2 and XCH_4 column averaging kernels for the COCCON Gobabeb and GOSAT observations. The colorbar denotes the SZA. For the COCCON instrument, SZAs from 0° to 85° are depicted, whereas for GOSAT only the averaging kernels for the actual measurements are shown, with SZAs approximately between 10° and 50°.



Figure 11. Correlation plots between coincident COCCON Gobabeb observations and GOSAT measurements over the gravel plains between 2016 and 2019. For this area GOSAT only performed M-gain soundings (red dots). The red solid line is the best fit line through all M-gain data points. The dotted black line is the 1:1 line. Error bars denote the 1 σ standard deviation of the hourly mean values for COCCON measurements and the measurement error for the GOSAT soundings.



Figure 12. Correlation plots between coincident COCCON Gobabeb observations and GOSAT measurements over the COCCON site between 2016 and 2019. For this area GOSAT performed M-gain (red dots) and H-gain (blue dots) soundings. The red solid line is the best fit line through all M-gain data points, the blue solid line is the best fit line through all H-gain data points and the black solid line is the best fit line through all data points. The dotted black line is the 1:1 line. Error bars denote the 1 σ standard deviation of the hourly mean values for COCCON measurements and the measurement error for the GOSAT soundings.



Figure 13. Same as Fig. 12, with GOSAT observations over the sand desert.



Figure 14. Column-averaged dry air mole fraction daily mean time series for XCO_2 and XCH_4 at Gobabeb, Namibia. COCCON measurements are shown as blue dots, CAMS model data as red and black dots. For COCCON, we show hourly pooled data, for CAMS we show 3-hourly model output for XCO_2 and 6-hourly model output for XCH_4 .



Figure 15. XCO_2 correlation plots between coincident COCCON Gobabeb observations and CAMS model data. The left panel shows the OCO-2 assimilated model data, the right panel shows the in situ assimilated model data. Note that the OCO-2 assimilated data is available until 2019 and the in situ assimilated data is available until 2018. Error bars denote the 1 σ standard deviation of the hourly mean values for COCCON measurements.



Figure 16. XCH₄ correlation plots between coincident COCCON Gobabeb observations and CAMS model data. The left panel shows the model data assimilated with in situ and GOSAT data, the right panel shows the in situ assimilated model data. Error bars denote the 1 σ standard deviation of the hourly mean values for COCCON measurements.



Figure 17. Global map showing OCO-2 assimilated CAMS a posteriori surface carbon fluxes for 16 February 2017 12 UTC.



Figure 18. NOAA HYSPLIT backward trajectory ensemble simulations on 16 February 2017. The endpoint of the backward trajectories is the COCCON Gobabeb station, 5000 m above ground level. The colors and symbols are used to make the different trajectories of the ensemble distinguishable.



Figure 19. Same as Fig. 18, but for the TCCON Reunion Island station.



Figure 20. Same as Fig. 18, but with the endpoint of the backward trajectories at 1000 m above ground level.



Figure A1. Same as Fig. 4, but for calibration measurements performed between February 2018 and February 2019.



Figure A2. Comparisons of intraday XCO_2 and XCH_4 variability for one day in 2015, 2016, 2017, 2018 and 2019 between the COCCON station Gobabeb (blue dots) and the TCCON station Reunion Island (black dots).