



Quantitative comparison of measured and simulated O₄ absorptions for one day with extremely low aerosol load over the tropical Atlantic

5 Thomas Wagner¹, Steffen Dörner¹, Steffen Beirle¹, Sebastian Donner¹, Stefan Kinne²

¹Satellite Remote Sensing Group, Max Planck Institute for Chemistry, Mainz, Germany

²Max-Planck Institute for Meteorology, Hamburg, Germany

Correspondence to: Thomas Wagner (thomas.wagner@mpic.de)

Abstract. In this study we compare measured and simulated O₄ absorptions for conditions of extremely low aerosol optical depth (between 0.034 to 0.056 at 360nm) on one day during a ship cruise in the tropical Atlantic. For such conditions, the uncertainties related to imperfect knowledge of aerosol properties don't significantly affect the comparison results. We find that the simulations underestimate the measurements by 15% to 20%. Even for simulations without any aerosols the measured O₄ absorptions are still systematically higher than the simulation results. The observed discrepancies can not be explained by uncertainties of the measurements and simulations and thus indicate a fundamental inconsistency between simulations and measurements.

1 Introduction

Remote sensing measurements of the atmospheric absorption of the oxygen dimer (O₂)₂ are often used to derive properties of aerosols and clouds. The atmospheric concentration of (O₂)₂ (in the following referred to as O₄) varies only slightly with temperature, pressure and humidity variations (aside the dependence on altitude). Thus deviations from the O₄ absorptions for clear sky conditions indicate changes of the atmospheric radiative transfer, e.g. due to clouds and aerosols. In recent years, inconsistencies between the measured atmospheric O₄ absorption and radiative transfer simulations were detected for Multi-AXis-DOAS (MAX-DOAS) observations. MAX-DOAS instruments measure scattered sun light under different, mostly slant elevation angles (Hönninger and Platt, 2002). Several studies found that a scaling factor (SF<1) had to be applied to the observed atmospheric O₄ absorptions in order to bring them into agreement with radiative transfer simulations. Other studies, however, did not find the need to apply such a scaling factor (e.g. Ortega et al., 2016; see also discussion in Wagner et al., 2019, and references therein). One major difficulty in the quantitative interpretation of these comparisons is that usually the atmospheric aerosol properties are not well known (e.g. the vertical extinction profile and/or the optical properties). And even if they were known, it is still a challenge to accurately represent them in atmospheric radiative transfer simulations.

In this study we mainly minimise these difficulties by using atmospheric observations in the presence of very low aerosol loads. During a ship campaign across the tropical Atlantic, very low aerosol optical depth (AOD) was observed on one day (2 May 2019). At 360 nm (the wavelength at which we analyse the atmospheric O₄ absorption), the AOD ranged from 0.034 to 0.056, which is an order of magnitude lower than the optical depth of molecular Rayleigh scattering.

Like in previous studies, we compare the observed atmospheric O₄ absorption with the results of radiative transfer simulations. Information about the aerosol properties is derived from sun photometer measurements in combination with ceilometer measurements. Also in our study, considerable uncertainties about the aerosol vertical profile and the aerosol optical properties exist. However, these uncertainties are less important for the interpretation of the comparison results than in previous studies because of the low AOD, and we find large discrepancies between the measured and simulated O₄ absorptions.

The paper is organised as follows: In section 2, an overview on the ship campaign and the instruments used in this study is given. Sections 3 to 5 describe the spectral analysis, the cloud classification, and the calculation of the O₄ profile. In section



6, the radiative transfer simulations and the extraction of the aerosol extinction profiles are presented. Section 7 presents the
45 comparison results, and section 8 the summary and conclusions.

2 Overview on the ship campaign and the instruments used in this study

The MAX-DOAS measurements were carried out during a cruise (MSM82/2) of the German research vessel RV Maria S.
50 Merian (<https://www.ldf.uni-hamburg.de/merian.html>) from Montevideo (Uruguay) to Las Palmas (Spain) from 26 April
2019 to 14 May 2019 (see Fig. 1). More details on the ship cruise MSM82/2 can be found in Krastel et al. (2019). In this
study, we focus on one day with particularly low AOD (2nd May), which is marked in Fig. 1.

The MAX-DOAS instrument was mounted above the ship's bridge at about 20m altitude above sea level. The telescope was
aligned in the driving direction of the ship (Fig. 2).

55

2.1 MAX-DOAS instrument

The MAX-DOAS instrument is a so-called Tube MAX-DOAS instrument which was developed and built by the electronic
workshop of the Max Planck Institute for Chemistry in Mainz (Donner, 2016). It consists of two major parts: the telescope
60 unit and the spectrometer unit. The telescope unit is mounted outside on the railing of the ship. The spectrometer unit is
located inside the ship. Besides the spectrometer it also contains a peltier cooling element which stabilises the spectrometer
temperature at 15 °C. Both units are connected via a quartz glass fibre bundle and electric cables. The telescope unit is
equipped with a gyroscope to stabilise the elevation angles by continuously adjusting the motor position with an accuracy of
 $\pm 0.1^\circ$.

65 The spectrometer is an Avantes ULS2048x64-USB2. It covers the spectral range from 299.4 nm to 463.1 nm with a spectral
resolution between 0.52 and 0.54 nm as described by the full width half maximum (FWHM). Spectra are measured with an
integration time of 1 min at the following elevation angles: -2° , -1° , -0.5° , 0° , 0.5° , 1° , 2° , 3° , 4° , 5° , 6° , 8° , 10° , 15° , 30° ,
 90° . Note that in this study only measurements with positive elevation angles are used. One elevation sequence is completed
within about 21 min. Dark current and offset spectra are taken during night and are used to correct the measured spectra
70 before the spectral analysis.

2.2 Sun photometer

A MICROTOPS II sunphotometer provided atmospheric totals on aerosol and water vapor. The instrument, when directed
75 towards the sun (in a handheld operation), captures via diodes the solar intensity in five sub-spectral bands near wavelengths
of 380, 440, 670, 870 and 940nm. In combination with the larger reference solar intensity at the top of the atmosphere -
using time and (GPS-provided) position data - sun-photometer measurements define the atmospheric attenuation at these
solar sub-spectral bands. Four spectral bands (near 380, 440, 670 and 870 nm) sample in trace-gas poor regions, while one
spectral band (near 940nm) is strongly affected by water vapor absorption. In the absence of clouds, the solar attenuations in
80 the four trace-gas poor bands can be linked to aerosol - after (surface air pressure defined) contributions from air-molecule
(Rayleigh) scattering have been removed. Hereby, the aerosol associated attenuations are quantified by the (vertically
normalized) aerosol optical depth (AOD). As the instrument offers AOD values simultaneously at four different solar
wavelengths, the typical aerosol particle size is revealed and even AOD contributions from sub-micrometer (mainly from
pollution and wildfire) and super-micrometer size aerosol particles (mainly from dust and seasalt) can be distinguished. The
85 determination of the atmospheric water vapor is based on the differential absorption between 870 and 940 nm attenuation
data. Any quality measurement usually relies on many repeated samples in order to identify and remove poor data associated



with sun-view contamination by clouds and/or inaccurate orientations of the instrument into the sun (with is done manually with the help of a pointing device). NASA's Aeronet sub-group of the Maritime Aerosol Network (MAN, Smirnov et al., 2009) provided the calibrated instrument for the cruise and also stores cruise data at
90 https://aeronet.gsfc.nasa.gov/new_web/cruises_new/Maria_Merian_19_0.html.

2.3 Ceilometer

The Jenoptik 15K ceilometer of the MPI-M is a simple laser system operating at 1064nm, at an invisible trace-gas free near-
95 IR wavelength. Laser impulses are sent upward into the atmosphere and based on strength and delay of backscattered return signals altitude positions for atmospheric aerosol and clouds are derived. Due to their stronger backscatter at optically thicker media, such as clouds, overhead cloud base altitudes are well captured. However, as laser light strongly attenuates in optically thicker media, no information above a cloud base is possible. Vertical profiles of aerosol for clouds-free views (and below clouds) are possible up to about 7km in altitude during the night but only up to about 4km in altitude during the day,
100 due to scattering noise by sunlight. No useful aerosol profiling is possible near the surface (e.g. lower 300m), because signal sender and receiver are not the identical location. Recorded ceilometer data of the cruise are accessible via an anonymous ftp-site at ftp://ftp-projects.zmaw.de/aerocom/ships/ceilometer_MSM/.

3 Spectral analysis

105 The spectral analysis is performed following the settings suggested by Wagner et al. (2019). The spectral range from 352 to 387 nm is chosen, which contains two O₄ absorption bands. The details of the analysis are given in Table 1. Fig. A1 (left) presents an example of the spectral analysis as used in this study. In addition to the other cross sections, also a H₂O cross section (Polyansky et al. 2018) is included. The reason for including a H₂O cross section as well as the effect of including a
110 second O₄ cross section are discussed in appendix A1.

The results of the spectral analysis represent the integrated trace gas concentration along the atmospheric light path, the so-called slant column density (SCD). For O₄ the SCD is expressed with respect to the square of the O₂ concentration (see Greenblatt et al., 1990). Thus, the unit of the O₄ SCD is molec²/cm⁵. For the analysis of the measured spectra, a so-called Fraunhofer reference spectrum is used. In this study, the Fraunhofer reference spectrum is calculated as the average of the
115 zenith spectra before and after the chosen elevation sequence, weighted by the time of the selected measurement from that elevation sequence. Before performing the spectral analysis, these sequential Fraunhofer reference spectra are fitted to a 'universal' Fraunhofer reference spectrum (29 April 13:43, SZA: 44.8°, elevation angle: 90°) to transfer the spectral calibration of the universal Fraunhofer reference spectrum to the sequential Fraunhofer reference spectra. The universal Fraunhofer reference spectrum was calibrated using a high resolved solar spectrum.

120 Since the Fraunhofer reference spectrum also contains atmospheric trace gas absorptions, the output of the spectral analysis represents the difference between the SCDs of the selected non-zenith spectrum and the Fraunhofer reference spectrum, the so-called differential SCD (or dSCD).

The typical fit error of the derived O₄ dSCD is between 2·10⁴¹ molec²/cm⁵ and 4·10⁴¹ molec²/cm⁵. Depending on the magnitude of the retrieved O₄ dSCD this corresponds to relative errors between 1 and 4 %.

125

4 Cloud detection using the MAX-DOAS measurements

Although during most of the afternoon on 2 May clear sky conditions prevailed, also some scattered clouds were present. They were e.g. detected by the ceilometer in zenith direction (see Fig. 3). In order to derive information about possible cloud



130 contamination for the individual MAX-DOAS measurements, the MAX-DOAS measurements themselves were used for the
detection of cloud contamination, similar as in Wagner et al., (2014, 2016). Figure A3 in the appendix shows the time series
of the retrieved O₄ dSCDs on 2 May for the different elevation angles. During the morning the O₄ dSCDs show strong
variability caused by the presence and variability of clouds as also seen in the ceilometer data (Fig. 3). During the afternoon,
for most of the time, smooth variations of the O₄ dSCDs are found indicating clear sky conditions. However, for some times
135 and elevation angles, also small systematic deviations (usually reductions) of the O₄ dSCDs occur, which are caused by
scattered clouds. During periods without any cloud contamination, the temporal variability of the retrieved O₄ dSCDs is
rather small (scatter of the O₄ dSCDs is typically $\leq 5 \cdot 10^{41}$ molec²/cm⁵). Measurements with deviations $> 10^{42}$ molec²/cm⁵
compared to the extrapolated O₄ SCDs from the smooth (cloud-free) neighboring measurements are thus flagged as cloud-
contaminated. From the selected 11 elevation sequences during the mainly cloud-free periods in the afternoon of 2 May, 7
140 are found to be completely free of cloud contamination.

5 Calculation of the O₄ profile and O₄ VCD

The O₄ height profile and VCD for 2 May 2019 are calculated from vertical profiles of temperature and pressure. Also the
145 effect of the atmospheric humidity is accounted for. For the profiles of temperature, pressure and atmospheric humidity we
used the results from the ECMWF ERA-Interim data set (Berrisford et al., 2011) for 2 May 2019. From the temperature and
pressure profiles the air concentration [air] is calculated. Then the O₂ concentration [O₂] is derived according to the
following equation:

$$150 \quad [O_2] = [air] \cdot M_{O_2} \cdot (1 - M_{H_2O}) \quad (1)$$

Here M_{H₂O} is the mixing ratio of water vapour taken from the ERA interim data. For the mixing ratio of O₂ (M_{O₂}) a value of
21% is assumed. The O₄ concentration is then represented by the square of the O₂ concentration (Greenblatt et al., 1990). To
derive the O₄ VCD, the O₄ concentration is vertically integrated between the surface and 30 km with a vertical resolution of
155 20 m.

The temperature and pressure from the ECMWF ERA-Interim data set at the surface are also compared to the in situ
measurements on the ship. It is found that the ECMWF temperature is slightly lower (-0.7 K) and the ECMWF pressure is
slightly higher (+2 hPa) than the corresponding in situ measurements, see Fig. A4 in the appendix. Therefore, we repeated
our calculations of the O₄ profiles by shifting the ECMWF values for the whole profiles by +0.7 K and - 2 hPa. The
160 resulting change of the O₄ VCD is rather small (+0.3 %). The derived O₄ VCD for the modified profile is (1.245 ± 0.25)
 $\cdot 10^{43}$ molec²/cm⁵.

To estimate the uncertainty of the derived O₄ VCD the temperature and pressure of the whole profiles are varied by ± 2 K and
 ± 2 hPa, respectively. The resulting changes of the O₄ VCDs are ± 1.5 % and ± 0.9 %, respectively. In addition, assuming an
uncertainty of the atmospheric humidity profile of 30% leads to an uncertainty of the derived O₄ VCD of 0.9 %. Thus, we
165 estimate the total uncertainty of the O₄ VCD to ± 2 %.

Finally, a subtle detail should be mentioned: the integration of the O₄ VCD was performed starting from sea level, while the
instrument was located about 20 m above sea level. This rather small difference would result in a reduction of the O₄ VCD
by 0.4 %. However, this effect is considered in exactly the same way in the radiative transfer simulations, where the
instrument was also put at an altitude of 20 m, while the O₄ AMFs are calculated for the O₄ column starting from sea level.
170 Thus it is a consistent procedure to use the O₄ VCD integrated from sea level for the conversion of the measured O₄ dSCDs
into O₄ dAMFs.



6 Radiative transfer simulations

175 O₄ dSCDs are calculated using the full spherical Monte Carlo radiative transfer model MCARTIM (Deuschmann et al. 2011). For the simulations the profiles of temperature, pressure, and O₄ as described in section 5 are used. The vertical resolution was set to 20 m close to the surface and increases with altitude (see Table 2). The surface albedo was set to 0.05. The simulations were performed for the exact SZA and relative azimuth angles of the individual measurements. From the obtained O₄ AMFs, the corresponding O₄ dAMFs are calculated by subtracting the simulated O₄ AMFs for the zenith
180 viewing direction. To achieve best consistency with the measurements, for the simulation of the zenith measurements (interpolated between the zenith observations before and after the sequence) the SZA and relative azimuth angle for the exact time of the non-zenith measurements are also used for the simulations of the zenith measurements. The temporal evolution of the SZA and relative azimuth angle for 2 May are shown in Fig. A5 in the appendix.

It should be noted that it is important to use a consistent treatment of the SZA and relative azimuth angles in the simulations
185 and measurement analyses. Especially the choice of the Fraunhofer reference spectra is important. If e.g. either zenith measurements before or after the selected elevation sequence are used as reference spectra, systematic deviations of the retrieved O₄ dSCDs of up to 10% can occur (see Fig. A6 in the appendix).

O₄ dAMFs are simulated for two aerosol extinction profiles as well as for a pure Rayleigh atmosphere. For the extraction of the aerosol extinction profiles, the observations by the sun photometer and the ceilometer were used (see section 6.1). For the
190 simulations including aerosols, the phase function is represented by a Henyey-Greenstein (HG) parameterisation with an asymmetry parameter of 0.68. The single scattering albedo was set to 0.95. Variations of these properties lead to changes of the simulated O₄ dSCDs by up to ± 3 % (see Fig. A7 in the appendix). These rather low uncertainties are related to the low AOD on 2 May 2019. For measurements with higher aerosol loads, the corresponding uncertainties are usually much larger (e.g. Wagner et al., 2019). Here it should be noted that the HG phase function model is a rather simplified approximation for
195 true aerosol phase functions. Thus especially for measurements with small scattering angles (e.g. around noon on 2 May 2019) the uncertainties of the RTM simulations might be larger.

6.1 Extraction of the aerosol extinction profiles

200 Figure 4 presents the hourly averaged and range corrected ceilometer backscatter profiles for three periods in the afternoon on 2 May 2019 without cloud contamination. In a first order approximation, these backscatter profiles are proportional to the aerosol extinction. Thus together with the total AOD from the sun photometer measurements, the aerosol extinction profiles can be determined. However, ceilometer measurements are affected by several instrumental limitations, which complicate the direct conversion to aerosol extinction profiles:

205 a) Due to the missing overlap between the outgoing beam and the field of view of the detector, the sensitivity of the ceilometer is very low for altitudes below 500 m. Thus for this altitude range, no information on the aerosol extinction can be derived from the ceilometer measurements.

b) In spite of the long averaging period, still strong noise appears for altitudes above 3 km.

Due to these limitations, the ceilometer profiles can only be used for a restricted altitude range. In the following we used the
210 ceilometer profiles for the altitude range between 500 m and about 7 to 9 km. Between 500 m and 3000 m, averages for 100 m layers are calculated. Below 500 m, the values at 500 m are either set constant for the layer below, or are linearly extrapolated from the ceilometer data between 500 m and 800 m (similar as in Wagner et al., 2019). Since between 3 km and 10 km the noise increases strongly, a third order polynomial was fitted to the ceilometer data in that height range. The polynomial values are used for the altitude range for which positive values are obtained. Between 7 and 9 km the polynomial



215 values for the three profiles cross zero. Above these altitudes, the the profile values are set to zero. These extraction steps are illustrated in Fig. A8 in the appendix.

Before the backscatter profiles are normalised with the total AODs measured by the sun photometer, the stratospheric part of the total aerosol profile has to be added. This step is usually not important, because in more polluted areas the total AOD is clearly dominated by the tropospheric part. However, for our study, the total AOD is so low that the stratospheric part
220 constitutes a substantial fraction (up to 25 %) of the total AOD. Thomason et al. (2018) report the stratospheric AODs in the Tropics at 525 nm to be about 0.005 to 0.006. Assuming an Angström exponent of 2 the corresponding AOD at 360 nm is estimated to around 0.011 and 0.013. In the following we used a value of 0.012.

This value is subtracted from the total AOD (Fig. 3) measured by the sun photometer. Then the tropospheric aerosol profiles (as described above, see also Fig. A8) are normalised by the resulting tropospheric AOD. Finally, the stratospheric extinction
225 profile is added to the normalised tropospheric aerosol extinction profiles. For the stratospheric extinction profile we used a simplified shape with an AOD of 0.012. Here it is important to note that the details of the extinction profile in the upper troposphere and stratosphere are not critical. For example, the simulated O₄ dAMFs using aerosol profiles with or without the stratospheric part are almost the same. The final aerosol extinction profiles used for the RTM simulations are shown in Fig. 4.

230

6.2 Calculation of effective temperatures for the O₄ absorption

Since the temperature of the troposphere decreases with altitude, and the O₄ absorption cross section depends on temperature, the retrieved O₄ dSCDs might deviate from the true O₄ dSCDs (the integrated O₄ concentration along the
235 atmospheric light paths), because only one O₄ cross section for a fixed temperature is used in the spectral analysis. Thus, before the O₄ dAMFs from the measured spectra are compared to those from the radiative transfer simulations, the effect of the temperature dependence of the O₄ absorption has to be investigated.

The effective temperature of the O₄ measurements is calculated according to:

$$240 \quad T_{\text{eff},\alpha} = \frac{\sum_z [O_4]_z \cdot (bAMF_{z,\alpha} - bAMF_{z,90^\circ}) \cdot T_z}{\sum_z [O_4]_z \cdot (bAMF_{z,\alpha} - bAMF_{z,90^\circ})} \quad (2)$$

Here $[O_4]_z$ represents the O₄ concentration at altitude z , $bAMF_{z,\alpha}$ the box-AMF for elevation angle α at altitude z , and T_z the temperature at altitude z . $T_{\text{eff},\alpha}$ is the effective temperature for the measured O₄ dSCD at elevation angle α .

Equation 2 is applied for each individual measurement, the results are shown in Fig. A9. The effective temperatures range
245 from 276 K to 299 K. They depend systematically on the elevation angle and SZA. Measurements at low elevation angles are most sensitive for the layers near the surface, at which the highest temperatures occur. Measurements at high SZA (towards the end of the considered time period) have higher sensitivities for higher atmospheric layers with colder temperatures. Both dependencies are well represented by the results shown in Fig. A9.

To correct the effect of the temperature dependence, the correction factors presented in Fig. 13 in Wagner et al. (2019) are
250 applied to the O₄ dSCDs retrieved with the O₄ cross section for 293 K. The corrected O₄ dSCDs differ by up to a few percent (between -2 % and +7 %) from the original O₄ dSCDs. In Fig. A10 in the appendix the effect of the temperature correction is shown for two selected elevation sequences. For the comparison with the radiative transfer simulations the temperature-corrected O₄ dSCDs (or dAMFs) are used.

255 7 Comparison results



7.1 Direct comparison between measurements and RTM results

260 In Fig. 6 the O_4 dAMFs derived from the MAX-DOAS measurements are compared to those obtained from the radiative transfer simulations for elevation sequences not affected by clouds (similar comparisons for the sequences with cloud-contaminated measurements are shown in Fig. A11 in the appendix).

265 In the left part of Fig. 6, the results from radiative transfer simulations without aerosols are shown. Here, for almost all cases, the measured O_4 dAMFs are systematically smaller than the simulated O_4 dAMFs. This is an important finding, because especially for the low elevation angles, the presence of aerosol scattering leads to a decrease of the O_4 dSCDs. Thus the simulations for a pure Rayleigh atmosphere represent an upper limit of the achievable O_4 dSCDs. Only for cloudy cases with a high probability for multiple scattering events higher O_4 dSCDs could occur, but such conditions can be ruled out here because of the absence of thick and vertically extended clouds. Thus the overestimation of the simulated O_4 dSCDs for a pure Rayleigh atmosphere by the measured O_4 dSCDs indicates a fundamental inconsistency between measurements and simulations. Similar results are found for the elevation sequences with cloud contamination (Fig. A11 in the appendix).

270 In the right part of Fig. 6, simulation results for the aerosol profiles extracted in section 6.1 are shown. Note the separate y-axes for the simulated O_4 dAMFs on the right side, for which the maxima are chosen to achieve best agreement between the measured and simulated O_4 dAMFs. The exact values of the axis maxima were determined by fitting the measured O_4 dAMFs to the simulated O_4 dAMFs for elevation angles $>4^\circ$. For these elevation angles the simulation results for the different profile shapes below 500m) are almost the same. Good qualitative agreement between measurements and simulation is found, especially for the aerosol profiles with constant extinction below 500 m. However, the absolute values differ strongly. The ratios between measured and simulated O_4 dAMFs are found to be between 0.8 and 0.86. Again, similar results are found for the elevation sequences with cloud contamination (Fig. A11 in the appendix).

275 The scaling factors derived from this comparison between measured and simulated O_4 dSCDs are presented as blue data points in Fig. 7.

280

7.2 Profile inversion with MAPA

We also applied our profile inversion algorithm, the Mainz profile algorithm (MAPA, Beirle et al., 2019), to the measured O_4 dAMFs. For that purpose, a new MAPA LUT had to be created, because the lowest AOD in the original LUT (0.05) is larger than all AODs observed on 2 May 2020. The new LUT includes AOD values from zero to 0.1 in steps of 0.02. MAPA provides the option to apply a fixed user-defined scaling factor, or to determine a scaling factor yielding best match between forward model and measurement during profile inversion.

285 In Fig. A12 in the appendix the retrieved extinction profiles are shown for different scaling factors. Here it should be noted that the individual measurements (not the sequences) with cloud contamination were skipped before the profile inversion. Only profiles with either 'valid' or 'warning' flags are shown (profiles with 'error' flags are not shown). In Fig. A13 in the appendix the retrieved AODs for the different scaling factors are compared to the tropospheric AODs from the sun photometer measurements (stratospheric AOD of 0.012 was subtracted). Also the RMS between the measured and simulated O_4 dAMFs are shown (right). The colour of the MAX-DOAS inversion results indicates the quality of the profile inversion.

290 Most valid profiles are obtained for scaling factors between 0.80 and 0.90, or for a free fitted (variable) scaling factor. For the inversions with larger scaling factors, rather high RMS are found. For most cases, the retrieved AODs are smaller than those measured by the sun photometer. However, here it should be noted that for these low aerosol extinctions, the information content of the measurements is probably too low to constrain the aerosol extinction profiles, especially for high altitudes.



300 The obtained scaling factors are shown in Fig. 7. Overall good agreement between both comparison methods is found. For
all elevation sequences, values of the scaling factor < 1 are found. For the direct comparison, the difference from unity is
mostly larger than 15 % and can thus not be explained by the uncertainties of the measurements and simulations.

8 Conclusions

305

We compared measured and simulated O_4 absorptions for one day with very low aerosol optical depth. For such conditions,
the uncertainties caused by imperfect knowledge of the aerosol properties play a smaller role than for comparison under
more polluted conditions.

310 One important result of the comparison was that for all measurements, the observed O_4 absorption was higher than the
simulation results for an atmosphere without aerosols. In the absence of optically thick clouds, the simulated O_4 dAMFs for
an atmosphere without aerosols constitutes an upper limit, since especially for the low elevation angles the inclusion of
aerosols leads to a decrease of the O_4 absorption. The observed discrepancies thus indicate a fundamental inconsistency
between simulations and measurements.

315 The measured O_4 absorptions are also compared to simulations including aerosol extinction profiles. The aerosol extinction
profiles were constrained by measurements of the sun photometer, the ceilometer and a climatology of stratospheric aerosols.
Again, a large discrepancy was found for the absolute values. However, for the relative dependence of the O_4 dAMFs on the
elevation angle good agreement could be achieved. For each elevation sequence, the ratio of simulated and measured O_4
dAMFs was calculated. For that purpose the elevation angles $> 4^\circ$ were used, for which the O_4 dAMFs are almost insensitive
to the profile shape in the lower atmospheric layers. For all elevation sequences, ratios of 0.85 or less were found. Similar
320 ratios were also obtained from the application of our profile inversion algorithm (MAPA) to the measurements. The
observed discrepancies cannot be explained by the uncertainties of measurements and/or simulations. Here it is important to
note that in the spectral analysis, we explicitly corrected for the (small) temperature dependence of the atmospheric O_4
absorption.

325 Our results indicate that something fundamental is missing/wrong in either the radiative transfer simulations or the spectral
analysis of the atmospheric O_4 absorptions. We did not find a clear reason for the discrepancies. One possible reason for the
discrepancies could be a systematically too small O_4 absorption cross section.

Author contributions

330 T. Wagner performed the measurements, data analysis and prepared the manuscript. S. Dörner prepared the MAX-DOAS
instrument and extracted the ERA interim data. S. Dörner and S. Donner contributed to the MAX-DOAS operation and data
analysis. S. Beirle performed the MAPA profile inversions. S. Kinne operated the sun photometer and ceilometer.

335 Acknowledgements

The scientific party of RV MARIA S. MERIAN Cruise MSM82/2 gratefully acknowledges the very friendly and most
effective cooperation with Captain Maaß and his crew. Their great flexibility and their perfect technical assistance
substantially contributed to make this cruise a scientific success. We also appreciate the valuable support by the Leitstelle
340 Deutsche Forschungsschiffe (German Research Fleet Coordination Centre) at the University of Hamburg. The expedition
was funded by the Deutsche Forschungsgemeinschaft.



References

- 345 Bogumil, K., J. Orphal, T. Homann, S. Voigt, P. Spietz, O. C. Fleischmann, A. Vogel, M. Hartmann, H. Bovensmann, J. Frerik and J. P. Burrows, Measurements of Molecular Absorption Spectra with the SCIAMACHY Pre-Flight Model: Instrument Characterization and Reference Data for Atmospheric Remote-Sensing in the 230-2380 nm Region, *J. Photochem. Photobiol. A.*, 157, 167-184, 2003.
- 350 Beirle, S., Dörner, S., Donner, S., Remmers, J., Wang, Y., and Wagner, T.: The Mainz profile algorithm (MAPA), *Atmos. Meas. Tech.*, 12, 1785–1806, <https://doi.org/10.5194/amt-12-1785-2019>, 2019.
- Berrisford, P., Dee, DP, Poli, P, Brugge, R, Fielding, M, Fuentes, M, Kållberg, PW, Kobayashi, S, Uppala, S, Simmons, A, The ERA-Interim archive Version 2.0, ERA Report Series No.1 Version 2.0 (<https://www.ecmwf.int/node/8174>), 2011.
- Chen, Z., Bhartia, P. K., Torres, O., Jaross, G., Loughman, R., DeLand, M., Colarco, P., Damadeo, R., and Taha, G.:
- 355 Evaluation of the OMPS/LP stratospheric aerosol extinction product using SAGE III/ISS observations, *Atmos. Meas. Tech.*, 13, 3471–3485, <https://doi.org/10.5194/amt-13-3471-2020>, 2020.
- Deutschmann, T., Beirle, S., Frieß, U., Grzegorski, M., Kern, C., Kritten, L., Platt, U., Pukite, J., Wagner, T., Werner, B., and Pfeilsticker, K.: The Monte Carlo Atmospheric Radiative Transfer Model McArtim: Introduction and Validation of Jacobians and 3D Features, *J. Quant. Spectrosc. Ra.*, 112, 1119–1137, doi:10.1016/j.jqsrt.2010.12.009, 2011.
- 360 Donner, S.: Mobile MAX-DOAS measurements of the tropospheric formaldehyde column in the Rhein-Main region, Master's thesis, University of Mainz, <http://hdl.handle.net/11858/00-001M-0000-002C-EB17-2>, 2016.
- Greenblatt G.D., Orlando, J.J., Burkholder, J.B., and Ravishankara, A.R.: Absorption measurements of oxygen between 330 and 1140 nm, *J. Geophys. Res.*, 95, 18577-18582, 1990.
- Hönninger, G. and Platt, U.: Observations of BrO and its vertical distribution during surface ozone depletion at Alert,
- 365 *Atmospheric Environment*, 36, 2481 – 2489, 2002.
- Krastel, S., RV Maria S. Merian-Cruise MSM82/2, short cruise report, German Research Fleet Coordination Centre, University of Hamburg (<https://www.lfd.uni-hamburg.de/merian/wochenberichte/wochenberichte-merian/msm82-2-msm84/msm82-2-scr.pdf>), 2019.
- Lampel, J., Pöhler, D., Polyansky, O. L., Kyuberis, A. A., Zobov, N. F., Tennyson, J., Lodi, L., Frieß, U., Wang, Y., Beirle,
- 370 S., Platt, U., and Wagner, T.: Detection of water vapour absorption around 363 nm in measured atmospheric absorption spectra and its effect on DOAS evaluations, *Atmos. Chem. Phys.*, 17, 1271-1295, <https://doi.org/10.5194/acp-17-1271-2017>, 2017.
- Ortega, I., Berg, L. K., Ferrare, R. A., Hair, J. W., Hostetler, C. A., and Volkamer, R.: Elevated aerosol layers modify the O₂-O₂ absorption measured by ground-based MAX-DOAS, *J. Quant. Spectrosc. Ra.*, 176, 34–49,
- 375 <https://doi.org/10.1016/j.jqsrt.2016.02.021>, 2016.
- Polyansky, O.L., A.A. Kyuberis, N.F. Zobov, J. Tennyson, S.N. Yurchenko and L. Lodi ExoMol molecular line lists XXX: a complete high-accuracy line list for water, *Monthly Notices of the Royal Astronomical Society*, 480 (2), 2597-2608, 2018.
- Smirnov, A., B. N. Holben, I. Slutsker, D. M. Giles, C. R. McClain, T. F. Eck, S. M. Sakerin, A. Macke, P. Croot, G.
- 380 Zibordi, P. K. Quinn, J. Sciare, S. Kinne, M. Harvey, T. J. Smyth, S. Piketh, T. Zielinski, A. Proshutinsky, J. I. Goes, N. B. Nelson, P. Larouche, V. F. Radionov, P. Goloub, K. Krishna Moorthy, R. Matarrese, E. J. Robertson, and F. Jourdin
- Maritime Aerosol Network as a component of Aerosol Robotic Network, *J. Geophys. Res.*, 114, D06204, doi:10.1029/2008JD011257, 2009.
- Thalman, R. and Volkamer, R.: Temperature dependent absorption cross-sections of O₂-O₂ collision pairs between 340 and
- 385 630 nm and at atmospherically relevant pressure, *Phys. Chem. Chem. Phys.*, 15, 15371, doi:10.1039/c3cp50968k, 2013.



- Thomason, L. W., Ernest, N., Millán, L., Rieger, L., Bourassa, A., Vernier, J.-P., Manney, G., Luo, B., Arfeuille, F., and Peter, T.: A global space-based stratospheric aerosol climatology: 1979–2016, *Earth Syst. Sci. Data*, 10, 469–492, <https://doi.org/10.5194/essd-10-469-2018>, 2018.
- Vandaele, A. C., C. Hermans, P. C. Simon, M. Carleer, R. Colin, S. Fally, M.-F. Mérienne, A. Jenouvrier, and B. Coquart, 390 Measurements of the NO₂ Absorption Cross-section from 42000 cm⁻¹ to 10000 cm⁻¹ (238-1000 nm) at 220 K and 294 K, *J. Quant. Spectrosc. Radiat. Transfer*, 59, 171-184, 1997.
- Wagner, T., Beirle, S., and Deutschmann, T.: Three-dimensional simulation of the Ring effect in observations of scattered sun light using Monte Carlo radiative transfer models, *Atmos. Meas. Tech.*, 2, 113-124, 2009.
- Wagner, T., Apituley, A., Beirle, S., Dörner, S., Friess, U., Remmers, J., and Shaiganfar, R.: Cloud detection and 395 classification based on MAX-DOAS observations, *Atmos. Meas. Tech.*, 7, 1289-1320, doi:10.5194/amt-7-1289-2014, 2014.
- Wagner, T., Beirle, S., Remmers, J., Shaiganfar, R., and Wang, Y.: Absolute calibration of the colour index and O₄ absorption derived from Multi AXis (MAX-) DOAS measurements and their application to a standardised cloud classification algorithm, *Atmos. Meas. Tech.*, 9, 4803-4823, doi:10.5194/amt-9-4803-2016, 2016.
- Wagner, T., Beirle, S., Benavent, N., Bösch, T., Chan, K. L., Donner, S., Dörner, S., Fayt, C., Frieß, U., García-Nieto, D., 400 Gielen, C., González-Bartolome, D., Gomez, L., Hendrick, F., Henzing, B., Jin, J. L., Lampel, J., Ma, J., Mies, K., Navarro, M., Peters, E., Pinardi, G., Puentedura, O., Puķīte, J., Remmers, J., Richter, A., Saiz-Lopez, A., Shaiganfar, R., Sihler, H., Van Roozendael, M., Wang, Y., and Yela, M.: Is a scaling factor required to obtain closure between measured and modelled atmospheric O₄ absorptions? An assessment of uncertainties of measurements and radiative transfer simulations for 2 selected days during the MAD-CAT campaign, *Atmos. Meas. Tech.*, 12, 2745–2817, [https://doi.org/10.5194/amt-12-2745-](https://doi.org/10.5194/amt-12-2745-2019) 405 2019, 2019.
- Wang, Y., Beirle, S., Hendrick, F., Hilboll, A., Jin, J., Kyuberis, A. A., Lampel, J., Li, A., Luo, Y., Lodi, L., Ma, J., Navarro, M., Ortega, I., Peters, E., Polyansky, O. L., Remmers, J., Richter, A., Puentedura, O., Van Roozendael, M., Seyler, A., Tennyson, J., Volkamer, R., Xie, P., Zobov, N. F., and Wagner, T.: MAX-DOAS measurements of HONO slant column densities during the MAD-CAT campaign: inter-comparison, sensitivity studies on spectral analysis settings, and error 410 budget, *Atmos. Meas. Tech.*, 10, 3719-3742, <https://doi.org/10.5194/amt-10-3719-2017>, 2017.
- Wang, Y., Apituley, A., Bais, A., Beirle, S., Benavent, N., Borovski, A., Bruchkouski, I., Chan, K. L., Donner, S., Drosoglou, T., Finkenzeller, H., Friedrich, M. M., Frieß, U., Garcia-Nieto, D., Gómez-Martín, L., Hendrick, F., Hilboll, A., Jin, J., Johnston, P., Koenig, T. K., Kreher, K., Kumar, V., Kyuberis, A., Lampel, J., Liu, C., Liu, H., Ma, J., Polyansky, O. L., Postlyakov, O., Querel, R., Saiz-Lopez, A., Schmitt, S., Tian, X., Tirpitz, J.-L., Van Roozendael, M., Volkamer, R., 415 Wang, Z., Xie, P., Xing, C., Xu, J., Yela, M., Zhang, C., and Wagner, T.: Inter-comparison of MAX-DOAS measurements of tropospheric HONO slant column densities and vertical profiles during the CINDI-2 campaign, *Atmos. Meas. Tech.*, 13, 5087–5116, <https://doi.org/10.5194/amt-13-5087-2020>, 2020.

420

425

430



435

Tables

Table 1: Settings for the DOAS analysis of O₄

Parameter	Value, Remark / Reference
Spectral range	352 – 387 nm
Degree of DOAS polynomial	5
Degree of intensity offset polynomial	2
Fraunhofer reference spectrum	Interpolated between 90° measurement before and after each elevation sequence
Wavelength calibration	Fit to high resolution solar spectrum using Gaussian slit function
Shift / squeeze	The measured spectrum is shifted and squeezed against all other spectra
Ring spectrum 1	Normal Ring spectrum calculated from measured zenith spectrum
Ring spectrum 2	Ring spectrum 1 multiplied by λ^{-4} (Wagner et al. (2009))
O ₃ cross section	223 K, Bogumil et al. (2003)
NO ₂ cross section	294 K, Vandaele et al. (1997)
H ₂ O cross section	293 K, Polyansky et al. (2018)
O ₄ cross section	293 K, Thalman and Volkamer (2013)

440

Table 2: Vertical resolution used for the radiative transfer simulations

Altitude range [km]	Vertical resolution [km]
0 - 0.5	0.02
0.5 - 2	0.1
2 - 12	0.2
12 - 25	1
25 - 45	2
45 - 100	5

445

450

455

460



465
Figures

470





Fig. 1: Left: Ship route from Montevideo to Las Palmas. The blue circle indicates the location of the measurements used in this study. Right: Aerosol optical depth at 380 nm measured with a hand-held sun photometer.

475


480



485


Fig. 2: The position and viewing direction of the MAX-DOAS instrument on the RV Maria S. Merian during the ship cruise (ship drawing taken from <https://briese-research.de/research-department/research-vessels/rv-maria-s-merian>).

490



495 Fig. 3: Top: AOD at 360 nm measured from the hand-held sun photometer. The data were extrapolated from the measurements at 380 nm using the Angström coefficient calculated from 380 nm and 440 nm. Bottom: Range-corrected ceilometer backscatter profile at 1064 nm.

500




505 **Fig. 4: Hourly averaged and range corrected ceilometer backscatter profiles for 3 periods on 2 May 2019 without cloud contamination. The thin lines represent the raw data. The dotted lines represent the smoothed profiles (averages over 100 m).**

510

515

520

525



530

Fig. 5: Complete aerosol extinction profiles for the three time periods without clouds after all corrections are applied. The green curves represent the profiles with linear extrapolation below 500 m; the red curves represent profiles with constant values below 500 m.

535











540


545

550



555






560

Fig. 6: Comparison of the measured and simulated O_4 dAMFs for selected elevation sequences without cloud contamination. In the left part, the measured O_4 dAMFs are compared to simulations for a pure Rayleigh atmosphere. In the right part they are compared to simulation results including aerosols (two profiles with either constant or linearly extrapolated aerosol extinction below 500 m). Note that in the right part separate y-axes on the right sides are used for the simulation results. The maxima of the right y-axes are chosen to achieve best agreement between the measured and simulated O_4 dAMFs (see text).

565

570



575

Fig. 7: Scaling factors derived from the direct comparison between the measured and simulated O_4 dSCDs (blue) and from the MAPA profile inversion (green) for all elevation sequences shown in Fig. 6 and Fig. A11. Many of the measurements of the two last elevation sequences are affected by clouds.

580



585 Appendix A1 Effect of including a H₂O cross section or a second O₄ cross section on the retrieved O₄ dSCDs

H₂O cross section

Recent studies found evidence for substantial atmospheric H₂O absorptions in measured spectra in the UV (Lampel et al.,
590 2017, Wang et al., 2017, 2020). These absorptions are usually rather small, but especially for measurement conditions with
high atmospheric humidity the inclusion of a H₂O cross section in the spectral analysis can be useful.

Fig. A1 presents examples of the spectral analysis with either a H₂O cross section included or excluded. A clear H₂O
absorption signal is found around 363 nm. The H₂O dSCDs retrieved at 363 nm agree reasonably well ($r^2=0.63$) with those
retrieved at 442 nm (see Fig. A2) with a similar slope (2.07) as presented in Lampel et al. (2017) who found a slope of 2.39.

595 If the H₂O cross section is not included in the analysis, a systematic structure appears in the residual. Thus, in this study, a
water vapour cross section (Polyansky et al. 2018) is included in the spectral analysis. Here it should be noted that compared
to other locations, the water vapour absorption during the ship cruise was rather high because most of the measurements
were carried out under conditions of high atmospheric temperature and humidity. At other, colder locations, the impact of
the H₂O absorption might be negligible.

600 Although the H₂O absorption is clearly found in the spectral analysis, the effect of including a H₂O cross section or not on
the retrieved O₄ dSCDs is still rather small. If a H₂O cross section is included, the retrieved O₄ dSCDs are about 2.5 % larger
than without a H₂O cross section included.

O₄ cross section at low temperature


605

We also investigated the effect of including a second O₄ cross section for low temperature (203 K). Before using this cross
section in the fit it was orthogonalised with respect to the O₄ cross section at 293 K. Including the additional O₄ cross section
leads to only small changes of the retrieved O₄ dSCD of about -1.5 %. Here it is interesting to note that the retrieved O₄
dSCDs for the O₄ cross section at low temperature were negative and the absolute values much smaller ($<2 \cdot 10^{43}$ molec²/cm⁵)

610 than those at high temperature ($<6 \cdot 10^{43}$ molec²/cm⁵). The largest negative O₄ dSCDs for the O₄ cross section at low
temperature were found indicating that the effective atmospheric temperatures decrease with elevation angle (see section
6.2). Also, the correlation between both O₄ dSCDs ($r^2=0.20$) is very low. Thus we conclude that the measured spectra do not
contain significant O₄ absorptions at low temperatures. For the interpretation of this finding it should be noted that low
temperatures exist only at higher atmospheric layers. The O₄ absorptions at these layers mostly cancel out in the spectral
analysis, because the light paths of the measured spectra and the Fraunhofer reference spectra at these layers are very similar.

615 This explains that the retrieved O₄ absorptions at cold temperatures are very small. To further confirm this hypothesis, we
calculated the effective temperatures for the O₄ absorptions on 2 May 2019 (see section 6.2) and found them to be very close
to the temperature of the high temperature O₄ cross section (293 K). Based on these findings, the O₄ results in this study are
retrieved without including a second O₄ cross section at low temperature. It should, however, be noted that for measurements

620 at other locations and seasons including a second O₄ cross section in the spectral analysis might be meaningful.



625 Fig. A1: Fit results for a spectrum taken on 2 May 2019, 13:14:50, at an elevation angle of 1° (SZA: 33.6°). Left: results if a H₂O cross section is included in the spectral analysis; right: results if no H₂O cross section is included in the spectral analysis. The black lines represent the fitted cross section, the red lines indicate the residual (bottom) or the residual plus the fitted cross section.

630




Fig. A2: Correlation plot of the H₂O dSCDs retrieved at 363 nm versus those retrieved at 442 nm for the whole ship cruise. The regression line is fitted assuming that the H₂O dSCDs retrieved at 442 nm have no error.



635

Appendix A2 Additional Figures

640




Fig. A3: Time series of the retrieved O_4 dSCD on 2 May 2019 for the different elevation angles. During the afternoon, for most of the time smooth variations are found. However, for some times and elevation angles systematic deviations of the O_4 dSCDs occur, which are caused by scattered clouds.

645






Fig. A4: Diurnal variation of the surface pressure and temperature from in situ measurements on the ship. The green box indicates the period of the MAX-DOAS measurements used in this study. The corresponding values from the ECMWF model simulations are 1012.8 hPa and 299.8 K, respectively.



655


Fig. A5: Top: Variation of the latitude and longitude of the ship position during 2 May 2019. Bottom: Corresponding variation of the SZA and relative azimuth angle (RAA).

660



665


Fig. A6: Effect of using different Fraunhofer reference spectra for the analysis of individual elevation sequences. Shown are the ratios of the obtained O_4 dSCDs for different selections versus those for Fraunhofer reference spectra interpolated between the zenith measurements before and after the selected elevation sequence. Before: zenith measurement before the sequence is used; After: zenith measurement after the sequence is used; Average before and after: the average of the zenith measurements before and after the sequence is used.



670


Fig. A7: Effect of different phase functions and single scattering albedos on the O_4 dSCDs. Shown are the ratios for simulations with variations of asymmetry parameter (AP) and single scattering albedo (SSA) versus simulations using the standard settings (AP= 0.68, SSA: 0.95). The results are for SZA of 64.5° and RAA of 87.7° (around 18:00 on 2 May 2019). The results for other SZA/RAA combinations during the afternoon of 2 May 2019 are similar.

675



680

Fig. A8: The light blue data show the original backscatter profile averaged between 14:00 and 15:00. The blue dots show the smoothed (with a 100m kernel) profile, which are used between 500 m and 3 km. Below 500m either constant or linearly extrapolated data (see text) are used. Between 3 km and 10 km a third order polynomial is fitted to the raw data. The polynomial values are used between 3 km and the altitude at which they become negative. Above the values are set to zero.



685

Fig. A9: Effective temperatures calculated for the individual measurements according to equation 2.

690





Fig. A10: Effect of the temperature correction for two selected elevation sequences.

695

700




705



710

Fig A11: Comparison of the measured and simulated O_4 dAMFs for five elevation sequences with few cloud-contaminated measurements (indicated by the red arrows). In the left part, the measured O_4 dAMFs are compared to simulations for a pure Rayleigh atmosphere. In the right part they are compared to simulation results including aerosols (two profiles with either constant or linearly extrapolated aerosol extinction below 500 m). Note that in the right part separate y-axes on the right sides are used for the simulation results. The maxima of the right y-axes are chosen to achieve best agreement between the measured and simulated O_4 dAMFs (see text).


715



720

Fig A12: Extinction profiles retrieved with MAPA for the selected elevation sequences for different scaling factors. Only profiles for inversions with ,valid' or ,warning' flags are shown.

725



730

Fig A13: Right: Comparison of the retrieved AOD for the MAPA profile inversions with different scaling factors (squares) and the (tropospheric) AOD observed by the sun photometer (blue lines). Right: RMS between the measured and fitted O₄ dAMFs. The colours indicate the quality flags for the individual profile inversions.

735