

# Ash particle refractive index model for simulating the brightness temperature spectrum of volcanic ash clouds from satellite infrared sounder measurements

5 Hiroshi Ishimoto<sup>1</sup>, Masahiro Hayashi<sup>1</sup>, Yuzo Mano<sup>1</sup>

<sup>1</sup>Meteorological Research Institute, Tsukuba, 305-0052, Japan

*Correspondence to:* Hiroshi Ishimoto (hiroishi@mri-jma.go.jp)

**Abstract.** Using data from the Infrared Atmospheric Sounding Interferometer (IASI) measurements of volcanic ash clouds (VACs) and radiative transfer calculations, we attempt to simulate the measured brightness temperature spectra (BTS) of volcanic ash aerosols in the infrared region. In particular, the dependence on the ash refractive index (RI) model is investigated. We apply 21 RI models for the optical properties of ash particles, including recently published models. The results of BTS simulations are evaluated using the root mean square (RMS) of the brightness temperature difference between measurements and simulations for channels in the wavenumber range of 750–1400 cm<sup>-1</sup>. The results of numerical simulations for 156 pixels of IASI measurements for ash plumes from seven volcanoes indicated that some newly established RI models have smaller RMS than those obtained with the conventional ash RI model. For some eruption events, the RI models deduced from the chemical composition of ash samples for the same volcanic eruption events simulate the measured BTS well. The results suggest that infrared sounder measurements and precise BTS simulations of the VACs contribute to estimating the appropriate RI model in satellite VAC analysis.

## 20 1. Introduction

For continuous near-real-time monitoring of volcanic ash clouds (VACs) during both day and night, satellite observations obtained using infrared sensors are particularly useful (Mackie and Watson, 2014; Mackie et al., 2016; Clarisse and Prata, 2016). Silicate has a characteristic absorption band around 10 μm due to the stretching vibration of the Si-O bond; thus, VACs containing fine ash particles can be easily detected and distinguished from water or ice clouds by measuring the brightness temperature difference between two channels in this absorption band, such as 10.8 μm and 12 μm (Wen and Rose, 1994; Prata and Grant, 2001). Applying further constraints related to the atmospheric profile and microphysical properties of the ash particles, such as the particle size distribution, the physical properties of VACs (plume height, particle effective radius, optical depth, and the associated ash mass loading) have been determined using measurements from multiple infrared channels from

high-resolution satellite imagers and from satellite infrared sounders (e.g. Francis et al., 2012; Pavolonis et al., 2013; Clarisse et al., 2013; Ventress et al., 2016)). These retrieval methods are based on the results of radiative transfer calculations for the measured infrared brightness temperature of VACs under assumed atmospheric profiles and surface conditions; as such, the optical properties of the ash particles are essential. In particular, the complex refractive index (RI) is an important upstream parameter for estimating ash optical properties and their wavelength dependence.

A typical RI model over a wide range of infrared wavelengths, the ‘andesite’ model proposed by Pollack et al. (1973) (hereinafter PL1973) has been adopted for radiative transfer calculations for volcanic ash analysis. Although the PL1973 andesite model provides suitable approximations for the analysis of some VACs (Pavolonis et al., 2013), simulations of the infrared brightness temperature spectrum (BTS) of VACs based on radiative transfer calculations from the PL1973 andesite model often causes large discrepancies relative to calculations based on satellite infrared sounder measurements (Gangale et al., 2010; Newman et al., 2012; Ishimoto et al., 2016; Clarisse and Prata, 2016). For simulating a measured BTS to retrieve the physical properties of a VAC from satellite data, RI models that account for variations in the chemical composition of volcanic ash are essential. Optical properties of ash particles calculated using an appropriate RI model for an observed VAC support accurate and robust determination of the physical properties of the VAC in satellite image analysis. Furthermore, a good RI model leads to improvements in existing retrieval algorithms for volcanic SO<sub>2</sub> gas levels using the  $\nu_1$  SO<sub>2</sub> absorption band around 8.7  $\mu\text{m}$  ( $\sim 1152\text{ cm}^{-1}$ ) through accurate estimation of the ash contribution to the measured brightness temperature (Corradini et al., 2009; Clarisse et al., 2012).

In recent years, two RI datasets for volcanic ash over a broad infrared wavelength range have been proposed. One dataset is built from the results of laboratory experiments using ash sampled from several volcanoes (Reed et al., 2017, 2018). Based on the results of Reed et al. (2018), Prata et al. (2019) constructed another RI dataset parameterised using the SiO<sub>2</sub> content or ratio of non-bridging oxygens to tetrahedrally coordinated cations (NBO/T). If an appropriate RI model for analysing a measured VAC from satellite data is provided by the RI dataset of Reed et al. (2018), the RI model can be estimated based on the similarity with the rock type of the target volcano. Furthermore, if the SiO<sub>2</sub> content or NBO/T parameterisations of Prata et al. (2019) are effective, we can identify the RI model, for example, from the chemical composition of ash or pumice samples from ongoing or past eruption events. However, whether the selected RI model accurately reproduces satellite measurements via radiative transfer calculations remains unclear.

The most appropriate satellite measurement type to discuss such reproducibility is a BTS obtained using an infrared sounder. In this work, we attempt to simulate the observed BTS through radiative transfer calculations using the RI models from recently proposed RI datasets without compositional constraints for the ash aerosols. If the RI model obtained from infrared sounder measurements corresponds to that derived from SiO<sub>2</sub> content or NBO/T for a sample from the same volcano, the results are considered to support the validity of the RI model based on ash chemical composition. Volcanic ash particles in the atmosphere may have different optical properties compared to sampled ash particles due to modifications in the atmosphere, such as the condensation of ice, water, or sulphate on ash particles. Even pure ash particles identified from satellite infrared observations may have compositions different from those of ash or pumice samples (Carn and Krotkov, 2016). The particle sizes are

generally small for ash plumes, which cause the large negative brightness temperature difference between two infrared split-window channels. The estimated effective radii of the ash plumes are generally less than 5  $\mu\text{m}$  depending on the plume height and optical depth (Clarisse and Prata, 2016). For such fine ash particles, silicic micropumice, and small glass shards may remain in the atmosphere longer than bulk composition particles due to their density (Cashman and Rust, 2016). Thus, volcanic ash particles detected in infrared satellite measurements after long-distance transportation may have a more felsic composition than do ash particles sampled near a volcano. Conversely, the differences in the RI models used may not be statistically distinguishable from the reproduced BTS, because there are many unknowns and assumptions in the radiative transfer calculations. One of the objectives of this study was to investigate whether RI models that reproduce the observed BTS in the infrared region can be identified, and whether the compositional type (mafic or felsic) corresponding to that RI model is consistent with that of the volcanic ash sample or the type of volcano.

In this work, we use the time and places of VAC observation data from the Infrared Atmospheric Sounding Interferometer (IASI) on board the Meteorological Operation (MetOp) satellite to simulate measured BTS based on radiative transfer calculations. IASI has 8461 channels that range between 645 and 2760  $\text{cm}^{-1}$ , with a spectral sampling interval of 0.25  $\text{cm}^{-1}$ , and the ground resolution is 12 km at nadir (Hilton et al., 2012). In section 2, our numerical method for the radiative transfer calculations is explained. VAC parameters and RI models for the ash particles used in numerical simulations are discussed in section 3, and the selection of IASI pixels, which are used for analysis, is described in section 4. Results of case studies for some VACs are discussed in section 5.


## 2. Infrared spectral simulator Monochromatic-Based Cloud Radiation Model (MBCRM) for VAC analysis

To estimate VAC parameters and  $\text{SO}_2$  gas contents from satellite infrared sounder measurements, a fast radiative transfer model is needed that can calculate upward channel radiance at the top of the atmosphere (TOA), including multiple light scattering events among ash particles. Furthermore, we examine a number of RI models for volcanic ash to simulate the measured BTS. To meet these requirements, we developed a radiative transfer code, MBCRM, to simulate infrared sounder measurements. For the basic algorithm of MBCRM, we adopt the Atmospheric Radiative Transfer Simulator (Buehler et al., 2010). Upward radiance  $R_i$  of a sounder channel  $i$  at TOA is calculated as the weighted sum of  $N$  number of monochromatic radiances  $R_{v_j}$ .

$$R_i = \sum_{j=1}^N C_j R_{v_j} \quad (1)$$

where  $C_j$  is the weighting coefficient ( $1 = \sum C_j$ ). Using a diverse atmospheric profile dataset to compare the calculated  $R_i$  with values derived from exact integration of channel response functions using line-by-line calculations, the monochromatic wavenumber  $v_j$  and weight  $C_j$  were determined, as was the number  $N$ , using a simulated annealing technique. To simulate the radiance of the 8461 IASI channels, a total of 14,132 monochromatic wavenumber values were determined. Figure 1 shows the root-mean-square difference of the channel brightness temperature  $BT_i$  calculated from  $R_i$  for clear-sky conditions between


95 rigorous line-by-line calculations and MBCRM using the ECMWF 83 diverse atmospheric profiles dataset (Matricardi, 2008). We assume that the mixing ratios of the six atmospheric components ( $\text{CO}_2$ ,  $\text{N}_2\text{O}$ ,  $\text{CO}$ ,  $\text{CH}_4$ ,  $\text{O}_2$ , and  $\text{N}_2$ ) are fixed (the  $\text{CO}_2$  mixing ratio was set to 403.3 ppmv). For line-by-line calculations, the HITRAN 2012 database (Rothman et al., 2013) and same definitions of line parameters and correction formulas as those of LBLRTM ver. 12.4 (Clough et al., 2005) are applied, and MT-CKD2.5 (Mlawer et al., 2012) is used for the water vapour continuum. The radiances of IASI channels in Fig. 1 are  
 100 calculated by applying the IASI channel response function for the monochromatic line-by-line output with the wavenumber interval  $\sim 10^{-4} \text{ cm}^{-1}$ . The accuracy of MBCRM for calculating gas absorption was less than 0.4 K in wavenumber range  $650 \leq \nu \leq 2400 \text{ cm}^{-1}$ .



120 **Figure 1: The root-mean-square of the TOA brightness temperature for the IASI 7021 channels ( $645\text{--}2400\text{cm}^{-1}$ ) between line-by-line and MBCRM calculations. The satellite zenith angle  $0^\circ$  and surface emissivity 1.0 were applied, and the temperature of the atmosphere at surface pressure height was used as surface temperature. The brown line is the NEdT of IASI at 280 K (Hilton et al., 2012). The atmospheric profiles dataset is available from [https://nwp-saf.eumetsat.int/oldsite/deliverables/rtm/profile\\_datasets.html](https://nwp-saf.eumetsat.int/oldsite/deliverables/rtm/profile_datasets.html).**

125 A plane-parallel single homogeneous layer VAC was assumed for calculations of multiple scattering by ash particles. The values of  $\nu_j$  and  $C_j$  in Eq. (1) determined under clear-sky conditions can be applied to calculations in a cloudy atmosphere with high accuracy (Holl et al., 2012). We used the same formulation of Eq. (1) for radiance calculations in cloudy conditions

caused by volcanic ash. For monochromatic radiative transfer calculations at infrared wavelengths, we formulated an analytical expression of TOA radiance in which the terms for cloud multiple scattering were described by coefficients depending only on cloud geometry and cloud optical properties. A look-up-table (LUT) of coefficients for various ash cloud conditions was prepared for calculation of monochromatic radiance in advance using the discrete ordinate method with 64 streams. As a benchmark test for the calculations of TOA radiance for the atmosphere including the VAC layer, the simulations of IASI channel radiances between MBCRM and Radiative Transfer for the TIROS Operational Vertical Sounder (RTTOV) were compared using the optical properties of ‘new volcanic ash (VAPO)’ in RTTOV (Fig. 2).



**Figure 2: Comparisons of the IASI BTS simulated using MBCRM and RTTOV (ver.12.2) for clear sky ( $\tau_{11\mu\text{m}} = 0$ ) and cloudy ( $\tau_{11\mu\text{m}} = 1$ ) atmosphere. Optical depth was defined as that at wavelength  $11\ \mu\text{m}$  ( $\nu = 907.488\ \text{cm}^{-1}$ ). The average atmospheric profile in the ECMWF 52 diverse profile dataset (no. 52) and the satellite zenith angle  $0^\circ$  were used, and a sea surface emissivity model and surface temperature  $286.52\ \text{K}$  were applied. For cloudy atmosphere, a homogeneous aerosol layer between  $358.966\ \text{hPa}$  and  $459.712\ \text{hPa}$ , and the aerosol properties of the ‘new volcanic ash (VAPO)’ model in RTTOV (Vidot and Hocking, 2017) were considered. For RTTOV, result of 8 streams discrete-ordinate method (DOM) was plotted. For MBCRM calculations, we read the complex refractive index (RI) data of VAPO from the figure in the RTTOV-11 Science and Validation Report (Saunders et al., 2013).**

### 3. VAC parameters for radiative transfer calculations and RI models

165 Using LUTs for ash optical properties, numerical simulations of IASI measurements of VACs were conducted using MBCRM. A prolate spheroid shape with a 1:2 axis ratio and a log-normal size distribution (Hansen and Travis, 1974) with a geometric standard deviation of 0.74 (Pavolonis et al., 2013) were assumed for ash particles. The optical properties of the particles were calculated using the T-matrix method (Mishchenko et al., 2002) for particle effective radii  $r_{eff}$  between 0.1 and 20  $\mu\text{m}$  (particle size is defined as the radius of a sphere with equivalent volume). In satellite infrared sounder measurements, volcanic  
170 ash mainly affects TOA brightness temperature in the wavenumber range of  $\nu \leq 1400\text{cm}^{-1}$ , and LUTs of volcanic ash optical properties were prepared for  $640\text{ cm}^{-1} \leq \nu \leq 1400\text{ cm}^{-1}$  at an interval of  $\Delta\nu = 10\text{ cm}^{-1}$ . The optical properties of each IASI channel were estimated through linear interpolation of the LUT. TOA radiance and brightness temperature were calculated under the assumption that the ash plumes are in thermal equilibrium with the ambient air.

For RI models of volcanic ash particles, we used the datasets of Reed et al. (2018) and Prata et al. (2019) (hereinafter RE2018  
175 and PG2019). Additionally, we prepared datasets using two RI models, PLAND and PLRHY. These models were described as ‘andesite’ and ‘rhyolite (obsidian)’ models by PL1973. The RI models applied in this work are listed in Table 1, and the characteristics of the models are described below.

180

185

**Table 1: Abbreviations for the refractive index (RI) datasets and RI models. The RI models of Reed et al. (2018) (RE2018; RE010–RE080) are listed in alphabetical order and labelled with the name of the corresponding volcanic ash sample. Numbers used for the RI models based on the Prata et al. (2019) (PG2019) dataset (PG000–PG100) represent the ratio of non-bridging oxygens to tetrahedrally coordinated cations (NBO/T). The values of NBO/T and SiO<sub>2</sub> (wt.%) for the RI models of RE2018 were calculated  
190 from bulk components of the ash samples and derived by PG2019. PLAND and PLRHY are ‘andesite’ and ‘obsidian from Little Glass Mt. California’ by Pollack et al. (1973). The wavenumber dependence of each RI model is plotted in Figures 3–5.**

195

200

205

210

215

Dataset	RI model (original sample)	NBO/T	SiO <sub>2</sub>
RE2018 Reed et al. (2018)	RE010 (Askja)	0.13	72.35
	RE020 (Aso)	0.43	54.96
	RE030 (Eyjafjallajökull-a)	0.38	60.00
	RE040 (Eyjafjallajökull-b)	0.40	58.85
	RE050 (Grímsvötn)	0.75	50.33
	RE060 (Nisyros)	0.05	74.16
	RE070 (Spurr)	0.36	55.99
	RE080 (Tongariro)	0.17	64.13
PG2019 Prata et al. (2019)	PG100	1.00	-
	PG090	0.90	-
	PG080	0.80	-
	PG070	0.70	-
	PG060	0.60	-
	PG050	0.50	-
	PG040	0.40	-
	PG030	0.30	-
	PG020	0.20	-
	PG010	0.10	-
	PG000	0.00	-
PL1973 Pollack et al. (1973)	PLAND (andesite)		54.15
	PLRHY (rhyolite)		73.45

### 3.1 Refractive index dataset of Reed et al. (2018): RE2018


Using the assumptions of the Rayleigh continuous distribution of ellipsoids (CDE) scattering model for sampled ash particles and the Lorentz formulation for RI to ensure consistency with the Kramers–Kronig relationship, RE2018 constructed a spectral RI dataset of typical volcanic ash samples for wavelengths of 0.33–19  $\mu\text{m}$  (526 – 3030  $\text{cm}^{-1}$ ). RI data for eight ash samples from seven volcanoes (Askja, Aso, Eyjafjallajökull-(a) and -(b), Grímsvötn, Nisyros, Spurr, and Tongariro) are available from the Aerosol Refractive Index Archive (<http://eodg.atm.ox.ac.uk/ARIA/index.html>), as well as in the Supporting Information of PG2019 (<https://doi.org/10.1029/2018JD028679>). The spectral RIs of RE2018 over the wavenumber range of  $650 \text{ cm}^{-1} \leq \nu \leq 1400 \text{ cm}^{-1}$  are plotted in Figure 3. In the wavenumber range of  $700 \text{ cm}^{-1} \leq \nu \leq 800 \text{ cm}^{-1}$ , the RI models of RE2018 showed weak absorption features that were not clearly expressed in the andesite and rhyolite models of PL1973. In previous attempts to simulate sounder-observed BTS, the calculated BTS tended to exhibit a positive bias in this wavenumber range (Gangale et al., 2010; Clarisse et al., 2013; Ishimoto et al., 2016; Clarisse and Prata, 2016). With the RE2018 RI models, improvements in BTS fitting are expected, although the relationships of chemical composition and volcano type (mafic or

felsic) with the absorption index (imaginary part of the complex RI:  $k$ ) at  $700\text{ cm}^{-1} \leq \nu \leq 800\text{ cm}^{-1}$  remain unclear. All RI models based on the RE2018 dataset have relatively weak absorption peaks ( $k \leq 1.0$ ) and similar wavenumber dependence of  $k$  between 1110 and  $1250\text{ cm}^{-1}$ . The individual RI models of RE2018, RE020 (Aso), and RE080 (Tongariro) show a similar wavenumber dependence of  $k$ , although their NBO/T and  $\text{SiO}_2$  wt.% values differ significantly. In addition,  $k$  from RE050 (Grímsvötn) exhibits a broad feature that differs from those of the other RI models of RE2018.

235

240

245



**Figure 3: Real  $n$  (left) and imaginary  $k$  (right) parts of the RE2018 complex RI models ( $RI = n + i \cdot k$ ). For Eyjafjallajökull ash samples, the yellow solid and dashed lines indicate Eyjafjallajökull-(a) and -(b), respectively. See Table 1 for the abbreviations, NBO/T values, and  $\text{SiO}_2$  wt.% of each RI model.**

250

### 3.2 RI models parameterised by Prata et al. (2019): PG2019

Using the RE2018 RI dataset and the results of laboratory analysis of the chemical composition of ash samples, Prata et al. (2019) provided a new parameterisation for the ash RI, based on  $\text{SiO}_2$  content and NBO/T. According to Prata et al. (2019), NBO/T and  $\text{SiO}_2$  wt.% in the bulk composition of the ash samples exhibited stronger correlations with the derived RI over a broad spectral range compared to glass composition. Furthermore, parameterisation using NBO/T provides better results compared to using  $\text{SiO}_2$  wt.% at both visible and infrared wavelengths. In this study, we used 11 RI models parameterised using NBO/T values between 0 and 1, with a step size of 0.1. The  $(n, k)$  data for the corresponding NBO/T values were calculated using a supporting spreadsheet from Prata et al. (2019),

260



which is available from the publisher’s website. Compared to the RE2018 dataset, RI models using NBO/T parameterisation show simplified wavelength dependences in both the real and imaginary parts of the RI due to the use of linear regression. However, this simple parameterisation is useful for VAC retrieval based on satellite remote sensing, given that an appropriate RI model can be specified in advance based on the assumed chemical composition of the ash aerosols. In that context, it is important to verify how well PG2019 RI models reproduce the BTS of infrared sounder measurements.




Figure 4: Same as Figure 3 but for PG2019 RI models parameterised using NBO/T.

### 3.3 Andesite and rhyolite RI models of Pollack et al. (1973): PLAND and PLRHY

The two RI models PLAND and PLRHY are the andesite and rhyolite (obsidian from Little Glass Mt. California) models of PL1973, respectively. These RI models were used for comparisons to the results of BTS simulations using RI models based on the RE2018/PG2019 datasets.




Figure 5: Same as Figure 3 but for the PLAND (red) and PLRHY (blue) models. These RI models are the andesite and rhyolite models of PL1973. The refractive index ( $n, k$ ) values were calculated using dispersion parameters and the equations in PL1973.

#### 4. IASI ash plume data


To evaluate the RI models by comparing measured and simulated BTS, continuous data from an infrared sounder that covers the wavenumber range of 650–1400  $\text{cm}^{-1}$  are desirable, and VAC data collected by IASI were used in this work. The IASI Level-1C granule data were acquired from the National Oceanic and Atmospheric Administration Comprehensive Large Array-Data Stewardship System (<https://www.avl.class.noaa.gov/saa/products/>). As a primary condition for BTS analysis, the brightness temperature difference between 10.7  $\mu\text{m}$  (934.5  $\text{cm}^{-1}$ ) and 12.2  $\mu\text{m}$  (819.5  $\text{cm}^{-1}$ ) ( $\Delta BT_{split} \equiv BT_{10.7} - BT_{12.2}$ ) was used, and pixels with  $\Delta BT_{split} < -2$  K for VACs over the ocean were selected. For discriminating the results of BTS simulations from different RI models using radiative transfer calculations, VAC pixels with more negative values of  $\Delta BT_{split}$  are better, as the BTS of such pixels shows distinct wavenumber dependence due to absorption in the infrared window region. The value  $-2$  K was chosen as the minimum requirement for discriminating the difference in BTS between the results of different RI models. We excluded VAC data over land due to large uncertainties in spectral surface emissivity and surface temperature.

Our radiative transfer model MBCRM conducts forward calculations for a single layer of VAC with no contamination from meteorological clouds (MCs) consisting of water or ice particles. Because MC contamination in VAC pixels can cause large estimation errors for VAC parameters and RI models (Kylling et al., 2015), contaminated pixels must be excluded to ensure the validity of the estimated RI model based on the results of our BTS analysis. In the case of VAC covered with dense MC layers, most measured pixels are rejected due to the condition  $\Delta BT_{split} < -2$  K. On the other hand, if a portion of the pixel

area is covered by MC or if the VAC is located above the MC layers, the MC can cause a negative bias in the measured BTS and still meet  $\Delta BT_{split} < -2 K$ . As a result, large errors may occur for the estimation of VAC parameters under the assumption of a single VAC layer. Furthermore, the detailed wavelength dependence of the BTS for such contaminated pixels may differ from that of the pure VAC. For these reasons, we used VAC data from daytime IASI measurements, and pixels that were presumed to be affected by MC were excluded through reference to visible true-colour imagery for the same area from the Moderate Resolution Imaging Spectroradiometer (MODIS) on board the Terra and Aqua satellites. MODIS images were provided by the Level-1 and Atmosphere Archive & Distribution System Distributed Active Archive Centre, and are available online (<https://atmosphere-imager.gsfc.nasa.gov/images/11b-granules>). The visible true-colour image product of the Japanese geostationary meteorological satellite Himawari-8 Advanced Himawari Imager (AHI) (Bessho et al., 2016) was also used for volcanic eruptions since 2015 in the coverage area of Himawari-8 (images were downloaded from the Japan Aerospace Exploration Agency Himawari Monitor homepage: <https://www.eorc.jaxa.jp/ptree/index.html>). Because we used daytime IASI measurements that could be referenced to MODIS images, the number of VAC pixels available for our analysis was reduced significantly. Furthermore, the evaluation of IASI pixels using MODIS visible images does not always exclude MC contamination due to the time difference between IASI and MODIS measurements, as well as the difference in sensor spatial resolution. Therefore, we added another condition for IASI pixels in our analysis. From a simple approximation of the thermal emissions from sea surface and clouds, the observed BT at a satellite is usually lower than that without MC when the MC locates between the ash plume and sea surface, when the ash plume is not optically thick. Using the brightness temperature at wavenumber  $\nu_a$ , where the measured brightness temperature is maximised in the region of  $750 \text{ cm}^{-1} \leq \nu \leq 900 \text{ cm}^{-1}$  due to a local minimum of the absorption index  $k$ , the difference  $\Delta BT_{clr}$  between the brightness temperatures of clear sky  $BT_{clr}(\nu_a)$  and ash plumes  $BT_{obs}(\nu_a)$  was defined as  $[\Delta BT_{clr} \equiv BT_{clr}(\nu_a) - BT_{obs}(\nu_a)]$ . The value of  $\nu_a$  was derived from VAC measurements and  $BT_{clr}(\nu_a)$  was calculated using our forward model. Then, a threshold of  $\Delta BT_{clr}$  was set according to the value of  $\Delta BT_{split}$ . Figure 6 shows a plot of  $BT_{clr}(\nu_a)$  with respect to  $\Delta BT_{split}$  in 1026 IASI pixels of VAC measurements for the volcanic eruptions listed in Table 2. The pixels are selected based on the primary condition  $\Delta BT_{split} < -2K$  over ocean areas, including night-time data to increase the number of data points. The clear-sky brightness temperature  $BT_{clr}(\nu_a)$  was calculated using atmospheric profiles, sea surface temperature, and atmospheric pressure at the surface from the results of the global assimilation analysis (GANAL) using a four-dimensional variational method and the atmospheric state from the Global Spectral Model (GSM) in the operational numerical weather prediction system of the Japan Meteorological Agency (<https://www.jma.go.jp/jma/jma-eng/jma-center/nwp/outline2019-nwp/index.htm>). In Figure 6, pixels are discriminated between Eyjafjallajökull (blue) and other volcanic eruptions (red), and the two datasets in the plot show similar distribution patterns. If a pixel of pure VAC observation is contaminated with MC, the negative value of  $\Delta BT_{split}$  is likely to become smaller, while the value of  $\Delta BT_{clr}$  is likely to increase. Therefore, the upper right data in Figure 6 may have higher probabilities of MC contamination than the lower left data on the plot. In our retrieval analysis, a threshold value of  $\Delta BT_{clr}$  (K) was defined to reduce the fraction of MC-contaminated pixels:

$$\Delta BT_{clr.th} \equiv -2 \cdot \Delta BT_{split} + 6 \quad (2)$$

and  $\Delta BT_{clr} < \Delta BT_{clr.th}$  was applied as an additional retrieval condition. The coefficients of Eq. (2) were determined empirically, and approximately 30% of all pixels with  $\Delta BT_{split} < -2K$  were rejected due to this condition for the Eyjafjallajökull VAC shown in Figure 6. The aim of this condition is to reduce the fraction of MC-contaminated pixels in our analysis; this threshold cannot be used to determine whether an individual pixel is contaminated with MC. Even for a pure VAC of a single homogeneous layer,  $\Delta BT_{clr}$  may be greater than  $\Delta BT_{clr.th}$  depending on the VAC parameters. In addition to MC contamination, the negative value of  $\Delta BT_{split}$  decreases and  $\Delta BT_{clr}$  increases for VACs with larger particle sizes and optical depths, as shown in the scatter plot of Figure 6, essentially the same as when the split-window plotting method is used (Prata, 1989; Wen and Rose, 1994; Prata and Grant, 2001) with the vertical and horizontal axes switched. As a result, the retrieval condition of Eq. (2) selects pixels of optically thin VACs composed of relatively small particles.



**Figure 6: Estimated brightness temperature difference  $\Delta BT_{clr}$  for VAC pixels of  $\Delta BT_{split}$  measured with IASI. In total, 1026 pixels of  $\Delta BT_{split} < -2K$  from the volcanic eruptions listed in Table 2 were used. Data for Eyjafjallajökull eruptions (6–16 May 2010) are coloured blue. The dashed line indicates the threshold of  $\Delta BT_{clr}$  from Eq. (2) used for retrieval analysis in this work.**

For retrieval analysis, we selected up to 10 VAC pixels with large negative values of  $\Delta BT_{split}$  from a granular dataset of IASI measurements. Radiative transfer calculations and BTS simulations for the selected VAC pixels were conducted using assumptions for three VAC parameters: the pressure height of the plume top  $P_{top}$ , optical depth  $\tau_c$  at wavelength  $11 \mu m$  ( $\nu = 907.488 \text{ cm}^{-1}$ ), and effective radius of the ash particles  $r_{eff}$ , which were retrieval variables, and using the temperature/water

vapour profiles and sea surface temperature/pressure from GANAL as fixed values. We set the pressure height of the plume  
 395 base  $P_{base}$  to  $P_{base} = P_{top} + 100 \text{ hPa}$  for simplicity. For the ozone profile, the total column of ozone was treated as a retrieval  
 variable, and the relative values for the initial ozone profile from GANAL were fixed. A homogeneous volcanic  $\text{SO}_2$  gas layer  
 with a pressure difference of  $100 \text{ hPa}$  between the top and base was assumed, with the pressure height of the layer top and  
 total column used as variables. The estimated values of the retrieval variables were those for which the root mean square  
 (RMS) between measured and calculated brightness temperatures ( $BT^{obs}$  and  $BT^{cal}$ ) in the assumed channels of number  $N$  is  
 400 minimised. The RI models are primarily evaluated by comparing the RMS values of various RI models.

$$\text{RMS} = \sqrt{\frac{1}{N} \sum_{i=1}^N (BT_i^{obs} - BT_i^{cal})^2} \quad (3)$$

Retrieval calculations were conducted using the scattering properties for ash particles derived from the selected RI model and  
 data for IASI channels of  $650 \text{ cm}^{-1} \leq \nu \leq 1400 \text{ cm}^{-1}$  with the following procedure:

- a) VAC parameters ( $P_{top}$ ,  $r_{eff}$ ,  $\tau_c$ ) are estimated using  $BT^{obs}$  and  $BT^{cal}$  for channels excluding the ozone band of  
 405  $980 \text{ cm}^{-1} \leq \nu \leq 1080 \text{ cm}^{-1}$  and  $\text{SO}_2$  bands of  $1100 \text{ cm}^{-1} \leq \nu \leq 1210 \text{ cm}^{-1}$  and  $1320 \text{ cm}^{-1} \leq \nu \leq 1395 \text{ cm}^{-1}$ . By  
 changing the value of the VAC parameters  $P_{top}$  from  $100 \text{ hPa}$  to  $P_S - 100 \text{ hPa}$  ( $P_S$  is the surface pressure) in 12 steps,  $r_{eff}$   
 from  $0.1 \mu\text{m}$  to  $5.0 \mu\text{m}$  in 10 steps, and  $\tau_c$  from  $0.1$  to  $5.0$  in 15 steps, we calculate the radiance of all parameter combinations  
 and search for a set of ( $P_{top}$ ,  $r_{eff}$ ,  $\tau_c$ ) that makes the minimum RMS. Then, the minimum RMS is searched again at a smaller  
 step size for each VAC parameter within the range of the previous  $\pm 1$  steps. The set of ( $P_{top}$ ,  $r_{eff}$ ,  $\tau_c$ ) for the minimum  
 410 RMS is the final result for the VAC parameters for the specific RI model.
- b) The total columns of ozone and  $\text{SO}_2$  columns/top pressure are estimated using the channels  $980 \text{ cm}^{-1} \leq \nu \leq 1080 \text{ cm}^{-1}$   
 for ozone and  $1320 \text{ cm}^{-1} \leq \nu \leq 1395 \text{ cm}^{-1}$  for  $\text{SO}_2$ , assuming the VAC parameters ( $P_{top}$ ,  $r_{eff}$ ,  $\tau_c$ ) as fixed values. The  
 wavenumbers  $1100 \text{ cm}^{-1} \leq \nu \leq 1210 \text{ cm}^{-1}$  are excluded from estimation of the  $\text{SO}_2$  because the brightness temperature of  
 these channels is strongly influenced by the applied ash RI model. For ozone, the column amount was set to a retrieval  
 415 parameter and the value was estimated using the Gauss–Newton method. For volcanic  $\text{SO}_2$ , the layer top pressure is changed  
 from  $100 \text{ hPa}$  to  $P_S - 100 \text{ hPa}$  in the same way as for ash top pressure, and the column amount of  $\text{SO}_2$  gas is estimated by the  
 Gauss–Newton method for each  $\text{SO}_2$  layer. The final results of the  $\text{SO}_2$  parameters are the top height and column amount that  
 give the minimum RMS in wavenumbers  $1320 \text{ cm}^{-1} \leq \nu \leq 1395 \text{ cm}^{-1}$ .
- c) The RMS for the channels  $750 \text{ cm}^{-1} \leq \nu \leq 1400 \text{ cm}^{-1}$  is then calculated, considering the estimated VAC, ozone, and  $\text{SO}_2$   
 420 parameters as fixed. Although the range of  $650 \text{ cm}^{-1} \leq \nu < 750 \text{ cm}^{-1}$  is important for estimating the VAC parameters,  
 especially the top height of the ash plumes, we exclude this range to avoid RMS values related to error in the applied  
 atmospheric profiles and to clarify the difference between the RI models in the atmospheric window region.
- d) Processes a) to c) are conducted for all RI models.

The date and number of pixels in the selected IASI data for eruptions from seven volcanoes are listed in Table 2. The average  
 425 value of the wavenumber  $\bar{\nu}_a$  at the local maximum of brightness temperature is also listed. A TAS diagram based on the data

of tephra samples of volcanos from the literature is shown in Figure 7. In particular,  $\nu_a$  is related to the minimum of the absorption index  $k$  and has been reported as an important parameter for identifying ash aerosols from infrared sounder measurements (Gangale et al., 2010; Clarisse et al., 2013; Clarisse and Prata, 2016). Using a total of 21 RI models, which are listed in Table 1, the mean RMS of brightness temperature in the wavenumber range  $750 \text{ cm}^{-1} \leq \nu \leq 1400 \text{ cm}^{-1}$  was determined for each RI model. In the following sections, we discuss the results for the RI models determined from RMS values for the volcanic events in Table 2. The detailed results for RMS and VAC parameters ( $P_{top}$ ,  $r_{eff}$ ,  $\tau_c$ ), ash top height  $h_{top}$  estimated from  $P_{top}$  and the GANAL profile, and total column and top height of  $\text{SO}_2$  for each IASI pixel are provided in the Supplemental Material section. The mineral composition of ash varies across different eruptions and within the same eruption at different times (Prata and Lynch, 2019). In this study, however, we assumed that the ash aerosols ejected from a given volcano remains the same throughout the period of each eruption event in Table 2 and that the RI model is evaluated using the average of the RMS over all pixels for a given volcano. Even if its mean RMS was the smallest, we rejected an RI model as unrealistic if the converged values of  $r_{eff}$  reached the minimum size  $0.1 \mu\text{m}$  for many pixels in the retrieval analyses. In addition to the differences among RI models, the size of ash particles effectively changes the wavelength dependence of the simulated BTS (Clarisse and Prata, 2016; Ishimoto et al., 2016), and small values of  $r_{eff}$  tend to be estimated when a relatively mafic RI model is used with low  $\text{SiO}_2$  wt.% or large NBO/T values. The relationship between the estimated  $r_{eff}$  value and the RI model in the retrieval analysis is discussed in section 5.1.

In our radiative transfer calculations for the BTS simulations, we assumed that the VAC is distributed homogeneously within the area of the IASI pixel footprint ( $\sim 12 \text{ km}$  in the nadir direction). Note that the estimated VAC parameters might have large errors if the VAC is only partially distributed within the pixel area (VAC fraction  $< 1$ ). Of the three VAC parameters, ash top height was relatively sensitive to the assumed ash fraction and the homogeneous layer assumption tended to derive a lower height of  $P_{top}$ . Furthermore, the BTS simulations in this work include many potential error sources in addition to MC contamination, such as the error due to the monochromatic approximation of Eq. (1) in radiative transfer calculations for ash plumes, interpolation error of ash optical properties from the data table in MBCRM, applied atmospheric profiles and sea surface temperature/emissivity, and assumption of a homogeneous plane-parallel ash layer. Therefore, the results of the following analysis and discussion may change if different assumptions and numerical methods were applied for the ash plume analysis.

**Table 2: Volcanic eruption events used for BTS analysis of IASI measurements and examples of the RI models.** The RI models were selected through a comparison between measured and simulated BTS for IASI pixels on dates listed in the column ‘Date’. Plural RI models in the column ‘RI model’ indicate that the results of BTS fitting were almost the same. The average wavenumber  $\bar{\nu}_a$  is that at which the measured brightness temperature is maximised in the range of  $750 \text{ cm}^{-1} \leq \nu \leq 900 \text{ cm}^{-1}$ . For Calbuco eruptions, RI models for different ash plume locations, Calbuco-A and -B, are listed separately.

460

465

470


475

Name of volcano	Location of Volcano (Lat., Lon.)		Date (yyyymmdd)	IASI pixels	$\bar{\nu}_a(\text{cm}^{-1})$	RI model
Calbuco	-41.33	287.382	20150424	15 (A)	831.6	PG010 (RE040)
				10 (B)	825.7	PG060, RE030
Eyjafjallajökull	63.633	340.367	20100506-12	50	819.3	PG040, RE030
Grimsvötn	64.416	342.684	20110522-23	11	817.2	PG070, PG080, RE050, PLAND
Kelud	-7.93	112.308	20140214	4	860.4	PG000
Kirishimayama	31.934	130.862	20110127	9	830.4	RE080, RE020, PG010
Nishinoshima	27.247	140.874	20200731	18	827.3	RE070, PG070
Puyehue-Cordon	-40.59	287.883	20110606	39	859.9	PG000

480

485

490





495 **Figure 7: Total alkali–silica (TAS) diagram based on ash and pumice samples of the volcanos featured in Table 2. SiO<sub>2</sub> and Na<sub>2</sub>O+K<sub>2</sub>O wt.% values for each volcano were obtained from the literature as follows: Calbuco, Deguine et al. (2020); Eyjafjallajökull and Grímsvötn, Reed et al. (2018) and Prata et al. (2019); Kelud, Maeno et al. (2019); Kirishimayama, <https://www.gsj.jp/hazards/volcano/kirishima/2011/works-index.html>; Nishinoshima, [https://www.data.jma.go.jp/svd/vois/data/tokyo/STOCK/kaisetsu/CCPVE/shiryo/147/147\\_2-2.pdf](https://www.data.jma.go.jp/svd/vois/data/tokyo/STOCK/kaisetsu/CCPVE/shiryo/147/147_2-2.pdf); Puyehue-Cordón**  
500 **Caulle, Castro et al. (2013).**

**5. Refractive index models estimated through IASI measurement simulation**


Selected IASI measurements for the eruption events of seven volcanoes and **examples of the RI models** are listed in Table 2.  
505 Figure 8 shows ranking plots for the results of RMS analysis for each measurement pixel and Figure 9 shows the mean and standard deviation for each volcano. Ranking based on the smallest RMS was conducted for the 21 RI models. Detailed results for each RI model in each pixel, including the minimum RMS, estimated VAC parameters ( $P_{top}, h_{top}, r_{eff}, \tau_c$ ), and SO<sub>2</sub> **top height and** column amount, are provided in the Supplemental Material. From the scattered distribution of the ranking in Figure 8 and from the mean and standard deviation of the RMS in Figure 9, the basic features of BTS for the ash plumes measured  
510 by IASI can be reproduced **by several** RI models, although we chose **typical** RI models **1–4** in Table 2. Moreover, statistical discussions are difficult for some eruptions, such as Kelud and Kirishimayama, because of the small number of pixels available for the analysis. Therefore, **the selection of the RI models** used here may change depending on the measurement dataset and the **numerical conditions in the BTS simulations.**

The RI models from the PG2019 dataset were selected **for all** volcanic events. Furthermore, high-ranking RI models tended to  
515 converge at a specific value of NBO/T in the PG2019 dataset. This result suggests that NBO/T, which can be determined from the chemical composition of the ash, is an effective indicator of the appropriate RI model for satellite VAC analysis. On the other hand, the andesite model of PL1973 (PLAND) was not selected as the **high-ranking RI models**, except for the Grímsvötn eruptions. **This result** shows that the novel RI models of RE2018 and PG2019 are useful for simulating the measured BTS for various types of volcanic plume.

520 The results of VAC analysis for individual volcanoes are discussed in the following subsections. Typical results of the BTS simulations are shown in the figures, and the estimated VAC parameters, SO<sub>2</sub> column **and top height**, and RI model used for each simulation are summarised in Table 3.

525





565

570

575


580

585

590

595

Figure 8: Ranking plots showing the results of BTS simulations for the pixels used for the IASI ash plume measurements in Table 2. The ranking of the 21 RI models is in ascending order of the RMS values between the measured and simulated BTS. Results for different pixels are shown on different lines, which are ordered by the date and time of IASI granule measurements from top to bottom. The 11 RI models using the PG2019 dataset are arranged based on NBO/T values (0.0 to 1.0 in steps of 0.1 from right to left), while the 8 RI models using the RE2018 dataset and two RI models from PL1973 (PLAND/PLRHY) are arranged based on SiO<sub>2</sub> wt.% (Table 1). The RI model for the volcano (Table 2) was determined from the average of RMS values for the corresponding measurement pixels. A circle indicates that the BTS simulated using the RI model provides a good fit to the measured BTS, and other RI models are indicated by open triangles (△). The RI model with a solid triangle (▲) resulted in the smallest average of RMS values, but was rejected because many estimated  $r_{eff}$  values reach the assumed minimum size 0.1  $\mu\text{m}$ .



600

605 **Figure 9: Results of the RMS (filled circle) for each RI model averaged over the pixels of the eruptions in Table 2. The error bar indicates the standard deviation. The RI models are arranged in order from smallest to largest RMS from left to right and the corresponding RI models are listed on the right.**

610 **Table 3: Locations of IASI pixels, applied RI models, retrieved VAC parameters ( $h_{top}, r_{eff}, \tau_c$ ), and SO2 columns (in Dobson units) and top height (km) for BTS simulations of typical VAC (‘Figure No.’). BTS data for simulations and measurements are provided in the Supplemental Material.**

615

620

625


Figure No.	Volcano	Pixel location (lat., lon.) in deg.	RI model	Top height (km)	Effective radius ( $\mu\text{m}$ )	Optical thickness @11 $\mu\text{m}$	SO2	
							column (DU)	top (km)
10c	Eyjafjallajökull	(63.146N, 15.261W)	RE030	5.97	0.84	2.24	34.4	5.98
			PG040	6.23	1.18	2.08	54.4	6.23
10f	Eyjafjallajökull	(52.549N, 24.075W)	RE030	3.67	0.84	1.34	0.0	—
			PG040	4.30	1.30	0.96	0.0	—
12c	Grimsvötn	(61.390N, 21.484W)	RE050	1.67	0.76	1.15	0.0	—
			PG080	1.82	1.24	1.07	0.0	—
13c	Calbuco_A	(35.265S, 55.656W)	PG010	12.99	0.57	0.14	45.3	13.00
13f	Calbuco_B	(32.592S, 75.050W)	RE030	4.25	0.66	0.88	1.4	5.42
			PG060	4.47	0.52	0.80	1.3	5.59
14c	Kirishimayama	(31.079N, 136.500E)	PG010	2.23	1.57	1.34	0.0	—
			RE080	3.40	2.20	1.27	0.0	—
14f	Nishinoshima	(26.719N, 141.167E)	PG070	3.89	0.29	0.84	0.0	—
			RE070	3.78	0.39	0.88	0.0	—
15d	Kelut	(9.101S, 110.513E)	PG000	11.02	1.18	0.61	44.0	11.25
			PG010	11.97	0.93	0.49	46.2	12.09
16d	Puyehue-Cordon	(27.136S, 25.178W)	PG000	7.29	0.26	0.33	4.4	7.53
16g		(41.212S, 61.898W)	PG000	10.23	0.24	0.33	7.9	10.23

**5.1 Eyjafjallajökull**

630 In the case of the Eyjafjallajökull eruptions in 2010, RI models developed from the results of laboratory analysis of ash samples were provided by RE2018 and data of IASI measurements were obtained. Simulations of the measured BTS were conducted using 50 BTS datasets for IASI pixels that satisfied our retrieval conditions over the period of 6–12 May 2010. Overall, the RI

models of  $\text{NBO/T} = 0.3\text{--}0.6$  (PG030–PG060) for the PG2019 dataset and the RI model Eyjafjallajökull-(a) (RE030) using the RE2018 dataset showed good BTS fitting results (Fig. 9). Specifically, RE030 and PG040 gave similar values for the smallest of average RMS values over the 50 total pixels. Typical results of BTS analysis for the Eyjafjallajökull VAC are shown in Figure 10. For these IASI measurements, true-colour images from the MODIS product (Fig. 10a and d) taken near the IASI measurement time were obtained. Through comparison with the colour image shown in Figure 10a, the influence of MCs appears small in the vicinity of IASI pixels with highly negative  $\Delta TB_{split}$  values in Figure 10b. The results of BTS simulation with RE030 and PG040 were similar and in good agreement with the measured BTS, although some differences occurred for the estimated VAC parameters  $(h_{top}, r_{eff}, \tau_c)$  (Table 3). Because PG040 is parameterised by  $\text{NBO/T} = 0.4$  and this  $\text{NBO/T}$  value is close to that of the Eyjafjallajökull-(a) ash sample (0.38), the wavenumbers of the local minimum and local maximum of the absorption index  $k$  obtained using PG040 were similar to those of RE030 (Figs. 3 and 4). Thus, the results of the BTS simulations shown in Figure 10c arose from the similarity of the spectral pattern of  $k$  between PG040 and RE030. Another Eyjafjallajökull VAC was located more than 1000 km from the volcano on 9 May 2010 (Fig. 10d–f). In IASI measurements, pixels with highly negative values of  $\Delta TB_{split}$  are present in the area of  $51^\circ \text{N} \leq lat \leq 54^\circ \text{N}$  and  $20^\circ \text{W} \leq lon \leq 28^\circ \text{W}$  (circle in Fig. 10e); a faint VAC with sparse broken clouds was inferred in this area from the MODIS colour image, and an ash optical thickness of 0.15–0.20 at visible wavelengths was estimated by Ventress et al. (2016). Figure 10f shows the results of BTS simulation for one of the nine selected pixels in this area. As shown in Figure 10c, the BTS simulated using PG040 and RE030 agree well with the measured values, and the effect of MC contamination on the measured BTS appears minor. The average VAC parameter values over the nine pixels were  $(\bar{h}_{top}, \bar{r}_{eff}, \bar{\tau}_c) = (3.48 \text{ km}, 0.83 \mu\text{m}, 1.38)$  for PG040 and  $(\bar{h}_{top}, \bar{r}_{eff}, \bar{\tau}_c) = (3.31 \text{ km}, 0.70 \mu\text{m}, 1.57)$  for RE030. The values of  $h_{top}$  and  $r_{eff}$  are consistent with the mode values of the estimated  $h_{top}$  and  $r_{eff}$  histograms based on IASI data from the same day (Ventress et al., 2016). In our analysis of 50 total pixels showing the Eyjafjallajökull VAC from 6–12 May, the average value of the effective radius  $\bar{r}_{eff}$  was  $1.08 \mu\text{m}$  ( $0.95 \mu\text{m}$ ) with the PG040 (RE030) RI model, which agrees with the results from aircraft observations stating that  $r_{eff}$  was in the range of  $0.87\text{--}1.19 \mu\text{m}$  (Turnbull et al., 2012; Ventress et al., 2016).






**Figure 10: Examples of BTS analyses of the Eyjafjallajökull VAC on 6 May 2010 (a–c). (a) MODIS true-colour image at 11:55 UTC and (b) brightness temperature difference  $\Delta BT_{split}$  measured with IASI at 11:53 UTC. (c) BTS of IASI measurement at  $(lat, lon) = (63.146^\circ N, 15.261^\circ W)$  (red), and results of BTS simulations using the RI models RE030 (green) and PG040 (blue). (d–f) Same as (a–c), but for the VAC on 9 May 2010. The MODIS colour image is a composite from 12:25–12:30 UTC (d) and IASI  $\Delta BT_{split}$  is calculated at 12:32 UTC (e). The area of highly negative  $\Delta BT_{split}$  is indicated with circles in (d) and (e). (f) Measured and simulated BTS for the IASI pixel at  $(lat, lon) = (52.549^\circ N, 24.075^\circ W)$ . The retrieved VAC parameters and **SO2** for each BTS simulation are listed in Table 3.**

Using the retrieval results for 50 pixels of Eyjafjallajökull VAC measurements under various RI models (PG070, PG040, and PG010), the dependence of the RI model on the estimated VAC parameters  $(h_{top}, r_{eff}, \tau_c)$  was investigated (Fig. 11a–c). In particular, the results for  $r_{eff}$  showed a strong dependence on RI model selection (Fig. 11b). Within the PG2019 dataset, small

700 values of  $r_{eff}$  tend to be estimated for mafic RI models with large NBO/T. By contrast, relatively large  $r_{eff}$  values were  
 derived for felsic RI models. Similar dependence of RI models on estimated  $r_{eff}$  was also confirmed for our VAC retrievals  
 related to other volcanic events listed in Table 2. Using a pixel with small differences in retrieved  $h_{top}$  and  $\tau_c$  among the three  
 RI models (pixel no. 12), the results of BTS simulations with different  $r_{eff}$  values and with different RI models are plotted in  
 Figure 11d and e, respectively. If we assume that the RI model of PG040 is adequate for Eyjafjallajökull ash aerosols and that  
 705 the retrieval result  $r_{eff} \sim 1.0 \mu\text{m}$  using PG040 is close to the true value, a positive bias of the simulated BTS in the region of  
 $900 \text{ cm}^{-1} \leq \nu \leq 1230 \text{ cm}^{-1}$  occurs, except within the ozone band, when we use a more mafic RI model (PG060 and PG080)  
 than PG040 with the same  $r_{eff}$  value (Fig. 11e). The positive bias of the BTS is mitigated by applying a smaller value of  $r_{eff}$   
 and keeping the other VAC parameters constant in the BTS simulation, as shown in Fig. 11d. By contrast, a negative bias of  
 the BTS occurs in the same wavenumber range when we use a more felsic RI model (PG000 and PG020 in Fig. 11e), and  
 710 fitting of the calculated BTS improves when a larger  $r_{eff}$  value is used for the ash particles as in Fig. 11d. Thus, the retrieval  
 results of  $r_{eff}$  are sensitive to the wavenumber dependence of the particle absorption property around  $900 \text{ cm}^{-1} \leq \nu \leq$   
 $1230 \text{ cm}^{-1}$ . As such, the selection of the proper RI model is important for estimating  $r_{eff}$ . In our BTS analysis of the  
 Eyjafjallajökull VAC, PG040 and RE030 were **selected as** their fitting results were better than those of other RI models in  
 wavenumber regions other than  $900 \text{ cm}^{-1} \leq \nu \leq 1230 \text{ cm}^{-1}$  (in particular,  $750 \text{ cm}^{-1} \leq \nu \leq 900 \text{ cm}^{-1}$ ).

715

720



730





Figure 11: VAC parameters plume top height  $h_{top}$  (a), particle effective radius  $r_{eff}$  (b), and optical thickness  $\tau_c$  (c) estimated from our BTS analysis of 50 IASI pixels for Eyjafjallajökull VAC measurement. Observation dates corresponding to the serial number of the pixel are shown in the lower part of (a). (d) and (e) shows examples of measured and calculated BTS using IASI data for pixel no. 12 [7 May 11:32 UTC, ( $lat, lon$ ) = (61.669° N, 15.370° W)]. The green line in (d) and (e) indicates the BTS calculated using the PG040 RI model and the VAC parameters  $(h_{top}, r_{eff}, \tau_c) = (5.33 \text{ km}, 1.0 \text{ } \mu\text{m}, 1.50)$ , while the BTS calculated after changing the  $r_{eff}$  value are plotted in (d). (e) is the same as (d) but using PG000–PG080 RI models and the VAC parameters  $(h_{top}, r_{eff}, \tau_c) = (5.33 \text{ km}, 1.0 \text{ } \mu\text{m}, 1.50)$ .

## 5.2 Grímsvötn

For the Grímsvötn eruptions in May 2011, both the RI model from the laboratory experiments of RE2018 and IASI measurements of ash plumes were obtained. From IASI data for the Grímsvötn VAC, one pixel at 12:08 UTC on 22 May and 10 pixels at 11:47 UTC on 23 May were selected for our BTS analysis. The measured IASI pixels of  $\Delta BT_{split} < -2\text{K}$  on 23 May (Fig. 12b) were estimated to have low MC contamination based on the MODIS colour image taken at 12:05 UTC (Fig. 12a). Moreover, from this MODIS colour image, MCs in the area of the VAC were determined to be located at higher altitudes than the VAC (Prata et al., 2017b), suggesting that the MC-contaminated pixels were effectively excluded from the target

pixels in our BTS analysis. For VACs from Grímsvötn eruptions, the BTS based on IASI measurements was reported to produce good simulation results using the basalt RI model of PL1973 (Newman et al., 2012). The results of our BTS simulations showed that a small RMS was obtained for the PG070 and PG080 RI models using the PG2019 dataset, and RI models derived from data on Grímsvötn (RE050) and Spurr (RE070) ash samples listed in the RE2018 dataset, and the andesite model (PLAND) of PL1973, returned relatively small RMS values. Absorption features related to SO<sub>2</sub> gas were not present in the BTS of 11 analysed pixels, in accordance with the report of SO<sub>2</sub> gas separation from ash plumes (Prata et al., 2017b); thus, PG070-PG080, RE050, and PLAND were selected as the typical RI models based on their mean RMS scores. The estimate of NBO/T for the Grímsvötn ash sample was 0.74–0.75, which is the highest value within the RE2018 dataset (Prata et al., 2019). Because the PG2019 dataset was derived from NBO/T parameterisation using the RE2018 dataset, the wavelength dependence of PG070-PG080 is similar to that of RE050. As a result, similar VAC parameters were estimated for PG070-PG080 and RE050, and the average values over 11 pixels, i.e.,  $(\bar{h}_{top}, \bar{r}_{eff}, \bar{\tau}_c) = (1.59 \text{ km}, 1.47 \text{ }\mu\text{m}, 2.04)$   $[(\bar{h}_{top}, \bar{r}_{eff}, \bar{\tau}_c) = (1.56 \text{ km}, 1.44 \text{ }\mu\text{m}, 2.10)]$  for PG070 (PG080) and  $(\bar{h}_{top}, \bar{r}_{eff}, \bar{\tau}_c) = (1.60 \text{ km}, 0.93 \text{ }\mu\text{m}, 2.01)$  for RE050. This result for  $r_{eff}$  generally agrees with published results for a Grímsvötn ash sample (1.1  $\mu\text{m}$ ) (Reed et al., 2018). For plume top height, Moxnes et al. (2014) estimated  $h_{top} \leq 4 \text{ km}$  through the analysis of IASI data, and similar results for plume height were obtained using the Inversion Technique for Emission Modelling (Harvey et al., 2020). According to the measurements using Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) on board the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO), the top heights of the Grímsvötn VAC at lon. 60 – 62°N and lat. ~20°W on the same day were around 3 km, and the backscatter peak heights were less than 2 km (Taylor et al., 2019). Although our estimated values of  $h_{top}$  are smaller, they are generally consistent with previous research.

An example of the results from BTS simulations is shown in Figure 12c. Due to the small SO<sub>2</sub> contribution, the BTS in the range of  $1100 \text{ cm}^{-1} \leq \nu \leq 1210 \text{ cm}^{-1}$  is mainly driven by the optical properties of ash particles, and the difference in the BTS of PG080 and RE050 was small in this wavenumber range.





800 **Figure 12: Grímsvötn VAC on 23 May 2011, shown in a MODIS true-colour image taken at 12:05 UTC. (a),  $\Delta TB_{split}$  from IASI**  
measurements at 11:47 UTC (b), and the BTS of the IASI pixel at ( $lat, lon$ ) = (61.390° N, 21.484° W) (red) (c). The results of  
BTS simulations using RE050 (green) and PG080 (blue) are also shown in (c).

805

The selected RI models for simulating the observed BTS of the Eyjafjallajökull and Grímsvötn VACs had similar NBO/T values to those of ash samples from the same volcanic eruptions. This result indicates that the BTS of VACs obtained from satellite infrared sounder measurements is related to the spectrum of the ash RI and VAC parameters. Furthermore, it supports the proposals of Reed et al. (2018) and Prata et al. (2019) that an appropriate RI model for VAC retrieval from satellites can  
810 be determined using the NBO/T (or SiO<sub>2</sub> content) of ash samples. In addition, our results suggest that infrared sounder measurements for VAC may help to choose ash RI models under certain atmospheric and numerical conditions to perform exact radiative transfer calculations, provided that enough RI models have been prepared in advance.

815

### 5.3 Calbuco


In the series of Calbuco volcanic activity from 22 April 2015, IASI data for brightness temperature at 13:29–13:32 UTC on 24 April were used for our retrieval analysis. At this time, ash plumes with negative  $\Delta TB_{split}$  were observed over the Atlantic Ocean near the estuary of the La Plata River at latitudes of 35–38° S and longitudes of 55–56° W (Calbuco\_A: Fig. 13b), as  
820 well as over the Pacific Ocean near the west coast of Chile at 32–33° S and 73–76° W (Calbuco\_B: Fig. 13e). From the MODIS colour image taken at 13:35 UTC (Fig. 13a), it appeared that the effect of MCs on the BTS was small for pixels with large negative values of  $\Delta TB_{split}$  in Calbuco\_A. Although another VAC was confirmed around latitude 40° S in the MODIS image shown in Figure 13a, the IASI pixels had slightly negative or positive  $\Delta TB_{split}$  values, suggesting ice particles within or above the VAC layer. BTS analyses were conducted for 15 pixels with large negative  $\Delta TB_{split}$  values in Calbuco\_A. Small total  
825 RMS values were obtained from the PG010–PG030 RI models with the PG2019 dataset and RE040 with the RE2018 dataset, and the smallest RMS was obtained from RE040 (Figs. 8 and 9). The estimated  $r_{eff}$  values were small overall, and were less than 0.2  $\mu\text{m}$  for most pixels when relatively mafic RI models (PG030–PG100 of PG2019 dataset) were applied. For the RE040 RI model, low  $r_{eff}$  values of less than 0.2  $\mu\text{m}$  were obtained for most of the analysed pixels, and the smallest effective radius  $r_{eff} = 0.1 \mu\text{m}$  in our LUT was estimated for 8 of the 15 total pixels within Calbuco\_A. These small  $r_{eff}$  values are mainly  
830 related to the fitting of the BTS in the range of  $1070 \text{ cm}^{-1} \leq \nu \leq 1230 \text{ cm}^{-1}$ , as discussed in section 5.1 (Fig. 11). The area

of pixels with large negative  $\Delta BT_{split}$  in Figure 13b is ~1500 km from Calbuco volcano, and the particle size of the VAC is expected to be reduced due to depositional segregation during transportation in the atmosphere. Although a  $r_{eff} < 0.2 \mu\text{m}$  may not be too small for the ash particles present in this area, we considered the PG010 RI model, which had the second smallest mean RMS after RE040 and averaged  $r_{eff} \geq 0.2 \mu\text{m}$ , to be the better RI model in this study. As shown in Figure 13c, the absorption feature of  $\text{SO}_2$  gas is clearly apparent in the measured BTS, and the results of BTS simulations approximately agree with the measurements. However, the RMS values between measurements and simulations were relatively large for all pixels and systematic deviation from the measurements occurred even with PG010. For the retrieval calculations for the IASI pixels near Calbuco\_A, similar RMS values were obtained at different VAC top heights,  $h_{top} = 5 - 7 \text{ km}$  and  $h_{top} > 10 \text{ km}$ . According to the CALIOP measurements at 17:31–17:45 UTC of the same day, the height of the VAC near the Calbuco\_A region was 15–18 km. Therefore, we considered the retrieved VAC parameters when the estimated top height  $h_{top} > 10 \text{ km}$  (Table 3) to be closer to the true values.

MODIS observed the Calbuco\_B region at 15:00 UTC (Fig. 13d). We inferred the presence of an optically thin VAC in an area with little MCs, although the MODIS observation was made 1.5 hours after that of IASI. As  $\text{SO}_2$  absorption features were weak in the BTS from IASI measurements, we deduced that the measured BTS patterns arise mainly from the optical properties of the VAC. In BTS simulations for 10 pixels in Calbuco\_B, high-ranking RI models tended to have greater NBO/T in the PG2019 dataset compared with the results of Calbuco\_A (Fig. 8); results from the RI models PG040–PG070 and RE030 had small RMS values. The reason for this difference between the results of Calbuco\_A and Calbuco\_B has not been determined. According to the compositional analyses of (Romero et al., 2016) and (Deguine et al., 2020), Calbuco ash samples are of the basaltic andesite type with a  $\text{SiO}_2$  wt.% of 55–56.3, which is a similar or more mafic composition than Eyjafjallajökull ash samples. Our results for Calbuco\_B pixels are relatively consistent with the compositional analysis results; however, the PG040-PG070 RI had large RMS errors for Calbuco\_A pixels in our retrieval analysis.

We concluded that the measured BTS of the Calbuco\_A VAC were due to small, relatively felsic ash particles. Although we examined the contamination of ice clouds and internal mixing of sulphate and mafic ash particles using simple optical models in the preliminary calculations and the results were not successful; we cannot exclude the possibility that other factors explain the BTS of Calbuco\_A.

860




**Figure 13: Brightness temperature difference  $\Delta BT_{split}$  of Calbuco ash plumes measured by IASI at 13:29 UTC on 24 Apr 2015 over the Atlantic Ocean [Calbuco\_A, (b)], and the measured (red) and calculated (PG010, blue) BTS for the pixel at  $(lat, lon) = (35.265^\circ S, 55.656^\circ W)$  (c). MODIS visible-colour image of the same region at 13:25 UTC (a). (d–f) Same as (a–c) but for ash plumes over the Pacific Ocean [Calbuco\_B, marked with a circle in (e)] and the measured BTS at  $(lat, lon) = (32.592^\circ S, 75.050^\circ W)$ . Calculated BTS without SO<sub>2</sub> gas from the RE030 (green line) and PG060 (blue line) RI models are also shown. The measurement time of the MODIS colour image (d) was 15:00 UTC.**

## 5.4 Kirishimayama and Nishinoshima

During the Kirishimayama eruption on 26 January 2011, ash plumes were transported in the southeast direction over the Pacific Ocean, and IASI measured the VAC to the south of the Japanese main islands at 00:08 UTC on 27 January (Fig. 14b). The MODIS image taken at 01:05 UTC (Fig. 14a) suggests that the VAC in the region of highly negative  $\Delta BT_{split}$  is distributed above the streaks of MCs, and VACs with lower MC contamination are expected in locations between the MC streaks. Due to the 1-hour time difference between IASI and MODIS measurements, the pixels for analysis were determined using only IASI data. Among 30 pixels of  $\Delta BT_{split} < -2\text{K}$ , 21 pixels were rejected under the condition of Eq. (2), suggesting that many IASI pixels were contaminated with MCs. For the BTS of the remaining nine pixels, the RI models RE080, RE020, and PG010-PG020 provided small RMS values. The results of BTS simulations for RE080 and PG010 generally agree with measured values (Fig. 14c). However, some small differences between the measured and calculated BTS were confirmed, such as a negative bias in the RE080 simulation around  $920\text{ cm}^{-1}$ . From the plot ranking shown in Figure 8 for the PG2019 dataset, the Kirishimayama VAC at this time tended to fit the results of RI models with smaller NBO/T values than the Eyjafjallajökull and Grímsvötn ash plumes. This result is reasonable considering the reported  $\text{SiO}_2$  contents of the ash samples and the average local maximum of  $\bar{\nu}_a \approx 830\text{ cm}^{-1}$ , which is a greater wavenumber than the local maxima for Eyjafjallajökull and Grímsvötn (Table 2). However, the possibility of MC contamination of the BTS for all VAC pixels in this area cannot be excluded. Further precise analysis of the Kirishimayama VAC, including night-time measurements with no MC contamination, is necessary. An algorithm for detecting MC contamination in the VAC using infrared channels will be essential to solving this problem.

Compared to Kirishimayama, the VACs of Nishinoshima, which were observed by IASI at 00:08 (Fig. 14e) and 23:47 UTC on 31 July 2020, were easily identified by the AHI data on board the geostationary satellite Himawari-8 as having less MC contamination (Fig. 14d). In BTS simulations for 18 pixels with large negative values of  $\Delta BT_{split}$ , the RI models RE070 with the RE2018 dataset and PG050-PG080 with the PG2019 dataset returned small RMS values and small effective radius estimates ( $r_{eff} \leq 0.6\text{ }\mu\text{m}$ ) for all pixels. As shown in Figure 14f, the calculated BTS fit well to the measured BTS, especially at wavenumbers  $\nu \geq 1050\text{ cm}^{-1}$ ; however, systematic errors in the range of  $750\text{ cm}^{-1} \leq \nu \leq 1000\text{ cm}^{-1}$  were also observed. From the results of chemical analysis of volcanic ash samples from eruptions on 14 and 20 July 2020, it has been estimated that the bulk composition of Nishinoshima volcanic product has apparently changed from andesite (59–60  $\text{SiO}_2$  wt.%) to basaltic andesite (54–55  $\text{SiO}_2$  wt.%) during the 2019–2020 eruptions. Our result of relatively mafic RI models for the Nishinoshima VAC indicates that the BTS analysis might detect the compositional variation in the VAC from satellite measurements.



**Figure 14:** (a-c) Composite MODIS image of the Kirishimayama VAC at 01:05 UTC on 27 January 2011(a), and  $\Delta BT_{split}$  from IASI data at 00:08 UTC on the same day (b). Results of BTS simulations for the pixel at  $(lat, lon) = (31.079^\circ \text{ N}, 136.500^\circ \text{ E})$  are shown in (c), as well as the measured values. The RI models **PG010** (green) and **RE080** (blue) were used for the BTS simulations. For the Nishinoshima VAC on 31 July 2020, the visible true-colour image (d) is from the Himawari-8/AHI observation at 0:00–00:09 UTC, and  $\Delta TB_{split}$  is from IASI at 00:05–00:08 UTC (e). (f) Measured and calculated BTS for the pixel at  $(lat, lon) = (26.719^\circ \text{ N}, 141.167^\circ \text{ E})$ . The RI models **PG070** (green) and **RE070** (blue) were used for simulations.



For the VAC from the Kelud eruption event on 13 February 2014, Ishimoto et al. (2016) analysed data obtained by the Atmospheric Infrared Sounder (AIRS) on board the Earth Observing System Aqua polar-orbiting satellite and estimated that the spectral RI of the Kelud VAC was intermediate between the andesite model and the rhyolite (obsidian) model of PL1973. In their analysis, a mixture of these two RI models was assumed as the a priori RI and the spectral RI in the wavenumber range of  $\nu \geq 1100 \text{ cm}^{-1}$  was omitted due to missing spectral data between 1137 and 1216  $\text{cm}^{-1}$  from the AIRS measurements. Furthermore, MC contamination was not considered for pixels with highly negative  $\Delta BT_{split}$  values, although MC contamination may cause large errors in RI estimation. For the Kelud VAC at this time, Kylling (2016) reported that a mixture of ice clouds and ash plumes in an IASI pixel can explain the observed BTS from the PL1973 andesite model. In this work, retrieval analyses of the Kelud VAC were repeated using IASI pixels with less MC contamination, as estimated from visible images and from the condition of Eq. (2).

In the IASI data of the Kelud VAC observed at 02:08 UTC on 14 February (Fig. 15c), all pixels with  $\Delta BT_{split} < -2\text{K}$  in the region of  $8.0^\circ \text{ S} \leq lat \leq 9.5^\circ \text{ S}$  and  $106.5^\circ \text{ E} \leq lon \leq 109.5^\circ \text{ E}$  were rejected under the condition of Eq. (2), and four pixels (indicated by the circle in Fig. 15c) were retained for our retrieval analysis. This result suggests that most VAC measurements are affected by MC contamination, which is consistent with the conclusion of Kylling (2016) that IASI measures both ash and ice clouds in the same pixel. The area of the four retained pixels is presumed to be covered with thin ash plumes and have less MCs based on the MODIS true-colour image taken at 03:35 UTC (Fig. 15a) and the Himawari-7 visible monochromatic image taken at 02:00 UTC (Fig. 15b). The results of BTS simulations showed that the BTS of the Kelud VAC could be well simulated using relatively felsic RI models with small NBO/T values and the PG2019 dataset (Figs. 8 and 9), with PG000 and PG010 leading to the smallest RMS values (Fig. 15d). Although the measured BTS for pixels in the area of  $8.0^\circ \text{ S} \leq lat \leq 9.5^\circ \text{ S}$  and  $106.5^\circ \text{ E} \leq lon \leq 109.5^\circ \text{ E}$  are negatively biased due to MC contamination, their spectral features, such as the wavenumber of the local maximum ( $\nu_a$ ), were similar to those shown in Figure 15d (plots of BTS for the Kelud VAC are also shown in Clarisse and Prata, 2016). Whether these measured BTS can be simulated well by internal or external mixing between the andesite model of VAC and ice cloud remains unclear. In addition to internal mixing between ash and ice based on effective medium theory, a simple external mixing process between individual ash and ice layers was examined through preliminary calculations, and the BTS simulations were unsuccessful. We could not conclude that Kelud ash plumes were composed of felsic ash aerosols based on the chemical composition ( $\sim 55 \text{ SiO}_2 \text{ wt.}\%$ ) of the Kelud ash samples from the same eruption (Maeno et al., 2019). However, the measured BTS of the Kelud VACs in this eruption event can be well explained by assuming that ash particles in the Kelud VACs had a similar infrared RI feature to PG000 and PG010.

1000


1005

1010

1015

1020

1025



**Figure15: Kelud ash plumes on 14 February 2014. MODIS true-colour image taken at 03:35 UTC (a) and monochromatic Himawari-7 visible image taken at 02:00 UTC (b).  $\Delta BT_{split}$  from the IASI measurement at 02:08 UTC (c) and measured and calculated BTS for the pixel at  $(lat, lon) = (9.101^\circ S, 110.513^\circ E)$  (d) [indicated by circles in (a–c)].**

## 5.6 Puyehue-Cordón Caulle (PCC)

From the beginning of the eruption on 4 June 2011, large amounts of volcanic ash and SO<sub>2</sub> gas were ejected from PCC, and the plumes reached an altitude of 12–13 km. Subsequently, ash plumes remained at high latitudes in the Southern Hemisphere

for a long time (Theys et al., 2013; Klüser et al., 2013; Clarisse et al., 2013). BTS analysis was conducted for ash plumes observed over the Atlantic Ocean at 10:41–10:47 UTC and 12:29 UTC on 6 June. In the area of IASI measurements at 10:41–10:47 UTC (Fig. 16c), MODIS observations were made at 11:05 UTC (Fig. 16b) and 12:45 UTC (Fig. 16a). From these MODIS images, ash plumes in the area with highly negative values of  $\Delta TB_{split}$  based on IASI measurements were distributed above broken MCs, and BTS simulations were performed for 29 IASI pixels that met the condition of Eq. (2). In addition, 10 additional IASI pixels were selected in the area of latitude 40–42° S and longitude 60–63° W from the data obtained at 12:29 UTC to avoid low-level dense MCs (circles in Fig. 16e, f). As shown in the ranking plot in Figure 8, almost all pixels have similar spectral features, and smaller mean RMS values were obtained for RI models with smaller NBO/T values in the PG2019 dataset. The smallest RMS was obtained for PG000 among all RI models. In accordance with the results of BTS analysis conducted by Newman et al. (2012), felsic RI models agreed well with the results of compositional analysis of ash sampled by Castro et al. (2013). Comparing BTS from simulations and measurements, the simulation results obtained using PG000 differ significantly from the observations (Fig. 16d, g). Although the simulation results exhibit a good fit to measurements in the wavenumber region of 820–1000  $\text{cm}^{-1}$  and at wavenumbers greater than 1130  $\text{cm}^{-1}$ , a large difference in brightness temperature at wavenumbers less than 820  $\text{cm}^{-1}$  suggests an excess of relative absorption in the PG000 RI model. Moreover, the simulated BTS values were significantly larger than the measurements between 1070 and 1130  $\text{cm}^{-1}$ , particularly for pixels in the area shown in Figure 16c. According to the results of VAC parameter retrieval using PG000, the effective radius  $r_{eff}$  was less than 0.5  $\mu\text{m}$  for most pixels,  $r_{eff}$  tended to decrease as distance from PCC increased, and most of the retrieval results showed  $r_{eff} \leq 0.3 \mu\text{m}$  for pixels around  $(lat, lon) = (27^\circ \text{ S}, 25^\circ \text{ W})$  (circle in Fig. 16c), which is 4500–4600 km from PCC. Considering a decrease in large particles due to deposition over a long transport period in the atmosphere,  $r_{eff} \leq 0.3 \mu\text{m}$  may be realistic in the case of PCC. Ash plumes from PCC eventually circled the Southern Hemisphere three times. Very small ash particles may increase the longevity of PCC ash plumes (Carn and Krotkov, 2016). According to the results of measurements by CALIOP on-board CALIPSO, PCC ash plumes are mainly composed of fine-mode non-spherical ash particles with a low sulphate contribution (Vernier et al., 2013; Prata et al., 2017a). For BTS in the range of 1070–1200  $\text{cm}^{-1}$ , the RI model PLRHY provides a better fit than PG000 (data not shown). This result suggests that an RI model with a stronger absorbing feature in the index  $k$  around 1050  $\text{cm}^{-1}$  than that of PG000 is better for explaining the measured BTS of PCC ash plumes.




1065

1070

1075

1080

1085



1090

Figure 16. (a–c) MODIS true-colour images at (a) 12:45 UTC and (b) 11:05 UTC, and (c) IASI  $\Delta BT_{split}$  at 10:41–10:47 UTC for Puyehue-Cordón Caulle ash plumes on 6 June 2011. (d) Results of BTS simulation with the PG000 RI model for the pixel at  $(lat, lon) = (27.136^\circ S, 25.178^\circ W)$ . (e–g) MODIS image at (e) 14:25 UTC and (f) IASI  $\Delta BT_{split}$  at 12:29 UTC. (g) Results of BTS simulation for the pixel at  $(lat, lon) = (41.212^\circ S, 61.898^\circ W)$ .

## 6. Summary and discussions

Using infrared brightness temperature data from IASI measurements of volcanic ash clouds (VACs) and radiative transfer calculations, we investigate whether refractive index (RI) models for ash particles could reproduce the measured brightness temperature spectra (BTS) in the infrared window region. We applied a total of 21 RI models to BTS simulations, including

1095

eight RI models using the Reed et al. (2018) dataset (RE2018), 11 RI models using the Prata et al. (2019) dataset (PG2019), and two additional RI models based on the andesite and rhyolite models of Pollack et al. (1973) (PL1973). For the eruption events of seven volcanoes, 156 daytime IASI pixels containing apparent volcanic ash features ( $\Delta TB_{split} \leq -2K$ ) with low meteorological clouds (MCs) contamination were selected with reference to MODIS and Himawari-8/AHI visible-colour images and an additional brightness temperature condition. The BTS simulation for each pixel included the retrieval of VAC parameters ( $h_{top}, r_{eff}, \tau_c$ ), SO<sub>2</sub> column content and top height, and the root mean square (RMS) between the measured and simulated BTS for each RI model was calculated. The results of the RI models were ranked based on the smallest RMS values for each pixel. Typical BTS simulation results were discussed for the ash plumes of some volcanoes.

As shown in Fig. 9, the RMS and its standard deviation between the simulation and observed BTS for each RI model are relatively large. Furthermore, the differences in the derived mean RMS between the RI models are relatively small because we individually estimated the VAC parameters and SO<sub>2</sub> for each RI model in the BTS fittings procedure. Therefore, it is difficult to determine a specific RI model from statistically insufficient BTS measurements, and the RI models proposed here are not always statistically significant and they can vary depending on the measurement dataset and the assumed atmospheric state.

With that in mind, the findings obtained from the analyses are summarised below.

Using RI models based on new RI datasets (RE2018, PG2019), the fitting of brightness temperature simulation for VACs in the atmospheric window region was improved. In particular, RI models using the PG2019 dataset performed as well as RE2018 dataset. This suggests that the NBO/T parameterisation of Prata et al. (2019) is effective for RI modelling at infrared window wavelengths.

In our BTS simulations, RI models for Eyjafjallajökull (RE030, RE040) with the RE2018 dataset and RI models parameterised using similar NBO/T values to those of Eyjafjallajökull ash samples (PG030, PG040) represented the measured BTS of Eyjafjallajökull ash plumes with good accuracy. Such consistency for mafic RI models was also observed for Grímsvötn ash plumes. The correlation between ash samples and ash plume measurements based on RI for the same volcanic event indicates that hyperspectral sounder measurements may help the selection of the RI model for ash aerosols, in addition to the information on NBO/T and SiO<sub>2</sub> wt.% data obtained from compositional analysis of ash samples.

The RI models estimated from BTS simulations of IASI measurements were not always consistent with models deduced from the reported chemical compositions of ash samples. For ash plumes released from Calbuco (Calbuco\_A) and Kelud, the RI models estimated from IASI measurements were more felsic (or had lower NBO/T) than the rock type or SiO<sub>2</sub> wt.% of tephra samples of the corresponding volcano. Furthermore, some RI models were excluded from acceptable RI models despite good BTS fitting results because the retrieved effective radii were too small. Negative values of brightness temperature difference  $\Delta TB_{split}$  are greater for ash particles of smaller sizes, and therefore our analysis is biased toward ash plumes with small  $r_{eff}$ . In addition, the retrieved  $r_{eff}$  values tended to be smaller when mafic RI models were used compared to the results of felsic RI models. This difference is due to fitting of the measured BTS in the wavenumber range of  $950 \text{ cm}^{-1} \leq \nu \leq 1230 \text{ cm}^{-1}$ , excluding the ozone absorption band. An effective radius of less than 0.2  $\mu\text{m}$  is too small for fresh ash plumes shortly after

1130 eruption, but may be realistic for ash plumes that have undergone long-distance transport in the atmosphere. Moreover, the condensation of volcanic sulphate on pure ash particles during transport might alter the inherent optical properties of the ash plumes.

We used the conventional andesite model (PLAND) by Pollack et al. (1973) for comparisons in our BTS analysis. Except for the Grímsvötn ash plumes, PLAND did not always perform well among the RI models for IASI pixels in this study. This result indicates that substantial improvement in satellite retrieval results for VACs can be expected from replacing the conventional andesite model with a proper RI model based on the RE2018 and PG2019 datasets. We explored the reproducibility of measured BTS over the entire wavenumber range of 650–1400 cm<sup>-1</sup>, including bands for CO<sub>2</sub>, water vapour, ozone, and SO<sub>2</sub>, and noted that the estimated RI models may not produce good results for VAC parameters in retrievals from multi-channel satellite imagers using certain infrared wavelength channels.

1140 In this study, we used MODIS and Himawari-8 daytime visible images for rough estimation of MC contamination in IASI measurement areas. For that reason, the number of IASI pixels available for analysis was significantly reduced, and statistical evaluation of the retrieval results for some volcanic events was difficult. To achieve a sufficient number of analyses, thorough evaluation of MC contamination of night-time VAC data from infrared sounder measurements is necessary, and methods for such evaluation should be explored in future research. In recent years, plans for mounting infrared sounders on geostationary satellites have been undertaken. High-frequency observations of distinct volcanic plumes increase the number of BTS data available, and precise retrieval is expected with further improvements in VAC analysis to combine infrared sounder and high-resolution imager measurements.

We tested 21 RI models in this article, and conspicuous discrepancies in measured and calculated BTS at specific wavenumber ranges were confirmed in all RI models for some volcanic events. These discrepancies may be resolved by applying precise atmospheric profiles, such as temperature and humidity profiles, and surface temperature in the radiative transfer calculations. Nevertheless, the number of RI models may still be insufficient for simulating all BTS patterns of VACs for various types of volcanoes. Recently, another RI dataset based on laboratory analysis of ash samples from six volcanoes was published by Deguine et al. (2020). A set of RI models, in which the silica content and porosity as well as glass-to-crystal ratio were considered, was also published by Piontek et al. (2021). These data are available from the journal's website. We would like to examine these novel RI models in the future.

## References

- Bessho, K., Date, K., Hayashi, M., Ikeda, A., Imai, T., Inoue, H., Kumagai, Y., Miyakawa, T., Murata, H., Ohno, T., Okuyama, A., Oyama, R., Sasaki, Y., Shimazu, Y., Shimoji, K., Sumida, Y., Suzuki, M., Taniguchi, H., Tsuchiyama, H., Uesawa, D., Yokota, H. and Yoshida, R.: An Introduction to Himawari-8/9 -Japan's New-Generation Geostationary Meteorological Satellites, J. Meteorol. Soc. Japan. Ser. II, 94(2), 151–183, doi:10.2151/jmsj.2016-009, 2016.

- Buehler, S. A., John, V. O., Kottayil, A., Milz, M. and Eriksson, P.: Efficient radiative transfer simulations for a broadband infrared radiometer—Combining a weighted mean of representative frequencies approach with frequency selection by simulated annealing, *J. Quant. Spectrosc. Radiat. Transf.*, 111(4), 602–615, doi:10.1016/j.jqsrt.2009.10.018, 2010.
- 1165 Carn, S. A. and Krotkov, N. A.: Ultraviolet Satellite Measurements of Volcanic Ash, in *Volcanic Ash*, edited by S. Mackie, K. Cashman, H. Ricketts, A. Rust, and M. Watson, pp. 217–231, Elsevier., 2016.
- Cashman, K. and Rust, A.: Volcanic Ash: Generation and Spatial Variations, in *Volcanic Ash*, edited by S. Mackie, K. Cashman, H. Ricketts, A. Rust, and M. Watson, pp. 5–22, Elsevier., 2016.
- 1170 Castro, J. M., Schipper, C. I., Mueller, S. P., Militzer, A. S., Amigo, A., Parejas, C. S. and Jacob, D.: Storage and eruption of near-liquidus rhyolite magma at Cordón Caulle, Chile, *Bull. Volcanol.*, 75(4), 702, doi:10.1007/s00445-013-0702-9, 2013.
- Clarisse, L. and Prata, F.: Infrared Sounding of Volcanic Ash, in *Volcanic Ash*, edited by S. Mackie, K. Cashman, H. Ricketts, A. Rust, and M. Watson, pp. 189–215, Elsevier., 2016.
- 1175 Clarisse, L., Hurtmans, D., Clerbaux, C., Hadji-Lazaro, J., Ngadi, Y. and Coheur, P.-F.: Retrieval of sulphur dioxide from the infrared atmospheric sounding interferometer (IASI), *Atmos. Meas. Tech.*, 5(3), 581–594, doi:10.5194/amt-5-581-2012, 2012.
- Clarisse, L., Coheur, P.-F., Prata, F., Hadji-Lazaro, J., Hurtmans, D. and Clerbaux, C.: A unified approach to infrared aerosol remote sensing and type specification, *Atmos. Chem. Phys.*, 13(4), 2195–2221, doi:10.5194/acp-13-2195-2013, 2013.
- 1180 Clough, S. A., Shephard, M. W., Mlawer, E. J., Delamere, J. S., Iacono, M. J., Cady-Pereira, K., Boukabara, S. and Brown, P. D.: Atmospheric radiative transfer modeling: a summary of the AER codes, *J. Quant. Spectrosc. Radiat. Transf.*, 91(2), 233–244, doi:10.1016/j.jqsrt.2004.05.058, 2005.
- Corradini, S., Merucci, L. and Prata, A. J.: Retrieval of SO<sub>2</sub> from thermal infrared satellite measurements: Correction procedures for the effects of volcanic ash, *Atmos. Meas. Tech.*, 2(1), 177–191, doi:10.5194/amt-2-177-2009, 2009.
- 1185 Deguine, A., Petitprez, D., Clarisse, L., Gudmundsson, S., Outes, V., Villarosa, G. and Herbin, H.: Complex refractive index of volcanic ash aerosol in the infrared, visible, and ultraviolet, *Appl. Opt.*, 59(4), 884, doi:10.1364/AO.59.000884, 2020.
- Francis, P. N., Cooke, M. C. and Saunders, R. W.: Retrieval of physical properties of volcanic ash using Meteosat: A case study from the 2010 Eyjafjallajökull eruption, *J. Geophys. Res. Atmos.*, 117(D20), 1–14, doi:10.1029/2011JD016788, 2012.
- 1190 Gangale, G., Prata, A. J. and Clarisse, L.: The infrared spectral signature of volcanic ash determined from high-spectral resolution satellite measurements, *Remote Sens. Environ.*, 114(2), 414–425, doi:10.1016/j.rse.2009.09.007, 2010.
- Hansen, J. E. and Travis, L. D.: Light scattering in planetary atmospheres, *Space Sci. Rev.*, 16(4), 527–610, doi:10.1007/BF00168069, 1974.
- 1195 Harvey, N. J., Dacre, H. F., Webster, H. N., Taylor, I. A., Khanal, S., Grainger, R. G. and Cooke, M. C.: The Impact of Ensemble Meteorology on Inverse Modeling Estimates of Volcano Emissions and Ash Dispersion Forecasts: Grímsvötn 2011, *Atmosphere (Basel)*, 11(10), 1022, doi:10.3390/atmos11101022, 2020.

- Hilton, F., Armante, R., August, T., Barnet, C., Bouchard, A., Camy-Peyret, C., Capelle, V., Clarisse, L., Clerbaux, C., Coheur, P.-F., Collard, A., Crevoisier, C., Dufour, G., Edwards, D., Faijan, F., Fourrié, N., Gambacorta, A., Goldberg, M., Guidard, V., Hurtmans, D., Illingworth, S., Jacquinet-Husson, N., Kerzenmacher, T., Klaes, D., Lavanant, L., Masiello, G., Matricardi, M., McNally, A., Newman, S., Pavelin, E., Payan, S., Péquignot, E., Peyridieu, S., Phulpin, T., Remedios, J., Schlüssel, P., Serio, C., Strow, L., Stubenrauch, C., Taylor, J., Tobin, D., Wolf, W. and Zhou, D.: Hyperspectral Earth Observation from IASI: Five Years of Accomplishments, *Bull. Am. Meteorol. Soc.*, 93(3), 347–370, doi:10.1175/BAMS-D-11-00027.1, 2012.
- Holl, G., Buehler, S. A., Mendrok, J. and Kottayil, A.: Optimised frequency grids for infrared radiative transfer simulations in cloudy conditions, *J. Quant. Spectrosc. Radiat. Transf.*, 113(16), 2124–2134, doi:10.1016/j.jqsrt.2012.05.022, 2012.
- Ishimoto, H., Masuda, K., Fukui, K., Shimbori, T., Inazawa, T., Tuchiya, H., Ishii, K. and Sakurai, T.: Estimation of the refractive index of volcanic ash from satellite infrared sounder data, *Remote Sens. Environ.*, 174, 165–180, doi:10.1016/j.rse.2015.12.009, 2016.
- Klüser, L., Erbetseder, T. and Meyer-Arnek, J.: Observation of volcanic ash from Puyehue–Cordón Caulle with IASI, *Atmos. Meas. Tech.*, 6(1), 35–46, doi:10.5194/amt-6-35-2013, 2013.
- Kylling, A.: Ash and ice clouds during the Mt Kelud February 2014 eruption as interpreted from IASI and AVHRR/3 observations, *Atmos. Meas. Tech.*, 9(5), 2103–2117, doi:10.5194/amt-9-2103-2016, 2016.
- Kylling, A., Kristiansen, N., Stohl, A., Buras-Schnell, R., Emde, C. and Gasteiger, J.: A model sensitivity study of the impact of clouds on satellite detection and retrieval of volcanic ash, *Atmos. Meas. Tech.*, 8(5), 1935–1949, doi:10.5194/amt-8-1935-2015, 2015.
- Mackie, S. and Watson, M.: Probabilistic detection of volcanic ash using a Bayesian approach, *J. Geophys. Res. Atmos.*, 119(5), 2409–2428, doi:10.1002/2013JD021077, 2014.
- Mackie, S., Cashman, K., Ricketts, H., Rust, A. and Watson, M.: Introduction, in *Volcanic Ash*, edited by S. Mackie, K. Cashman, H. Ricketts, A. Rust, and M. Watson, pp. 1–3, Elsevier., 2016.
- Maeno, F., Nakada, S., Yoshimoto, M., Shimano, T., Hokanishi, N., Zaennudin, A. and Iguchi, M.: A sequence of a plinian eruption preceded by dome destruction at Kelud volcano, Indonesia, on February 13, 2014, revealed from tephra fallout and pyroclastic density current deposits, *J. Volcanol. Geotherm. Res.*, 382, 24–41, doi:10.1016/j.jvolgeores.2017.03.002, 2019.
- Matricardi, M.: The generation of RTTOV regression coefficients for IASI and AIRS using a new profile training set and a new line-by-line database, *ECMWF Tech. Memo.*, (May), 2008.
- Mishchenko, M. I., Travis, L. and Lacis, A. a: *Scattering, Absorption, and Emission of Light by Small Particles*, Cambridge, pp445, 2002.
- Mlawer, E. J., Payne, V. H., Moncet, J.-L., Delamere, J. S., Alvarado, M. J. and Tobin, D. C.: Development and recent evaluation of the MT\_CKD model of continuum absorption, *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 370(1968), 2520–2556, doi:10.1098/rsta.2011.0295, 2012.

- 1230 Moxnes, E. D., Kristiansen, N. I., Stohl, A., Clarisse, L., Durant, A., Weber, K. and Vogel, A.: Separation of ash and sulfur dioxide during the 2011 Grímsvötn eruption, *J. Geophys. Res. Atmos.*, 119(12), 7477–7501, doi:10.1002/2013JD021129, 2014.
- Newman, S. M., Clarisse, L., Hurtmans, D., Marengo, F., Johnson, B., Turnbull, K., Havemann, S., Baran, A. J., O’Sullivan, D. and Haywood, J.: A case study of observations of volcanic ash from the Eyjafjallajökull eruption: 2. Airborne and  
1235 satellite radiative measurements, *J. Geophys. Res. Atmos.*, 117(D20), 1–19, doi:10.1029/2011JD016780, 2012.
- Pavolonis, M. J., Heidinger, A. K. and Sieglaff, J.: Automated retrievals of volcanic ash and dust cloud properties from upwelling infrared measurements, *J. Geophys. Res. Atmos.*, 118(3), 1436–1458, doi:10.1002/jgrd.50173, 2013.
- Piontek, D., Hornby, A. J., Voigt, C., Bugliaro, L. and Gasteiger, J.: Determination of complex refractive indices and optical properties of volcanic ashes in the thermal infrared based on generic petrological compositions, *J. Volcanol. Geotherm. Res.*, 411, 107174, doi:10.1016/j.jvolgeores.2021.107174, 2021.  
1240
- Pollack, J. B., Toon, O. B. and Khare, B. N.: Optical properties of some terrestrial rocks and glasses, *Icarus*, 19(3), 372–389, doi:10.1016/0019-1035(73)90115-2, 1973.
- Prata, A. J.: Infrared radiative transfer calculations for volcanic ash clouds, *Geophys. Res. Lett.*, 16(11), 1293–1296, doi:10.1029/GL016i011p01293, 1989.
- 1245 Prata, A. J. and Grant, I. F.: Retrieval of microphysical and morphological properties of volcanic ash plumes from satellite data: Application to Mt Ruapehu, New Zealand, *Q. J. R. Meteorol. Soc.*, 127(576), 2153–2179, doi:10.1002/qj.49712757615, 2001.
- Prata, A. T., Young, S. A., Siems, S. T. and Manton, M. J.: Lidar ratios of stratospheric volcanic ash and sulfate aerosols retrieved from CALIOP measurements, *Atmos. Chem. Phys.*, 17(13), 8599–8618, doi:10.5194/acp-17-8599-2017, 2017a.
- 1250 Prata, F. and Lynch, M.: Passive Earth Observations of Volcanic Clouds in the Atmosphere, *Atmosphere (Basel)*, 10(4), 199, doi:10.3390/atmos10040199, 2019.
- Prata, F., Woodhouse, M., Huppert, H. E., Prata, A., Thordarson, T. and Carn, S.: Atmospheric processes affecting the separation of volcanic ash and SO<sub>2</sub> in volcanic eruptions: Inferences from the May 2011 Grímsvötn eruption, *Atmos. Chem. Phys.*, 17(17), 10709–10732, doi:10.5194/acp-17-10709-2017, 2017b.
- 1255 Prata, G. S., Ventress, L. J., Carboni, E., Mather, T. A., Grainger, R. G. and Pyle, D. M.: A New Parameterization of Volcanic Ash Complex Refractive Index Based on NBO/T and SiO<sub>2</sub> Content, *J. Geophys. Res. Atmos.*, 124(3), 1779–1797, doi:10.1029/2018JD028679, 2019.
- Reed, B. E., Peters, D. M., McPheat, R., Smith, A. J. A. and Grainger, R. G.: Mass extinction spectra and size distribution measurements of quartz and amorphous silica aerosol at 0.33–19  $\mu\text{m}$  compared to modelled extinction using Mie, CDE, and T-matrix theories, *J. Quant. Spectrosc. Radiat. Transf.*, 199, 52–65, doi:10.1016/j.jqsrt.2017.05.011, 2017.  
1260
- Reed, B. E., Peters, D. M., McPheat, R. and Grainger, R. G.: The Complex Refractive Index of Volcanic Ash Aerosol Retrieved From Spectral Mass Extinction, *J. Geophys. Res. Atmos.*, 123(2), 1339–1350, doi:10.1002/2017JD027362, 2018.

- Romero, J. E., Morgavi, D., Arzilli, F., Daga, R., Caselli, A., Reckziegel, F., Viramonte, J., Díaz-Alvarado, J., Polacci, M.,  
 1265 Burton, M. and Perugini, D.: Eruption dynamics of the 22–23 April 2015 Calbuco Volcano (Southern Chile): Analyses  
 of tephra fall deposits, *J. Volcanol. Geotherm. Res.*, 317(April 2015), 15–29, doi:10.1016/j.jvolgeores.2016.02.027,  
 2016.
- Rothman, L. S., Gordon, I. E., Babikov, Y., Barbe, A., Chris Benner, D., Bernath, P. F., Birk, M., Bizzocchi, L., Boudon, V.,  
 Brown, L. R., Campargue, A., Chance, K., Cohen, E. A., Coudert, L. H., Devi, V. M., Drouin, B. J., Fayt, A., Flaud, J.-  
 1270 M., Gamache, R. R., Harrison, J. J., Hartmann, J.-M., Hill, C., Hodges, J. T., Jacquemart, D., Jolly, A., Lamouroux, J.,  
 Le Roy, R. J., Li, G., Long, D. A., Lyulin, O. M., Mackie, C. J., Massie, S. T., Mikhailenko, S., Müller, H. S. P.,  
 Naumenko, O. V., Nikitin, A. V., Orphal, J., Perevalov, V., Perrin, A., Polovtseva, E. R., Richard, C., Smith, M. A. H.,  
 Starikova, E., Sung, K., Tashkun, S., Tennyson, J., Toon, G. C., Tyuterev, V. G. and Wagner, G.: The HITRAN2012  
 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transf.*, 130, 4–50, doi:10.1016/j.jqsrt.2013.07.002, 2013.
- 1275 Saunders, R., Hocking, J., Rundle, D., Rayer, P., Matricardi, M., Geer, A., Lupu, C., Brunel, P. and Vidot, J.: Rttov-11  
 Science and Validation Report. [online] Available from: [https://nwpsaf.eu/deliverables/rtm/docs\\_rttov11/rttov11\\_svr.pdf](https://nwpsaf.eu/deliverables/rtm/docs_rttov11/rttov11_svr.pdf),  
 2013.
- Taylor, I. A., Carboni, E., Ventress, L. J., Mather, T. A. and Grainger, R. G.: An adaptation of the CO<sub>2</sub> slicing technique for  
 the Infrared Atmospheric Sounding Interferometer to obtain the height of tropospheric volcanic ash clouds, *Atmos.*  
 1280 *Meas. Tech.*, 12(7), 3853–3883, doi:10.5194/amt-12-3853-2019, 2019.
- Theys, N., Campion, R., Clarisse, L., Brenot, H., van Gent, J., Dils, B., Corradini, S., Merucci, L., Coheur, P.-F., Van  
 Roozendael, M., Hurtmans, D., Clerbaux, C., Tait, S. and Ferrucci, F.: Volcanic SO<sub>2</sub> fluxes derived from satellite data: a  
 survey using OMI, GOME-2, IASI and MODIS, *Atmos. Chem. Phys.*, 13(12), 5945–5968, doi:10.5194/acp-13-5945-  
 2013, 2013.
- 1285 Turnbull, K., Johnson, B., Marengo, F., Haywood, J., Minikin, A., Weinzierl, B., Schlager, H., Schumann, U., Leadbetter, S.  
 and Woolley, A.: A case study of observations of volcanic ash from the Eyjafjallajökull eruption: 1. In situ airborne  
 observations, *J. Geophys. Res. Atmos.*, 117(D20), 1–19, doi:10.1029/2011JD016688, 2012.
- Ventress, L. J., McGarragh, G., Carboni, E., Smith, A. J. and Grainger, R. G.: Retrieval of ash properties from IASI  
 measurements, *Atmos. Meas. Tech.*, 9(11), 5407–5422, doi:10.5194/amt-9-5407-2016, 2016.
- 1290 Vernier, J.-P., Fairlie, T. D., Murray, J. J., Tupper, A., Trepte, C., Winker, D., Pelon, J., Garnier, A., Jumelet, J., Pavolonis,  
 M., Omar, A. H. and Powell, K. A.: An Advanced System to Monitor the 3D Structure of Diffuse Volcanic Ash Clouds,  
*J. Appl. Meteorol. Climatol.*, 52(9), 2125–2138, doi:10.1175/JAMC-D-12-0279.1, 2013.
- Vidot, J. and Hocking, J.: Note on RTTOV v12 unit conversions for gases , clouds and aerosols. [online] Available from:  
[https://www.nwpsaf.eu/site/download/documentation/rtm/docs\\_rttov12/rttov\\_gas\\_cloud\\_aerosol\\_units.pdf](https://www.nwpsaf.eu/site/download/documentation/rtm/docs_rttov12/rttov_gas_cloud_aerosol_units.pdf), 2017.
- 1295 Wen, S. and Rose, W. I.: Retrieval of sizes and total masses of particles in volcanic clouds using AVHRR bands 4 and 5, *J.*  
*Geophys. Res.*, 99(D3), 5421, doi:10.1029/93JD03340, 1994.