

Laboratory evaluation of the scattering matrix of ragweed, ash, birch and pine pollen towards pollen classification

Danaël Cholleton^{1,2}, Émilie Bialic², Antoine Dumas², Pascal Kaluzny², Patrick Rairoux¹, Alain Miffre¹

¹University of Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622, VILLEURBANNE, France

²TERA Sensor, ZI Rousset, 296 Avenue Georges Vacher, 13790, Rousset, France

Correspondence to: Alain Miffre (alain.miffre@univ-lyon1.fr)

Abstract. Pollen are nowadays recognized as one of the main atmospheric particles affecting public human health as well as the Earth's climate. In this context, an important issue concerns our ability to detect and differentiate among the existing pollen taxa. In this paper, the potential differences that may exist in light scattering by four of the most common pollen taxa, namely ragweed, birch, pine and ash, are analysed in the framework of the scattering matrix formalism at two wavelengths simultaneously (532 and 1064 nm). Interestingly, our laboratory experimental error bars are precise enough to show that these four pollen, when embedded in ambient air, exhibit different spectral and polarimetric light scattering characteristics, in the form of ten scattering matrix elements (five per wavelength), which allow identifying each separately. To end with, a simpler light scattering criterion is proposed for classifying among the four considered pollen by performing a principal component (PC) analysis, that still accounts for more than 99 % of the observed variance. We thus believe this work may open new insights for future atmospheric pollen detection.

1 Introduction

Pollen are biological aerosols impacting public health (Schaffner et al., 2020) and the Earth's climate (Li et al., 2013). The economical cost of pollen allergy on public health is impressive and has been estimated to reach up to 151 billion euros in Europe in 2014 (Lake et al., 2017). Moreover, this cost is expected to increase as the prevalence of allergies in the global population is increasing worldwide. Apart from this socio-economical cost, pollen may locally influence the Earth's radiative forcing, by increasing the IR downwelling flux (Spänkuch et al., 2000) and by acting as cloud-condensation nuclei (Pope, 2010). Indeed, *Ambrosia* pollen concentrations are expected to increase by 400 % in the following decades (Hamaoui-Laguel et al., 2015) for *Ambrosia* is an invasive plant. To our knowledge, no similar projection exists for other taxa. Based on observed evolutions over the 1994-2010 period however, (Zhang et al., 2015) reported a 42 % increase in the birch pollen concentration in the US. Due to climate change, which increases the global temperature and CO₂ atmospheric concentrations, the length of the pollen season should extend (Bielory et al., 2012). Moreover, the geographical repartition of pollen plants is also expected to extend (Ziska et al., 2011). These health and climatic impacts mainly depend on the involved pollen species, which are numerous and induce various allergic reactions. To better quantify the impact of pollen on human health and climate, and as

underscored by (Crouzy et al., 2016), reliable measurements and forecasts are required as well as a reliable method for pollen identification and classification. At present, the most widely used methodology is that developed by (Hirst, 1952) seventy years ago, consisting of sampling then precipitating atmospheric pollen grains on an adhesive substrate prior to the microscopic observation of the deposited pollen grains. Though rather accurate, this historical methodology suffers from a too long time duration, with characteristic times ranging from a few hours to a week, much higher than the characteristic time for the impact of pollen on human health. Another limitation of the historical methodology is that the pollen grains are counted one by one and identified by a scientist, which is time-consuming and may be operator dependent (Comtois et al., 1999). In this context, deep learning recently improved the identification of pollen grains through microscopy (Schaefer et al., 2021) by automating the procedure to differentiate among existing pollen. Besides, new methodologies based on interferometry, light-scattering or laser-induced fluorescence have been developed. In the latter, the fluorescence spectrum and its life-time have been used from the UV to the NIR spectral range to identify pollen (Pan et al., 2011; Kiselev et al., 2013). Likewise, image recognition on the scattering pattern of pollen grains have been investigated, as described by (Šaulienė et al., 2019), and holographic images are also used (Giri et al., 2019; Sauvageat et al., 2020; Kemppinen et al., 2020) as an identification methodology. Finally, light scattering by pollen has been studied in laboratory in aqueous solutions by (Bickel and Stafford, 1980) or when pollen are deposited on a substrate or an holder by (Surbek et al., 2011; Iwai, 2013; Raman et al., 2013; Nouri et al., 2018). Concerning airborne pollen, (Matsuda and Kawashima, 2018; Holler et al., 2016) studied forward and side scattering while polarization-dependent features of light scattering were first studied by our group by evaluating the scattering matrix of ragweed pollen in laboratory ambient air at near backscattering angles (Cholleton et al., 2020). More recently, (Gómez Martín et al., 2021) evaluated the scattering matrix of cypress pollen far from the backscattering angle. Also, the depolarization ratio of several pollen has been studied in field by (Cao, 2010; Bohlmann et al., 2018; Sicard et al., 2021) using lidar measurements. Nevertheless, light scattering by pollen embedded in ambient air still remains a complex topic, mostly due to the complexity in size and shape of the pollen taxa. In particular, it is not a priori granted that pollen differing in size and shape exhibit different light scattering characteristics. Indeed, as underscored in (Cholleton et al., 2020), no analytical light scattering numerical simulation exists for such complex objects so that a remaining issue is to predict how these grains scatter light. Therefore, a controlled-laboratory experiment is coveted to address light scattering by pollen with a precision allowing to identify each pollen.

In this paper, to complement the above airborne pollen studies, the scattering matrix of several airborne pollen taxa is for the first time presented. We focus on ragweed, ash and birch pollen, which are allergenic, and currently monitored in several countries in North America and in Europe. Also, pine pollen is studied in complement as strong pine pollen events have been reported in the literature: (Spänkuch et al., 2000) reported that a high pine pollen concentration increased the down welling infrared flux up to 8 times the monthly means. Our main selection criteria for choosing our pollen samples was then the geographical location (Europe, North America) and the allergenic character or / and the climatic impact of the pollen taxon. Still, other pollen taxa could be likewise studied by applying our new measurement technique, provided that the same accuracy

65 is experimentally achieved. Extension of this work to all pollen taxa in a given geographical region is however beyond the scope of our contribution. The laboratory experiment is carried out at two wavelengths simultaneously ($\lambda_{VIS} = 532$ nm and $\lambda_{IR} = 1064$ nm) and provides five scattering matrix elements per pollen taxa and per wavelength. Interestingly, our experimental error bars are precise enough to reveal the existing differences in light scattering by these four pollen, thus allowing to identify each separately. Ten scattering matrix elements are provided per pollen (five per wavelength). Hence, 70 compared with our previous contribution (Cholleton et al., 2020), three more pollen are here studied (ash, birch and pine) which allows discussing on airborne pollen classification using polarization-resolved light scattering. As an output, a principal component analysis is applied to provide a simpler light-scattering criterion for classifying among these four pollen. The classification can then be visualized on a simple 2D-plane representation. The novelty of the paper is hence two-fold. Firstly, precise light scattering characteristics of ragweed, ash, birch and pine are revealed in the form of ten scattering matrix elements 75 (five per pollen per wavelength). Secondly, an analysis is proposed towards a pollen classification. The paper first presents the studied pollen samples, then details our laboratory methodology to precisely evaluate the scattering matrix of these pollen. From that, the evaluation of their scattering matrix is presented, and a principal component analysis is proposed as an outlook to help classifying among these pollen.

2 Pollen samples

80 Figure 1 shows scanning electron microscopic images of the four studied pollen species, namely ragweed, ash, birch and pine. These microscopic images allow characterizing the size and the shape of these pollen: each pollen exhibits a very characteristic shape with comprising grains presenting a quasi-monodisperse size distribution. The four considered species clearly differ in size and shape, from nearly spherical with a $20\ \mu\text{m}$ volume equivalent diameter for ragweed to highly irregular shapes with a diameter larger than $60\ \mu\text{m}$ for pine. Small-scale features exist at each pollen surface, such as granularity or apertures, which 85 show the great complexity of these microscopic objects. These specific sizes and shapes are used for pollen recognition as extensively detailed in palynological databases (PalDat, 2021), as recalled in the introduction.

2.1 Ragweed

Ragweed or *Ambrosia artemisiifolia* pollen is amongst the most allergenic pollen in Europe and Northern America (Smith et al., 2013) with 15.8 million persons sensitized in Europe (Schaffner et al., 2020). Ragweed pollen induces particular allergic 90 reactions twice more often as other pollen (Dahl et al., 1999) with an annual economic cost of 7 billion euros in Europe (Schaffner et al., 2020). The ragweed season usually lasts from August to October and this duration is getting longer with expected increasing mean temperatures. Hence, ragweed pollen has been extensively studied in the literature (Smith et al., 2013; Hamaoui-Laguel et al., 2015; Zink et al., 2017) and extensive efforts are made for the observations and forecasts of its concentration (Prank et al., 2013). Ragweed pollen grains exhibit a characteristic shape, that bears some resemblance with 95 coronavirus virions (Petrov, 2020), are nearly but non spherical, with a mean volume-equivalent diameter of 21 micrometres.

Its membrane exhibit three apertures (2 micrometres long), regular spikes (*echinus*) of about 1 micrometre length and a perforate structure, i.e. covered with holes of about 100 nanometres diameter.

2.2 Ash


We here consider ash (*Fraxinus americana*) pollen which is a relevant source of allergenic reactions in North America. In Europe, ash pollen has been underestimated for a long time as its bloom season overlaps with that of birch (Imhof et al., 2014). Hence, *Fraxinus excelsior*, highly present in Europe, may also be considered as an outlook of this work, provided that the same experimental accuracy is achieved. Moreover, it presents a high cross-reactivity from allergens from other plant species such as birch pollen, as underscored by (Niederberger et al., 2002). Its blooming season lasts from March to May. Ash pollen grains have an irregular shape, with a reticulate exine, i.e. ornamental elements arranged as a network of ridges with gaps of the micron scale, and 3-4 colpi, i.e. thin elongated apertures regularly distributed along its equatorial region. Its volume equivalent diameter is about 31 micrometres.

2.3 Birch

Alongside with ragweed, birch (*Betula pendula*) pollen is one of the most impacting in central and northern Europe (Biedermann et al., 2019). It also exhibits a cross-reactivity with other allergens. Depending on its location, the birch pollen season starts from March and lasts until late June. Birch pollen grains are irregularly shaped, with sunken interpectoral areas when dried. Their mean volume equivalent diameter is about 28 to 29 micrometres. They present 3 protruding pores, i.e. three 2 µm large holes on its polar view. Birch pollen has been one off the first to be integrated in global pollen forecasts (Siljamo et al., 2013), using dispersion models.

2.4 Pine

In contrary to the previous taxa, pine (*Pinus strobus*) pollen rarely cause allergies. Moreover, this pollen has been studied by optical diffraction tomography (Kim et al., 2018) to highlight its structure and morphological features, and evaluate its refractive index. At last, (Sicard et al., 2021) recently studied its vertical distribution in the atmosphere using lidar-derived profiles. As shown Fig. 1(d), pine pollen is the largest of the studied samples with a mean diameter of 69 micrometers. Its shape largely differs from that of other pollen, as two *sacci*, i.e. two large air-filled bladders are attached to the center part (Schwendemann et al., 2007). The refractive index of pine at 532 nm has been estimated between 1.5 and 1.54 (Kim et al., 2018), while its imaginary part remains undocumented.



125 Figure 1 : Scanning electron microscopy (SEM) images of the studied pollen taken at iLM: (a) ragweed (*Ambrosia artemisiifolia*), (b) ash (*Fraxinus americana*), (c) birch (*Betula pendula*), (d) pine (*Pinus strobus*). The scale bar is 5 μm on all pictures. Pollen grains were suspended in ambient air and deposited on an adhesive substrate for SEM observations.

2.5 Pollen suspension

The studied pollen samples were supplied by Stallergenes Greer. Dry pollen powder has been suspended in the light scattering volume using a solid aerosol generator supplied with dried compresses air ($\text{RH} < 10\%$). The size distribution and concentration of the generated airborne ragweed pollen grains was measured using an Aerodynamic Particle Sizer (APS). As a consistency check, the retrieved ragweed pollen size distribution was found in agreement with the size specified by the supplier and also with above SEM observations issued from the state-of-the-art literature (PalDat, 2021). These commercial pollen grains may differ from that of atmospheric pollen grains. However, our Figure 1 microscopic images did not exhibit differences in size nor shape compared with fresher pollens. Otherwise, the most recent literature does not report any changes in the pollen light scattering properties over a ten days period (Miki and Kawashima, 2021).

3 Methodology

3.1 Light scattering by pollen


We consider elastic light scattering by an ensemble of pollen grains suspended in ambient air with a polarized electromagnetic incident radiation of wavelength λ . For the pollen samples are nonspherical (see Fig. 1), the polarization state of the scattered radiation may differ from that of the incident radiation. To describe the polarization states of the incident and scattered waves, the Stokes vector formalism (Mishchenko et al., 2002) is applied. The polarization state of the incident and scattered radiation is then described by a Stokes vector $S = [I, Q, U, V]^T$ where I, Q, U, V respectively relate to the total intensity, the linear degree of polarization in the scattering plane, and at 45° from it, and the degree of circular polarization. In the far-field single scattering approximation and assuming random orientation, we learn from light scattering textbooks (Mishchenko et al., 2002) that the incident and scattered Stokes vectors S_0 and S relate by the so-called scattering matrix :

$$\begin{bmatrix} 1 & f_{12}^\lambda & 0 & 0 \\ f_{12}^\lambda & f_{22}^\lambda & 0 & 0 \\ 0 & 0 & f_{33}^\lambda & f_{34}^\lambda \\ 0 & 0 & -f_{34}^\lambda & f_{44}^\lambda \end{bmatrix} \quad (1)$$

where the scattering matrix elements f_{ij}^λ ($i, j = 1 - 4$) have been normalized with respect to the scattering phase function so that f_{ij}^λ at most equals unity. f_{22}^λ and f_{33}^λ characterize how a linear polarization state is preserved during light scattering by the considered pollen while f_{44}^λ relates if circular polarization is preserved during light scattering by the pollen grains. Off-diagonal element f_{12}^λ quantifies the amount of non-polarized light after scattering by pollen, while the other off-diagonal element f_{34}^λ traduces the ability of pollen to convert a linearly polarized radiation to a circular polarization state throughout light scattering. The scattering matrix elements describe how the polarization state of the incident radiation is modified during light scattering by the studied pollen grains. These matrix elements hence depend on the shape of the pollen grains which we here investigate.

155 3.2 Laboratory experimental set-up for pollen light scattering

The specific size and shape of each pollen taxon is addressed by considering the Fig. 2 laboratory experiment, which has proven efficiency for ragweed pollen detection (Cholleton et al., 2020). We here recall its main characteristics for the sake of clarity before discussing on its applicability to the detection of other pollen in the next paragraph. This laboratory polarimeter operates at the near backscattering angle $\theta = (177.5 \pm 0.2)^\circ$ and at two wavelengths $\lambda_{VIS} = 532$ nm and $\lambda_{IR} = 1064$ nm simultaneously. A reflecting polarizing beam-splitter cube (PBC) is used so that it is the s-component of the scattered radiation that is measured by the photodetector. This experimental set-up has been validated on spherical particles (Cholleton et al., 2020) for which the scattering matrix can be analytically computed by applying the Mie theory, by measuring the corresponding size distribution with particle optical sizers.



165 Figure 2: Scheme of the laboratory experimental set-up of the $(\lambda_{VIS}, \lambda_{IR})$ polarimeter from (Cholleton et al., 2020). The 177.5° scattering angle has been exaggerated to ease the reading. $\mathbf{k}_0(\lambda)$ and $\mathbf{k}(\lambda)$ respectively stand for the incident and scattered wavevectors at wavelength λ . Pollen grains are embedded in laboratory ambient air as described Section 2.5. ψ is the angle between the fast axis of the quarter-wave plate (QWP) and the scattering plane.

170 Following Fig. 2, the detected pollen scattered intensity can be obtained by considering the successive Mueller matrices encountered by the incident radiation. After a few calculations (Cholleton et al., 2020), we get at wavelength λ :

$$I_i^\lambda(\psi) = I_0^\lambda \times [a_i^\lambda - b_i^\lambda \sin(2\psi) - c_i^\lambda \cos(4\psi) - d_i^\lambda \sin(4\psi)] \quad (2)$$

175 where subscript i stands for incident polarization state while I_0^λ is a proportionality constant that accounts for the incident laser power and the electro-optics efficiency. The pollen scattering matrix elements are retrieved from Eq. (2) by adjusting the detected scattered intensity as a function of the angle ψ to retrieve the $a_i^\lambda, b_i^\lambda, c_i^\lambda$ and d_i^λ coefficients, which only depend on the scattering matrix elements. To gain in accuracy, the polarization state of the scattered radiation is analysed for several incident polarization states, obtained by rotating an analyser (QWP), whose position is labelled by the angle ψ . To retrieve all
180 the pollen matrix elements, the polarization state of the scattered radiation is analysed for three successive incident polarization states, namely (p) , $(45 +)$ and (RC) . To give an example, a (RC) -polarized incident radiation allows retrieving $f_{12}^\lambda, f_{34}^\lambda$ and f_{44}^λ , as established in (Cholleton et al., 2020). Adding a $(45 +)$ -polarized incident radiation allows evaluating f_{33}^λ in addition. Finally, f_{22}^λ is retrieved from a (p) -polarized incident radiation. In summary, the following set of equations, established in (Cholleton et al., 2020), is applied to retrieve the pollen scattering matrix elements from adjusted coefficients $a_i^\lambda, b_i^\lambda, c_i^\lambda$ and
185 d_i^λ :

$$f_{12}^\lambda = 2c_{RC}^\lambda / (a_{RC}^\lambda + c_{RC}^\lambda) \quad (3a)$$

$$f_{34}^\lambda = 2d_{RC}^\lambda / (a_{RC}^\lambda + c_{RC}^\lambda) \quad (3b)$$

$$f_{44}^\lambda = -b_{RC}^\lambda / (a_{RC}^\lambda + c_{RC}^\lambda) \quad (3c)$$

$$f_{33}^\lambda = 2d_{45+}^\lambda / (a_{45+}^\lambda + c_{45+}^\lambda) \quad (3d)$$

$$f_{22}^\lambda = [f_{12}^\lambda \times (c_p^\lambda - a_p^\lambda) + 2c_p^\lambda] / (a_p^\lambda + c_p^\lambda) \quad (3e)$$

3.3 Applicability of this set-up to the detection of other pollen

To study the sensitivity of our methodology to other pollen, three different Taxa are considered, characterized by the following
190 set of scattering matrix elements:

- a spherical pollen called *Taxon 1*, with normalized scattering matrix elements [$f_{22}^\lambda = 1, f_{33}^\lambda = 1, f_{44}^\lambda = -1, f_{12}^\lambda = 0, f_{34}^\lambda = 0$]. *Taxon 1* is to our knowledge an hypothetical pollen since pollen are nonspherical (Hesse, 2009). It is here however considered for several pollen exhibits an overall spherical shape. Also, the spherical shape is the ideal shape to be detected and can hence be considered as a reference case for polarimetric studies.
- a nonspherical pollen called *Taxon 2* with scattering matrix elements [$f_{22}^\lambda = 0.47, f_{33}^\lambda = -0.41, f_{44}^\lambda = -0.30, f_{12}^\lambda = 0.02, f_{34}^\lambda = 0.06$]. As to be seen in Section 4, dedicated to our laboratory results, these scattering matrix elements are that of ragweed pollen at wavelength λ_{IR} .

- another nonspherical pollen called *Taxon 3*, exhibiting close but different scattering matrix elements compared with *Taxon 2*, namely $[f_{22}^\lambda = 0.38, f_{33}^\lambda = -0.38, f_{44}^\lambda = -0.17, f_{12}^\lambda = 0.02, f_{34}^\lambda = 0.01]$. As to be seen in Section 4, dedicated to our laboratory results, these scattering matrix elements are that of pine pollen at wavelength λ_{IR} .

To fix ideas, we plotted in Fig. 3 the variation of I_i^λ as a function of the ψ -angle of the analyser at a given wavelength λ , for *Taxon 1, 2, 3* at the three successive incident polarization states (p), (45°), (RC). Interestingly, each pollen *Taxon* is associated with a specific light scattering curve, showing the sensitivity of our methodology. From *Taxon 1* to 3, the (p) -polarization curve $I_p^\lambda(\psi)$ exhibits varying minima to be related to variations in the f_{22}^λ -scattering element. The same conclusions can be drawn with the $I_{RC}^\lambda(\psi)$ curve minima, which are determined by the f_{44}^λ scattering matrix element. If the $I_{45^\circ}^\lambda(\psi)$ curve is π -periodic whatever the considered *taxon*, its extrema, which depend on $f_{12}^\lambda, f_{34}^\lambda$ and f_{33}^λ , are also *Taxon*-dependent. As a result, the proposed methodology can potentially be used to differentiate pollen exhibiting different scattering matrix elements. For that however, the scattering matrix elements should be retrieved with precision, as discussed below.




Figure 3: Numerical simulation of the detected scattered light intensity $I_i^\lambda(\psi)$ as a function of the ψ -angle of the analyser for the three considered case studies of pollen taxa : **Taxon 1** (dotted lines), **Taxon 2** (solid lines), **Taxon 3** (dashed lines) at the three successive incident polarization states ($i = (p), (45^\circ), (RC)$) from left to right. Each curve is normalized to unity (i.e. $I_0^\lambda \times a_i^\lambda = 1$) so that any change in the scattered light intensity be due to polarimetric considerations.

3.4 Scattering matrix elements retrieval accuracy

Taxon 2 can then be distinguished from *Taxon 3* using polarimetric light scattering if their corresponding Fig. 3 curves can be distinguished within our experimental error bars. To give an example of our ability to distinguish two pollen taxa, to a 1 %-variation in the f_{44}^λ scattering matrix element corresponds a 1 %-variation in the detected scattered light intensity. For *Taxon 2* and *3* exhibit f_{44}^λ -values differing by almost 50 %, the required precision can be reached. Moreover, in our experiment, the evaluation of the $a_i^\lambda, b_i^\lambda, c_i^\lambda$ and d_i^λ coefficients, and hence of the scattering matrix elements (see Eq. 2), relies on a large number

of measurements data points (180 points per fitted curve). Special care has indeed been taken to specify our experimental error bars on the retrieved scattering matrix elements by considering both statistical and systematic errors. Polarization cross-talks in the analyser have been minimized by aligning a second polarizing beam-splitter cube in the detector. Wavelength cross-talks are also fully negligible, which is key for revealing the spectral dependence of the scattering matrix elements, for laser-line selective interference filters are used in the light detector. Also, the incident polarization state may slightly differ from the (p) , $(45 +)$, (RC) polarization states. To quantify this uncertainty, we considered the following incident Stokes vector $S_0 = [1, 2\chi, 2\omega]$, where 2χ and 2ω represent deviations from the (p) incident polarization state (a similar discussion is drawn for $(45 +)$ and (RC) polarization states). At first order in 2χ and 2ω , the uncertainties on the scattering matrix elements $[f_{33}^\lambda, f_{44}^\lambda, f_{12}^\lambda, f_{34}^\lambda]$ are $[2\omega f_{34}^\lambda, 2\chi f_{34}^\lambda, 2\omega f_{22}^\lambda, 2\chi f_{33}^\lambda]$ while the matrix element f_{22}^λ remains unperturbed. As f_{ij}^λ elements at most equal unity, the uncertainties can be majored by their upper limit 2χ or 2ω , depending on the considered scattering matrix element. From a practical point of view, 2χ is evaluated by recording the scattered light intensity by complementary incident polarization states $(45 +)$ and $(45 -)$. Our calculations indeed show that $2\chi = a_{45+}/(a_{45+} + c_{45+}) - a_{45-}/(a_{45-} + c_{45-})$. Besides, 2ω is evaluated by recording the scattering curve corresponding to incident polarization states (RC) and (LC) since $2\omega = a_{RC}/(a_{RC} + c_{RC}) - a_{LC}/(a_{LC} + c_{LC})$. Finally, we used the outputs of a paper by Mishchenko (Mishchenko et al., 2007) to check that the single scattering approximation was rather safe in our experiment where $k_1 \langle d \rangle \gg 30$ (k_1 is the wave vector in the surrounding medium and $\langle d \rangle$ is the average inter-pollen distance), while particle volume concentration remained lower than 1 %, hence considering a tenuous media. Statistical errors due to potential fluctuations in the grain number concentration in the scattering volume are also accounted for by normalizing the detected intensity by that of a photodetector placed at 170° scattering angle, which depends on the pollen grains number concentration. Indeed, the scattered light intensity at 170° scattering angle is proportional to the pollen grain concentration as this detector is polarization insensitive. As a result, statistical errors due to potential fluctuations in the pollen grains number concentration are removed by considering the ratio of the two intensities at these two scattering angles. As a conclusion, from the measurements of the scattered intensity at wavelength λ_{VIS} and λ_{IR} , the above detailed method allows to precisely retrieve the scattering matrix with error bars that are sufficiently low to distinguish pollen Taxa through polarimetric light scattering, as detailed in the Section 4 below, dedicated to our laboratory results.

4 Laboratory results

4.1 Detected scattered light intensity by ragweed, ash, birch and pine

Figure 4 displays the detected scattered light intensity by our four pollen samples (ragweed, ash, birch, pine) at wavelength λ_{VIS} (in green) and λ_{IR} (in red) and at the three incident polarization states (p) , $(45 +)$, (RC) for a complete rotation of the analyser labelled by the angle ψ . The uncertainty on the incident polarization state has been evaluated by applying the


methodology presented in Section 3.4, thus comparing (45 +) and (RC) polarization state to their complementary states (45 –) and (LC): at both wavelengths, both 2χ and 2ω at most equals 0.01. To minimize statistical errors, each data point results from an average of four measurements each composed of a sequence of 100 laser shots. In Fig. 4, the error bar affecting each data point is then the standard deviation of these four sequences. Moreover, the reproducibility of the observed minima during a full rotation of the analyser indicates that the shape of each considered pollen remained constant during the acquisition. Let us first discuss on the scattered light intensity by ragweed pollen. The (*p*)-polarization curve exhibits non-zero minima, showing that ragweed pollen grains are non-spherical, in agreement with the SEM observation. In the (45 +)-polarization curve, the two successive local minima are not equal at wavelength λ_{IR} (see for example, the first and second minima of ragweed pollen at wavelength λ_{IR}), which proves that $f_{34}^{\lambda_{IR}}$ is not null, as opposed to wavelength λ_{VIS} where, within our error bars, no significant differences occur in the successive local minima for ragweed for example. Moreover, the light scattering curves differ for the three other pollen (ash, birch, pine). For instance, pine pollen, whose successive local minima are identical in the (45 +)-polarization curve, exhibits a near zero f_{34}^{λ} -value at both wavelengths as detailed below in Section 4.2. Likewise, the (RC) polarization curve of the ash pollen is nearly constant, which relates to a f_{44}^{λ} -value of the matrix element close to zero.

265 4.2 Retrieved scattering matrix elements for ragweed, ash, birch and pine


The Fig. 4 experimental data points are adjusted with Eq. (2) to retrieve the coefficients a_i^{λ} to d_i^{λ} (see Fig. 4 fitted curves). The agreement between our measurement data points and the Eq. (2) adjustment is notable. Therefore, we can apply the scattering matrix formalism to evaluate f_{ij}^{λ} at each wavelength by applying Eqs. (3), as presented in Fig. 5 with detailed numerical values given in Table 1 (wavelength λ_{VIS}) and in Table 2 (wavelength λ_{IR}). We first focus on the comparison of retrieved $f_{ij}^{\lambda_{VIS}}$ between the four pollen before discussing on the spectroscopic variations of these matrix elements. For our four pollen are nonspherical, all the retrieved $f_{22}^{\lambda_{VIS}}$ -values differ from unity. More interestingly, within our experimental error bars, each pollen exhibits a different $f_{22}^{\lambda_{VIS}}$ matrix element. Likewise, within our error bars, other diagonal matrix elements $f_{33}^{\lambda_{VIS}}$ and $f_{44}^{\lambda_{VIS}}$, which are also not null and differ from one to another, specifically describe light scattering by each pollen. The $f_{44}^{\lambda_{VIS}}$ element is clearly higher for ragweed and birch, compared with ash and pine. Hence, among our set of four pollen, a set of $f_{ij}^{\lambda_{VIS}}$ elements can be unequivocally linked to a single pollen taxon. Off-diagonal elements $f_{12}^{\lambda_{VIS}}$ and $f_{34}^{\lambda_{VIS}}$ are nearly equal to zero within our error bars. Regarding now wavelength λ_{IR} , the same conclusions can be drawn with $f_{22}^{\lambda_{IR}}$, $f_{33}^{\lambda_{IR}}$ and $f_{44}^{\lambda_{IR}}$, i.e. this set of matrix elements unequivocally relates to a single pollen taxon. Interestingly, while $f_{12}^{\lambda_{IR}}$ remains near-zero, $f_{34}^{\lambda_{IR}}$ is not null for ragweed, ash and birch pollen. As underlined by (Bickel and Stafford, 1980), this matrix element may be relevant for the study of pollen particles. The diagonal elements are generally larger at wavelength λ_{IR} , which indirectly means easier to evaluate with a higher precision. The considered pollen exhibit a rather pronounced spectral variation except birch for which the matrix elements remain constant within error bars. The spectral dependence however remains complex to interpret, as the scattering matrix elements of pollen depend on the complex refractive index of pollen grains and the literature on pollen

refractive indices is rather sparse, with, to our knowledge, no evaluation of their imaginary part and no spectral measurement at wavelengths λ_{VIS} and λ_{IR} . The fact that the pine pollen is not symmetrical along its axis and has a distinct size, texture and shape is indeed responsible for the observed differences in the retrieved scattering matrix elements for pine compared with other taxa. Our spectral and polarimetric light scattering methodology is indeed sensitive to the size and to the shape of each pollen taxon and the achieved precision in the scattering matrix retrieval allows accounting for these specific size and shape features, which allow identifying each pollen separately.

290



295 Figure 4: Detected scattered light intensity by ragweed, ash, birch and pine pollen as a function of the of the ψ -angle of the analyser for the three successive incident polarization states (p), (45+) and (RC) at wavelength λ_{VIS} (in green) and wavelength λ_{IR} (in red). The detected light intensity is normalized so that $\mathbf{a}_i \times \mathbf{I}_0^\lambda = \mathbf{1}$, as detailed Section 3.3. A break has been inserted to ease the reading. Each measurement data point is obtained by repeating 400 lasers shots and the uncertainty is calculated by following the methodology described in Section 3.4.



300 Figure 5: Retrieved scattering matrix elements of ragweed, ash, birch and pine at 177.5° scattering angle at wavelengths λ_{VIS} and λ_{IR} , with corresponding uncertainties evaluated by applying the methodology presented in Section 3. For each pollen taxon, the evaluation of the scattering matrix has been repeated with different samples: each evaluated scattering matrix lies within presented error bars.

Table 1 : Retrieved scattering matrix elements of ragweed, ash, birch and pine at 177.5° scattering angle at wavelength λ_{VIS} .

Pollen	$f_{22}^{\lambda_{VIS}}$	$f_{33}^{\lambda_{VIS}}$	$f_{44}^{\lambda_{VIS}}$	$f_{12}^{\lambda_{IR}}$	$f_{34}^{\lambda_{IR}}$
Ragweed	0.38 ± 0.03	-0.33 ± 0.03	-0.18 ± 0.01	0.00 ± 0.02	-0.01 ± 0.02
Ash	0.30 ± 0.02	-0.29 ± 0.01	0.02 ± 0.02	0.00 ± 0.02	-0.01 ± 0.01
Birch	0.44 ± 0.03	-0.41 ± 0.04	-0.20 ± 0.02	0.00 ± 0.03	-0.02 ± 0.03
Pine	0.24 ± 0.02	-0.23 ± 0.02	-0.04 ± 0.01	0.01 ± 0.02	-0.01 ± 0.02

Table 2: Retrieved scattering matrix elements of ragweed, ash, birch and pine at 177.5° scattering angle at wavelength λ_{IR} .

Pollen	$f_{22}^{\lambda_{IR}}$	$f_{33}^{\lambda_{IR}}$	$f_{44}^{\lambda_{IR}}$	$f_{12}^{\lambda_{IR}}$	$f_{34}^{\lambda_{IR}}$
Ragweed	0.47 ± 0.01	-0.41 ± 0.01	-0.30 ± 0.01	0.02 ± 0.01	0.06 ± 0.01
Ash	0.43 ± 0.01	-0.33 ± 0.02	-0.05 ± 0.01	0.02 ± 0.02	0.06 ± 0.01
Birch	0.48 ± 0.03	-0.42 ± 0.05	-0.20 ± 0.02	0.01 ± 0.03	0.08 ± 0.03
Pine	0.38 ± 0.01	-0.38 ± 0.01	-0.17 ± 0.01	0.02 ± 0.01	0.01 ± 0.01

4.3 Towards pollen identification using principal component analysis

As above explained, each of the four considered pollen exhibits its own set of ten scattering elements (five per wavelength), which allow identifying them among the considered set of four pollen. Pine can however be unequivocally identified using $f_{34}^{\lambda IR}$ only as being the only taxon for which $f_{34}^{\lambda IR}$ is null. Also, $f_{22}^{\lambda IR}$ may allow distinguishing pine from other taxa, however not ragweed from birch. Likewise, within our experimental uncertainties, $f_{12}^{\lambda VIS}$ and $f_{34}^{\lambda VIS}$ do not allow differentiating among the four considered pollen. Consequently, for pollen identification through polarized light scattering, the complete set of scattering matrix elements must be generally taken into account, giving rise to the forty elements given in Tables 1 and 2. It however remains challenging and at least non intuitive to identify a pollen taxon based on these set of intricate forty scattering matrix elements. In the literature, decision trees have been applied for classification but overlaps exist due to experimental uncertainties, which limit the application of this classification methodology. Rather, we here propose to reduce the dimensionality of our system to a 2D-representation plane while taking into account experimental uncertainties. Therefore, the goal of this paragraph is to provide a simpler light scattering criterion for classifying among the four considered pollen, based on a 2D-representation plane. To reduce the dimension of our four pollen dataset, a principal component analysis (*PCA*) has been performed. A Linear Discriminant Analysis (*LDA*) may appear more suited for labelled classes. However, as published by (Martinez and Kak, 2001), *PCA* might outperform *LDA* when the number of samples per class is small, and in our methodology, each class (pollen taxon) is represented by a single point as being representative of the distinct size and shape of each pollen taxon. Applying a *LDA* would require having more than one single point per class. *PCA* is a statistical method for dimension reduction and feature extraction (Jolliffe, 2005). There, each pollen is described by its two principal components, hereafter called PC_1 and PC_2 , which form an orthogonal basis of two uncorrelated components. Each measured scattering pattern can then be defined in terms of a combination of the *PC*s. Interestingly, besides the advantage of a dimensionality reduction, the *PCA* maximizes the variance in the (PC_1, PC_2) representation of the dataset. For that, we considered the explained variance, which measures the proportion to which a mathematical model accounts for the dispersion of a given data set. In our case, a threshold of 99 % explained variance is reached by taking into account only two principal components. As a result, Fig. 6 presents the projection of each pollen taxon in the new (PC_1, PC_2) -orthogonal basis. The newly obtained coordinates PC_1 and PC_2 of each pollen account for our ten retrieved scattering matrix elements (five per wavelength per pollen) and the corresponding error bars in Fig. 6 have been evaluated by propagating uncertainties on each f_{ij}^{λ} . The explained variance of PC_1 is 78.8 % while that of PC_2 reaches 20.3 % so that the dimension reduction still accounts for 99.1 % of our experimental variance. Based on Fig. 6, a classification criterion among our four pollen dataset is to assign to each pollen taxon the area delimited by its error bar. To each measured scattering pattern (i.e. each set of ten evaluated f_{ij}^{λ} matrix elements) then corresponds a single point in the (PC_1, PC_2) -basis, which allows identifying the pollen if the evaluated point lies in one of the Fig. 6 coloured areas, which provides a simple criterion for classifying among the four considered pollen, based on a 2D-

representation plane. When considering each wavelength ($\lambda_{VIS}, \lambda_{IR}$) separately, the PCA still allows identifying a simple light-scattering criterion to differentiate each taxon, with a precision depending on the achieved accuracy in the retrieved scattering matrix elements at the considered wavelength. This methodology can interestingly be extended to other pollen taxa provided that precise laboratory measurements of its f_{ij}^λ matrix elements are first accurately carried out, as performed in Section 3 for ragweed, ash, birch and pine. Still as is, our methodology allows identifying non equivocal regions in the *PCA* basis, due to the sensitivity and accuracy of our laboratory work. Hence, our evaluation of pollen scattering matrix at two wavelengths, when associated with a principal component analysis, is a step towards identifying non-equivocal polarized light scattering characteristics of pollen, that work for the set of following taxa: ragweed, ash, birch and pine.




Figure 6: Projection of the ten f_{ij}^λ scattering matrix elements (five per wavelength) for each pollen on the new (PC_1, PC_2)-orthogonal basis using principal component analysis. Error bars have been evaluated by propagating the uncertainties on each f_{ij}^λ scattering matrix element. Coloured regions correspond to the range of (PC_1, PC_2) where the pollen taxon can be unequivocally identified among our set of four pollen.

5 Conclusion and outlooks

In a context where pollen are highly impacting atmospheric bioaerosols with a high socio-economical cost, new methodologies are required for monitoring and differentiating pollen. In this paper, the ability of polarized light scattering to differentiate a set of four pollen (ragweed, ash, birch and pine) is experimentally studied in laboratory at two wavelengths. Each pollen taxon having a very specific size and shape, light scattering by these complex-shaped particles is challenging and cannot be easily numerically simulated. In this context, a controlled laboratory experiment, relying on the robust scattering matrix formalism, has been carried out, with special emphasis on the required precision to identify different spectral and polarimetric characteristics of the four considered pollen. This laboratory experiment provides precise retrievals of the scattering matrix

elements of each pollen taxon in the form of ten matrix elements (five per wavelength). Hence, a set of ten retrieved matrix
360 elements has been drawn per pollen. Interestingly, within our experimental uncertainties, the four considered pollen, when
embedded in ambient air, exhibit clearly different light scattering characteristics, which allowed to differentiate between each
pollen taxon from the set of forty matrix elements (five per wavelength per pollen). Finally, to reduce the dimensionality of
our system to a 2D-representation plane while taking into account our experimental uncertainties, a principal component
analysis is here proposed. Though the dimensionality is reduced, this projection of each pollen taxon in the new (PC_1, PC_2) -
365 orthogonal basis interestingly accounts for more than 99 % of our experimental variance, which allow to provide a simple
criterion for classifying among the four considered pollen, based on a 2D-representation plane. There are multiple outlooks to
this work. To remain focused on laboratory work, which is currently in short supply, the evaluation of the scattering matrix
for a larger set of pollen taxon should be considered. If the four chosen pollen taxa are among the most impacting pollen, this
work could be extended to other pollen by following our methodology. Any pollen taxon can in principle be studied and
370 identified by applying our new methodology: our experimental error bars are indeed very low, so does the probability for two
different pollen taxa to exhibit the same ten scattering matrix elements (five per wavelength). Laboratory intense work is
however required for extending this work to other species, which is far beyond the scope of this contribution, aimed at
introducing this new measurement technique. We may expect our methodology to be applicable to other sets of pollen taxa,
provided that the same accuracy is experimentally achieved. The laboratory experiment should then be carried out. As well,
375 extension of this work to other wavelengths, to pollen mixtures, or to other scattering angles is also an interesting outlook of
this work.

Author contribution

Danaël Cholleton: Formal analysis, Investigation, Software, Visualization, Writing - original draft, Writing - review & editing.
Emilie Bialic: Writing - original draft, Project administration. **Antoine Dumas:** Writing - original draft, Project
380 administration. **Pascal Kaluzny:** Project administration, Supervision. **Patrick Rairoux:** Conceptualization, Writing - original
draft, Project administration, Supervision. **Alain Miffre:** Conceptualization, Formal analysis, Investigation, Methodology,
Supervision, Writing - original draft, Writing - review & editing

Competing interests

The authors declare that they have no conflict of interest.

385 Acknowledgements

CNRS is acknowledged for financial support and the Lyon Centre Technologique des Microstructures for their help in electronic microscopy.

References

- PalDat: <https://www.paldat.org/>, last access: 21 June 2021.
- 390 Bickel, W. S. and Stafford, M. E.: Biological Particles as Irregularly Shaped Scatterers, in: *Light Scattering by Irregularly Shaped Particles*, Springer, Boston, MA, 299–305, https://doi.org/10.1007/978-1-4684-3704-1_35, 1980.
- Biedermann, T., Winther, L., Till, S. J., Panzner, P., Knulst, A., and Valovirta, E.: Birch pollen allergy in Europe, *Allergy*, 74, 1237–1248, <https://doi.org/10.1111/all.13758>, 2019.
- 395 Bielory, L., Lyons, K., and Goldberg, R.: Climate Change and Allergic Disease, *Curr. Allergy Asthma Rep.*, 12, 485–494, <https://doi.org/10.1007/s11882-012-0314-z>, 2012.
- Bohlmann, S., Filioglou, M., Giannakaki, E., Shang, X., Saarto, A., and Komppula, M.: Characterization of atmospheric pollen with active remote sensing in Finland, in: *Geophysical Research Abstracts*, 1, 2018.
- Cao: Lidar polarization discrimination of bioaerosols, *Opt. Eng.*, 49, 116201, <https://doi.org/10.1117/1.3505877>, 2010.
- 400 Cholleton, D., Bialic, E., Dumas, A., Kaluzny, P., Rairoux, P., and Miffre, A.: Laboratory evaluation of the (VIS, IR) scattering matrix of complex-shaped ragweed pollen particles, *J. Quant. Spectrosc. Radiat. Transf.*, 254, 107223, <https://doi.org/10.1016/j.jqsrt.2020.107223>, 2020.
- Comtois, P., Alcazar, P., and Neron, D.: Pollen counts statistics and its relevance to precision, *Aerobiologia*, 10, 1999.
- Crouzy, B., Stella, M., Konzelmann, T., Calpini, B., and Clot, B.: All-optical automatic pollen identification: Towards an operational system, *Atmos. Environ.*, 140, 202–212, <https://doi.org/10.1016/j.atmosenv.2016.05.062>, 2016.
- 405 Dahl, Å., Strandhede, S.-O., and Wihl, J.-Å.: Ragweed – An allergy risk in Sweden?, *Aerobiologia*, 15, 293–297, <https://doi.org/10.1023/A:1007678107552>, 1999.
- Giri, R., Morello, C., Heinson, Y. W., Kemppinen, O., Videen, G., Videen, G., Videen, G., and Berg, M. J.: Generation of aerosol-particle light-scattering patterns from digital holograms, *Opt. Lett.*, 44, 819–822, <https://doi.org/10.1364/OL.44.000819>, 2019.
- 410 Gómez Martín, J. C., Guirado, D., Frattin, E., Bermudez-Edo, M., Cariñanos Gonzalez, P., Olmo Reyes, F. J., Nousiainen, T., Gutiérrez, P. J., Moreno, F., and Muñoz, O.: On the application of scattering matrix measurements to detection and identification of major types of airborne aerosol particles: Volcanic ash, desert dust and pollen, *J. Quant. Spectrosc. Radiat. Transf.*, 271, 107761, <https://doi.org/10.1016/j.jqsrt.2021.107761>, 2021.
- 415 Hamaoui-Laguel, L., Vautard, R., Liu, L., Solmon, F., Viovy, N., Khvorostyanov, D., Essl, F., Chuine, I., Colette, A., Semenov, M. A., Schaffhauser, A., Storkey, J., Thibaudon, M., and Epstein, M. M.: Effects of climate change and seed dispersal on airborne ragweed pollen loads in Europe, *Nat. Clim. Change*, 5, 766–771, <https://doi.org/10.1038/nclimate2652>, 2015.
- Hesse, M. (Ed.): *Pollen terminology: an illustrated handbook*, Springer, Wien ; New York, 261 pp., 2009.
- Hirst, J. M.: An automatic volumetric spore trap, *Ann. Appl. Biol.*, 39, 257–265, <https://doi.org/10.1111/j.1744-7348.1952.tb00904.x>, 1952.

- Holler, S., Fuerstenau, S. D., and Skelsey, C. R.: Simultaneous two-color, two-dimensional angular optical scattering patterns from airborne particulates: Scattering results and exploratory analysis, *J. Quant. Spectrosc. Radiat. Transf.*, 178, 167–175, <https://doi.org/10.1016/j.jqsrt.2016.01.009>, 2016.
- 420 Imhof, K., Probst, E., Seifert, B., Regenass, S., and Schmid-Grendelmeier, P.: Ash pollen allergy: reliable detection of sensitization on the basis of IgE to Ole e 1, *Allergo J. Int.*, 23, 78–83, <https://doi.org/10.1007/s40629-014-0010-8>, 2014.
- Iwai, T.: Polarization Analysis of Light Scattered by Pollen Grains of *Cryptomeria japonica*, *Jpn. J. Appl. Phys.*, 52, 062404, <https://doi.org/10.7567/JJAP.52.062404>, 2013.
- 425 Jolliffe, I.: Principal Component Analysis, in: *Encyclopedia of Statistics in Behavioral Science*, American Cancer Society, <https://doi.org/10.1002/0470013192.bsa501>, 2005.
- Kemppinen, O., Laning, J. C., Mersmann, R. D., Videen, G., and Berg, M. J.: Imaging atmospheric aerosol particles from a UAV with digital holography, *Sci. Rep.*, 10, 16085, <https://doi.org/10.1038/s41598-020-72411-x>, 2020.
- 430 Kim, G., Lee, S., Shin, S., and Park, Y.: Three-dimensional label-free imaging and analysis of Pinus pollen grains using optical diffraction tomography, *Sci. Rep.*, 8, <https://doi.org/10.1038/s41598-018-20113-w>, 2018.
- Kiselev, D., Bonacina, L., and Wolf, J.-P.: A flash-lamp based device for fluorescence detection and identification of individual pollen grains, *Rev. Sci. Instrum.*, 84, 033302, <https://doi.org/10.1063/1.4793792>, 2013.
- 435 Lake, I. R., Jones, N. R., Agnew, M., Goodess, C. M., Giorgi, F., Hamaoui-Laguel, L., Semenov, M. A., Solomon, F., Storkey, J., Vautard, R., and Epstein, M. M.: Climate Change and Future Pollen Allergy in Europe, *Environ. Health Perspect.*, 125, 385–391, <https://doi.org/10.1289/EHP173>, 2017.
- Li, Y., Steiner, A., and Solmon, F.: Investigating the direct climatic forcing of pollen and subpollen particles, 2013, A11B-0020, 2013.
- Martinez, A. M. and Kak, A. C.: Pca versus Ida, *IEEE Trans. Pattern Anal. Mach. Intell.*, 23, 228–233, 2001.
- Matsuda, S. and Kawashima, S.: Relationship between laser light scattering and physical properties of airborne pollen, *J. Aerosol Sci.*, 124, 122–132, <https://doi.org/10.1016/j.jaerosci.2018.07.009>, 2018.
- 440 Miki, K. and Kawashima, S.: Estimation of pollen counts from light scattering intensity when sampling multiple pollen taxa – establishment of an automated multi-taxa pollen counting estimation system (AME system), *Atmospheric Meas. Tech.*, 14, 685–693, <https://doi.org/10.5194/amt-14-685-2021>, 2021.
- Mishchenko, M. I., Travis, L. D., and Lacis, A. A.: *Scattering, Absorption, and Emission of Light by Small Particles*, Cambridge University Press, 492 pp., 2002.
- 445 Mishchenko, M. I., Liu, L., and Videen, G.: Conditions of applicability of the single-scattering approximation, *Opt. Express*, 15, 7522, <https://doi.org/10.1364/OE.15.007522>, 2007.
- Niederberger, V., Purohit, A., Oster, J. P., Spitzauer, S., Valenta, R., and Pauli, G.: The allergen profile of ash (*Fraxinus excelsior*) pollen: cross-reactivity with allergens from various plant species, *Clin. Exp. Allergy*, 32, 933–941, <https://doi.org/10.1046/j.1365-2222.2002.01369.x>, 2002.
- 450 Nouri, S. A., Gregory, D. A., and Fuller, K.: Development of an angle-scanning spectropolarimeter: Preliminary results, *J. Quant. Spectrosc. Radiat. Transf.*, 206, 342–354, <https://doi.org/10.1016/j.jqsrt.2017.11.024>, 2018.
- Pan, Y.-L., Hill, S. C., Pinnick, R. G., House, J. M., Flagan, R. C., and Chang, R. K.: Dual-excitation-wavelength fluorescence spectra and elastic scattering for differentiation of single airborne pollen and fungal particles, *Atmos. Environ.*, 45, 1555–1563, <https://doi.org/10.1016/j.atmosenv.2010.12.042>, 2011.

- 455 Petrov, D.: Photopolarimetric properties of coronavirus model particles: Spike proteins number influence, *J. Quant. Spectrosc. Radiat. Transf.*, 248, 107005, <https://doi.org/10.1016/j.jqsrt.2020.107005>, 2020.
- Pope, F. D.: Pollen grains are efficient cloud condensation nuclei, *Environ. Res. Lett.*, 5, 044015, <https://doi.org/10.1088/1748-9326/5/4/044015>, 2010.
- 460 Prank, M., Chapman, D. S., Bullock, J. M., Belmonte, J., Berger, U., Dahl, A., Jäger, S., Kovtunen, I., Magyar, D., Niemelä, S., Rantio-Lehtimäki, A., Rodinkova, V., Sauliene, I., Severova, E., Sikoparija, B., and Sofiev, M.: An operational model for forecasting ragweed pollen release and dispersion in Europe, *Agric. For. Meteorol.*, 182–183, 43–53, <https://doi.org/10.1016/j.agrformet.2013.08.003>, 2013.
- Raman, P., Fuller, K. A., and Gregory, D. A.: Polarization signatures of airborne particulates, *Opt. Eng.*, 52, 074106, <https://doi.org/10.1117/1.OE.52.7.074106>, 2013.
- 465 Šaulienė, I., Šukienė, L., Daunys, G., Valiulis, G., Vaitkevičius, L., Matavulj, P., Brdar, S., Panic, M., Sikoparija, B., Clot, B., Crouzy, B., and Sofiev, M.: Automatic pollen recognition with the Rapid-E particle counter: the first-level procedure, experience and next steps, *Atmospheric Meas. Tech. Discuss.*, 1–33, <https://doi.org/10.5194/amt-2018-432>, 2019.
- Sauvageat, E., Zeder, Y., Auderset, K., Calpini, B., Clot, B., Crouzy, B., Konzelmann, T., Lieberherr, G., Tummon, F., and Vasilatou, K.: Real-time pollen monitoring using digital holography, *Atmospheric Meas. Tech.*, 13, 1539–1550, <https://doi.org/10.5194/amt-13-1539-2020>, 2020.
- 470 Schaefer, J., Milling, M., Schuller, B. W., Bauer, B., Brunner, J. O., Traidl-Hoffmann, C., and Damialis, A.: Towards automatic airborne pollen monitoring: From commercial devices to operational by mitigating class-imbalance in a deep learning approach, *Sci. Total Environ.*, 796, 148932, <https://doi.org/10.1016/j.scitotenv.2021.148932>, 2021.
- Schaffner, U., Steinbach, S., Sun, Y., Skjøth, C. A., Weger, L. A. de, Lommen, S. T., Augustinus, B. A., Bonini, M., Karrer, G., Šikoparija, B., Thibaudon, M., and Müller-Schärer, H.: Biological weed control to relieve millions from Ambrosia allergies in Europe, *Nat. Commun.*, 11, 1–7, <https://doi.org/10.1038/s41467-020-15586-1>, 2020.
- 475 Schwendemann, A. B., Wang, G., Mertz, M. L., McWilliams, R. T., Thatcher, S. L., and Osborn, J. M.: Aerodynamics of saccate pollen and its implications for wind pollination, *Am. J. Bot.*, 94, 1371–1381, <https://doi.org/10.3732/ajb.94.8.1371>, 2007.
- Sicard, M., Jorba, O., Ho, J. J., Izquierdo, R., De Linares, C., Alarcón, M., Comerón, A., and Belmonte, J.: Measurement report: Characterization of the vertical distribution of airborne Pinus pollen in the atmosphere with lidar-derived profiles: a modelling case study in the region of Barcelona, NE Spain, *Aerosols/Remote Sensing/Troposphere/Physics (physical properties and processes)*, <https://doi.org/10.5194/acp-2021-235>, 2021.
- 480 Siljamo, P., Sofiev, M., Filatova, E., Grewling, L., Jäger, S., Khoreva, E., Linkosalo, T., Ortega Jimenez, S., Ranta, H., Rantio-Lehtimäki, A., Svetlov, A., Veriankaite, L., Yakovleva, E., and Kukkonen, J.: A numerical model of birch pollen emission and dispersion in the atmosphere. Model evaluation and sensitivity analysis, *Int. J. Biometeorol.*, 57, 125–136, <https://doi.org/10.1007/s00484-012-0539-5>, 2013.
- 485 Smith, M., Cecchi, L., Skjøth, C. A., Karrer, G., and Šikoparija, B.: Common ragweed: A threat to environmental health in Europe, *Environ. Int.*, 61, 115–126, <https://doi.org/10.1016/j.envint.2013.08.005>, 2013.
- Spänkuch, D., Döhler, W., and Güldner, J.: Effect of coarse biogenic aerosol on downwelling infrared flux at the surface, *J. Geophys. Res. Atmospheres*, 105, 17341–17350, <https://doi.org/10.1029/2000JD900173>, 2000.
- 490 Surbek, M., Esen, C., Schweiger, G., and Ostendorf, A.: Pollen characterization and identification by elastically scattered light, *J. Biophotonics*, 4, 49–56, <https://doi.org/10.1002/jbio.200900088>, 2011.
- Zhang, X., Rao, R., Huang, Y., Mao, M., Berg, M. J., and Sun, W.: Black carbon aerosols in urban central China, *J. Quant. Spectrosc. Radiat. Transf.*, 150, 3–11, <https://doi.org/10.1016/j.jqsrt.2014.03.006>, 2015.

Zink, K., Kaufmann, P., Petitpierre, B., Broennimann, O., Guisan, A., Gentilini, E., and Rotach, M. W.: Numerical ragweed pollen forecasts using different source maps: a comparison for France, *Int. J. Biometeorol.*, 61, 23–33, <https://doi.org/10.1007/s00484-016-1188-x>, 2017.

- 495 Ziska, L., Knowlton, K., Rogers, C., Dalan, D., Tierney, N., Elder, M. A., Filley, W., Shropshire, J., Ford, L. B., Hedberg, C., Fleetwood, P., Hovanky, K. T., Kavanaugh, T., Fulford, G., Vrtis, R. F., Patz, J. A., Portnoy, J., Coates, F., Bielory, L., and Frenz, D.: Recent warming by latitude associated with increased length of ragweed pollen season in central North America, *Proc. Natl. Acad. Sci.*, 108, 4248–4251, <https://doi.org/10.1073/pnas.1014107108>, 2011.