

Synergetic use of IASI profile and TROPOMI total column level 2 methane retrieval products

Matthias Schneider¹, Benjamin Ertl^{1,2}, Qiansi Tu^{1,a}, Christopher J. Diekmann^{1,b}, Farahnaz Khosrawi¹, Amelie N. Röhling¹, Frank Hase¹, Darko Dubravica¹, Omaira E. García³, Eliezer Sepúlveda³, Tobias Borsdorff⁴, Jochen Landgraf⁴, Alba Lorente⁴, André Butz⁵, Huilin Chen⁶, Rigel Kivi⁷, Thomas Laemmel^{8,c}, Michel Ramonet⁸, Cyril Crevoisier⁹, Jérôme Pernin⁹, Martin Steinbacher¹⁰, Frank Meinhardt¹¹, Kimberly Strong¹², Debra Wunch¹², Thorsten Warneke¹³, Coleen Roehl¹⁴, Paul O. Wennberg¹⁴, Isamu Morino¹⁵, Laura T. Iraci¹⁶, Kei Shiomi¹⁷, Nicholas M. Deutscher¹⁸, David W. T. Griffith¹⁸, Voltaire A. Velazco^{18,d}, and David F. Pollard¹⁹

¹Institute of Meteorology and Climate Research (IMK-ASF), Karlsruhe Institute of Technology, Karlsruhe, Germany

²Steinbuch Centre for Computing (SCC), Karlsruhe Institute of Technology, Karlsruhe, Germany

³Izaña Atmospheric Research Center, Agencia Estatal de Meteorología (AEMET), Santa Cruz de Tenerife, Spain

⁴Earth Science Group, SRON Netherlands Institute for Space Research, Utrecht, The Netherlands

⁵Institute of Environmental Physics, Heidelberg University, Heidelberg, Germany

⁶Center for Isotope Research, University of Groningen, Groningen, The Netherlands

⁷Space and Earth Observation Centre, Finnish Meteorological Institute, Sodankylä, Finland

⁸Laboratoire des Sciences du Climat et de l'Environnement (LSCE), CEA, 91191 Gif sur Yvette, France

⁹LMD/IPSL, CNRS, Ecole polytechnique, University Paris-Saclay, Palaiseau, France

¹⁰Swiss Federal Laboratories for Materials Science and Technology (EMPA), Dübendorf, Switzerland

¹¹Air Monitoring Network, Federal Environment Agency (UBA), Langen, Germany

¹²Department of Physics, University of Toronto, Toronto, Canada

¹³Institute of Environmental Physics, University of Bremen, Bremen, Germany

¹⁴California Institute of Technology, Pasadena, CA, USA

¹⁵Satellite Observation Center, National Institute for Environmental Studies (NIES), Tsukuba, Ibaraki, Japan

¹⁶Atmospheric Science Branch, NASA Ames Research Center, Moffett Field, CA, USA

¹⁷Japan Aerospace Exploration Agency (JAXA), Tsukuba, Japan

¹⁸Centre for Atmospheric Chemistry, School of Earth, Atmospheric and Life Sciences, Faculty of Science, Medicine and Health, University of Wollongong, Wollongong, Australia

¹⁹National Institute of Water and Atmospheric Research Ltd (NIWA), Lauder, New Zealand

^anow at: School of Mechanical Engineering, Tongji University, Shanghai, China

^bnow at: Telespazio Germany GmbH, Darmstadt, Germany

^cnow at: Laboratory for the Analysis of Radiocarbon with AMS (LARA) Department of Chemistry, Biochemistry and Pharmaceutical Sciences (DCBP) & Oeschger Centre for Climate Change Research (OCCR), University of Bern, Switzerland

^dnow at: Deutscher Wetterdienst (DWD), Albin-Schwaiger-Weg 10, Hohenpeissenberg, Germany

Correspondence: M. Schneider

(matthias.schneider@kit.edu)

Abstract. The thermal infrared nadir spectra of IASI (Infrared Atmospheric Sounding Interferometer) are successfully used for retrievals of different atmospheric trace gas profiles. However, these retrievals offer generally reduced information about the lowermost tropospheric layer due to the lack of thermal contrast close to the surface. Spectra of scattered solar radiation observed in the near and/or short wave infrared, for instance by TROPOMI (TROPOspheric Monitoring Instrument) offer higher

5 sensitivity near ground and are used for the retrieval of total column averaged mixing ratios of a variety of atmospheric trace gases. Here we present a method for the synergetic use of IASI profile and TROPOMI total column level 2 retrieval products. Our method uses the output of the individual retrievals and consists of linear algebra a posteriori calculations (i.e. calculation after the individual retrievals). We show that this approach has strong theoretical similarities to applying the spectra of the different sensors together in a single retrieval procedure, but with the substantial advantage of being applicable to data generated
10 with different individual retrieval processors, of being very time efficient, and of directly benefiting from the high quality and most recent improvements of the individual retrieval processors.

We demonstrate the method exemplarily for atmospheric methane (CH_4). We perform a theoretical evaluation and show that the a posteriori combination method yields a total column averaged CH_4 product (XCH_4) that conserves the good sensitivity of the corresponding TROPOMI product while merging it with the high quality upper tropospheric and lower stratospheric
15 (UTLS) CH_4 partial column information of the corresponding IASI product. As consequence, the combined product offers additional sensitivity for the tropospheric CH_4 partial column, which is not provided by the individual TROPOMI nor the individual IASI product. The theoretically predicted synergetic effect is verified by comparisons to CH_4 reference data obtained from collocated XCH_4 measurements at 14 globally distributed TCCON (Total Carbon Column Observing Network) stations, CH_4 profile measurements made by 36 individual AirCore soundings, and tropospheric CH_4 data derived from continuous
20 ground-based in-situ observations made at two nearby Global Atmospheric Watch (GAW) mountain stations. The comparisons clearly demonstrate that the combined product can reliably detect the actual variations of atmospheric XCH_4 , CH_4 in the UTLS, and CH_4 in the troposphere. A similar good reliability for the latter is not achievable by the individual TROPOMI and IASI products.

1 Introduction

25 Measurements from different ground- or satellite-based sensors target at the observations of the same atmospheric parameters (e.g. the same trace gases), but with different characteristics (e.g. sensitivities for different vertical regions). Often the different sensors use different observation geometries (limb scanning, nadir, solar light reflected at the Earth's surface) and/or different spectral regions (e.g. UV/vis, near infrared, thermal infrared, microwave). Dedicated experts and efforts are needed to develop retrieval techniques that are specifically optimized for an individual sensor. An algorithm that uses coincident measurements
30 of all the different sensors for a multispectral approach (synergetic use of level 1 data) for the optimal estimation of the atmospheric state would well exploit the synergies of the different observation geometries and spectral regions and thus allows for detecting the atmospheric state in more detail than achievable by individual optimal estimation retrievals.

There is a variety of studies investigating the multispectral synergism when retrieving atmospheric trace gases from space. Examples of theoretical studies using synthetic thermal infrared and UV spectra for a simulated synergistic retrieval of atmospheric ozone (O_3) are Landgraf and Hasekamp (2007); Worden et al. (2007); Cuesta et al. (2013); Costantino et al. (2017).
35 These studies considered the thermal infrared spectra of TES (Tropospheric Emission Spectrometer) and IASI (as well as its successor IASI - New Generation), and UV spectra of OMI (Ozone Monitoring Instrument) and GOME-2 (Global Ozone

Monitoring Experiment - 2 as well as its successor UVNS – Ultraviolet Visible Near-infrared Shortwave-infrared) and are complemented by studies with real spectra (e.g. Cuesta et al., 2013; Fu et al., 2013). Another example of a study with real spectra is Luo et al. (2013), who examine the combination of the TES thermal nadir spectra with the MLS (Microwave Limb Sounder) microwave limb spectra for a synergetic retrieval of atmospheric carbon monoxide (CO) profiles. All the different studies clearly show that the synergetic use of the measured spectra results in an increased sensitivity with respect to the targeted trace gases.

However, the development of these multispectral retrievals requires experts in different remote sensing techniques to work closely together. Furthermore, as soon as measurements from a new sensor become available (or as soon as sensors are modified/improved) such multispectral processors have to be adapted accordingly, i.e. continuous collaborative retrieval developments are required. While this is certainly possible, it might be not the most efficient way, in particularly considering the steadily increasing amount of available satellite data products (level 2 retrieval products). The optimal synergetic exploitation of the already available level 2 retrieval products would be much less computationally expensive than running dedicated multispectral retrievals. Such synergetic combination of level 2 products is the topic of this paper.

There are already several examples of a level 2 product fusion discussed in literature (the following list is not intended to be complete): Worden et al. (2015) combine the thermal and near infrared level 2 products of methane (CH_4) of TES and GOSAT (Greenhouse gas mOnitoring SATellite), respectively, by performing approximative calculations and with a focus on monthly mean data. Data aggregation is necessary due to the reduced temporal and horizontal coverage of TES and GOSAT and their imperfect collocation. Cortesi et al. (2016) combine the thermal infrared MIPAS-STR (Michelson Interferometer for Passive Atmospheric Sounding - STRatospheric aircraft) and microwave MARSCHALS (Millimetre-wave Airborne Receivers for Spectroscopic Characterisation in Atmospheric Limb Sounding) aircraft-based remote sensing products of O_3 , nitric acid (HNO_3), water vapour (H_2O), and atmospheric temperature (applying the so-called Measurement-Space-Solution data fusion method of Ceccherini et al., 2009). Another example is Warner et al. (2014), who use a Kalman filter for combining the CO data products of AIRS (Atmospheric Infrared Sounder) – available for a large horizontal area, but with weak vertical details – and TES (and MLS) – available with detailed vertical information, but only for very localised areas.

Here, we present a method for fusing the available level 2 CH_4 profile product of IASI and the XCH_4 (total column averaged methane) product of TROPOMI (Tropospheric Monitoring Instrument) by means of a Kalman filter approach. Our objective is a data product with improved vertical profile information (determine tropospheric CH_4 independently from CH_4 at higher altitudes, which is not possible by IASI or TROPOMI data alone) by synergetically exploiting the different vertical sensitivities of the two products. The method allows a computationally very efficient generation of global daily maps of the combined data product and only needs the individually retrieved states, averaging kernels and noise covariances provided by the respective remote sensing experts in the context of their standard retrieval work. The proposed method can be used flexibly for combining measurement information of different satellite sensors and is in particularly interesting for combining a profile product with a total column product. The method has strong theoretical similarities to a dedicated combined optimal estimation retrieval that uses the combined IASI and TROPOMI spectra as input (synergetic use of level 1 data).

The reliable and global detection of tropospheric CH_4 independently from CH_4 at higher altitudes can lead to an improved understanding of the CH_4 cycle. Respective data allow a more direct investigation of the CH_4 boundary layer source and sink signals than total column averaged mixing ratios (XCH_4) provided globally for instance by GOSAT (e.g. Parker et al., 2020) or TROPOMI (Lorente et al., 2021). This is because XCH_4 signals are strongly affected by vertical shifts of the tropopause altitude, i.e. their use for investigating CH_4 absorption and release at ground depends on the correct consideration of the tropopause altitude by model simulations (Pandey et al., 2016).

This manuscript is organised as follows. Section 2 briefly presents the used IASI and TROPOMI products (generated by two individual retrievals). Section 3 presents the equations for the optimal a posteriori combination of the two independent retrieval outputs (level 2 product combination), and performs a theoretical evaluation of the individual and combined products. Section 4 validates the total column and tropospheric and UTLS (upper tropospheric/lower stratospheric) partial column products obtained by the individual IASI and TROPOMI retrievals and by the a posteriori combination by an inter-comparison to reference data from TCCON, AirCore, and GAW. Section 5 discusses the global consistency of the products and shows global maps. Section 6 resumes the results of our study and briefly discusses upcoming possibilities. Furthermore, in Appendix A we give a brief overview on retrieval theory and in Appendix B we discuss the theory of our a posteriori combination method and show that the method has strong similarities to performing a full multispectral optimal estimation retrieval. Appendix C introduces the operator for transferring logarithmic scale differentials into linear scale differentials. Appendix D presents the operators used for converting vertical profile data into total and partial column data. Appendix E examines the dislocation error, i.e. to what extent the temporal and spatial dislocation of the IASI and TROPOMI observations (the two sensors are on two different satellites having different orbits) impacts the combined data product. Appendix F explains how we assess the comparability of the satellite products with the reference data and reveals the reasonable agreement between the characteristics of the satellite products and the results of the validation study.

2 Satellite data

In this section we briefly present the satellite data products that are used for the combination procedure. These are the XCH_4 data obtained from the analysis of the near and short wave infrared (SWIR and NIR) spectra measured by TROPOMI and the CH_4 profiles derived from IASI thermal nadir (TIR) spectra. In addition, we explain the criteria used for collocating the two satellite observations.

2.1 RemoTeC TROPOMI XCH_4

The TROPOMI XCH_4 data used in this study are generated by the RemoTeC algorithm (Butz et al., 2011), which is used for the operational processing of Sentinel 5 Precursor/TROPOMI XCH_4 data (Hu et al., 2016; Hasekamp et al., 2019; Landgraf et al., 2019). Here we work with data of the operational processing algorithm version 2.2.0 (which has been presented and validated in Lorente et al., 2021). The TROPOMI output files provide the XCH_4 data together with the used a priori data (constructed from simulations of the global chemistry-transport model TM5, Krol et al., 2005), the column averaging kernels,

and the error values. Here we work with all TROPOMI data that pass the standard quality filtering (TROPOMI output variable
105 `qa_value` must be equal to 1.0, which means a filtering according to Table A1 of Lorente et al., 2021). In addition, we
remove observations over ground covered by snow (which show a high bias as discussed in Lorente et al., 2021) by requiring
a blended albedo (A_b , calculated from the NIR, A_{NIR} , and SWIR, A_{SWIR} , albedos according to Wunch et al., 2011b, as
 $A_b = 2.4A_{\text{NIR}} - 1.13A_{\text{SWIR}}$) of smaller than 0.85.

2.2 MUSICA IASI CH₄ profiles

110 As IASI CH₄ data product we use the data generated by the retrieval processor MUSICA (MUlti-platform remote Sensing of
Isotopologues for investigating the Cycle of Atmospheric water, a European Research Council project between 2011 and 2016).
The MUSICA IASI data full retrieval product encompasses trace gas profiles of H₂O, the HDO/H₂O ratio, N₂O, CH₄, and
HNO₃. The data have been validated in several previous studies (Schneider et al., 2016; Borger et al., 2018; García et al., 2018),
and it has been shown that the CH₄ product can detect the CH₄ signals originating in the upper troposphere/lower stratosphere
115 particularly well. MUSICA IASI data using processor versions 3.2.1 and 3.3.0/1 are currently available for the 2014 to 2021
period and are presented in Schneider et al. (2022). This MUSICA IASI data set is best suited for a posteriori data reuse (e.g.
Diekmann et al., 2021), because in addition to the retrieved trace gas profiles it contains full information on retrieval settings (a
priori states and constraints) and on averaging kernel and error covariance matrices. In order to ensure highest MUSICA IASI
data quality, here we require the flag variable `musica_fit_quality_flag` to be set to 3 (the spectral fit of the MUSICA
120 IASI retrieval has a good quality and the spectral residuals are close to the instrumental noise level). Furthermore, we only use
MUSICA IASI data for which the flag variable `eumetsat_cloud_summary_flag` is set to 1, which guarantees that the
IASI instrumental field of view is cloud-free.

A particularity of the MUSICA IASI processor is that the trace gas inversions are performed on a logarithmic scale. In
Appendix B of Schneider et al. (2022) it is shown that the MUSICA IASI retrieval can be considered as a moderately non-
125 linear problem, in particular if the differentials (averaging kernels and covariances) are used on the logarithmic scale. In the
following equations we take special care on the correct usage of the corresponding logarithmic scale differentials. Nevertheless,
all equations are also applicable for retrievals made on linear scale by replacing in the following the operator \mathbf{L} (which is
introduced in Appendix C) by the identity operator.

2.3 Collocation of TROPOMI and IASI observations

130 As temporal collocation criterion we use 6 hours, for a valid horizontal collocation the centres of the TROPOMI and IASI
ground pixels must be closer than 50 km, and the difference between the ground pressure at the TROPOMI and IASI ground
pixels must be within 50 hPa. Generally multiple TROPOMI/IASI ground pixel pairs fulfill the aforementioned criteria. In
such case we use the pair with the smallest distance metric. This metric is defined as the Euclidean distance that considers a
norm of 12 hours for the temporal distance, a norm of 50 km for the horizontal distance, and a norm of 5 hPa for the vertical
135 distance. The possible small difference in the TROPOMI and IASI ground pixel pressures is taken into account by correcting
the TROPOMI XCH₄ values according to Appendix B of Sha et al. (2021).

3 Presentation and characterisation of the combination method

3.1 Calculation of the combined state vector

For this study we use the CH₄ a priori profile as provided by the TROPOMI product as the common a priori for all products (these are simulations of the global chemistry-transport model TM5, Krol et al., 2005). For this purpose we modify the MUSICA IASI product and bring it in line with the TROPOMI a priori profile choice by applying Eq. (B13).

For updating the IASI CH₄ profile product using the TROPOMI XCH₄ observation we apply a Kalman filter and obtain the combined CH₄ state as:

$$\hat{\mathbf{x}}_C^l = \hat{\mathbf{x}}_I^l + \mathbf{L}^{-1} \mathbf{m} [\hat{\mathbf{x}}_T^* - \mathbf{a}_T^{*T} \hat{\mathbf{x}}_I] - \mathbf{L}^{-1} \mathbf{m} (\mathbf{w}^{*T} - \mathbf{a}_T^{*T}) \mathbf{x}_a. \quad (1)$$

Here the vector $\hat{\mathbf{x}}_I$ and scalar $\hat{\mathbf{x}}_T^*$ are the MUSICA IASI CH₄ profile and the TROPOMI XCH₄ column averaged products. The row vector \mathbf{a}_T^{*T} is the total column averaged mixing ratio kernel of the TROPOMI product interpolated to the vertical grid used by the MUSICA IASI processor and the row vector \mathbf{w}^{*T} is the operator for converting mixing ratio vertical profiles into total column averaged mixing ratios (for details on the interpolation see Appendix D). The state vector $\hat{\mathbf{x}}_C^l$ represents the logarithmic scale combined CH₄ profile product (i.e. the MUSICA IASI CH₄ data updated with the TROPOMI XCH₄ observation). The superscript ‘*l*’ used with $\hat{\mathbf{x}}_C^l$ and $\hat{\mathbf{x}}_I^l$ indicates the use of the logarithmic scale. Here and in the following we will mark scalars, vectors or matrix operators that are in logarithmic scale by the superscript ‘*l*’. The matrix \mathbf{L} is the operator for the transformation of differentials or small changes (as given by averaging kernels or error covariances) from the logarithmic to the linear scale (for more details see Appendix C).

The column vector \mathbf{m} is the Kalman gain operator and it is given by:

$$\mathbf{m} = \mathbf{L} \mathbf{S}_{\hat{\mathbf{x}}_I}^l \mathbf{L}^T \mathbf{a}_T^* (\mathbf{a}_T^{*T} \mathbf{L} \mathbf{S}_{\hat{\mathbf{x}}_I}^l \mathbf{L}^T \mathbf{a}_T^* + S_{\hat{\mathbf{x}}_T, n}^*)^{-1}, \quad (2)$$

with the matrix $\mathbf{S}_{\hat{\mathbf{x}}_I}^l$ and the scalar $S_{\hat{\mathbf{x}}_T, n}^*$ being the logarithmic scale a posteriori covariance of the MUSICA IASI CH₄ product and the noise error variance of the TROPOMI XCH₄ product, respectively. The vector operator \mathbf{a}_T^* is the transpose of the TROPOMI column averaging kernel, i.e. $\mathbf{a}_T^* = (\mathbf{a}_T^{*T})^T$.

Except for the logarithmic scale transformation, the Eqs. (1) and (2) are analogous to Eqs. (B9) and (B10). As demonstrated in Appendix B this kind of Kalman filter application has a strong similarity to an optimal estimation retrieval that uses a combined IASI and TROPOMI measurement vector (synergetic use of level 1 data). The application of this Kalman filter is possible because the MUSICA IASI data are provided with full information on a priori states, constraints, error covariances, and averaging kernels (Schneider et al., 2022), and because the TROPOMI data are provided together with their a priori state, averaging kernel, and retrieval noise error (Lorente et al., 2021).

The Kalman Gain according to Eq. (2) describes how differences between the MUSICA IASI and TROPOMI XCH₄ product are used to update the MUSICA IASI CH₄ profile. An example for a Kalman Gain operator is depicted in Fig. 1. It shows that a positive difference of +1 ppb of $[\hat{\mathbf{x}}_T^* - \mathbf{a}_T^{*T} \hat{\mathbf{x}}_I]$ will lead to a combined CH₄ profile product that has been modified with respect to the MUSICA IASI CH₄ product by almost +3 ppb in the lowermost troposphere, by about −0.5 ppb at 10 km, and by about +1.5 ppb above 20 km.




Figure 1. Visualisation of a Kalman Gain operator for optimally combining TROPOMI XCH₄ data with MUSICA IASI CH₄ profile data. This is the column vector \mathbf{m} according to Eq. (2). The example shown is for a late summer atmosphere (27 September 2018) over Central Europe.

170 3.2 Vertical resolution and representativeness

In this section we compare the vertical resolution and representativeness of the individual retrieval products with those achieved when combining the two retrieval products. According to Eq. 1 the averaging kernels for the combined data product can be calculated as:

$$\mathbf{A}_C^1 = \mathbf{A}_I^1 + \mathbf{L}^{-1} \mathbf{m} (\mathbf{a}_T^{*T} - \mathbf{a}_T^{*T} \mathbf{L} \mathbf{A}_I^1 \mathbf{L}^{-1}) \mathbf{L}. \quad (3)$$

175 Here \mathbf{A}_I^1 and \mathbf{A}_C^1 are the logarithmic scale averaging kernels of the MUSICA IASI CH₄ product and of the combined product (the MUSICA IASI CH₄ product after being updated with the information provided by the TROPOMI XCH₄ product), respectively. These are the kernels for the profile products represented in nal (nal : number of atmospheric levels) levels, i.e. they are matrices of dimension $nal \times nal$. Logarithmic scale kernels are also called fractional or relative averaging kernels (e.g. Keppens et al., 2015).

180 Figure 2 depicts the rows of typical averaging kernels for the MUSICA IASI product (Fig. 2a) and the combined data product (Fig. 2b). Adding the information provided by TROPOMI clearly improves the sensitivity in the lower troposphere: for the MUSICA IASI product the lower tropospheric kernels generally peak at the upper limit of the lower troposphere (at about 5 km a.s.l.). For the combined product these peak values are obtained at significantly lower altitudes (at about 2 km a.s.l.). In the UTLS we see no significant difference between the kernels.

185 In this work we focus on the total column and the partial columns between the surface and 6 km a.s.l. (the tropospheric partial column) and between 6 km a.s.l. and 20 km a.s.l. (the UTLS partial column). The total and partial column kernels are calculated from \mathbf{A}_I^1 and \mathbf{A}_C^1 by their transformation on linear scale (see Appendix C) and the vertical resampling as explained in Appendix D. Figure 3 depicts the total and partial column kernels corresponding to the row kernels of Fig. 2.




Figure 2. Logarithmic scale row kernels for (a) the MUSICA IASI and (b) the combined product for the same late summer observations as used in the context of Figs. 1 and 3. The symbols mark the kernel values at the nominal altitude.




Figure 3. Total column amount and partial column amount kernels corresponding to the TROPOMI, MUSICA IASI, and combined product for the same late summer observation as used in Figs. 1 and 2: (a) total column amount kernels; (b) lower tropospheric partial column amount kernels, surface - 6 km a.s.l.; (c) upper tropospheric/lower stratospheric (UTLS) partial column amount kernels, 6 - 20 km a.s.l.

Total column amount kernels are available for all three products (see Fig. 3a): the TROPOMI, the MUSICA IASI, and the
 190 combined product. The TROPOMI kernel is close to unity for all altitudes, documenting the good sensitivity for CH_4 at all altitudes. The combined total column amount kernel is even closer to unity than the respective TROPOMI kernel, which means that the combined retrieval product does also well reflect the actual atmospheric total column amounts. The MUSICA IASI




Figure 4. Time series of the degree of freedom for signal (DOFS, example for Central Europe). Black squares: TROPOMI (please note that only the total column data are made available); Red dots: MUSICA IASI; blue crosses: combined product.

kernel has relatively low values in the lower troposphere and above 15 km, only in the UTLS region the kernel values are between 0.75 and 1.25. This means that MUSICA IASI can actually not well detect total column amounts, because it lacks
 195 sensitivity in the lower troposphere. The altitude regions where the MUSICA IASI product has reduced sensitivities are the regions where TROPOMI's total column information has the strongest impact on the combined product (see Fig. 1).

Partial column amount kernels are only available for profile products, i.e. the MUSICA IASI and the combined product (MUSICA IASI updated with information from TROPOMI). Figure 3b shows tropospheric partial column amount kernels. For the MUSICA IASI product we observe values that are generally lower than 0.5. The highest values are achieved around 6 km
 200 a.s.l., i.e. at the upper boundary of the vertical layer we defined as the tropospheric partial column. The kernel of the combined product shows a good sensitivity with peak values of almost 0.95 at 2.5 km a.s.l. and values above 0.75 for almost all altitudes between the surface and 6 km a.s.l.

UTLS partial column amount kernels are depicted in Fig. 3c. The values are closest to unity for the altitudes that we attributed to the UTLS layer (altitudes between 6 km and 20 km a.s.l.). There is almost no difference between the MUSICA IASI and
 205 the combined kernels, meaning that the information provided by TROPOMI has almost no effect on the UTLS partial column, which is because the MUSICA IASI product is already very sensitive to this altitude region.

The example kernels document that the combined product allows for detecting tropospheric CH_4 largely independent from CH_4 in the UTLS, which is not possible by the IASI product alone. Figure 4 shows a time series of the degree of freedom for signal (DOFS: it is calculated as the trace of the averaging kernels and is a measure for the profiling capability, Rodgers,
 210 2000). It documents that the combination of TROPOMI with IASI improves the profiling capability of IASI rather consistently throughout all seasons. Here we also show the DOFS values of the TROPOMI retrieval, but please note that only the total column data are made available, i.e. there is no profile information in the provided TROPOMI CH_4 data product.

An optimal estimation retrieval updates the a priori knowledge with information provided by a measurement. The a posteriori uncertainty is the uncertainty achieved by optimally combining the a priori knowledge (captured by the inverse of the a priori
 215 covariance matrix, i.e. \mathbf{S}_a^{-1}) with the measurement. As shown in Appendix A the a posteriori uncertainty covariance is the




Figure 5. Time series of the representativeness error (example for Central Europe). Black squares: TROPOMI; red dots: MUSICA IASI; blue crosses: combined product. (a) Total column; (b) tropospheric partial column; (c) UTLS partial column.

sum of the noise covariance and the representativeness error covariance (called "smoothing error" covariance in Rodgers, 2000). According to Eq. (A7) the representativeness error matrix is calculated from the averaging kernel (\mathbf{A}) and the a priori covariance (\mathbf{S}_a) as

$$\mathbf{S}_{\hat{x},r}^1 = (\mathbf{A}^1 - \mathbf{I})\mathbf{S}_a^1(\mathbf{A}^1 - \mathbf{I})^T, \quad (4)$$

220 with \mathbf{I} being the identity operator. By using the kernels \mathbf{A}_T^1 and \mathbf{A}_C^1 we can calculate the representativeness error for the MUSICA IASI and the combined product, respectively. The resampling of $\mathbf{S}_{\hat{x},r}^1$ on total and partial columns is made according to Eq. (D7). For the TROPOMI total column averaged mixing ratios we can calculate the representativeness error by $(\mathbf{w}^{*T} - \mathbf{a}_T^{*T})\mathbf{S}_a(\mathbf{w}^{*T} - \mathbf{a}_T^{*T})^T$. For more details see Appendix D.

Figure 5 depicts the representativeness error relative to the retrieved values for the total column, the tropospheric and UTLS
 225 partial columns. Shown are time series for measurements over Central Europe, which confirm the observations made in the context of the example kernels of Fig. 3: for the total column (Fig. 5a) the representativeness error on the TROPOMI and the combined products are rather small and can be neglected, i.e. both products can detect total column signals. In contrast the MUSICA IASI representativeness error is much larger and the respective data do not well represent the total column, i.e. provide no independent observations of the total column. Concerning partial column products (Fig. 5b and c) we can
 230 compare the MUSICA IASI and the combined product (the TROPOMI product has no information on the vertical distribution). The tropospheric MUSICA IASI partial column has a significant representativeness error (and a seasonal cycle with highest values of about 3% in winter). In the combined product this error is throughout all seasons generally smaller than 1%. In the UTLS both the MUSICA IASI and combined products are well representative for the actual atmospheric methane concentration signals (representativeness error is generally between 0.5 and 1%). In summary, TROPOMI only provides total column data.
 235 With IASI alone we can well detect signals in the UTLS, but not in the lower troposphere. The detection of signals in both altitude regions independently from the a priori information is only possible by using the combined product.




Figure 6. Time series of estimated relative noise error for the retrieved products (example for Central Europe). Colours are as in Fig. 5. (a) Total column; (b) tropospheric partial column; (c) UTLS partial column.

3.3 Retrieval noise error

After documenting the representativeness error in the previous subsection, here we investigate the retrieval noise error. We compare the retrieval noise errors of the individual retrieval products with those achieved when combining the two retrieval products. According to Eq. (1) we can calculate the retrieval noise covariance matrix for the combined data product by

$$\mathbf{S}_{\hat{\mathbf{x}}_{C,n}}^1 = (\mathbf{I} - \mathbf{L}^{-1} \mathbf{m} \mathbf{a}_T^{*T}) \mathbf{L} \mathbf{S}_{\hat{\mathbf{x}}_{I,n}}^1 \mathbf{L}^T (\mathbf{I} - \mathbf{L}^{-1} \mathbf{m} \mathbf{a}_T^{*T})^T + (\mathbf{L}^{-1} \mathbf{m}) S_{\hat{\mathbf{x}}_{T,n}}^* (\mathbf{L}^{-1} \mathbf{m})^T. \quad (5)$$

Here $\mathbf{S}_{\hat{\mathbf{x}}_{I,n}}^1$ is the retrieval noise covariance matrix of the MUSICA IASI retrieval. The error covariances resampled to the total and partial columns are then determined according to Appendix D. Figure 6 shows the retrieval noise errors (which are the square root values of the error variances) relative to the retrieved values for the total column and the tropospheric and UTLS partial columns.

The errors for the total columns (Fig. 6a) are generally below 0.2% for the TROPOMI product. For the MUSICA IASI product they are rather stable at about 0.6%. Concerning the combined product the retrieval noise error is very similar to the retrieval noise error of the TROPOMI data.

For the tropospheric partial columns (Fig. 6b) the error is in general above 1% for the MUSICA IASI product and below 1% for the combined product. For the UTLS partial columns (Fig. 6c) we observe an error of generally below 1% and no significant difference between the MUSICA IASI and the combined data products. This suggests that the error in the combined product is dominated by the error in the MUSICA IASI data, which reveals the very limited impact of the TROPOMI data on the combined UTLS data product.




Figure 7. Time series of estimated relative dislocation error for the combined product due to temporal and spatial dislocation of the TROPOMI and IASI satellite ground pixels (example for Central Europe). (a) Total column; (b) tropospheric partial column; (c) UTLS partial column.

3.4 Dislocation error

255 As mentioned in Sect. 2.3 we allow for small dislocations between the TROPOMI and IASI observations of up to 6 hours and 50 km. As derived in Appendix E the dislocation error covariance matrix is calculated by

$$\mathbf{S}_{\mathbf{x}_{C,d1}}^1 = \mathbf{A}_{C,d1}^1 \mathbf{S}_{\Delta,d1}^1 \mathbf{A}_{C,d1}^{1T}, \quad (6)$$

where $\mathbf{A}_{C,d1}^1$ is the dislocation kernel and $\mathbf{S}_{\Delta,d1}^1$ is the covariance matrix for the CH_4 dislocation uncertainty, whose main characteristics are visualised in Figs. E1 and E2. The low entries of the dislocation kernel at low altitudes (typical example see 260 Fig. E4), reduces the impact of the spatial and temporal dislocation on the total and tropospheric partial columns of combined product.

Over Central Europe we estimate an error in the combined product due to the dislocations between IASI and TROPOMI as shown in Fig. 7. For the total column the error is below 0.1% and for the tropospheric and UTLS partial columns it is generally below 0.8%. If compared to the noise error (see Fig. 6), the dislocation error is of secondary importance. Details on 265 the estimation of these dislocation errors and examples for other locations are given in Appendix E.

4 Validation

In this section we empirically evaluate the quality of the TROPOMI, MUSICA IASI, and combined products by their inter-comparison to different reference data products. As reference for the total column averaged mixing ratio (X_{CH_4}) we use TCCON (Total Carbon Column Observing Network, Wunch et al., 2011a) ground-based remote sensing data from 14 sites 270 located in different climate zones. As reference for the total and the partial columns we use in-situ profiles measured by the AirCore system (Karion et al., 2010) at two geophysically different European locations. Furthermore, we use in-situ data measured at two nearby Central European Global Atmospheric Watch (GAW) mountain stations.




Figure 8. Maps showing the location of the reference measurements used for the validation. (a) Global map with the 14 TCCON stations (more detailed information on these sites is given in Table 1). (b) Map indicating the areas accepted for valid horizontal collocations in the surroundings of the European reference stations: TCCON sites and the 100 km collocation radius (red crosses and circles), AirCore sites and the spatial collocation circles with 500 km radius (blue stars and circles), and GAW sites and the collocation circle with 150 km radius (grey dots and circle).

Figure 8 depicts the geographical location of the reference observations. Figure 8a shows that the considered TCCON stations are distributed around the globe (more detailed information on these sites is given in Table 1). Figure 8b gives details on the different European reference sites and the areas accepted for a valid collocation. For collocation with TCCON the satellite ground pixels should fall within a circle around the stations with a radius of 100 km (red crosses and circles). For the comparison to the GAW data the collocation circle has a radius of 150 km (grey circle) and is centred in the middle of the two GAW stations (Jungfraujoch in Switzerland and Schauinsland in South-western Germany indicated by the grey dots). For the comparison with AirCore we relax the radius of the collocation circle to 500 km in order to achieve a sufficient high number of coincidences between AirCore and satellite observations. The two AirCore sites (Trainou in France and Sodankylä in Finland) and the collocation circles are indicated by the blue stars and circles.

Appendix F reveals that the following validation results are in reasonable agreement with sensitivities and errors of the different satellite data products as shown in Sects. 3.2 - 3.4.

4.1 TCCON XCH₄

We use TCCON ground-based remote sensing data from 14 sites located in different climate zones representative for high, mid and low latitudes. Details on the locations of these sites, the respective data amounts, and references are given in Table 1. We use the TROPOMI a priori setting for the comparison between the ground-based TCCON and the satellite-based remote sensing products. For this purpose the TCCON product is adjusted to the TROPOMI a priori settings according to Eq. (B13), which ensures the usage of the same a priori data for all the remote sensing products. As spatial collocation criteria we require the TROPOMI and IASI ground pixels to be located within 100 km of the TCCON station (where we consider the viewing direction of the TCCON spectrometer by using as location, the TCCON's line of sight {latitude,longitude} at 5 km altitude). Differences in the satellite and TCCON ground pressures are taken into account by correcting the TCCON XCH₄

Table 1. Locations of TCCON sites, amount of the satellite data compared to TCCON, and references. "Number" gives the total number of single satellite footprint collocations and "Days" the number of days with collocations.

Station (ID)	Location	Period	Number	Days	References
Eureka (EUR)	80.1°N; 86.4°W; 610 m a.s.l.	06/2020 - 06/2020	271	5	Strong et al. (2019)
Sodankylä (SOD)	67.4°N; 26.6°E; 190 m a.s.l.	05/2018 - 09/2020	17453	107	Kivi et al. (2014), Kivi and Heikkinen (2016)
East Trout Lake (ETL)	54.4°N; 105.0°W; 500 m a.s.l.	02/2018 - 08/2020	8549	122	Wunch et al. (2018)
Karlsruhe (KAR)	49.1°N; 8.4°E; 120 m a.s.l.	02/2018 - 11/2020	21208	167	Hase et al. (2015)
Orleans (ORL)	48.0°N; 2.1°E; 130 m a.s.l.	02/2018 - 09/2020	16094	128	Warneke et al. (2019)
Park Falls (PAR)	46.0°N; 90.3°W; 440 m a.s.l.	11/2017 - 12/2020	7808	87	Wennberg et al. (2017)
Rikubetsu (RIK)	43.5°N; 143.8°E; 380 m a.s.l.	11/2017 - 09/2019	730	25	Morino et al. (2018)
Lamont (LAM)	36.6°N; 97.5°W; 320 m a.s.l.	12/2017 - 12/2020	21548	171	Wennberg et al. (2016)
Edwards (EDW)	35.0°N; 118.9°W; 700 m a.s.l.	06/2018 - 12/2020	36821	438	Iraci et al. (2016)
Saga (SAG)	33.2°N; 130.3°E; 10 m a.s.l.	11/2017 - 12/2020	916	67	Kawakami et al. (2014)
Burgos (BUR)	18.5°N; 120.7°E; 40 m a.s.l.	11/2018 - 03/2020	37	11	Velazco et al. (2017)
Darwin (DAR)	12.5°S; 130.9°E; 40 m a.s.l.	06/2018 - 04/2020	2841	59	Griffith et al. (2014a)
Wollongong (WOL)	34.4°S; 150.9°E; 30 m a.s.l.	03/2018 - 06/2020	1657	86	Griffith et al. (2014b)
Lauder (LAU)	45.0°S; 169.7°E; 610 m a.s.l.	11/2017 - 12/2020	1437	115	Sherlock et al. (2014), Pollard et al. (2019)

values according to Appendix B of Sha et al. (2021). For collocation with respect to time, we use as TCCON reference the median XCH₄ value calculated from all TCCON data measured within 2 hours of the TROPOMI observation. Furthermore, we require stable conditions for atmospheric CH₄. This is achieved by performing the comparisons only, if there are at least 3 individual TCCON observations that fulfil the collocation criterion and if the timestamps of these observations have a 1 σ standard deviation of 1 hour and the 1 σ standard deviation of the respective XCH₄ data is smaller than 0.5%.

In Fig. 9 the TROPOMI, MUSICA IASI, and combined XCH₄ products are compared to the TCCON XCH₄ data. The crosses represent the daily mean data and the filled symbols in the background show all data corresponding to all individual valid collocations (between all single pixel satellite observations and individual TCCON observations). Figure 9a-c show time series of the differences with respect to the TCCON references. The daily mean data have error bars, which is the 1 σ standard deviation of the data used for calculating the daily mean.

Statistics in form of the median of the difference and the scatter around the median difference are given in each panel (for statistics using daily mean data in black fonts and for statistics using all individual valid collocations in grey fonts). We use here the median in order to be less affected by outliers. For the same reason we use as metric for the scatter the half inter-percentile range between the 15.9 and 84.1 percentiles (hIPR68.2, which is analogous to the 1 σ standard deviation in case of a pure




Figure 9. Comparison of the different XCH₄ satellite products with TCCON XCH₄ data from 14 globally representative stations (the different colours correspond to the stations as shown in Fig.8a). Data for all individual coincidences are plotted in the background as squares and daily mean data are depicted as crosses with error bars representing the 1σ standard deviation (daily means are only calculated if there are at least 3 observations per day): (a)-(c) time series of the differences. (d)-(f) Correlations between TCCON and satellite data (the black line is the one-to-one diagonal). (g)-(i) Correlations between the a priori free TCCON data (Δ_a TCCON = XCH₄(TCCON) – XCH₄(apriori)) and the a priori free satellite data (Δ_a Satellite = XCH₄(Satellite) – XCH₄(apriori)). The inserted text reports median and scatter (hIPR68.2, a-c) and the coefficients of determination, the slope, and the intercept of the robust linear regression model (R^2 , m , and b , d-i). Black and grey fonts represent the values for the daily mean data and for data from all individual collocations, respectively.

Gaussian distribution). Concerning TROPOMI (Fig. 9a) we observe a good agreement. For daily mean data as well as for the statistics based on all individual differences, the median difference is within 0.1% and the scatter lies below 0.7%.

A similar good agreement and low values for median difference and scatter is also achieved for the combined product (Fig. 9c). For the MUSICA IASI product (Fig. 9b) we have reduced sensitivity in the lower troposphere (see Figs. 3 and 5). Because of uncertainties in the a priori assumptions the agreement with the TCCON XCH₄ data is weaker (uncertainties in the a priori assumption are less well detected, see Fig. F2). We observe no significant systematic negative or positive difference for the satellite versus TCCON comparisons, i.e. the satellite data sets seem to be in good absolute agreement with TCCON. In general the observed scatter values are within the range that can be expected from the data uncertainties and the data comparability (for more details see Appendix F).

Figure 9d-f depicts the correlation plots. In order to reduce the effect of outliers, we apply a robust linear regression model (the iteratively reweighted least-squares algorithm with Tukey's bisquare weight function and the respective tuning parameter set to the commonly used value of 4.685). For daily mean data the obtained coefficients of determination (R^2) are larger than 80% for the TROPOMI and the combined product. The slope of the obtained linear regression line is very close to unity. When considering all individual coincidences the R^2 values are about 70%. The error bars on the daily mean data are the 1σ standard deviations of the data used for calculating the daily mean. For the MUSICA IASI product, we observe a similar good correlation as for the TROPOMI and the combined products. However, concerning the MUSICA IASI data, part of the common signal might be due to the a priori on which the MUSICA IASI total column product is not independent (the MUSICA IASI data have a reduced sensitivity, i.e. an increased representativeness error, see Fig. 5a).

Figure 9g-i reveals the information gained by the satellite data with respect to the a priori data. It depicts the correlation between the collocated a priori free TCCON data ($\Delta_a\text{TCCON} = \text{XCH}_4(\text{TCCON}) - \text{XCH}_4(\text{apriori})$) and the a priori free satellite data ($\Delta_a\text{Satellite} = \text{XCH}_4(\text{Satellite}) - \text{XCH}_4(\text{apriori})$). It shows the same data as in 9d-f but with the a priori knowledge removed. We find that the TROPOMI and the combined data product adds a significant amount of information to the a priori knowledge (R^2 values for the respective linear correlations of above 32%). This information gain is much smaller in the case of the MUSICA IASI data, which confirms that the good correlation as observed in Fig. 9e is to a large extent due to the good quality of the a priori data.

4.2 Air-Core in-situ CH₄ profiles

We use the AirCore balloon borne in-situ measurements (Karion et al., 2010) as the reference for CH₄ total columns as well as for the CH₄ vertical distribution. The AirCore system samples the vertical distribution of CH₄ with a much better vertical resolution than the satellite remote sensing systems. For this reason we can generate an AirCore profile (\hat{x}_{AC}) that has the same vertical sensitivity and resolution characteristics as the remote sensing data. According to Eqs. (A2) and (A4) for the MUSICA IASI and the combined retrieval data we can write:

$$\hat{x}_{AC}^l = x_a^l + \mathbf{A}^1(x_{AC}^l - x_a^l). \quad (7)$$

Here \mathbf{A}^1 and x_a^l are the logarithmic scale averaging kernels and the logarithmic scale a priori state of the satellite retrieval, respectively, x_{AC}^l is the measured logarithmic scale AirCore profile regridded to the atmospheric model grid used for the satellite retrievals. The resampling of these data on total and partial columns is made with the linear scale data according to

Table 2. Locations of AirCore sites and amount of satellite data compared to AirCore. "Number" gives the total number of single satellite footprint collocations, "Days" the number of days with collocations, and "AC number" the number of collocated AirCore profiles.

Station	Location	Period	Number	Days	AC number
Sodankylä	67.4°N; 26.6°E	04/2018 - 08/2020	16326	14	14
Trainou	48.0°N; 2.1°E	01/2018 - 11/2020	18458	17	22

Eq. (D5). For the TROPOMI total column averaged mixing ratios we calculate the adjusted AirCore total column averaged mixing ratio (a scalar) by $\hat{x}_{AC}^* = \mathbf{w}^{*T} \mathbf{x}_a + \mathbf{a}_T^{*T} (\mathbf{x}_{AC} - \mathbf{x}_a)$. For more details see Appendix D.

As spatial collocation criteria we require that the ground pixels of the TROPOMI and IASI measurement fall within a circle with a radius of 500 km around the mean horizontal location of the AirCore system when sampling between the 450 and 550 hPa pressure levels. The temporal collocation requirements for both satellite observations is 6 hours. AirCore data are typically not available close to the ground and above the burst altitude of the balloon (approximately 25 hPa). At low altitudes we extend the profile with the concentrations closest to the ground. At high altitudes we extend the profile with the TM5 model data, with a smooth transition between the measured values and the modelled data.

Table 2 gives an overview of the satellite data amount compared to the AirCore profiles measured at Trainou (France, 48.0°N, 2.1°E) and Sodankylä (Finland, 67.4°N, 26.6°E). In total we have 36 individual AirCore profiles measured on 31 different days for which collocated satellite observations exist. The total number of collocated single pixel satellite observations is 34784. We estimate that the AirCore data can serve as reliable references for the validation of the total column as well as for the validation of the tropospheric and UTLS partial columns (see Appendix F).

The comparison between the satellite and the AirCore XCH₄ data is shown in Fig. 10. The differences of collocated measurements are shown in Figs. 10a-c. The agreement between the different satellite products and AirCore is good: the scatter around the median difference is low and similar to the comparison with TCCON. Furthermore, we observe no significant bias in none of the satellite data, which demonstrates the good consistency between the RemoTeC TROPOMI and MUSICA IASI XCH₄ data.

Figure 10d-f depicts the correlation between the satellite and the adjusted AirCore data. Here we apply the same iteratively reweighted least-squares as in Sect. 4.1. The obtained R^2 values are high (for all products above 60% for daily mean data), although a bit lower than the R^2 value achieved for the correlation with TCCON data; however, we have to consider that the amplitude in the analysed total column signals is much smaller in the AirCore data set (data from two northern hemispheric sites only) if compared to the TCCON data set (data from 14 globally distributed sites). As for the comparison to TCCON we examine the correlation between the a priori free reference data ($\Delta_a \text{AirCore} = \text{XCH}_4(\text{AirCore}) - \text{XCH}_4(\text{apriori})$) and the a priori free satellite data ($\Delta_a \text{Satellite} = \text{XCH}_4(\text{Satellite}) - \text{XCH}_4(\text{apriori})$). These correlations are visualised in Fig. 10g-i. We find reasonable correlation for the daily mean TROPOMI and combined data products, but no significant correlation for the daily mean IASI product. This indicates that the correlation as observed between the IASI and the adjusted AirCore data in




Figure 10. Comparison of the different satellite XCH₄ products with adjusted AirCore XCH₄ measured at Trainou (black) and Sodankylä (blue). Data for all individual coincidences are shown in the background as pale crosses and daily averages are depicted as crosses with error bars (daily means are only calculated if there are at least 3 observations per day, which is the case on 29 days of the total 31 days with AirCore observations). (a)-(c) Time series of the differences (error bars represent the daily 1σ standard deviation of the difference). (d)-(f) Correlation between AirCore and satellite data (the black line is the one-to-one diagonal, x-axis error bars represent the mean uncertainty estimated for the AirCore data – according to Eq. (F3) – and y-axis error bars the daily 1σ standard deviation of the satellite data, respectively). (g)-(i) Correlation between the a priori free AirCore and satellite data (error bars as in d-f). The inserted text reports median and scatter (hIPR68.2, a-c) and the coefficients of determination, the slope, and the intercept of the robust linear regression model (R^2 , m , and b , d-i). Black and grey fonts represent the values for the daily mean data and for data from all individual collocations, respectively.

Fig. 10e is mainly due to the a priori data, i.e. IASI adds almost no information with respect to XCH₄ to what is already known by the a priori model. These findings are in line with the vertical resolution and representativeness analyses of Sect. 3.2.




Figure 11. Same as Fig. 10, but for the comparisons of tropospheric partial column averaged CH₄ AirCore and satellite products (IASI and combined).

Figure 11 presents the comparison between the AirCore and satellite tropospheric partial column CH₄ data. The differences between the satellite and the AirCore data are depicted in Fig. 11a and b. If compared to the total column data the agreement worsens a bit (increased median difference and scatter). Nevertheless, the agreement is still good and close to what can be expected from the uncertainty and the comparability of the different data (see Appendix F). Concerning the daily mean data
 375 the combined product has a median difference and hIPR68.2 scatter of below about 0.9%. These values increase to about 1.25% for the IASI product. These results might indicate a weak systematic bias in the MUSICA IASI lower tropospheric partial columns.




Figure 12. Same as Fig. 11, but for comparisons of UTLS partial column averaged CH₄ data.

Figure 11c and d show the correlation plots. In particular for the combined product we observe a reasonable correlation (R^2 of about 26% for daily mean data obtained by using the robust linear regression model). For the IASI product the correlation strength is reduced (R^2 of about 18% for daily mean data). Furthermore, we have to consider that the IASI product has a rather limited tropospheric sensitivity (see Sect. 3.2), which means that a large part of the observed correlation is due to the a priori data: according to Eq. (7) for low entries in \mathbf{A}^1 the variability in the satellite data as well as in the adjusted AirCore data is determined by the variability in the a priori (\mathbf{x}_a^l). This is confirmed by Figs. 11e and f, which show the correlations after removing the a priori data. We observe still a good correlation for the combined product (R^2 of about 44% and regression line slope of 0.93 for daily mean data), but only a weak correlation for the IASI daily mean data (R^2 of about 11%). This clearly

documents the importance of combining IASI and TROPOMI in order to be sensitive to and reliably detect tropospheric CH₄ variations.

Concerning the UTLS partial column we find a very good agreement between the adjusted AirCore data and the IASI and combined satellite data products (see Fig. 12a and b): median difference calculated from the daily mean data are about -0.3% and the scatter values are within about 0.5% . We find no indication of a bias in the satellite data product. The scatter observed between the AirCore and satellite data is even better than what we estimate from the data uncertainty and the data comparability analysis (see Appendix F). Figure 12c and d show that in the UTLS the AirCore and satellite data are strongly correlated (for daily mean data and when using the robust linear regression model, we get R^2 values of up to about 82% and regression line slopes of very close to unity). In this altitude region the MUSICA IASI and the combined products have a very good sensitivity (see Sect. 3.2). This means that the entries in \mathbf{A}^1 of Eq. (7) are large and any deviation between the a priori and the actual CH₄ concentrations in the UTLS are well captured by the adjusted AirCore and satellite data products. Nevertheless, the correlation strength observed for the a priori free data (Fig. 12e and f) is relatively weak (R^2 values of 20% - 23% for daily mean data). This indicates that the a priori model does generally capture well the actual variation of the CH₄ concentration in the UTLS above France and northern Scandinavia.

4.3 GAW surface in-situ CH₄ measurements

At many globally distributed sites atmospheric trace gas in-situ measurements are made continuously within the Global Atmospheric Watch (GAW, <https://community.wmo.int/activity-areas/gaw>) programme. Appendix A of Sepúlveda et al. (2014) presents a method for filtering common signals in night time CH₄ data of the two nearby mountain GAW stations Jungfraujoch (46.5°N , 8.0°E , 3580 m a.s.l.) and Schauinsland (47.9°N , 7.9°E , 1205 m a.s.l.). Data were retained as common signals when deviations of observations (after correction for vertical gradient, i.e. application of an offset, and a temporal shift in the annual cycles) at both sites were below a certain threshold. Sepúlveda et al. (2014) showed that the common signals are well representative for a broader layer in the lower free troposphere. Here we follow this approach and use the mean of the Jungfraujoch and Schauinsland CH₄ mixing ratio – whenever identified as a common signal – as a validation reference for the remote sensing data in South-western Germany and Northern Switzerland (indicated by the grey circle in Fig. 8b). We assume that the signals obtained from this GAW data filtering are well representative for the tropospheric partial column averaged mixing ratios (surface - 6 km a.s.l.) and compare these data directly to different satellite products as a fully independent data set: we do not adjust the data to a common a priori data usage as in Sects. 4.1, because the in-situ data represent absolute measurements and do not depend on any a priori information. Furthermore, we do not adjust sensitivities as in Sect. 4.2 (see Eq. (7)), which means that we validate here also the sensitivities of the products.

In order to be able to compare TROPOMI data to the GAW data, we calculate from the TROPOMI XCH₄ data a proxy ($f_{\text{T}}(\text{TROPOMI})$) that represents the tropospheric column averaged mixing ratios:

$$\begin{aligned} \text{troXCH}_4(\text{TROPOMI}) &\approx f_{\text{T}}(\text{TROPOMI}) \\ &= \frac{X_{\text{air}}}{\text{troX}_{\text{air}}} [\text{XCH}_4(\text{TROPOMI}) - \text{XCH}_4(\text{apriori})] + \text{troXCH}_4(\text{apriori}). \end{aligned} \quad (8)$$




Figure 13. Comparison of GAW measurements made at Jungfraujoch and Schauinsland with the TROPOMI tropospheric CH_4 proxy product according to Eq. 8 and the IASI and the combined tropospheric CH_4 products. Data for all individual coincidences are shown in the background as squares and daily averages are depicted as crosses with error bars representing the daily 1σ standard deviations (daily means are only calculated if there are at least 3 observations per day). (a)-(c) Time series of differences. (d)-(f) Correlation between GAW and satellite data (the black line is the one-to-one diagonal). (g)-(i) Correlations between the difference of GAW and a priori data ($\Delta_a \text{GAW} = \text{troXCH}_4(\text{GAW}) - \text{troXCH}_4(\text{apriori})$) and the a priori free satellite data ($\Delta_a \text{Satellite} = \text{troXCH}_4(\text{Satellite}) - \text{troXCH}_4(\text{apriori})$). The inserted text reports median and scatter (hIPR68.2, a-c) and the coefficients of determination, the slope, and the intercept of the robust linear regression model (R^2 , m , and b , d-i). Dark and pale coloured fonts represent the values for the daily mean data and for data from all individual collocations, respectively.

In Eq. (8) X_{air} and troX_{air} are the dry air total and tropospheric partial columns, respectively, and $\text{troXCH}_4(\text{apriori})$ is the tropospheric column averaged CH_4 a priori. In the case that the CH_4 a priori in the UTLS is of very good quality, this proxy is well representative for the tropospheric CH_4 variations.

Figure 13 shows the comparison with the different satellite products. For the tropospheric proxy product calculated from the XCH₄ product of TROPOMI we observe no systematic difference and a scatter of the daily mean differences of within 1.3% (Fig. 13a). However, the correlation is rather weak (from the robust linear regression model we get R^2 values of about 10% and regression line slope m of below 0.5, see Fig. 13d), which might suggest that this proxy is affected by signals in the UTLS, where CH₄ values are dominated by shifts of the tropopause height.

For the MUSICA IASI tropospheric partial column averaged mixing ratio product (Fig. 13b and e) we observe a smaller median difference than for the comparison with the TROPOMI tropospheric proxy CH₄ data, but at the same time an increased scatter. The R^2 values are larger than for the correlation of TROPOMI proxy data; however, we have to be careful, because in the lower troposphere the MUSICA IASI CH₄ data have a limited sensitivity (see Fig. 5b). This means that the respective data are significantly affected by the a priori assumptions and the observed correlation might actually be due to a correlation with the tropospheric a priori data. This is confirmed by Fig. 13h, which shows the correlation after removing the a priori data. Then the correlation strength is weaker if compared to the data that include the a priori information (R^2 decreases from about 25% to 20% and from about 20% to 10% for correlations with daily mean and all individual data, respectively).

The combined product has a good sensitivity in the troposphere (see Fig. 5b), i.e. the respective partial column averaged mixing ratio product is practically independent from the a priori assumptions. We find a good agreement and correlation between the GAW data and the combined products as illustrated in Fig. 13c and f: for instance, for daily mean data the difference and scatter is $+0.28\% \pm 1.05\%$, the R^2 value is about 37%, and the regression line slope very close to 1.0. This demonstrates that the combined product can reliably capture actual tropospheric CH₄ variations independently from the UTLS CH₄ variations and from the a priori assumption. The latter is confirmed by Fig. 13i, which shows the correlation after removing the tropospheric a priori information. We observe that the good correlation remains even after removing the a priori information (for daily mean data the R^2 value is about 39% and the regression line slope close to 1.0). A similar good correlation is not achieved by the TROPOMI tropospheric proxy and the IASI product.

5 Global data

5.1 Discussion on global data consistency

The TCCON and AirCore comparisons of Sects. 4.1 and 4.2 suggest that the combined total column and UTLS partial column products have no significant bias with respect to reference data. However, there might be a weak bias in the troposphere (see discussions in the context of Fig. 11). In general we have to consider that the study on biases in the profile data is limited to the two sites where AirCore references are available: Sodankylä in northern Scandinavia and Trainou in France. In this section, we argue that it is reasonable to assume similar insignificant or low biases also for other locations.

According to Eq. (A2) a varying error in the a priori state together with a poor sensitivity (i.e. an averaging kernel being very different from an identity matrix) can cause a varying bias. If the error in the a priori state is latitudinally dependent the bias will also be latitudinally dependent. Similarly, a systematic error source (like an error in a spectroscopic parameter) can have a variable impact on the remote sensing product, if the sensitivity is variable. If the sensitivity has a dependency on




Figure 14. Latitudinal dependency of the overall mean values obtained at the 14 TCCON and two AirCore observation sites. Grey colour represents the TCCON and AirCore reference data and black, red and blue colours the TROPOMI, MUSICA IASI, and combined satellite data, respectively: (a) for total columns (XCH_4) and (b) for tropospheric and UTLS partial columns. The error bars on the AirCore data describe the variability range due to the AirCore data treatment – according to Eq. (7) – with the different averaging kernels of the TROPOMI, MUSICA IASI and combined data product.

455 latitude, a systematic error source can thus also cause a latitudinally dependent bias. In this context, variabilities (e.g. latitudinal dependencies) of biases are likely for a low or variable sensitivity. In contrast, inconsistencies in the bias are unlikely in case of a high and constant sensitivity (as observed in Fig. 5 for the total column and tropospheric and UTLS partial columns of the combined data product).

Figure 14 depicts the overall mean total and partial column values obtained at the 14 TCCON and and two AirCore ob-
 460 servation sites. For total column data (Fig. 14a) we achieve a good latitudinal coverage by the TCCON observation sites and can investigate possible latitudinal inconsistencies in the satellite data products. We find that the TROPOMI and the combined satellite data product capture a latitudinal dependency that is similar to the dependency as seen in the TCCON data.

Figures 3a and 5a reveal that for the TROPOMI and the combined XCH_4 products the sensitivities are very high and stable, in contrast to the MUSICA IASI data product, which has a relatively weak and seasonally (and supposed latitudinally) varying
 465 sensitivity. This explains that in Fig. 14a the latitudinal dependency of the MUSICA IASI XCH_4 data is different from the TCCON data. Table 3 resumes the statistics made with the overall mean XCH_4 values obtained for the 14 TCCON observation sites. For the TROPOMI and the combined data product the 1σ standard deviation calculated from the mean difference of the 14 stations is about 0.4%. A standard linear least squares fit results in R^2 values of almost 100% and regression line slope values of close to unity, which confirms the very good latitudinal data consistency of the TROPOMI and combined data products. The
 470 MUSICA IASI XCH_4 product shows poorer performance with regard to the values of standard deviation and R^2 , which is in line with its weak and varying sensitivity.

A similar study of the latitudinal consistency of the partial column data products is compromised by the lack of profile references for low latitudes and southern hemispheric sites (Fig. 14b). Nevertheless, because the combined product has a rather high and constant sensitivity for the tropospheric as well as the UTLS partial column (see Figs. 3 and 5), we expect – as for
 475 XCH_4 – a good latitudinal consistency, i.e. a bias at low and/or southern latitudes that is similarly insignificant or low as the biases observed at for two AirCore station in the middle and high northern latitudes.

Table 3. Statistics based on the comparisons between satellite and TCCON data of the overall mean XCH₄ values obtained for the 14 TCCON sites of Table 1.

Product	Difference (mean±std)	R^2	Slope (m)	Intercept (b)
TROPOMI	-0.04%±0.41%	94.2%	1.09	-174 ppb
MUSICA IASI	+0.71%±0.67%	79.2%	0.88	+237 ppb
Combined	+0.02%±0.41%	94.2%	1.09	-173 ppb

5.2 Example of global maps

The proposed synergetic use method needs no extra retrievals, and is thus computationally very efficient. This makes it ideal for combining the large TROPOMI and IASI data sets on global scale. Figure 15 shows monthly mean global maps ($1^\circ \times 1^\circ$ resolution) of TROPOMI XCH₄ data and the tropospheric and UTLS partial column CH₄ data of the combined product. The maps are generated from about 1.62 million and 3.77 million individual data points in January and July 2020, respectively. These are the data that remain after requiring collocation of the quality filtered IASI and TROPOMI data according to Sect. 2.

TROPOMI alone only reports the XCH₄ data (Fig. 15a and b). We observe low XCH₄ values at high latitudes. The lowest values are encountered in the summertime southern hemisphere. The highest XCH₄ values are present between northern low and middle latitudes. Here Fig. 15a and b shows the TROPOMI data, the XCH₄ data of the combined product are very similar.

The combined product offers the most reliable tropospheric partial columns. Respective maps are shown in Fig. 15c and d. We observe partial column averaged CH₄ mixing ratios that are almost monotonically increasing from south to north. In northern hemispheric winter (January 2020) this gradient is significantly stronger than in northern hemispheric summer (July 2020). The latitudinal patterns of tropospheric CH₄ is significantly different from the respective patterns of XCH₄, which might indicate to an extra potential of this tropospheric CH₄ data when investigating the CH₄ sources and sinks.

Figure 15e and f shows the respective maps of the UTLS partial columns (here we depict the combined data product, the respective MUSICA IASI data are very similar). We observe highest partial column averaged mixing ratios at low latitudes (in January 2020 around the equator and in July 2020 in the northern subtropics). The mixing ratios are lowest in high latitudes. This latitudinal pattern is in line with the tropopause height, which increases from high to low latitudes.

495 6 Summary and outlook

We present a method for a synergetic combination of the IASI vertical profile and TROPOMI total column level 2 retrieval products. It is computationally very efficient, because it is based on simple linear algebra calculations that works with the output already available from individual IASI and TROPOMI retrievals. Nevertheless, theoretically it approximates closely to a computationally expensive multispectral retrieval, that use the TROPOMI and IASI level 1 data (see Appendix B). We apply




Figure 15. Global maps with $1^\circ \times 1^\circ$ (longitude \times latitude) resolution of monthly mean data for January and July 2020. (a)+(b) XCH₄ as observed by TROPOMI. (c)+(d) Tropospheric partial columns of CH₄ as obtained by the combined product. (e)+(f) UTLS partial columns of CH₄ as obtained by the combined product.

500 the method to CH₄ level 2 products. By providing a compilation with all important equations we support its application to other data products.

We theoretically examine the sensitivity, vertical resolution, and errors of the individual TROPOMI and IASI products and of the combined product. The TROPOMI product consists of reliable total column CH₄ data, but does not offer information on the vertical distribution. The IASI product offers some information on the vertical distribution and has best sensitivity in
 505 the UTLS region, but lacks sensitivity in the lower troposphere and in consequence to the total column. We show that the combined product combines both strengths: it is a reliable reference for the total column and also for the UTLS partial column.

In addition, we found as a clear synergetic effect that the combined product is theoretically able to distinguish variations of CH_4 that take place in the troposphere from variations at higher altitudes (it is a reliable reference for the tropospheric partial columns). We empirically demonstrate the functionality of the synergetic use method by comparing the different satellite CH_4 products to reference data of TCCON, AirCore and GAW.

The TCCON data offer good references for XCH_4 . In this study we use data from 14 stations covering different climate regions in the northern and southern hemisphere. For the TROPOMI and the combined data products, which are well sensitive for XCH_4 , we get an agreement with the TCCON data within about 0.7% (the agreement is slightly poorer with the IASI satellite product due its reduced sensitivity).

AirCore offers XCH_4 references as well as references for the vertical distribution of CH_4 . For this study 36 individual AirCore profiles measured at two sites in the northern hemispheric high and middle latitudes are available. Concerning XCH_4 , the comparisons to AirCore data confirm the results obtained by the comparison to TCCON data, and in addition demonstrate a very good consistency between the TROPOMI and the IASI product. Concerning CH_4 in the UTLS – where the MUSICA IASI and the combined data product are well sensitive – we find that both products agree well with the respective AirCore references (agreement within 0.7%).

The validation study with the TCCON and AirCore references shows that the total column and the UTLS partial column of the combined product has almost the same good quality as the respective products of TROPOMI and MUSICA IASI. This allows two conclusions: firstly, the assumption of the moderate non-linearity – required for a reliable functionality of the level 2 product combination according to Eq. (1) – is valid and secondly, the combined product's tropospheric data are also of good quality (good total column and UTLS data quality is an indirect proof of a good tropospheric data quality).

The good quality of the combined product in the troposphere is in addition directly proven by the comparison to tropospheric reference data. We find an agreement of the daily mean tropospheric AirCore and combined product data within 1%. This validation result is confirmed by the statistically very robust comparison with CH_4 data observed continuously at two nearby GAW stations (the collocated GAW reference data cover all seasons for more than 3 years and represent more than 186 different days). The GAW and the combined product's data capture very similar tropospheric CH_4 short term variabilities and seasonal cycles. Similar good agreements are not achieved by comparisons to the individual MUSICA IASI or TROPOMI data products, i.e. we empirically and directly prove the synergetic effect of the level 2 product combination.

The proposed method takes benefit from the outputs generated by the dedicated individual TROPOMI and IASI retrievals, it needs no extra retrievals, and is thus computationally very efficient. This is ideal for an operational combination of IASI and TROPOMI products in an efficient and sustained manner. This has a particular attraction, because IASI and TROPOMI successor instruments will be jointly aboard the upcoming Metop (Meteorological operational) Second Generation satellites (guaranteeing observations from the 2020s to the 2040s). IASI and TROPOMI successor instruments will have globally-distributed and perfectly-collocated observations (over land) in the order of several hundred thousands per day, for which a combined product can be generated in a computationally very efficient way.

540 *Data availability.* Access to the MUSICA IASI data is provided via <http://www.imk-asf.kit.edu/english/musica-data.php>. The TROPOMI XCH₄ data used in this study are available for download at ftp://ftp.sron.nl/open-access-data-2/TROPOMI/tropomi/ch4/14_14_Lorente_et_al_2020_AMTD/. TCCON data are made available via the TCCON Data Archive, hosted by CaltechDATA, California Institute of Technology, California (USA), <http://tccodata.org>. For Trainou AirCore data please contact Michel Ramonet (michel.ramonet@Isce.ipsl.fr) and for Sodankylä AirCore data please contact Huilin Chen (huilin.chen@rug.nl). The GAW surface in-situ data are available via the World Data Centre for Greenhouse Gases (WDCGG), <https://gaw.kishou.go.jp/search/>.

Appendix A: Basics on retrieval theory

In this appendix, we give a brief overview on the theory of optimal estimation remote sensing methods and follow the notation as recommended by the TUNER activity (von Clarmann et al., 2020), which is closely in line with the notation used by Rodgers (2000). The overview focuses on the equations that are important for our work, i.e. the a posteriori combination of two independently retrieved optimal estimation remote sensing products. For a more detailed and general insight into the theory of optimal estimation remote sensing methods we refer to Rodgers (2000).

Atmospheric remote sensing instruments measure radiance spectra (written as state vector \mathbf{y}), which can be well simulated by models (\mathbf{F}) whenever the actual atmospheric state (the vector \mathbf{x}) is known. Using the a priori atmospheric state vector \mathbf{x}_a we can linearise and write:

$$555 \quad \mathbf{F}(\mathbf{x}) - \mathbf{F}(\mathbf{x}_a) = \Delta\mathbf{y} = \mathbf{K}(\mathbf{x} - \mathbf{x}_a). \quad (\text{A1})$$

Here, \mathbf{K} is the Jacobian matrix, i.e. derivatives that capture how the measurement vector (the measured radiances) will change for changes of the atmospheric state (the atmospheric state vector \mathbf{x}). A remote sensing retrieval inverts Eq. (A1) and provides an estimation of the difference between the atmospheric state and the a priori atmospheric state. For a moderately non-linear problem (according to Chapter 5 of Rodgers, 2000), the retrieved optimal estimation product ($\hat{\mathbf{x}}$) can be written as:

$$560 \quad \hat{\mathbf{x}} - \mathbf{x}_a = \mathbf{G}\Delta\mathbf{y} = \mathbf{G}[\mathbf{K}(\mathbf{x} - \mathbf{x}_a)]. \quad (\text{A2})$$

\mathbf{G} is the gain matrix and realises the inversion from the measurement domain (radiances) to the domain of the atmospheric states. It consists of derivatives that capture how the retrieved atmospheric state vector will change for changes in the measurement vector:

$$565 \quad \begin{aligned} \mathbf{G} &= (\mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K} + \mathbf{S}_a^{-1})^{-1} \mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \\ &= \mathbf{S}_a \mathbf{K}^T (\mathbf{K} \mathbf{S}_a \mathbf{K}^T + \mathbf{S}_{\mathbf{y},n})^{-1}, \end{aligned} \quad (\text{A3})$$

with $\mathbf{S}_{\mathbf{y},n}$ and $\mathbf{R} = \mathbf{S}_a^{-1}$ being the retrieval's noise covariance and the constraint matrices (in a strict optimal estimation sense, the constraint matrix is the inverse of the a priori covariance matrix \mathbf{S}_a), respectively. The equivalence of both lines in Eq. (A3) is demonstrated in Chapter 4.1 of Rodgers (2000), where the first line is called the n -form and the second line the m -form.

The averaging kernel

$$570 \quad \mathbf{A} = \mathbf{G}\mathbf{K}, \quad (\text{A4})$$

is an important component of a remote sensing retrieval, because according to Eq. (A2) it reveals how changes of the actual atmospheric state vector \boldsymbol{x} affect the retrieved atmospheric state vector $\hat{\boldsymbol{x}}$.

A valuable diagnostic quantity is the a posteriori covariance matrix, which can be calculated as follows:

$$\mathbf{S}_{\hat{\boldsymbol{x}}} = (\mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K} + \mathbf{S}_{\mathbf{a}}^{-1})^{-1}. \quad (\text{A5})$$

575 The linearised formulation of the retrieval solution according to (A2) is very useful for the analytic characterisation of the product. The retrieval state's noise error covariance matrix for noise can be analytically calculated as:

$$\begin{aligned} \mathbf{S}_{\hat{\boldsymbol{x}},n} &= \mathbf{G} \mathbf{S}_{\mathbf{y},n} \mathbf{G}^T \\ &= \mathbf{S}_{\hat{\boldsymbol{x}}} \mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K} \mathbf{S}_{\hat{\boldsymbol{x}}}, \end{aligned} \quad (\text{A6})$$

580 where $\mathbf{S}_{\mathbf{y},n}$ is the covariance matrix for noise on the measured radiances \mathbf{y} . The second line of Eq. (A6) is obtained by substituting \mathbf{G} by $\mathbf{S}_{\hat{\boldsymbol{x}}} \mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1}$ according to Eqs. (A3) and (A5). The representativeness error reveals the deficit of the retrieval product in representing the actual variations of the state vector \boldsymbol{x} . In Chapter 3 of Rodgers (2000) it is called the smoothing error and can be calculated as (with \mathbf{I} being the identity matrix):

$$\begin{aligned} \mathbf{S}_{\hat{\boldsymbol{x}},r} &= (\mathbf{A} - \mathbf{I}) \mathbf{S}_{\mathbf{a}} (\mathbf{A} - \mathbf{I})^T \\ &= \mathbf{S}_{\hat{\boldsymbol{x}}} \mathbf{S}_{\mathbf{a}}^{-1} \mathbf{S}_{\hat{\boldsymbol{x}}}. \end{aligned} \quad (\text{A7})$$

585 The second line of Eq. (A7) is obtained using:

$$\mathbf{S}_{\hat{\boldsymbol{x}}} = (\mathbf{I} - \mathbf{A}) \mathbf{S}_{\mathbf{a}}, \quad (\text{A8})$$

which in turn follows from Eqs. (A3) - (A6).

Using Eqs. (A5) - (A7) reveals that the a posteriori covariance is the sum of the noise error covariance and the representativeness error covariance:

$$590 \quad \mathbf{S}_{\hat{\boldsymbol{x}}} = \mathbf{S}_{\hat{\boldsymbol{x}},n} + \mathbf{S}_{\hat{\boldsymbol{x}},r}. \quad (\text{A9})$$

Appendix B: Theory on the optimal combination of retrieval data products

In this section, we discuss an optimal estimation retrieval that uses a combined measurement vector (two measurements from different instruments). First we show that the retrieval output of two profile retrievals performed on the same vertical grid can be used in a way that yields to the same results as performing a retrieval with the combined measurement vector. Then we present an approach for combining the outputs of a retrieval that provides profiles and another retrieval that provides column data. We show that the combination of profile and column data can be realised in a computationally efficient manner via a Kalman filter. Finally, we discuss the validity of the methods and the requirements on the individual retrieval products.

B1 Inversion of a combined measurement vector

According to Eqs. (A2), (A3), and (A5) the retrieval product obtained from measurement \mathbf{y} can be written as:

$$600 \quad \hat{\mathbf{x}} - \mathbf{x}_a = (\mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K} + \mathbf{S}_a^{-1})^{-1} \mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K} (\mathbf{x} - \mathbf{x}_a). \quad (\text{B1})$$

In the case of two individual measurements (measurement 1 and 2), we obtain from using a combined measurement vector $\{\mathbf{y}_1, \mathbf{y}_2\}$:

$$\begin{aligned} \hat{\mathbf{x}} - \mathbf{x}_a &= (\mathbf{K}_1^T \mathbf{S}_{\mathbf{y}_1,n}^{-1} \mathbf{K}_1 + \mathbf{K}_2^T \mathbf{S}_{\mathbf{y}_2,n}^{-1} \mathbf{K}_2 + \mathbf{S}_a^{-1})^{-1} (\mathbf{K}_1^T \mathbf{S}_{\mathbf{y}_1,n}^{-1} \mathbf{K}_1 + \mathbf{K}_2^T \mathbf{S}_{\mathbf{y}_2,n}^{-1} \mathbf{K}_2) (\mathbf{x} - \mathbf{x}_a) \\ &= (\mathbf{S}_{\hat{\mathbf{x}}_1}^{-1} + \mathbf{S}_{\hat{\mathbf{x}}_2}^{-1} - \mathbf{S}_a^{-1})^{-1} (\mathbf{K}_1^T \mathbf{S}_{\mathbf{y}_1,n}^{-1} \mathbf{K}_1 + \mathbf{K}_2^T \mathbf{S}_{\mathbf{y}_2,n}^{-1} \mathbf{K}_2) (\mathbf{x} - \mathbf{x}_a), \end{aligned} \quad (\text{B2})$$

605 where $\mathbf{S}_{\mathbf{y}_1,n}$ and $\mathbf{S}_{\mathbf{y}_2,n}$ are the respective measurement noise covariances, \mathbf{K}_1 and \mathbf{K}_2 the respective Jacobians and $\mathbf{S}_{\hat{\mathbf{x}}_1}$ and $\mathbf{S}_{\hat{\mathbf{x}}_2}$ the respective a posteriori covariances. The second line follows from Eq. (A5). According to Eqs. (A3) - (A5) we can substitute $\mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K} (\mathbf{x} - \mathbf{x}_a)$ by $\mathbf{S}_{\hat{\mathbf{x}}}^{-1} (\hat{\mathbf{x}} - \mathbf{x}_a)$ and write Eq. (B2) as

$$\hat{\mathbf{x}} - \mathbf{x}_a = (\mathbf{S}_{\hat{\mathbf{x}}_1}^{-1} + \mathbf{S}_{\hat{\mathbf{x}}_2}^{-1} - \mathbf{S}_a^{-1})^{-1} [\mathbf{S}_{\hat{\mathbf{x}}_1}^{-1} (\hat{\mathbf{x}}_1 - \mathbf{x}_a) + \mathbf{S}_{\hat{\mathbf{x}}_2}^{-1} (\hat{\mathbf{x}}_2 - \mathbf{x}_a)]. \quad (\text{B3})$$

Using Eq. (B3) we can realise an optimal combination of the two retrieval products that only needs the a priori covariance, 610 the a posteriori covariances, and the two retrieval products. The Jacobians are not needed. This combination is mathematically equivalent to using the Jacobians of a combined measurement vector $\{\mathbf{y}_1, \mathbf{y}_2\}$, i.e. within a linear subspace (validity of moderate non-linearity according to Chapter 5 of Rodgers, 2000) it is equivalent to a synergetic use of level 1 data in form of a multispectral retrieval.

B2 Combining profile and column data products

615 Equation (B3) requires two retrieval results on the same vertical grid and can be used to combine two profile products. Here we will develop a method for combining a profile and a column data product. For a column retrieval we can write in analogy to Eq. (A1)

$$\Delta x^* = \mathbf{a}^{*T} (\mathbf{x} - \mathbf{x}_a), \quad (\text{B4})$$

where \mathbf{a}^{*T} is the column averaged mixing ratio according to Appendix D2. Equation (B4) poses an inverse problem of the 620 same kind as Eq. (A1) and in order to optimally estimate a profile from an available column product we can apply the same solution approach as in Eqs. (A2) and (A3). A similar application of this approach is also presented in Sect. 4.2 of Rodgers and Connor (2003). For the application here we substitute in Eq. (A3) \mathbf{K} by \mathbf{a}^{*T} and $\mathbf{S}_{\mathbf{y},n}$ by the scalar $S_{\hat{\mathbf{x}},n}^*$ (the noise error variance of the column data product) and get the profile

$$\begin{aligned} \hat{\mathbf{x}} - \mathbf{x}_a &= (\mathbf{a}^* S_{\hat{\mathbf{x}},n}^{*-1} \mathbf{a}^{*T} + \mathbf{S}_a^{-1})^{-1} \mathbf{a}^* S_{\hat{\mathbf{x}},n}^{*-1} \mathbf{a}^{*T} (\mathbf{x} - \mathbf{x}_a) \\ 625 \quad &= (\mathbf{a}^* S_{\hat{\mathbf{x}},n}^{*-1} \mathbf{w}^{*T} \mathbf{S}_{\hat{\mathbf{x}}} \mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K} + \mathbf{S}_a^{-1})^{-1} \mathbf{a}^* S_{\hat{\mathbf{x}},n}^{*-1} \mathbf{w}^{*T} \mathbf{S}_{\hat{\mathbf{x}}} \mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K} (\mathbf{x} - \mathbf{x}_a). \end{aligned} \quad (\text{B5})$$

For the second line of Eq. (B5) we use $\mathbf{a}^{*T} = \mathbf{w}^{*T} \mathbf{A}$ – according to Eq. (D6) – and $\mathbf{A} = \mathbf{S}_{\hat{\mathbf{x}}} \mathbf{K}^T \mathbf{S}_{\mathbf{y},n}^{-1} \mathbf{K}$ – according to Eqs. (A3) - (A5). We write this second line to discuss similarities with Eq. (B1). The comparison of Eq. (B5) with Eq. (B1)

reveals that for a retrieval providing only a column product, the Jacobian information provided by \mathbf{K} is vertically aggregated according to the operator $\mathbf{a}^* S_{\hat{x},n}^{*-1} \mathbf{w}^{*T} \mathbf{S}_{\hat{x}}$. The term $\mathbf{S}_{\hat{x}}$ is the vertically resolved a posteriori covariance, which exist for a
630 retrieval that internally inverts profiles, but only distributes the column products; however, it is only an internal measure of the retrieval, and actually not available.

Instead of the term of Eq. (B4) we now invert the term $\Delta x^* = \mathbf{a}_2^{*T} (\mathbf{x} - \hat{\mathbf{x}}_1)$, i.e. we replace \mathbf{x}_a by the profile product $\hat{\mathbf{x}}_1$ of a first retrieval (retrieval 1) on the right side of (B4) and use \mathbf{a}_2^{*T} and $S_{\hat{x}_2,n}^*$ for the column averaging kernel and the noise error variance of a second retrieval (retrieval 2), respectively. Here and in the following, retrieval 1 is the profile retrieval and
635 retrieval 2 the retrieval that provides only column products. The solution can easily be achieved by substituting in (B5) \mathbf{S}_a by $\mathbf{S}_{\hat{x}_1}$, which is the a posteriori covariance of retrieval 1:

$$\hat{\mathbf{x}} - \hat{\mathbf{x}}_1 = (\mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{a}_2^{*T} + \mathbf{S}_{\hat{x}_1}^{-1})^{-1} \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{a}_2^{*T} (\mathbf{x} - \hat{\mathbf{x}}_1). \quad (\text{B6})$$

We modify Eq. (B6) by using $\hat{\mathbf{x}}_1 = \mathbf{A}_1 (\mathbf{x} - \mathbf{x}_a) + \mathbf{x}_a$:

$$\begin{aligned} \hat{\mathbf{x}} - \mathbf{x}_a &= \mathbf{A}_1 (\mathbf{x} - \mathbf{x}_a) + (\mathbf{S}_{\hat{x}_1}^{-1} + \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{a}_2^{*T})^{-1} \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{a}_2^{*T} (\mathbf{I} - \mathbf{A}_1) (\mathbf{x} - \mathbf{x}_a) \\ 640 \quad &= (\mathbf{S}_{\hat{x}_1}^{-1} + \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{a}_2^{*T})^{-1} [\mathbf{S}_{\hat{x}_1}^{-1} \mathbf{A}_1 + \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{a}_2^{*T}] (\mathbf{x} - \mathbf{x}_a) \\ &= (\mathbf{S}_{\hat{x}_1}^{-1} + \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{a}_2^{*T})^{-1} [\mathbf{S}_{\hat{x}_1}^{-1} (\hat{\mathbf{x}}_1 - \mathbf{x}_a) + \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} (\hat{x}_2^* - \mathbf{w}^{*T} \mathbf{x}_a)]. \end{aligned} \quad (\text{B7})$$

In the third line of Eq. (B7) we use the column product $\hat{x}_2^* = \mathbf{a}_2^{*T} (\mathbf{x} - \mathbf{x}_a) + \mathbf{w}^{*T} \mathbf{x}_a$. Similarly to Eq. (B3) we can generate a combined product without the need of the Jacobian matrices. The combination is possible by using the profile and the column product ($\hat{\mathbf{x}}_1$ and \hat{x}_2^* , respectively) together with the a posteriori covariance of the profile product and the noise error and
645 averaging kernel of the column product.

If we substitute in the second line of Eq. (B7) $\mathbf{S}_{\hat{x}_1}^{-1}$ by $\mathbf{K}_1^T \mathbf{S}_{y_1,n}^{-1} \mathbf{K}_1 + \mathbf{S}_a^{-1}$ – according to Eq. (A5), \mathbf{a}_2^{*T} by $\mathbf{w}^{*T} \mathbf{A}_2$ – according to Eq. (D6), and then \mathbf{A}_2 by $\mathbf{S}_{\hat{x}_2} \mathbf{K}_2^T \mathbf{S}_{y_2,n}^{-1} \mathbf{K}_2$ – according to Eqs. (A3) - (A5), we get:

$$\begin{aligned} \hat{\mathbf{x}} - \mathbf{x}_a &= (\mathbf{K}_1^T \mathbf{S}_{y_1,n}^{-1} \mathbf{K}_1 + \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{w}^{*T} \mathbf{S}_{\hat{x}_2} \mathbf{K}_2^T \mathbf{S}_{y_2,n}^{-1} \mathbf{K}_2 + \mathbf{S}_a^{-1})^{-1} \\ &[\mathbf{K}_1^T \mathbf{S}_{y_1,n}^{-1} \mathbf{K}_1 + \mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{w}^{*T} \mathbf{S}_{\hat{x}_2} \mathbf{K}_2^T \mathbf{S}_{y_2,n}^{-1} \mathbf{K}_2] (\mathbf{x} - \mathbf{x}_a). \end{aligned} \quad (\text{B8})$$

This equation has strong similarities to the first line of Eq. (B2), i.e. the retrieval product obtained when using the combined measurement vector $\{\mathbf{y}_1, \mathbf{y}_2\}$. The only difference is that in Eq.(B8) the information provided by Jacobian \mathbf{K}_2 is vertically aggregated according to the operator $\mathbf{a}_2^* S_{\hat{x}_2,n}^{*-1} \mathbf{w}^{*T} \mathbf{S}_{\hat{x}_2}$.

B3 Linear Kalman filter

Here we show that the approach developed in Appendix B2 is equivalent to a Kalman filter. An important application of a
655 Kalman filter (Kalman, 1960; Rodgers, 2000) is data assimilation in the context of atmospheric modelling. There, the filter operates sequentially in different time steps. Kalman filter data assimilation methods determine the analysis state ($\hat{\mathbf{x}}^a$) by optimally combining the background (or forecast) state ($\hat{\mathbf{x}}^b$) with the information as provided by a new observation ($\hat{\mathbf{x}}^o$):

$$\hat{\mathbf{x}}^a = \hat{\mathbf{x}}^b + \mathbf{M}[\hat{\mathbf{x}}^o - \mathbf{H}\hat{\mathbf{x}}^b] \quad (\text{B9})$$

Optimal means here that the uncertainties of both, the background state and the observation, are correctly taken into account
660 by the Kalman gain matrix (\mathbf{M}):

$$\mathbf{M} = \mathbf{S}_{\hat{\mathbf{x}}^b} \mathbf{H}^T (\mathbf{H} \mathbf{S}_{\hat{\mathbf{x}}^b} \mathbf{H}^T + \mathbf{S}_{\hat{\mathbf{x}}^o, n})^{-1}, \quad (\text{B10})$$

with $\mathbf{S}_{\hat{\mathbf{x}}^b}$ and $\mathbf{S}_{\hat{\mathbf{x}}^o, n}$ being the uncertainty covariances of background state and the new measurement, respectively. The matrix \mathbf{H} is the measurement forward operator, which maps the background domain into the measurement domain.

By rearranging the n -form of (B6) as the m -form – in analogy to Eq. (A3) – and by using again $\hat{\mathbf{x}}_1 = \mathbf{A}_1(\mathbf{x} - \mathbf{x}_a) + \mathbf{x}_a$ and
665 $\hat{x}_2^* = \mathbf{a}_2^{*T}(\mathbf{x} - \mathbf{x}_a) + \mathbf{w}^{*T} \mathbf{x}_a$ we get

$$\begin{aligned} \hat{\mathbf{x}} &= \hat{\mathbf{x}}_1 + \mathbf{S}_{\hat{\mathbf{x}}_1} \mathbf{a}_2^* (\mathbf{a}_2^{*T} \mathbf{S}_{\hat{\mathbf{x}}_1} \mathbf{a}_2^* + S_{\hat{x}_2, n}^*)^{-1} \mathbf{a}_2^{*T} (\mathbf{x} - \hat{\mathbf{x}}_1) \\ &= \hat{\mathbf{x}}_1 + \mathbf{S}_{\hat{\mathbf{x}}_1} \mathbf{a}_2^* (\mathbf{a}_2^{*T} \mathbf{S}_{\hat{\mathbf{x}}_1} \mathbf{a}_2^* + S_{\hat{x}_2, n}^*)^{-1} [\hat{x}_2^* - \mathbf{a}_2^{*T} \hat{\mathbf{x}}_1 - (\mathbf{w}^{*T} \mathbf{x}_a - \mathbf{a}_2^{*T} \mathbf{x}_a)] \\ &= \hat{\mathbf{x}}_1 + \mathbf{m} (\hat{x}_2^* - \mathbf{a}_2^{*T} \hat{\mathbf{x}}_1) - \mathbf{m} (\mathbf{w}^{*T} \mathbf{x}_a - \mathbf{a}_2^{*T} \mathbf{x}_a) \end{aligned} \quad (\text{B11})$$

with

$$670 \quad \mathbf{m} = \mathbf{S}_{\hat{\mathbf{x}}_1} \mathbf{a}_2^* (\mathbf{a}_2^{*T} \mathbf{S}_{\hat{\mathbf{x}}_1} \mathbf{a}_2^* + S_{\hat{x}_2, n}^*)^{-1}. \quad (\text{B12})$$

Disregarding the term that accounts for the a priori information ($\mathbf{m}(\mathbf{w}^{*T} \mathbf{x}_a - \mathbf{a}_2^{*T} \mathbf{x}_a)$), the Eqs. (B11) and (B12) are the same as the Kalman filter Eqs. (B9) and (B10): retrieval 1 provides the background state and retrieval 2 the new observation. Compared to Eqs. (B7) and (B8) the form of Eq. (B11) has the advantage that no matrices have to be inverted only the scalar ($\mathbf{a}_2^{*T} \mathbf{S}_{\hat{\mathbf{x}}_1} \mathbf{a}_2^* + S_{\hat{x}_2, n}^*$).

675 We have shown that Eq. (B11) is mathematically the same as Eq. (B7) and Eq. (B8). The latter is in turn very similar to the synergetic use of level 1 data in form of a multispectral retrieval as discussed in the context of Eq. (B3).

B4 Discussion and requirements

In the Appendices B2 and B3, we assume the usage of the same a priori for the two individual retrievals. Since generally two individually performed retrievals use two different a priori settings we have to perform an a priori adjustment. Using the a priori
680 of retrieval 2 as the reference ($\mathbf{x}_{2,a} = \mathbf{x}_a$), we can adjust the output of retrieval 1 by (see Eq. (10) of Rodgers and Connor, 2003):

$$\hat{\mathbf{x}}_1' = \hat{\mathbf{x}}_1 + (\mathbf{A}_1 - \mathbf{I})(\mathbf{x}_{1,a} - \mathbf{x}_{2,a}), \quad (\text{B13})$$

where $\mathbf{x}_{1,a}$ is the a priori used by retrieval 1.

For a combination according to Eq. (B3) we need retrieval 1 and 2 outputs obtained by using the same constraint (the inverse
685 of the a priori covariance \mathbf{S}_a). This has to be accounted for before applying Eq. (B3), by adjusting the constraint according to the formalism as presented in Chapter 10.4 of Rodgers (2000) or Sect. 4.2 of Rodgers and Connor (2003). By applying Eq. (B7) or the Kalman filter according to Eq. (B11) the common constraint is automatically set to the constraint of the retrieval 1 product and no extra modification is necessary.

The synergetic combination of remote sensing profile and column products according to Eq. (B7) or Eq. (B11) is possible, whenever: (1) the two remote sensing observations are made at the same time and detect the same location, (2) the problems is moderately non-linear (according to Chapter 5 of Rodgers, 2000), and (3) the individual retrieval output as listed by Eq. (B7) or Eq. (B11) is made available: for the profile retrieval, we need the a posteriori covariances ($\mathbf{S}_{\hat{x}}$, which might also be reconstructed from \mathbf{A} and $\mathbf{R} = \mathbf{S}_a^{-1}$ according to Eq. (A8)), the averaging kernels (\mathbf{A}), and the retrieved and a priori state vectors (\hat{x} and x_a , respectively). For the column retrieval, we need the noise variances (the scalar $S_{\hat{x},n}^*$), the column averaging kernels (the row vector a^{*T}), the column product (\hat{x}_2^*), and the a priori column data ($w^{*T} x_a$), respectively.

Appendix C: Operator for transformation between linear and logarithmic scales

Linear scale differentials and logarithmic scale differentials are related by $\partial x = x \partial \ln x$. For transforming differentials or covariances of a state vector with dimension nal (nal : number of atmospheric levels) from logarithmic to linear scale we define the $nal \times nal$ diagonal matrix \mathbf{L} :

$$\mathbf{L} = \begin{pmatrix} \hat{x}_1 & 0 & \cdots & 0 \\ 0 & \hat{x}_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \hat{x}_{nal} \end{pmatrix}. \quad (C1)$$

Here \hat{x}_i is the value of the i th element of the retrieved state vector (i.e. in case of an atmospheric CH_4 state vector the CH_4 mixing ratios retrieved at the i th model level).

Approximatively, a logarithmic scale averaging kernel matrix \mathbf{A}^1 can then be expressed in the linear scale as:

$$\mathbf{A} \approx \mathbf{L} \mathbf{A}^1 \mathbf{L}^{-1}. \quad (C2)$$

This is here an approximation, because on the right side the operator \mathbf{L} should contain the actual instead of the retrieved mixing ratios. It is a valid approximation as long as the a priori is reasonable and there is no large bias in the retrieval data.

Similarly a logarithmic scale covariance matrix \mathbf{S}^1 can be approximately expressed in the linear scale as:

$$\mathbf{S} \approx \mathbf{L} \mathbf{S}^1 \mathbf{L}^T. \quad (C3)$$

Here the approximation is because $\Delta x \approx x \Delta \ln x$.

Appendix D: Operators for column data

This appendix explains the calculation of operators for partial (and total) column data. Although some sections are similar to Appendix C of Schneider et al. (2022) we think it is a very useful reference here, because it facilitates the reproducibility of our results.

For converting mixing ratio profiles into amount profiles we set up a pressure weighting operator \mathbf{Z} , as a diagonal matrix
 715 with the following entries:

$$Z_{i,i} = \frac{\Delta p_i}{g_i m_{\text{air}} \left(1 + \frac{m_{\text{H}_2\text{O}}}{m_{\text{air}}} \hat{x}_i^{\text{H}_2\text{O}}\right)}. \quad (\text{D1})$$

Using the pressure p_i at atmospheric grid level i we set $\Delta p_1 = \frac{p_2 - p_1}{2} - p_1$, $\Delta p_{nal} = p_{nal} - \frac{p_{nal} - p_{nal-1}}{2}$, and $\Delta p_i = \frac{p_{i+1} - p_i}{2} - \frac{p_i - p_{i-1}}{2}$ for $1 < i < nal$. Furthermore, g_i is the gravitational acceleration at level i , m_{air} and $m_{\text{H}_2\text{O}}$ the molecular mass of dry air and water vapour, respectively, and $\hat{x}_i^{\text{H}_2\text{O}}$ the retrieved or modelled water vapour mixing ratio at level i .

720 We define an operator \mathbf{W}^T for resampling fine gridded atmospheric amount profiles into coarse gridded atmospheric partial column amount profiles. It has the dimension $c \times nal$, where c is the number of the resampled coarse atmospheric grid levels and nal , the number of atmospheric levels of the original fine atmospheric grid. Each line of the operator has the value '1' for the levels that are resampled and '0' for all other levels:

$$\mathbf{W}^T = \begin{pmatrix} 1 & \cdots & 1 & 0 & \cdots & \cdots & \cdots & \cdots & 0 \\ 0 & \cdots & 0 & 1 & \cdots & 1 & 0 & \cdots & 0 \\ 0 & \cdots & \cdots & \cdots & \cdots & 0 & 1 & \cdots & 1 \end{pmatrix}. \quad (\text{D2})$$

725 In analogy we can define a row vector \mathbf{w}^T (with the dimension $1 \times nal$) with all elements having the value '1', which allows the resampling for the total column amounts.

D1 Column amounts

The kernel that describes how a change in the amount at a certain altitude affects the retrieved partial (or total) column amount can be calculated as:

$$730 \mathbf{A}' = \mathbf{W}^T \mathbf{Z} \mathbf{A} \mathbf{Z}^{-1}. \quad (\text{D3})$$

For the total column, we replace \mathbf{W}^T by \mathbf{w}^T and get the row vector \mathbf{a}'^T (dimension $1 \times nal$). This is the total column kernel provided by the TROPOMI data and it is typically written as \mathbf{a}^T . Figure 3 shows examples of such total and partial column amount kernels. The total column amount kernel can be interpolated to different altitude grids. For the applications in Sects. 3 and 4 we interpolate the TROPOMI total column amount kernel to the vertical grid used by the MUSICA IASI retrieval.

735 D2 Column averaged mixing ratios

We can also combine the operators \mathbf{Z} and \mathbf{W}^T for the calculation of a pressure weighted resampling operator by:

$$\mathbf{W}^{*T} = (\mathbf{W}^T \mathbf{Z} \mathbf{W})^{-1} \mathbf{W}^T \mathbf{Z}. \quad (\text{D4})$$

This operator resamples linear scale mixing ratio profiles into linear scale partial column averaged mixing ratio profiles. The respective total column operator \mathbf{w}^{*T} can be calculated in analogy to Eq. (D4) by replacing \mathbf{W}^T by \mathbf{w}^T

740 With operator \mathbf{W}^{*T} we can calculate a coarse gridded partial column averaged state $\hat{\mathbf{x}}^*$ from the fine gridded linear mixing ratio state $\hat{\mathbf{x}}$ by:

$$\hat{\mathbf{x}}^* = \mathbf{W}^{*T} \hat{\mathbf{x}}. \quad (\text{D5})$$

The kernels matrix of the partial column averaged mixing ratio state can then be calculated from the fine gridded linear scale kernel matrix (\mathbf{A}) by:

$$745 \quad \mathbf{A}^* = \mathbf{W}^{*T} \mathbf{A}. \quad (\text{D6})$$

This kernel describes how a change in the mixing ratio at a certain altitude affects the retrieved partial column averaged mixing ratio. Covariances of the partial column averaged mixing ratio state can be calculated from the corresponding covariance matrices of the fine gridded linear scale (\mathbf{S}) by:

$$\mathbf{S}^* = \mathbf{W}^{*T} \mathbf{S} \mathbf{W}^*. \quad (\text{D7})$$

750 The respective calculations for total column averaged mixing ratios can be made by replacing \mathbf{W}^{*T} by \mathbf{w}^{*T} . For the total column averaged mixing ratios the covariance is a simple variance (the scalar S^*) and the kernel has the dimension $1 \times \text{mol}$, i.e. it is a row vector \mathbf{a}^{*T} .

The total column amount kernel (\mathbf{a}_T^T) provided with the TROPOMI data set can be converted into a total column averaged mixing ratio kernel \mathbf{a}_T^{*T} by the following calculation (using Eqs. (D3), (D4), and (D6)):

$$755 \quad \mathbf{a}_T^{*T} = \mathbf{w}^{*T} \mathbf{A}_T = (\mathbf{w}^T \mathbf{Z} \mathbf{w})^{-1} \mathbf{a}_T^T \mathbf{Z}. \quad (\text{D8})$$

The total column averaged mixing ratio kernel \mathbf{a}_T^{*T} used in Sects. 3 and 4 is valid for the vertical grid used by the MUSICA IASI retrieval. It is calculated from the TROPOMI total column amount kernel (\mathbf{a}_T^T) provided in the TROPOMI output files according to Eq. (D8), after its interpolation onto the MUSICA IASI grid (see also Appendix D1).

Appendix E: Dislocation of TROPOMI and IASI

760 IASI is on an orbit with descending node equator crossing at 9:30 mean local solar time. TROPOMI is on an orbit with ascending node equator crossing at 13:30 mean local solar time. In this work we require a temporal collocation within at least six hours. This requirement causes the following typical time difference (IASI - TROPOMI) for observing the same location: at northern high latitudes -0.6 to $+3.7$ hours, at northern middle latitudes -3.3 to -2.2 hours, at the equator -4.5 to -3.5 hours, at southern middle latitudes -5.4 to -4.3 hours, and at southern high latitudes -5.9 to -4.5 hours. This means

765 that at all latitudes we find data that fulfil the temporal collocation requirements and that in the southern hemisphere the temporal collocation is typically larger than in the northern hemisphere. Furthermore, there are horizontal dislocations. In this work we use a horizontal collocation threshold of 50 km. In this appendix we estimate the impact of these spatial and temporal dislocations on the combined product.




Figure E1. Square root values of the diagonal entries of different dislocation covariance matrices ($S_{\Delta_{di}}^1$). These values are the root-mean-squares (RMS) of the difference between the reference methane profile (CAMS forecast for location 49.1°N and 8.4°E, corresponding to the location of Karlsruhe) and other forecasted profiles dislocated with respect to the reference: (a) horizontal dislocations; (b) temporal dislocations. The dashed black lines indicate the collocation threshold values used for valid combinations of IASI and TROPOMI.

E1 Uncertainty source

770 For investigating the spatial and temporal variability of the atmospheric CH₄ fields, we use the CAMS (Copernicus Atmospheric Monitoring service, <https://atmosphere.copernicus.eu/>) CH₄ forecast product at highest available resolution (≈ 9 km, Barré et al., 2021). By analysing the profiles forecasted for the same location but different timestamps, we can determine the temporal covariance of the vertical CH₄ fields. Similarly by analysing the profiles forecasted for the same timestamp but different locations we get the spatial covariance of the vertical CH₄ fields. The analyses are made with CAMS data between
 775 November 2017 and December 2020 for Central Europe in an area around Karlsruhe. The results are depicted in Figs. E1 to E3.

Figure E1 shows the root-mean-squares (RMS) of the difference between the forecasted reference methane profile and forecasted profiles that are dislocated with respect to the reference by different spatial distances and time differences. These are the square root values of the diagonal entries of the respective dislocation covariance matrices ($S_{\Delta_{di}}^1$, we use here the
 780 superscript ¹ for logarithmic scale, because we work with relative covariances: $\Delta \ln x \approx \Delta x/x$). The dashed black lines indicate our collocation threshold values used for the combination of TROPOMI and IASI (TROPOMI and IASI are only combined as long as the horizontal distance of their ground pixels is within 50 km and the time difference is within 6 hours). Naturally, the respective RMS values are increasing with increasing horizontal distance and time difference. The values are largest in a small layer close to the surface and in the stratosphere, but relatively small in the free troposphere. For a horizontal
 785 dislocation of 50 km the RMS value is about 2% very close to the surface, between 0.3 and 0.5% for the rest of the troposphere and then it increases again to about 2% above 25 km altitude. For a time difference of 6 hours the RMS value is about 2.5% in a very small layer above ground, 0.6-0.8% in the free troposphere below 10 km, and it reaches about 1.5% at 15 km and 3.5% at 30 km altitude.

Figure E2 reveals to what extent the dislocation uncertainties as shown in Fig. E1 are vertically correlated. Depicted are
 790 the vertical correlations for the example of a spatial dislocation of 50 km (Fig. E2a) and a temporal dislocation of 6 hours (Fig. E2b). We observe that for both spatial and temporal dislocations the vertical behavior of the vertical correlation length




Figure E2. Characteristics of the vertical dependencies covered by the matrices $\mathbf{S}_{\Delta_{dl}}^1$. Shown are the vertical correlation matrices for the difference between the reference CH_4 profile (location 49.1°N and 8.4°E , corresponding to the location of Karlsruhe) and profiles dislocated with respect to the reference: (a) horizontal dislocation of 50 km; (b) temporal dislocation of 6 hours.




Figure E3. Same as Fig. E1, but for column averaged data: total column, tropospheric partial column, and UTLS partial column.

(distance where correlation coefficient decays to 0.5) is similar. The vertical correlation lengths are rather short close to the surface (only 100-200 m). They are larger for higher altitudes: in the middle/upper troposphere and in the stratosphere they increase to about 1000 m and 6000 m, respectively.

795 The dislocation error for total and partial columns can be calculated by resampling the spatial and temporal dislocation covariance matrices according to Eq. (D7) (more details see Appendix D). The result of these resampling calculations are shown for the spatial dislocation in Fig. E3a and for the temporal dislocation in Fig. E3b. Naturally the dislocation uncertainties increase for increasing horizontal distance and time difference. For our horizontal collocation threshold values of 50 km the uncertainty (RMS value is used as the metric) is about 0.2% for the total column data. For our time difference collocation
800 threshold of 6 hours it is about 0.3% for the total column data. For the tropospheric and upper tropospheric / lower stratospheric partial columns the respective relative uncertainties values are slightly larger.

E2 Impact on the combined CH_4 product

For calculating the error in the combined profile due to the horizontal and spatial dislocation between IASI and TROPOMI we substitute \hat{x}_I in Eq. (1) by $\hat{x}_I + \mathbf{A}_I \Delta_{dl}$, where Δ_{dl} is the dislocation uncertainty of CH_4 as shown in Figs. E1 and E2. This




Figure E4. Example of dislocation kernel $\mathbf{A}_{C,dl}^1$ – calculated according to Eq. (E3) – for the same late summer observation as used in the context of Figs. 1 to 3.

805 results in a new term in Eq. (1) that gives the dislocation error in the combined profile:

$$\Delta_{dl} \hat{x}_C^l = (\mathbf{I} - \mathbf{L}^{-1} \mathbf{m} \mathbf{a}_T^{*T}) \mathbf{A}_I^1 \Delta_{dl}^l. \quad (\text{E1})$$

The respective error covariance matrix is

$$\mathbf{S}_{\hat{x}_C, dl}^1 = \mathbf{A}_{C, dl}^1 \mathbf{S}_{\Delta_{dl}}^1 \mathbf{A}_{C, dl}^{1T}, \quad (\text{E2})$$

810 where $\mathbf{S}_{\Delta_{dl}}^1$ is the covariance matrix for the CH_4 dislocation uncertainty whose main characteristics are visualised in Figs. E1 and E2. Here

$$\mathbf{A}_{C, dl}^1 = (\mathbf{I} - \mathbf{L}^{-1} \mathbf{m} \mathbf{a}_T^{*T}) \mathbf{A}_I^1 \quad (\text{E3})$$

is the dislocation averaging kernel. Figure E4 shows an example of this dislocation averaging kernel. For the altitudes where the dislocation uncertainty of CH_4 are largest (close to ground and above 20 km, see Fig. E1) the dislocation kernel has rather low values (i.e. there the combination procedure has only limited sensitivity to the dislocation uncertainty).

815 We calculate the dislocation error covariance matrices according to Eq. (E2) for different locations and then determine the corresponding total and partial column dislocation errors by summing up the temporal and spatial dislocation covariances and performing an subsequent resampling of the covariance matrices according to Eq. (D7) (more details see Appendix D). Figure E5 depicts this dislocation error in comparison to the noise error (respective resampling of the covariance matrices obtained by Eq. (5), see also Fig. 6). We focus here on three different latitudinal locations: Sodankylä (northern high latitudes), Darwin
820 (low latitudes), and Lauder (southern middle latitudes). We find that for the northern high latitude site (where horizontal and temporal dislocation are of similar importance) but also for the tropical and southern hemispheric middle latitude sites (where the temporal dislocation is dominating), the dislocation uncertainty is generally much smaller than the noise error.




Figure E5. Comparison of the dislocation error (due to the CH₄ dislocation uncertainty) and the noise error (an example of the typical temporal dependencies of the noise error is shown in Fig. 6). The comparison is depicted for a northern high latitude location (Sodankylä, violet crosses), a tropical location (Darwin, orange crosses), and a southern middle latitudinal location (Lauder, red crosses). (a) Total column product; (b) Tropospheric partial column product; (c) UTLS partial column product.

Appendix F: Data comparability

The satellite data products are representative for broad vertical layers of the atmosphere (see averaging kernels as shown in Figs. 2 and 3). Also the TCCON and AirCore reference data are sensitive to atmospheric CH₄ at different vertical regions. If we furthermore assume that the TCCON and the AirCore data offer a stable absolute calibration reference, their inter-comparisons with the satellite data as shown in Sect. 4 can in principle be used for empirically validating the characteristics (sensitivity and error) of the satellite data products. The level of agreement that can be expected between the reference data and the satellite products depends on the reliability of the references and the characteristics of the satellite data products. In the following Appendices F1 and F2 we estimate the reliability of the TCCON and AirCore data, respectively, to serve as reference for the satellite data products. Then in Appendix F3 we show that the results of the inter-comparison as shown in the context of Figs. 9 to 12 are in a reasonable agreement with the reliability of the references and the characteristics of the different satellite data products. This confirms the validity of the sensitivity of the satellite data products as shown in Sect. 3.2 and the validity of the errors of the satellite data as documented in Sect. 3.3.

835 F1 TCCON versus satellite

For estimating the reliability of the TCCON data as reference for the satellite data products we consider the TCCON retrieval noise errors, the incomparableness of TCCON and satellite data caused by their different averaging kernels, and the collocation mismatch between the TCCON and the satellite observations. The total column uncertainty variance (the scalar S_{ref}^*) for using the TCCON data as reference for the satellite data can be estimated by:

$$840 \quad S_{\text{ref}}^* = S_{\Delta\text{TC}}^* + (\mathbf{a}^{*T} - \mathbf{a}_{\text{TC}}^{*T}) \mathbf{S}_{\Delta\mathbf{a}} (\mathbf{a}^{*T} - \mathbf{a}_{\text{TC}}^{*T})^T + \mathbf{a}_{\text{TC}}^{*T} (\mathbf{S}_{\Delta\text{h}} + \mathbf{S}_{\Delta\text{t}}) \mathbf{a}_{\text{TC}}^*, \quad (\text{F1})$$




Figure F1. Comparison of the CH₄ state obtained from the TROPOMI a priori model TM5 (\mathbf{x}_{TM5}) and the collocated CAMS high resolution forecasts (\mathbf{x}_{CAMS}). (a) RMS of the relative differences; (b) Matrix showing the correlations of TM5-CAMS differences at different altitudes.

The first term (the scalar $S_{\Delta\text{TC}}^*$) is the TCCON retrieval error variance (the TCCON error is provided with the TCCON data is typically 1‰). The second term accounts for the different averaging kernels. The row vectors \mathbf{a}^{*T} and $\mathbf{a}_{\text{TC}}^{*T}$ are the total column averaged mixing ratio kernels of the satellite and the TCCON retrievals, respectively (calculated according to Appendix D). The matrix $\mathbf{S}_{\Delta\mathbf{a}}$ describes the uncertainty covariances of the used a priori data, and the matrices $\mathbf{S}_{\Delta\mathbf{h}}$ and $\mathbf{S}_{\Delta\mathbf{t}}$ the covariances for horizontal and temporal collocation mismatches.

For estimating $\mathbf{S}_{\Delta\mathbf{a}}$ we use the difference between the CH₄ state as modelled by TM5 (\mathbf{x}_{TM5}) and provided by the high resolution CAMS forecast (\mathbf{x}_{CAMS} , e.g., Barré et al., 2021). Figure F1 shows the results of these calculations for the surroundings of Karlsruhe documented by the RMS values of the differences in the vertical profiles (Fig. F1a) and the vertical correlation matrix of the differences (Fig. F1b). We estimate an uncertainty of the TM5 a priori model of about 6% close to the surface, about 2% up throughout the middle troposphere, a gradual increase to about 7.5% between the UTLS and about 23 km altitude, and a maximum value of about 27% in the stratosphere at about 30 km. The vertical correlation lengths (altitude range where correlation coefficient decreases to about 0.5) is a few hundred metres close to the surface, about 5000 m in the middle troposphere, about 2500 m in the UTLS, and about 7500 m in the stratosphere above 30 km altitude. We find that this relatively large disagreement between the TM5 a priori data and the high resolution forecast of CAMS are significantly influenced by inconsistencies between TM5 and CAMS in the years 2019 and 2020: after 2018 the TM5 model shows an increase of about 1% per year, but the CAMS high resolution forecast shows no significant increase.

Figure F2 shows the value of the term $(\mathbf{a}^{*T} - \mathbf{a}_{\text{TC}}^{*T})(\mathbf{x}_{\text{TM5}} - \mathbf{x}_{\text{CAMS}})$ for the different satellite data products, i.e. it reveals the uncertainty in the comparison with TCCON data due to differences in the averaging kernels and the a priori model uncertainty, which in Eq. (F1) is represented by the square root value of the term $(\mathbf{a}^{*T} - \mathbf{a}_{\text{TC}}^{*T})\mathbf{S}_{\Delta\mathbf{a}}(\mathbf{a}^{*T} - \mathbf{a}_{\text{TC}}^{*T})^T$. Because the TROPOMI and the TCCON kernels have both a similar good column sensitivity throughout the troposphere, the respective uncertainty is generally within 0.1% (see black squares in Fig. F2). The same is true for the validation of the total column of the combined product (see blue crosses in Fig. F2). For the validation of the total column of the MUSICA IASI product




Figure F2. Error in the comparison of TCCON and satellite products due to the a priori model error and the different column sensitivities of the TCCON product and the satellite products. These values are calculated as $(\mathbf{a}^{*T} - \mathbf{a}_{\text{TCCON}}^{*T})(\mathbf{x}_{\text{TM5}} - \mathbf{x}_{\text{CAM5}})$ and are represented in Eq. (F1) by the square root value of the term $(\mathbf{a}^{*T} - \mathbf{a}_{\text{TCCON}}^{*T})\mathbf{S}_{\Delta\mathbf{a}}(\mathbf{a}^{*T} - \mathbf{a}_{\text{TCCON}}^{*T})^T$.

this error is larger, because the total column sensitivity of IASI is significantly different from the respective sensitivity of the TCCON product and the other satellite products (see Fig. 3a). For the comparison of the IASI and TCCON total column data we estimate that the error due to the different sensitivities (of IASI and TCCON) can occasionally be even above 2% (see red dots in Fig. F2). This error is largest to the end of the time series, because then the TM5 a priori model error is largest (increasing difference between the TM5 model and the CAMS high resolution forecast after 2018).

The collocation mismatch covariances $\mathbf{S}_{\Delta\mathbf{h}}$ and $\mathbf{S}_{\Delta\mathbf{t}}$ are the linear scale versions of the matrices $\mathbf{S}_{\Delta\mathbf{d}_i}^1$ (characterised in Figs. E1 and E2) interpolated to the actual temporal and horizontal mismatch of the satellite and the TCCON measurements. The effect of this collocation mismatch on the comparison of the total columns (i.e. the term $\mathbf{a}_{\text{TCCON}}^{*T}(\mathbf{S}_{\Delta\mathbf{h}} + \mathbf{S}_{\Delta\mathbf{t}})\mathbf{a}_{\text{TCCON}}^*$) is estimated to be between 0.1% and 0.4%.

F2 AirCore versus satellite

Similar to the TCCON data we estimate the reliability of the AirCore profile data as reference for the satellite observations. For this estimation we consider an AirCore measurement noise covariance ($\mathbf{S}_{\Delta\text{AC},\mathbf{n}}$). It is calculated assuming an uncertainty for altitudes with AirCore CH_4 data of 0.3% (Karion et al., 2010) and the uncertainty according to $\mathbf{S}_{\Delta\mathbf{a}}$ from Sect. F1 for all other altitudes. The outer diagonal elements are determined by assuming the same vertical correlation as derived for $\mathbf{S}_{\Delta\mathbf{a}}$. In addition, we consider uncertainties in the height attribution, which is according to Wagenhäuser et al. (2021) below 10 m close to ground, about 200 m at 20 km a.s.l. and about 1 km at 27 km a.s.l. We construct a respective height attribution uncertainty covariance ($\mathbf{S}_{\Delta\text{AC},\mathbf{v}}$) by assuming a very strong correlation of the height attribution uncertainties between different altitude levels. The temporal and spatial collocation mismatch covariance between the AirCore and the satellite observations ($\mathbf{S}_{\Delta\mathbf{h}}$ and $\mathbf{S}_{\Delta\mathbf{t}}$, respectively) are calculated as described in Sect. F1.

All the aforementioned uncertainties are independent and we can calculate the total uncertainty as:

$$\mathbf{S}_{\Delta\text{AC}} = \mathbf{S}_{\Delta\text{AC},\mathbf{n}} + \mathbf{S}_{\Delta\text{AC},\mathbf{v}} + \mathbf{S}_{\Delta\mathbf{h}} + \mathbf{S}_{\Delta\mathbf{t}}. \quad (\text{F2})$$

The reliability of the AirCore data – after its adjustment according to Eq. (7) – as reference for the MUSICA IASI and combined
 885 satellite data can then be estimated by:

$$\mathbf{S}_{\text{ref}}^1 = \mathbf{A}^1 \mathbf{S}_{\Delta\text{AC}}^1 \mathbf{A}^{1T}. \quad (\text{F3})$$

Here and in Eq. (F2) the covariances are determined for the full vertical profile. Respective covariances for total or partial
 columns can be derived according to Appendix D. The reliability for the TROPOMI total column averaged mixing ratio data
 can be calculated by $S_{\text{ref}}^* = \mathbf{a}_T^{*T} \mathbf{S}_{\Delta\text{AC}} \mathbf{a}_T^*$.

890 In order to get a reasonable number of collocated AirCore data we relax the collocation criteria: we require a temporal collocation
 within 6 hours and a spatial collocation within 500 km (see Sect. 4.2). In particular loose spatial collocation requirement
 results in theoretically large collocation mismatch uncertainties. For instance, Fig. E3 reveals that a spatial mismatch of 400 km
 clearly dominates the temporal mismatch, whose threshold is set to 6 hours, but it is actually only in the southern hemisphere
 typically greater than 3 hours. The spatial mismatch uncertainty also dominates AirCore uncertainties due to measurement
 895 noise and uncertain height attribution, i.e. it is the term that mostly affects the comparability of the AirCore and satellite mea-
 surements. We estimate a spatial mismatch error that has to be considered for the AirCore satellite inter-comparison of about
 0.5% for the total column data and of about 0.6% for the tropospheric and UTLS partial column data.

F3 Summary

The sum of the uncertainty (co)variance of using TCCON or AirCore as the reference (S_{ref}^* or $\mathbf{S}_{\text{ref}}^1$, see Sects. F1 and F2,
 900 respectively) and the noise and dislocation error (co)variances of the satellite data products (see Sect. 3.3 and 3.4, respectively)
 gives the covariance that can be theoretically expected for the scatter between the TCCON or AirCore reference data and the
 satellite data products.

Figure F3 shows the correlations between the theoretically expected scatter (mean value of the scatter expected for the
 individual data points) and the actually observed scatter (the hIPR68.2 of the individual differences between the reference data
 and the satellite data products). Shown is one data point for the XCH₄ comparisons with the TCCON references, for the XCH₄
 905 comparisons with the AirCore references, and further data points for the comparisons of the tropospheric and UTLS partial
 columns with the AirCore references. A detailed contemplation suggests that the scatter observed for the total column data of
 the TROPOMI and the combined data products as well as the scatter observed for the tropospheric partial column data of the
 IASI and combined products are slightly larger than their theoretically expected counterparts. On the contrary, in the UTLS
 910 the scatter observed in the IASI and combined data products seems to be a bit smaller than the theoretically expected scatter
 values. However, the data points group reasonably well around the one-to-one diagonal, i.e. there is overall a good agreement
 between the theoretically expected scatter and the actually observed scatter. This means that the inter-comparison results as
 shown in Sect. 4 confirm the satellite data quality characterisation of Sects. 3.2 - 3.4.




Figure F3. Theoretically predicted and observed 1σ scatter for the comparison of single pixel satellite data with individual TCCON and AirCore reference data. Black, red and blue colours represent TROPOMI, MUSICA IASI, and combined satellite data, respectively. The squares and vertical crosses are for XCH_4 comparisons with TCCON and AirCore references, respectively. The diagonal crosses and stars are for tropospheric and UTLS partial column comparisons, respectively, with AirCore references.

Author contributions. Matthias Schneider developed the idea for the optimal a posteriori combination of the level 2 remote sensing products and he prepared the figures and the manuscript. Benjamin Ertl developed and performed the continuous MUSICA IASI data processing, where he was supported by Matthias Schneider, Christopher J. Diekmann, Farahnaz Khosrawi, Amelie N. Röhling, Omaira E. García, and Eliezer Sepúlveda. Frank Hase developed the PROFFIT-nadir retrieval code used for the MUSICA IASI processing. Quiansi Tu supported the use of the CAMS high resolution data. Tobias Borsdorff, Jochen Landgraf, Alba Lorente, and André Butz are responsible for the TROPOMI processing and made TROPOMI data available. Huilin Chen and Rigel Kivi are responsible for the AirCore profile measurements over Sodankylä. Thomas Laemmel, Michel Ramonet, Cyril Crevoisier, and Jérôme Pernin are responsible for the AirCore profile measurements over Trainou. Martin Steinbacher and Frank Meinhardt are responsible for the GAW data of Jungfraujoch and Schauinsland, respectively. Rigel Kivi, Darko Dubravica, Frank Hase, Kimberly Strong, Debra Wunch, Thorsten Warneke, Coleen Roehl, Paul O. Wennberg, Isamu Morino, Laura T. Iraci, Kei Shiomi, Nicholas M. Deutscher, David W. T. Griffith, Voltaire A. Velazco, and David F. Pollard are responsible for the TCCON data. All authors supported the generation of the final version of this manuscript.

Competing interests. The authors declare that they have no conflict of interest

Acknowledgements. We would like to thank our colleague Thomas von Clarmann for his strong support in revising the equations in the Appendices of this manuscript.

This research has largely benefit from funds of the Deutsche Forschungsgemeinschaft (provided for the two projects MOTIV and TEDDY with IDs/Geschäftszeichen 290612604/GZ:SCHN1126/2-1 and 416767181/GZ:SCHN1126/5-1, respectively) and from support by the Eu-

930 ropean Space Agency in the context the "Sentinel-5p+Innovation (S5p+I) - Water Vapour Isotopologues (H2O-ISO)" activities. Furthermore,
we acknowledge funds from the Ministerio de Economía y Competividad from Spain for the project INMENSE (CGL2016-80688-P).

Important part of this work was performed on the supercomputer ForHLR funded by the Ministry of Science, Research and the Arts
Baden-Württemberg and by the German Federal Ministry of Education and Research. We also acknowledge the contribution of Teide High-
Performance Computing facilities. TeideHPC facilities are provided by the Instituto Tecnológico y de Energías Renovables (ITER), S.A
935 (teidehpc.iter.es).

The TROPOMI data processing was carried out on the Dutch National e-Infrastructure with the support of the SURF cooperative. The
presented material contains modified Copernicus data (2017, 2019).

The Eureka TCCON measurements were made at the Polar Environment Atmospheric Research Laboratory (PEARL) by the Canadian
Network for the Detection of Atmospheric Change, primarily supported by NSERC, ECCC, and CSA. The East Trout Lake TCCON station
940 is supported by CFI, ORF, and NSERC. The Karlsruhe TCCON station has been supported by the German Bundesministerium für Wirtschaft
und Energie (BMWi) via DLR under grants 50EE1711A to E and by the Helmholtz Society via the research program ATMO. The Park Falls
and Lamont TCCON sites are supported by and would like to acknowledge NASA's Carbon Cycle Science Program (grant # NNX17AE15G)
and OCO-2/-3 projects (primary grant # NNN12AA01C), respectively. The Burgos and Rikubetsu TCCON sites are supported in part by
the GOSAT series project. Burgos is supported in part by the Energy Development Corp. Philippines. Funding for the Edwards TCCON
945 station is provided by NASA's Earth Science Division. NMD is funded by ARC Future Fellowship FT180100327. Darwin and Wollongong
TCCON stations are supported by ARC grants DP160100598, LE0668470, DP140101552, DP110103118 and DP0879468, and Darwin
through NASA grants NAG5-12247 and NNG05-GD07G. The Lauder TCCON programme is core funded by NIWA through New Zealand's
Ministry of Business, Innovation and Employment.

The Trainou AirCore measurements have been supported by CEA, CNES, UVSQ, IPSL and the EU H2020 RINGO project (GA no. 730944),
950 and are part of the French consortium for Aircore measurements (LMD, LSCE, GSMA, CNES). The Sodankylä TCCON and AirCore mea-
surements have been supported via the ESA FRM4GHG project (under the grant agreement no. ESA-IPLPOE-LG-cl-LE-2015-1129) and
the EU H2020 RINGO project.

The CH₄ observations at Jungfraujoch were established as part of the Swiss National Air Pollution Monitoring Network and are supported
through ICOS-CH, which is funded by the Swiss National Science Foundation and in-house contributions.

955 We acknowledge the support by the Deutsche Forschungsgemeinschaft and the Open Access Publishing Fund of the Karlsruhe Institute
of Technology.

References

- 960 Barré, J., Aben, I., Agustí-Panareda, A., Balsamo, G., Bousserez, N., Dueben, P., Engelen, R., Inness, A., Lorente, A., McNorton, J., Peuch, V.-H., Radnoti, G., and Ribas, R.: Systematic detection of local CH₄ anomalies by combining satellite measurements with high-resolution forecasts, *Atmospheric Chemistry and Physics*, 21, 5117–5136, <https://doi.org/10.5194/acp-21-5117-2021>, <https://acp.copernicus.org/articles/21/5117/2021/>, 2021.
- 965 Borger, C., Schneider, M., Ertl, B., Hase, F., García, O. E., Sommer, M., Höpfner, M., Tjemkes, S. A., and Calbet, X.: Evaluation of MUSICA IASI tropospheric water vapour profiles using theoretical error assessments and comparisons to GRUAN Vaisala RS92 measurements, *Atmospheric Measurement Techniques*, 11, 4981–5006, <https://doi.org/10.5194/amt-11-4981-2018>, <https://www.atmos-meas-tech.net/11/4981/2018/>, 2018.
- Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J.-M., Tran, H., Kuze, A., Keppel-Aleks, G., Toon, G., Wunch, D., Wennberg, P., Deutscher, N., Griffith, D., Macatangay, R., Messerschmidt, J., Notholt, J., and Warneke, T.: Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophysical Research Letters*, 38, <https://doi.org/https://doi.org/10.1029/2011GL047888>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011GL047888>, 2011.
- 970 Ceccherini, S., Raspollini, P., and Carli, B.: Optimal use of the information provided by indirect measurements of atmospheric vertical profiles, *Opt. Express*, 17, 4944–4958, <https://doi.org/10.1364/OE.17.004944>, <http://www.opticsexpress.org/abstract.cfm?URI=oe-17-7-4944>, 2009.
- 975 Cortesi, U., Del Bianco, S., Ceccherini, S., Gai, M., Dinelli, B. M., Castelli, E., Oelhaf, H., Woiwode, W., Höpfner, M., and Gerber, D.: Synergy between middle infrared and millimeter-wave limb sounding of atmospheric temperature and minor constituents, *Atmospheric Measurement Techniques*, 9, 2267–2289, <https://doi.org/10.5194/amt-9-2267-2016>, <https://amt.copernicus.org/articles/9/2267/2016/>, 2016.
- Costantino, L., Cuesta, J., Emili, E., Coman, A., Foret, G., Dufour, G., Eremenko, M., Chailleux, Y., Beekmann, M., and Flaud, J.-M.: Potential of multispectral synergism for observing ozone pollution by combining IASI-NG and UVNS measurements from the EPS-SG satellite, *Atmospheric Measurement Techniques*, 10, 1281–1298, <https://doi.org/10.5194/amt-10-1281-2017>, <https://www.atmos-meas-tech.net/10/1281/2017/>, 2017.
- 980 Cuesta, J., Eremenko, M., Liu, X., Dufour, G., Cai, Z., Höpfner, M., von Clarmann, T., Sellitto, P., Foret, G., Gaubert, B., Beekmann, M., Orphal, J., Chance, K., Spurr, R., and Flaud, J.-M.: Satellite observation of lowermost tropospheric ozone by multispectral synergism of IASI thermal infrared and GOME-2 ultraviolet measurements over Europe, *Atmospheric Chemistry and Physics*, 13, 9675–9693, <https://doi.org/10.5194/acp-13-9675-2013>, <https://www.atmos-chem-phys.net/13/9675/2013/>, 2013.
- 985 Diekmann, C. J., Schneider, M., Ertl, B., Hase, F., García, O., Khosrawi, F., Sepúlveda, E., Knippertz, P., and Braesicke, P.: The global and multi-annual MUSICA IASI {H₂O, δD} pair dataset, *Earth System Science Data*, 13, 5273–5292, <https://doi.org/10.5194/essd-13-5273-2021>, <https://essd.copernicus.org/articles/13/5273/2021/>, 2021.
- Fu, D., Worden, J. R., Liu, X., Kulawik, S. S., Bowman, K. W., and Natraj, V.: Characterization of ozone profiles derived from Aura TES and OMI radiances, *Atmospheric Chemistry and Physics*, 13, 3445–3462, <https://doi.org/10.5194/acp-13-3445-2013>, <https://acp.copernicus.org/articles/13/3445/2013/>, 2013.
- 990 García, O. E., Schneider, M., Ertl, B., Sepúlveda, E., Borger, C., Diekmann, C., Wiegeler, A., Hase, F., Barthlott, S., Blumenstock, T., Raffalski, U., Gómez-Peláez, A., Steinbacher, M., Ries, L., and de Frutos, A. M.: The MUSICA IASI CH₄ and N₂O products and their comparison to HIPPO, GAW and NDACC FTIR references, *Atmospheric Measurement Techniques*, 11, 4171–4215, <https://doi.org/10.5194/amt-11-4171-2018>, <https://www.atmos-meas-tech.net/11/4171/2018/>, 2018.

- Griffith, D. W., Deutscher, N. M., Velazco, V. A., Wennberg, P. O., Yavin, Y., Keppel-Aleks, G., Washenfelder, R. A.,
995 Toon, G. C., Blavier, J.-F., Paton-Walsh, C., Jones, N. B., Kettlewell, G. C., Connor, B. J., Macatangay, R. C., Roehl,
C., Ryzcek, M., Glowacki, J., Culfan, T., and Bryant, G. W.: TCCON data from Darwin (AU), Release GGG2014.R0,
<https://doi.org/10.14291/TCCON.GGG2014.DARWIN01.R0/1149290>, <https://data.caltech.edu/records/269>, 2014a.
- Griffith, D. W., Velazco, V. A., Deutscher, N. M., Paton-Walsh, C., Jones, N. B., Wilson, S. R., Macatangay, R. C., Ket-
tlewell, G. C., Buchholz, R. R., and Riggensbach, M. O.: TCCON data from Wollongong (AU), Release GGG2014.R0,
1000 <https://doi.org/10.14291/TCCON.GGG2014.WOLLONGONG01.R0/1149291>, <https://data.caltech.edu/records/291>, 2014b.
- Hase, F., Blumenstock, T., Dohe, S., Groß, J., and Kiel, M.: TCCON data from Karlsruhe (DE), Release GGG2014.R1,
<https://doi.org/10.14291/TCCON.GGG2014.KARLSRUHE01.R1/1182416>, <https://data.caltech.edu/records/278>, 2015.
- Hasekamp, O., Lorente, A., Hu, H., Butz, A., van de Brugh, J., and Landgraf, J.: Algorithm Theoretical Baseline Document for Sentinel-5
Precursor methane retrieval, <http://www.tropomi.eu/documents/atbd/>, 2019.
- 1005 Hu, H., Hasekamp, O., Butz, A., Galli, A., Landgraf, J., van de Brugh, J., Borsdorff, T., Scheepmaker, R., and Aben, I.: The operational
methane retrieval algorithm for TROPOMI, *Atmospheric Measurement Techniques*, 9, 5423–5440, [https://doi.org/10.5194/amt-9-5423-](https://doi.org/10.5194/amt-9-5423-2016)
2016, <https://amt.copernicus.org/articles/9/5423/2016/>, 2016.
- Iraci, L. T., Podolske, J. R., Hillyard, P. W., Roehl, C., Wennberg, P. O., Blavier, J.-F., Landeros, J., Allen, N., Wunch, D., Zavaleta, J.,
Quigley, E., Osterman, G. B., Albertson, R., Dunwoody, K., and Boyden, H.: TCCON data from Edwards (US), Release GGG2014.R1,
1010 <https://doi.org/10.14291/TCCON.GGG2014.EDWARDS01.R1/1255068>, <https://data.caltech.edu/records/270>, 2016.
- Kalman, R. E.: A New Approach to Linear Filtering and Prediction Problems, *Journal of Basic Engineering*, 82, 35–45,
<https://doi.org/10.1115/1.3662552>, <https://doi.org/10.1115/1.3662552>, 1960.
- Karion, A., Sweeney, C., Tans, P., and Newberger, T.: AirCore: An Innovative Atmospheric Sampling System, *Journal of Atmospheric and
Oceanic Technology*, 27, 1839 – 1853, <https://doi.org/10.1175/2010JTECHA1448.1>, [https://journals.ametsoc.org/view/journals/atot/27/](https://journals.ametsoc.org/view/journals/atot/27/11/2010jtecha1448_1.xml)
1015 [11/2010jtecha1448_1.xml](https://journals.ametsoc.org/view/journals/atot/27/11/2010jtecha1448_1.xml), 2010.
- Kawakami, S., Ohyama, H., Arai, K., Okumura, H., Taura, C., Fukamachi, T., and Sakashita, M.: TCCON data from Saga (JP), Release
GGG2014.R0, <https://doi.org/10.14291/TCCON.GGG2014.SAGA01.R0/1149283>, <https://data.caltech.edu/records/288>, 2014.
- Keppens, A., Lambert, J.-C., Granville, J., Miles, G., Siddans, R., van Peet, J. C. A., van der A, R. J., Hubert, D., Verhoelst, T., Delcloo,
A., Godin-Beekmann, S., Kivi, R., Stübi, R., and Zehner, C.: Round-robin evaluation of nadir ozone profile retrievals: methodology
1020 and application to MetOp-A GOME-2, *Atmospheric Measurement Techniques*, 8, 2093–2120, <https://doi.org/10.5194/amt-8-2093-2015>,
<https://www.atmos-meas-tech.net/8/2093/2015/>, 2015.
- Kivi, R. and Heikkinen, P.: Fourier transform spectrometer measurements of column CO₂ at Sodankylä, Finland, *Geoscientific Instrumenta-
tion, Methods and Data Systems*, 5, 271–279, <https://doi.org/10.5194/gi-5-271-2016>, <https://gi.copernicus.org/articles/5/271/2016/>, 2016.
- Kivi, R., Heikkinen, P., and Kyrö, E.: TCCON data from Sodankylä (FI), Release GGG2014.R0,
1025 <https://doi.org/10.14291/TCCON.GGG2014.SODANKYLA01.R0/1149280>, <https://data.caltech.edu/records/289>, 2014.
- Krol, M., Houweling, S., Bregman, B., van den Broek, M., Segers, A., van Velthoven, P., Peters, W., Dentener, F., and Bergamaschi, P.:
The two-way nested global chemistry-transport zoom model TM5: algorithm and applications, *Atmospheric Chemistry and Physics*, 5,
417–432, <https://doi.org/10.5194/acp-5-417-2005>, <https://acp.copernicus.org/articles/5/417/2005/>, 2005.
- Landgraf, J. and Hasekamp, O. P.: Retrieval of tropospheric ozone: The synergistic use of thermal infrared emission and ultraviolet reflectivity
1030 measurements from space, *Journal of Geophysical Research: Atmospheres*, 112, [https://doi.org/https://doi.org/10.1029/2006JD008097](https://doi.org/10.1029/2006JD008097),
<https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JD008097>, 2007.

- Landgraf, J., Butz, A., Hasekamp, O., Hu, H., , and aan de Brugh, J.: Sentinel 5 L2 Prototype Processors, Algorithm Theoretical Baseline Document: Methane Retrieval, 2019.
- 1035 Lorente, A., Borsdorff, T., Butz, A., Hasekamp, O., aan de Brugh, J., Schneider, A., Wu, L., Hase, F., Kivi, R., Wunch, D., Pollard, D. F., Shiomi, K., Deutscher, N. M., Velazco, V. A., Roehl, C. M., Wennberg, P. O., Warneke, T., and Landgraf, J.: Methane retrieved from TROPOMI: improvement of the data product and validation of the first 2 years of measurements, *Atmospheric Measurement Techniques*, 14, 665–684, <https://doi.org/10.5194/amt-14-665-2021>, <https://amt.copernicus.org/articles/14/665/2021/>, 2021.
- 1040 Luo, M., Read, W., Kulawik, S., Worden, J., Livesey, N., Bowman, K., and Herman, R.: Carbon monoxide (CO) vertical profiles derived from joined TES and MLS measurements, *Journal of Geophysical Research: Atmospheres*, 118, 10,601–10,613, <https://doi.org/https://doi.org/10.1002/jgrd.50800>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/jgrd.50800>, 2013.
- Morino, I., Yokozeki, N., Matsuzaki, T., and Horikawa, M.: TCCON data from Rikubetsu (JP), Release GGG2014.R2, <https://doi.org/10.14291/TCCON.GGG2014.RIKUBETSU01.R2>, <https://data.caltech.edu/records/957>, 2018.
- 1045 Pandey, S., Houweling, S., Krol, M., Aben, I., Chevallier, F., Dlugokencky, E. J., Gatti, L. V., Gloor, E., Miller, J. B., Detmers, R., Machida, T., and Röckmann, T.: Inverse modeling of GOSAT-retrieved ratios of total column CH₄ and CO₂ for 2009 and 2010, *Atmospheric Chemistry and Physics*, 16, 5043–5062, <https://doi.org/10.5194/acp-16-5043-2016>, <https://acp.copernicus.org/articles/16/5043/2016/>, 2016.
- Parker, R. J., Webb, A., Boesch, H., Somkuti, P., Barrio Guillo, R., Di Noia, A., Kalaitzi, N., Anand, J. S., Bergamaschi, P., Chevallier, F., Palmer, P. I., Feng, L., Deutscher, N. M., Feist, D. G., Griffith, D. W. T., Hase, F., Kivi, R., Morino, I., Notholt, J., Oh, Y.-S., Ohyama, H., Petri, C., Pollard, D. F., Roehl, C., Sha, M. K., Shiomi, K., Strong, K., Sussmann, R., Té, Y., Velazco, V. A., Warneke, T., Wennberg, P. O., and Wunch, D.: A decade of GOSAT Proxy satellite CH₄ observations, *Earth System Science Data*, 12, 3383–3412, <https://doi.org/10.5194/essd-12-3383-2020>, <https://essd.copernicus.org/articles/12/3383/2020/>, 2020.
- 1050 Pollard, D. F., Robinson, J., and Shiona, H.: TCCON data from Lauder (NZ), Release GGG2014.R0, <https://doi.org/10.14291/TCCON.GGG2014.LAUDER03.R0>, <https://data.caltech.edu/records/1220>, 2019.
- Rodgers, C.: *Inverse Methods for Atmospheric Sounding: Theory and Praxis*, World Scientific Publishing Co., Singapore, 2000.
- 1055 Rodgers, C. and Connor, B.: Intercomparison of remote sounding instruments, *J. Geophys. Res.*, 108, 4116–4129, <https://doi.org/10.1029/2002JD002299>, 2003.
- Schneider, M., Wiegeler, A., Barthlott, S., González, Y., Christner, E., Dyroff, C., García, O. E., Hase, F., Blumenstock, T., Sepúlveda, E., Mengistu Tsidu, G., Takele Kenea, S., Rodríguez, S., and Andrey, J.: Accomplishments of the MUSICA project to provide accurate, long-term, global and high-resolution observations of tropospheric {H₂O,δD} pairs – a review, *Atmospheric Measurement Techniques*, 9, 2845–2875, <https://doi.org/10.5194/amt-9-2845-2016>, <http://www.atmos-meas-tech.net/9/2845/2016/>, 2016.
- 1060 Schneider, M., Ertl, B., Diekmann, C. J., Khosrawi, F., Weber, A., Hase, F., Höpfner, M., García, O. E., Sepúlveda, E., and Kinnison, D.: Design and description of the MUSICA IASI full retrieval product, *Earth System Science Data*, 14, 709–742, <https://doi.org/10.5194/essd-14-709-2022>, <https://essd.copernicus.org/articles/14/709/2022/>, 2022.
- 1065 Sepúlveda, E., Schneider, M., Hase, F., Barthlott, S., Dubravica, D., García, O. E., Gomez-Pelaez, A., González, Y., Guerra, J. C., Gisi, M., Kohlhepp, R., Dohe, S., Blumenstock, T., Strong, K., Weaver, D., Palm, M., Sadeghi, A., Deutscher, N. M., Warneke, T., Notholt, J., Jones, N., Griffith, D. W. T., Smale, D., Brailsford, G. W., Robinson, J., Meinhardt, F., Steinbacher, M., Aalto, T., and Worthy, D.: Tropospheric CH₄ signals as observed by NDACC FTIR at globally distributed sites and comparison to GAW surface in situ measurements, *Atmospheric Measurement Techniques*, 7, 2337–2360, <https://doi.org/10.5194/amt-7-2337-2014>, <https://amt.copernicus.org/articles/7/2337/2014/>, 2014.

- Sha, M. K., Langerock, B., Blavier, J.-F. L., Blumenstock, T., Borsdorff, T., Buschmann, M., Dehn, A., De Mazière, M., Deutscher, N. M.,
1070 Feist, D. G., García, O. E., Griffith, D. W. T., Grutter, M., Hannigan, J. W., Hase, F., Heikkinen, P., Hermans, C., Iraci, L. T., Jeseck,
P., Jones, N., Kivi, R., Kumps, N., Landgraf, J., Lorente, A., Mahieu, E., Makarova, M. V., Mellqvist, J., Metzger, J.-M., Morino, I.,
Nagahama, T., Notholt, J., Ohyama, H., Ortega, I., Palm, M., Petri, C., Pollard, D. F., Rettinger, M., Robinson, J., Roche, S., Roehl, C. M.,
Röhling, A. N., Rousogonous, C., Schneider, M., Shiomi, K., Smale, D., Stremme, W., Strong, K., Sussmann, R., Té, Y., Uchino, O.,
1075 Velazco, V. A., Vigouroux, C., Vrekoussis, M., Wang, P., Warneke, T., Wizenberg, T., Wunch, D., Yamanouchi, S., Yang, Y., and Zhou,
M.: Validation of methane and carbon monoxide from Sentinel-5 Precursor using TCCON and NDACC-IRWG stations, *Atmospheric
Measurement Techniques*, 14, 6249–6304, <https://doi.org/10.5194/amt-14-6249-2021>, <https://amt.copernicus.org/articles/14/6249/2021/>,
2021.
- Sherlock, V., Connor, B., Robinson, J., Shiona, H., Smale, D., and Pollard, D. F.: TCCON data from Lauder (NZ), 125HR, Release
GGG2014.R0, <https://doi.org/10.14291/TCCON.GGG2014.LAUDER02.R0/1149298>, <https://data.caltech.edu/records/281>, 2014.
- 1080 Strong, K., Roche, S., Franklin, J. E., Mendonca, J., Lutsch, E., Weaver, D., Fogal, P. F., Drummond, J. R., Batchelor, R., and Lindenmaier,
R.: TCCON data from Eureka (CA), Release GGG2014.R3, <https://doi.org/10.14291/TCCON.GGG2014.EUREKA01.R3>, <https://data.caltech.edu/records/1171>, 2019.
- Velazco, V. A., Morino, I., Uchino, O., Hori, A., Kiel, M., Bukosa, B., Deutscher, N. M., Sakai, T., Nagai, T., Bagtasa, G., Izumi, T., Yoshida,
Y., and Griffith, D. W. T.: TCCON Philippines: First Measurement Results, Satellite Data and Model Comparisons in Southeast Asia,
1085 *Remote Sensing*, 9, <https://doi.org/10.3390/rs9121228>, <https://www.mdpi.com/2072-4292/9/12/1228>, 2017.
- von Clarmann, T., Degenstein, D. A., Livesey, N. J., Bender, S., Braverman, A., Butz, A., Compernelle, S., Damadeo, R., Dueck, S., Eriksson,
P., Funke, B., Johnson, M. C., Kasai, Y., Keppens, A., Kleinert, A., Kramarova, N. A., Laeng, A., Langerock, B., Payne, V. H., Rozanov,
A., Sato, T. O., Schneider, M., Sheese, P., Sofieva, V., Stiller, G. P., von Savigny, C., and Zawada, D.: Overview: Estimating and report-
ing uncertainties in remotely sensed atmospheric composition and temperature, *Atmospheric Measurement Techniques*, 13, 4393–4436,
1090 <https://doi.org/10.5194/amt-13-4393-2020>, <https://amt.copernicus.org/articles/13/4393/2020/>, 2020.
- Wagenhäuser, T., Engel, A., and Sitals, R.: Testing the altitude attribution and vertical resolution of AirCore measurements with a new spiking
method, *Atmospheric Measurement Techniques Discussions*, 2021, 1–18, <https://doi.org/10.5194/amt-2020-461>, <https://amt.copernicus.org/preprints/amt-2020-461/>, 2021.
- Warneke, T., Messerschmidt, J., Notholt, J., Weinzierl, C., Deutscher, N. M., Petri, C., and Grupe, P.: TCCON data from Orléans (FR),
1095 Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.ORLEANS01.R1>, <https://data.caltech.edu/records/1301>, 2019.
- Warner, J. X., Yang, R., Wei, Z., Carminati, F., Tangborn, A., Sun, Z., Lahoz, W., Attié, J.-L., El Amraoui, L., and Duncan, B.: Global carbon
monoxide products from combined AIRS, TES and MLS measurements on A-train satellites, *Atmospheric Chemistry and Physics*, 14,
103–114, <https://doi.org/10.5194/acp-14-103-2014>, <https://acp.copernicus.org/articles/14/103/2014/>, 2014.
- Wennberg, P. O., Wunch, D., Roehl, C. M., Blavier, J.-F., Toon, G. C., and Allen, N. T.: TCCON data from Lamont (US), Release
1100 GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.LAMONT01.R1/1255070>, <https://data.caltech.edu/records/279>, 2016.
- Wennberg, P. O., Roehl, C. M., Wunch, D., Toon, G. C., Blavier, J.-F., Washenfelder, R., Keppel-Aleks, G., Allen, N. T., and Ayers, J.:
TCCON data from Park Falls (US), Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.PARKFALLS01.R1>, <https://data.caltech.edu/records/295>, 2017.
- Worden, J., Liu, X., Bowman, K., Chance, K., Beer, R., Eldering, A., Gunson, M., and Worden, H.: Improved tropospheric ozone profile
1105 retrievals using OMI and TES radiances, *Geophysical Research Letters*, 34, <https://doi.org/https://doi.org/10.1029/2006GL027806>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006GL027806>, 2007.

- Worden, J. R., Turner, A. J., Bloom, A., Kulawik, S. S., Liu, J., Lee, M., Weidner, R., Bowman, K., Frankenberg, C., Parker, R., and Payne, V. H.: Quantifying lower tropospheric methane concentrations using GOSAT near-IR and TES thermal IR measurements, *Atmospheric Measurement Techniques*, 8, 3433–3445, <https://doi.org/10.5194/amt-8-3433-2015>, <https://amt.copernicus.org/articles/8/3433/2015/>, 2015.
- 1110 Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369, 2087–2112, <https://doi.org/10.1098/rsta.2010.0240>, <https://royalsocietypublishing.org/doi/abs/10.1098/rsta.2010.0240>, 2011a.
- 1115 Wunch, D., Wennberg, P. O., Toon, G. C., Connor, B. J., Fisher, B., Osterman, G. B., Frankenberg, C., Mandrake, L., O'Dell, C., Ahonen, P., Biraud, S. C., Castano, R., Cressie, N., Crisp, D., Deutscher, N. M., Eldering, A., Fisher, M. L., Griffith, D. W. T., Gunson, M., Heikkinen, P., Keppel-Aleks, G., Kyrö, E., Lindenmaier, R., Macatangay, R., Mendonca, J., Messerschmidt, J., Miller, C. E., Morino, I., Notholt, J., Oyafuso, F. A., Rettinger, M., Robinson, J., Roehl, C. M., Salawitch, R. J., Sherlock, V., Strong, K., Sussmann, R., Tanaka, T., Thompson, D. R., Uchino, O., Warneke, T., and Wofsy, S. C.: A method for evaluating bias in global measurements of CO₂ total columns from space, *Atmospheric Chemistry and Physics*, 11, 12 317–12 337, <https://doi.org/10.5194/acp-11-12317-2011>, <https://acp.copernicus.org/articles/11/12317/2011/>, 2011b.
- 1120 Wunch, D., Mendonca, J., Colebatch, O., Allen, N. T., Blavier, J.-F., Roche, S., Hedelius, J., Neufeld, G., Springett, S., Worthy, D., Kessler, R., and Strong, K.: TCCON data from East Trout Lake, SK (CA), Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.EASTTROUTLAKE01.R1>, <https://data.caltech.edu/records/362>, 2018.