

A kriging-based analysis of cloud Liquid Water Content using CloudSat data

Atmospheric Measurement Techniques

March 4, 2022

J.-M. Lalande^{1,*}, G. Bourmaud¹,
P. Minvielle², J.-F. Giovannelli¹
`jean-marie.lalande@meteo.fr`

¹IMS (Univ. Bordeaux, CNRS, Bordeaux INP), 33400 Talence, France

²CESTA, DAM, CEA, 33114 Le Barp, France

*Now at CNRM, Université de Toulouse, Météo-France, CNRS, Lannion, France

Contents

1	Reviewer 1	1
	Reviewer 1 Comment 1	1
	Reviewer 1 Comment 2	3
2	Reviewer 2	4

Abstract

We thank the two anonymous reviewers for their efforts and feedback.

1 Reviewer 1

This article proposed a statistical spatio-temporal kriging-based approach that is able to interpolate/predict from the dataset and provide uncertainties. The topic is of interest to the readership of this journal, this study is well planned, and the mathematics appears correct and keeps at an appropriate level. However, this paper still need to be improved with moderate revisions:

Reviewer 1 Comment 1

Kriging methods have been widely used in spatial and temporal interpolation for meteorological elements, in the meantime, many improved kriging methods have been proposed, such as Universal Kriging, Co-Kriging, Disjunctive Kriging and so on. Therefore, this paper should give a brief overview of these improved methods, and highlight the advance of this paper method compared with the existing methods.

Response

We agree with the reviewer that the manuscript would greatly benefit from a better overview of kriging methods applied to meteorological parameters, as well as more references about the kriging literature in general. In that sense, we propose to add the following paragraph in the manuscript (in the introduction) in order to address this comment:

A number of statistical methods dedicated to the analysis of spatial and spatio-temporal data have been developed over the years taking into account the spatial and/or temporal correlation of the observations Ripley (1981); Cressie (1993). Among them, the kriging estimator was initially introduced by Krige (1951), from which it takes its name, to estimate the gold distribution at the Witwatersrand reef complex in South Africa based on samples from boreholes. It was then formalized mathematically by Matheron (1963) in the context of mining geology. Afterwards, the kriging estimator spread to many other areas of sciences Wackernagel (2013) (hydrogeology, geotechnics, agronomy, air quality, fishery, epidemiology, water and soil pollution, noise, *etc.*). Best-known kriging techniques are Simple Kriging, which assumes stationarity of the 1st-order with a known mean and Ordinary Kriging, where the mean is unknown. Since its first development, kriging techniques have largely evolved, and a number of new kriging techniques have been developed Chiles and Delfiner (1999); Cressie (1993); Cressie and Wikle (2015). In the field of meteorology, the kriging estimator have mostly been used in order to estimate precipitation accumulation from rain gauges Nour et al. (2006); Belo-Pereira et al. (2011) and in combination with satellite-derived precipitation Jewell and Gaussiat (2015); Verdin et al. (2016); Varouchakis et al. (2021) as well as the estimation of aerosols concentration in the air from in-situ observations Park (2016). It has also been used for the estimation of temperature from in-situ measurements Heuvelink et al. (2012); Didari and Zand-Parsa (2018), from satellite observations Florio et al. (2004) or a combination of them Didari and Zand-Parsa (2018), and for the estimation of surface properties from remote measurements der Meer (2012); Zakeri and Mariethoz (2021).

The introduction has been slightly reorganized to include this paragraph. Moreover, as the reviewer stated, some of the extended kriging estimators include:

1. the Universal Kriging assumes a non-stationary random function at the 1st-order, but 2nd order stationary, hence the mean model is written such as $\mu(s) = \beta^t f(s)$ with parameters acting linearly with a set of linear and polynomial function such as $f(s) = [1, s, s^2, \dots]$ in the model. In that sense, our estimator is similar to the Universal Kriging estimator as our mean is a linear combination of cosinus functions. However, in order to estimate the phase parameters of the cosinus, which are non-linears, we had to modify the estimation approach used in Universal Kriging. Our contribution, then, rest upon a mean adapted to our problem that includes the periodicity of the observations.
2. Co-Kriging which is applied to the estimation of multivariate random variable that are supposedly statistically correlated. In our case we aim at estimating a single variables (Liquid Water Content) from spatio-temporal observations so that we don't need to resort to this approach. Future development could include the joint estimation of LWC/IWC. Co-kriging is an extension of ordinary kriging. Furthermore, co-kriging can become computationally intensive to use as it requires to estimate the covariance

for each variables as well as their cross-covariance, this becomes rapidly difficult when dealing with big amount of data (such as satellite observations).

3. Disjunctive Kriging is a kriging techniques that has been available for over 50 years and falls in the field of nonlinear geostatistics. This methods is dedicated to the estimation of some functions of the quantity of interest $f[\phi(x_0)]$ instead of $\phi(x_0)$. In that case considering linear combination of the observation is not sufficient and nonlinear kriging methods have been proposed. Nonlinear kriging was designed to face the more and more complicated mining evaluation problems, specifically, instead of estimating the proportion of ore in a block, the problem was to estimate if the block in question would exceed a threshold between ore and waste Rivoirard (1994). Disjunctive Kriging arises as the co-kriging of indicator functions that are used to express any function f of the quantity of interest. The original problem does not require to resort to such approach but we note that this estimator should be considered to solve other interesting scientific questions (*i.e.* estimate if a specific cloud is going to exceed some LWC threshold, classification problems, *etc.*)

A number of other kriging estimator has been developed over the years so that it is nearly impossible to give an exhaustive listing. Our kriging estimator is an adaptation of the Universal Kriging estimator to include a more complex mean for the quantity of interest. The two other kriging estimators cited by the reviewer (co-kriging and disjunctive kriging), while they could be investigated in our context (*i.e.*, CloudSat observations), serve different purposes and are not specifically adapted to our problem. However, in our perspective, we mention some other estimation problems (*i.e.*, joint estimation IWC/LWC, integrate different cloud types) for which these estimators should be considered (we slightly modified the Conclusion and Perspective part of the manuscript in that sense). Reviewing the literature, when we first got our hands on this study, we found out confusing the difference in treatment of the model parameters associated with the mean and those associated with the covariance. One of the starting point of our analysis, along with the specific application to the CloudSat observations, was to specify a unified treatment of the mean and the covariance parameters and then to estimate these parameters, with the MAP estimator, before applying the kriging equation. We believe such an approach gives a better estimation of the variance of the estimation error.

Reviewer 1 Comment 2

The experimental section is lack of adequate contrast experiments with other existing interpolation methods (especially the representative improved kriging methods).

Response

Our study was initially motivated to demonstrate the feasibility of the kriging estimator in the complex situation of the estimation of cloud liquid water content at spatial location not sampled by Cloudsat observations and to derive a generative model of the LWC distribution, including the uncertainties associated with estimated quantity of interest. Moreover, we also wanted to give an in-depth evaluation of the kriging estimator in order for any interested reader to be able to reproduce this specific experiment. A number of other studies have investigated the performance of kriging estimator in comparison with other interpolation/extrapolation methods Lam (1983); Caruso and Quarta (1998); Stein (1999), we believe

such comparison doesn't fall in the scope of this particular paper and would considerably lengthen the study at the cost of clarity.

Moreover, our philosophy was to go back to the basics in a context never treated before and designed a new version of the kriging estimator for our purposes. One of our main concern, in that sense, was to increase the interpretability of the model parameters by carefully choosing them from the exploratory analysis. The detailed discussion on the estimated parameters could hardly be extended to other interpolation or kriging methods as the model parameters would definitely be different and hardly comparable. Our opinion is that this should preferably be done in a different study.

2 Reviewer 2

In this article, the authors use a kriging method to interpolate measurements and predict cloud properties of LWC from CloudSat satellite retrievals. Because of the polar orbit and the fixed-nadir radar measurement, CloudSat provides the most spatiotemporally coarse measurements of any satellite cloud radar. This is a frequent impediment for users of CloudSat for climatologies, and many prior studies have dealt with this issue with comparatively simple methods, such as gaussian means and standard deviations. There is a clear benefit in an advanced statistical method that could provide well-described (that is, with numerical uncertainties) predictions of unsampled regions based off of neighboring retrievals, so I recommend this paper for publication.

The majority of this paper discusses the kriging method, application, and resulting optimal estimation. I do not have much of a background in Kriging, but it seems like the other reviewer has already gone in-depth with recommendations on this matter, so I will not add anything else. I do not find any other issues with the results.

Response

We thank the reviewer for his laudatory comments. As being asked, we refer him to the answer to the first reviewer.

References

Belo-Pereira, M., Dutra, E., and Viterbo, P. (2011). Evaluation of global precipitation data sets over the iberian peninsula. *Journal of Geophysical Research: Atmospheres*, 116(D20).

Caruso, C. and Quarta, F. (1998). Interpolation methods comparison. *Computers and Mathematics with Applications*, 35(12):109–126.

Chiles, J.-P. and Delfiner, P. (1999). *Geostatistics: modeling spatial uncertainty*. Wiley, New York.

Cressie, N. (1993). *Statistics for Spatial Data*. Wiley Series in Probability and Statistics. Wiley.

Cressie, N. and Wikle, C. (2015). *Statistics for Spatio-Temporal Data*. Wiley.

der Meer, F. V. (2012). Remote-sensing image analysis and geostatistics. *International Journal of Remote Sensing*, 33(18):5644–5676.

Didari, S. and Zand-Parsa, S. (2018). Enhancing estimation accuracy of daily maximum, minimum, and mean air temperature using spatio-temporal ground-based and remote-sensing data in southern iran. *International Journal of Remote Sensing*, 39(19):6316–6339.

Florio, E. N., Lele, S. R., Chang, Y. C., Stern, R., and Glass, G. E. (2004). Integrating avhrr satellite data and noaa ground observations to predict surface air temperature: a statistical approach. *International Journal of Remote Sensing*, 25(15):2979–2994.

Heuvelink, G., Griffith, D., Hengl, T., and Melles, S. (2012). Sampling design optimization for space-time kriging. *Spatio-Temporal Design: Advances in Efficient Data Acquisition*, pages 207–230.

Jewell, S. and Gaußiat, N. (2015). An assessment of kriging based rain-gauge-radar merging techniques. *Quarterly Journal of the Royal Meteorological Society*, 141.

Krige, D. (1951). A statistical approach to some basic mine valuation problems on the witwatersrand. *Journal of the Southern African Institute of Mining and Metallurgy*, 52(6):119–139.

Lam, N. (1983). Spatial interpolation methods: a review. *American Cartographer*, 10:129–149.

Matheron, G. (1963). Principles of geostatistics. *Economic Geology*, 58(8):1246–1266.

Nour, M., Smit, D., and Gamal El-Din, M. (2006). Geostatistical mapping of precipitation: implications for rain gauge network design. *Water Science and Technology*, 53(10):101–110.

Park, N.-W. (2016). Time-series mapping of pm10 concentration using multi-gaussian space-time kriging: A case study in the seoul metropolitan area, korea. *Advances in Meteorology*, 2016:1–10.

Ripley, B. D. (1981). *Spatial statistics*. Wiley New York.

Rivoirard, J. (1994). *Introduction to Disjunctive Kriging and Non-linear Geostatistics*. Spatial information systems. Clarendon Press.

Stein, M. L. (1999). *Interpolation of spatial data*. Springer Series in Statistics. Springer-Verlag, New York. Some theory for Kriging.

Varouchakis, E. A., Kamińska-Chuchmala, A., Kowalik, G., Spanoudaki, K., and Graña, M. (2021). Combining geostatistics and remote sensing data to improve spatiotemporal analysis of precipitation. *Sensors*, 21(9).

Verdin, A., Funk, C., Rajagopalan, B., and Kleiber, W. (2016). Kriging and local polynomial methods for blending satellite-derived and gauge precipitation estimates to support hydrologic early warning systems. *IEEE Transactions on Geoscience and Remote Sensing*, 54(5):2552–2562.

Wackernagel, H. (2013). *Multivariate Geostatistics: An Introduction with Applications*. Springer Berlin Heidelberg.

Zakeri, F. and Mariethoz, G. (2021). A review of geostatistical simulation models applied to satellite remote sensing: Methods and applications. *Remote Sensing of Environment*, 259:112381.

A kriging-based analysis of cloud Liquid Water Content using CloudSat data

Jean-Marie Lalande^{1,*}, Guillaume Bourmaud¹, Pierre Minvielle², and Jean-François Giovannelli¹

¹IMS (Univ. Bordeaux, CNRS, Bordeaux INP), 33400 Talence, France

²CESTA, DAM, CEA, 33114 Le Barp, France

*Now at CNRM, Université de Toulouse, Météo-France, CNRS, Lannion, France

Correspondence: Jean-Marie Lalande (jean-marie.lalande@meteo.fr)

Abstract. Spatiotemporal statistical learning has received increased attention in the past decade, due to spatially and temporally indexed data proliferation, especially collected from satellite remote sensing. In the mean time, observational studies of clouds are recognized as an important step to improve cloud representation in weather and climate models. Since 2006, the satellite CloudSat of NASA carries a 94 GHz cloud profiling radar and is able to retrieve, from radar reflectivity, microphysical parameter distribution such as water or ice content. The collected data is piled up with the successive satellite orbits of nearly two hours, leading to a large compressed database of 2 Tb (<http://cloudsat.atmos.colostate.edu/>).

These observations give the opportunity to extend the cloud microphysical properties beyond the actual measurement locations using an interpolation and prediction algorithm. In order to do so, we introduce a statistical estimator based on the spatiotemporal covariance and mean of the observations known as kriging. An adequate parametric model for the covariance and the mean is chosen from an exploratory data analysis. Beforehand, it is necessary to estimate the parameters of this spatiotemporal model; This is performed in a Bayesian setting. The approach is then applied to a subset of the CloudSat dataset.

Copyright statement. TEXT

1 Introduction

Clouds have a strong influence on weather and climate. They are a key element of Earth's hydrological cycle, bringing water from the air to the ground and from one region of the globe to another. They also dominate the energy budget of the Earth through their action on the exchange of solar and thermal radiation within the atmosphere and between the atmosphere, the hydrosphere, the land surface, the biosphere, and space. However, they still remain a major source of uncertainties in predicting the weather and climate change.

While the measurement of cloud occurrences and properties at useful spatial and temporal scales is notoriously difficult (Marshall and Davis, 2005; Stephens and Kummerow, 2007), the proliferation of satellite platforms in the last decades (Stephens et al., 2002; Eriksson et al., 2008; Wu et al., 2009) is fostering a number of new approaches. One such satellite, CloudSat, whose payload is a Cloud Profiling Radar (CPR), has been dedicated to measure the cloud vertical structure and microphys-

ical properties. It is part of the A-train constellation, who was originally set to comprise 7 satellites specifically designed to measure cloud and precipitation properties using different instruments. Since 2006 CloudSat has collected a large database of 25 cloud properties, globally and over an extended period of time, despite some malfunctions.

In this study, we propose to analyze statistically a part of this database in order to perform interpolation and prediction of cloud properties. This kind of spatio-temporal estimation can be of major importance for the assessment of satellite cloud attenuation (Lyras et al., 2016), including Global Positioning System (GPS) radio occultation (Yang and Zou, 2012). It is required for improving the representation of cloud systems in numerical weather prediction (Bodas-Salcedo et al., 2008; Chen 30 et al., 2011), e.g. for data assimilation (Qu et al., 2018). It can be involved in the assessment of an aircraft icing detection system (Vivekanandan et al., 2001), the design of a satellite communication system (Khan et al., 2012), the systematic comparison with other cloud products from different instruments/satellites, *etc.*

As observations from different instruments are likely not collected at the same spatiotemporal positions, the procedure of interpolation-prediction usually involves regridding selected slots of data. A number of statistical methods dedicated to the 35 analysis of spatial and spatio-temporal data have been developed over the years taking into account the spatial and/or temporal correlation of the observations (Ripley, 1981; Cressie, 1993). In this study, we propose to use an approach based on a kriging estimator for the interpolation/prediction problem. The kriging estimator was initially introduced by (Krige, 1951), from which it takes its name, to estimate the gold distribution at the Witwatersrand reef complex in South Africa based on samples from boreholes. It was then formalized mathematically by (Matheron, 1963) in the context of mining geology. Afterwards, the 40 kriging estimator spread to many other areas of sciences (Wackernagel, 2013) (hydrogeology, geotechnics, agronomy, air quality, fishery, epidemiology, water and soil pollution, noise, *etc.*). Best-known kriging techniques are the Simple Kriging, which assumes stationarity of the 1st-order with a known mean, the Ordinary Kriging, where the mean is unknown and to be estimated, and the Universal Kriging, which assumes non-stationarity of the 1st-order. Since its first development, 45 kriging techniques have largely evolved, and a number of new kriging techniques have been developed (Chiles and Delfiner, 1999; Cressie, 1993; Cressie and Wikle, 2015). In the field of meteorology, the kriging estimator have primarily been used to estimate precipitation accumulation from rain gauges (Nour et al., 2006; Belo-Pereira et al., 2011) and in combination with satellite-derived precipitation (Jewell and Gaussiat, 2015; Verdin et al., 2016; Varouchakis et al., 2021) as well as aerosols 50 concentration in the air from in-situ observations (Park, 2016). It has also been used for the estimation of temperature from in-situ measurements (Heuvelink et al., 2012; Didari and Zand-Parsa, 2018), from satellite observations (Florio et al., 2004) or a combination of them (Didari and Zand-Parsa, 2018) as well as for the estimation of surface properties from remote 55 measurements (der Meer, 2012; Zakeri and Mariethoz, 2021).

Our approach is based on a second-order analysis of cloud Liquid Water Content (LWC) obtained from Level 2B product of the CloudSat ground segment data. We perform a detailed analysis of those properties and propose an adequate parametric model for their mean and covariance. The model parameters are obtained by maximizing the a posteriori probability density 55 which comprises a likelihood term expressing how the model fits the observation conditionnally to the presence of clouds. Finally we apply this model in the context of interpolation and prediction of CloudSat observations. To our knowledge, it is the first time the kriging estimator have been employed to infer cloud microphysical properties using CloudSat observations :

– We provide a comprehensive mathematical description of the kriging estimator. It allows us to mention that the parameters of the mean and covariance functions are commonly treated differently in the kriging literature. In our approach, 60 parameters can enter both linearly and non-linearly in the mean, which makes it more flexible in accommodating the trend of the observations. Thus, we propose a global treatment of all the parameters of the model as the standard Universal Kriging is not compatible with non-linearity.

– We propose a model for the mean and covariance functions and thoroughly justify each choice we make (*e.g.*, stationarity assumption, homogeneity assumption, *etc.*)

65 – We perform a detailed analysis of both the estimated model parameters and the resulting interpolated/predicted cloud LWC.

2 Data and Mathematical Description

2.1 The CloudSat data and the CPR instrument

CloudSat has been flying in formation in the A-train with other satellites including *Aqua*, *Aura* and CALIPSO. CloudSat 70 payload consists of a 94 GHz cloud-profiling radar (CPR) that was specifically designed to sense cloud-sized particles (*i.e.*, cloud ice, snow, cloud droplets and light rain).

It was declared operational on June 2-nd 2006 and has been flying in the A-train until February 22-nd 2018. It follows a Sun-synchronous orbit with an approximately 1330-LT equatorial crossing time. Since 2011 and a battery malfunction, it provides 75 observations only during daytime. The satellite visits the same position of the globe after a period of 16 days corresponding to 233 orbits. Each orbit is achieved in about 1 hour and 58 minutes. The CloudSat radar samples profiles at 625 kHz and has an along-track velocity of approximately 7 km/s, which corresponds to a profile measured every 0.16 second with an along-track displacement of approximately 1.1 km. Each profile has 125 vertical bins of 240 m thickness.

In this study, we use the level 2B Cloud Water Content product (2B-CWC-RO) that contains retrieved estimates of cloud 80 Liquid and Ice Water Content, effective radius, and related quantities for each radar backscattered reflectivity profile (Austin et al., 2009). We focus on a region centered over Europe and under-sampled at a rate of 1/50 (*cf.* Fig. 1). In this region, we select data from June 16-th 2006 to June 14-th 2015 constituting a set of 239087 profiles. The CPR observations are classified according to their cloud types in the level 2B-CLDCLASS product (*cf.* Fig. 2). After analysis of the cloud types distribution inside the considered region, we decided to focus on the LWC labeled as Altostratus as they constitute a good balance between the amount of available data and the computational feasibility. Moreover, altostratus are less fractionated than other cloud 85 types, thereby their physical properties are more continuous.




Figure 1. CloudSat ground track representing 14 orbits (a), zoom in on the European zone under study (b).




Figure 2. Sample of CloudSat cloud classification over a period of 15 days. Clouds are classified according to Clear (no clouds), Cs (Cumulus), As (Altocstratus), Ac (Altocumulus), St (Stratus), Sc (Stratocumulus), Cu (Cirrus), Ns (Nimbostratus), DC (Deep Convective).

2.2 Mathematical description of the data

2.2.1 Background

We are considering the scalar function Φ of three spatial variables (x, y, z) and a temporal variable t mapping elements from D into \mathbb{R} :

$$90 \quad \Phi : D = [0, 2\pi] \times [-\pi/2, \pi/2] \times \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R} \quad (1)$$

$$s = (x, y, z, t) \mapsto \Phi(x, y, z, t),$$

where $s \in D$ is spatiotemporal localization, x, y represent respectively the longitude and latitude at the Earth's surface and z is the altitude. The function Φ can be modeled as a deterministic or stochastic quantity. The stochastic nature of the function Φ can be introduced to model some inner variability of the physical phenomenon under study or an incomplete knowledge of the

phenomenon itself. The complete description of a stochastic process requires the construction of the joint probability density function for the continuous variables in space and time (Chonavel and Ormrod, 2002; Gaetan and Guyon, 2008; Brockwell and Davis, 2009), which can be cumbersome in a high-dimensional space. To alleviate this burden, it is possible to resort to the description of the mean and covariance of the random function. This proves very efficient when the distribution associated to the random function is not too complex (for instance when the random function is Gaussian, the mean and covariance are sufficient to completely describe the distribution).

100 In this study, we assume that the mean and covariance exist and represent sufficiently well the distribution of the random function Φ . The mean is a function of the spatiotemporal localization s :

$$m_\Phi(s) = E[\Phi(s)],$$

where $E[\cdot]$ denotes the expectation of the random variable. The covariance is a function of 4 spatiotemporal variables and 4 spatiotemporal shift variables, noted $\delta = (\delta_x, \delta_y, \delta_z, \delta_t)$. Thus we have,

$$105 \quad R_\Phi(s, \delta) = E[(\Phi(s) - m_\Phi(s))(\Phi(s + \delta) - m_\Phi(s + \delta))].$$

The covariance function describes the spatiotemporal dependency of the stochastic function Φ . Based on these definitions, it is useful to introduce certain stationary model to characterize the degree of homogeneity of the random function Φ .

2.2.2 Stationarity

In order to infer the distribution of the random function Φ , the structure of the spatiotemporal process is of great importance.

110 In this sense, stationary model enable to structure the spatiotemporal variability of the random function Φ . The process under study is said to be strictly stationary if the distributions $[\Phi(s_1), \dots, \Phi(s_k)]$ and $[\Phi(s_1 + \delta), \dots, \Phi(s_k + \delta)]$ are identical for all $\delta, s_1, \dots, s_k \in D$, and for all $k \in \mathbb{N}$. This is a very restrictive condition which is not usually satisfied in real-life applications. The weak stationarity stipulates that:

- the random function Φ is 1-st order stationary when its expected value does not depend on the localization s :

$$115 \quad m_\Phi(s) = \mu$$

- the random function Φ is 2-nd order stationary when its covariance function depends only on the lag vector δ between 2 localizations s and $s' = s + \delta$:

$$R_\Phi(s, \delta) = R_\Phi(\delta), \text{ for all } s \text{ and } \delta.$$

2.2.3 Observations and uncertainties

120 We denote the successive spatiotemporal positions $s_n = (x_n, y_n, z_n, t_n)$, $n = 1, \dots, N$ and the targeted quantity Φ at these positions:

$$\Phi_n = \Phi(s_n), \quad n = 1, \dots, N.$$

We model the actual observations of the targeted quantity Φ as corrupted by an additive noise related to the measurement uncertainty. The observation at position s_n is then denoted Ψ_n and is written:

125 $\Psi_n = \Phi_n + B_n,$

where Ψ_n , Φ_n and B_n are random variables. We denote the values associated with a realization of these random variables by ψ_n , φ_n and b_n . Thus, a realization of the random variable Ψ at the position s_n is written:

$$\psi_n = \varphi_n + b_n, \quad (2)$$

and additionally, we write:

130 $\psi = \varphi + b,$

where $\psi = [\psi_1, \dots, \psi_N]$, $\varphi = [\varphi_1, \dots, \varphi_N]$ and $b = [b_1, \dots, b_N]$ are the collections of the related N quantities.

2.2.4 Objectives

The objective of this work is to determine the value φ_0 of the quantity of interest φ at a given location $s_0 = (x_0, y_0, z_0, t_0)$. This is an estimation problem that is tackled by designing an estimate $\hat{\varphi}_0$ of the quantity of interest φ_0 from the data ψ described

135 in the previous section. On the one hand, spatially, this is an interpolation problem since the point x_0, y_0, z_0 is usually not on the grid of observations. On the other hand, temporally, this is a prediction problem since t_0 is naturally positioned in the future. The considered approach is to statistically learn from the large amount of available data both to interpolate/predict the targeted quantity and to quantize the uncertainty associated with the estimated value. In the next section, we introduce the kriging estimator, that is specifically designed to perform such a task.

140 **3 The kriging estimator**

The goal is to construct a statistical estimator $\hat{\Phi}_0$

$$\begin{aligned} \hat{\Phi}_0 : \mathbb{R}^N &\rightarrow \mathbb{R} \\ \psi &\mapsto \hat{\varphi}_0 = \hat{\Phi}_0(\psi) \end{aligned}$$

in order to estimate φ_0 from the observations ψ . This section describes the strategy for the construction of the kriging estimator

which has been widely used in geostatistics. This framework has been developed in a handful of handbooks (Cressie, 1993;

145 Chiles and Delfiner, 1999; Diggle et al., 2003) and later on by (Montero et al., 2015), in the case of a known mean and covariance. Our development differs slightly from the standards of geostatistical literature in two ways.

150 1. The parameters of the mean function $m_\Phi(s)$ usually act linearly and consequently their estimation can be performed inside the kriging estimator while the estimation of the covariance parameters is classically performed in a previous offline procedure. Instead, in order to design a model with a higher capacity, we consider both a non-linear mean and a non-linear covariance functions and estimate their parameters in an offline procedure.

2. Our approach is fully parametric and consequently relies on a parametric covariance function instead of a variogram (Banerjee et al., 2014).

Thus, in the following development, we consider that the mean and covariance functions are known and defer the (offline) estimation of their parameters to the next section.

155 We consider a linear estimator of the form:

$$\hat{\Phi}_0(\Psi) = \mathbf{a}^t \Psi + a_0, \quad (3)$$

where $\Psi = [\Psi_1, \dots, \Psi_N]$ are the observations, $\mathbf{a} = [a_1, \dots, a_N]$ are scalar coefficients and N is the number of available observations. This can be alternatively expressed in terms of realizations of the random variables as:

$$\varphi_0 = \hat{\Phi}_0(\psi) = \mathbf{a}^t \psi + a_0. \quad (4)$$

160 Our goal is to determine the coefficients \mathbf{a} and a_0 . The strategy is to minimize an error between the estimated quantity $\hat{\Phi}_0$ and the true value Φ_0 . We choose the Mean Squared Error (MSE),

$$\mathcal{E}(\mathbf{a}, a_0) = \mathbb{E}[(\hat{\Phi}_0 - \Phi_0)^2]. \quad (5)$$

The estimator $\hat{\Phi}$, defined by (3)-(4) with:

$$(\mathbf{a}, a_0)^{\text{opt}} = \arg \min_{\mathbf{a}, a_0} \mathcal{E}(\mathbf{a}, a_0),$$

165 and the corresponding estimator is the so-called Minimum Mean Square Error estimator (MMSE). The value a_0^{opt} minimizing (5) satisfies

$$\frac{\partial \mathcal{E}}{\partial a_0} \Big|_{a_0^{\text{opt}}} = 0$$

giving,

$$a_0^{\text{opt}} = m_{\Phi_0} - \mathbf{a}^t \mathbf{m}_{\Phi}. \quad (6)$$

170 Then, plugin (6) into (5) and expanding leads to:

$$\mathcal{E}(\mathbf{a}) = \mathcal{E}(\mathbf{a}, a_0^{\text{opt}}) = \mathbf{a}^t \mathbf{R}_{\Psi} \mathbf{a} - 2\mathbf{a}^t \mathbf{r}_{\Psi \Phi_0} + \text{var}[\Phi_0] \quad (7)$$

where we introduced the following notations:

$$\mathbf{R}_{\Psi} = \mathbb{E}[(\Psi - \mathbf{m}_{\Phi})(\Psi - \mathbf{m}_{\Phi})^t]$$

$$\mathbf{r}_{\Psi \Phi_0} = \mathbb{E}[(\Phi_0 - m_{\Phi_0})(\Psi - \mathbf{m}_{\Phi})]$$

$$\text{var}[\Phi_0] = \mathbb{E}[(\Phi_0 - m_{\Phi_0})^2],$$

\mathbf{R}_Ψ is the covariance matrix of the observations, $\mathbf{r}_{\Psi\Phi_0}$ the covariance between the observations and the quantity of interest, and $\text{var}[\Phi_0]$ is the variance of the quantity of interest. The vector \mathbf{a}^{opt} that minimizes (7) satisfies

$$\frac{\partial \mathcal{E}}{\partial \mathbf{a}} \bigg|_{\mathbf{a}^{\text{opt}}} = \mathbf{0},$$

175 and that leads to:

$$\mathbf{a}^{\text{opt}} = \mathbf{R}_\Psi^{-1} \mathbf{r}_{\Psi\Phi_0}.$$

Moreover, modeling noise \mathbf{B} independent of the quantity of interest Φ , we have

$$\mathbf{R}_\Psi = \mathbf{R}_\Phi + \mathbf{R}_B \quad \text{and} \quad \mathbf{r}_{\Psi\Phi_0} = \mathbf{r}_{\Phi\Phi_0}, \quad (8)$$

so that,

$$180 \quad \mathbf{a}^{\text{opt}} = (\mathbf{R}_\Phi + \mathbf{R}_B)^{-1} \mathbf{r}_{\Phi\Phi_0}. \quad (9)$$

Finally, we deduce the estimated value $\hat{\varphi}_0$ of the quantity of interest φ at position s_0 by inserting equations (9) and (6) in (4):

$$\hat{\varphi}_0 = \mathbf{r}_{\Phi\Phi_0}^t (\mathbf{R}_\Phi + \mathbf{R}_B)^{-1} (\psi - \mathbf{m}_\Phi) + m_\Phi(s_0). \quad (10)$$

Additionally, the MSE of the estimated value is given by:

$$\mathcal{E}(\mathbf{a}^{\text{opt}}, \mathbf{a}_0^{\text{opt}}) = \text{var}[\Phi_0] - \mathbf{r}_{\Phi\Phi_0}^t (\mathbf{R}_\Phi + \mathbf{R}_B)^{-1} \mathbf{r}_{\Phi\Phi_0}, \quad (11)$$

185 which, in this context, equals to the variance of the estimation error $\text{var}[\hat{\Phi}_0 - \Phi_0]$ because the bias is equal to zero. This estimator is known as the kriging estimator in the geostatistical literature (Montero et al., 2015). In a slightly different form, this estimator is also the one of Kalman and Wiener in the field of filtering.

4 Construction of a model for the mean and the covariance

4.1 Time-altitude and spectral analysis

190 In order to construct an adequate model for the mean and the covariance, we start with an exploratory analysis to extract some raw characteristics from the time-altitude observations represented in Fig. 3. We compute the biased empirical covariance for the 2-dimensional case $(\delta t, \delta z)$ and the 1-dimensional case at a fixed altitude of $z = 6.711$ km (cf. Fig. 4). This shows 1 year periodicity of the LWC observations in the time component, while the altitude component decreases to 0 for $\delta z = 1$ km. The blue spots located at $\delta z = +/ - 6$ km are due to zero-padding above and below the cloudy profile that introduces an artificial correlation. Another way of looking at the data ψ is to compute the empirical power spectral density $\hat{S}_\psi(f)$, i.e., the squared modulus of the Fourier transform of the observations ψ (a.k.a. periodogram) (cf. Fig. 5). There is clearly a peak at 1 year $^{-1}$. For some altitudes, there are lower peaks at 2 year $^{-1}$ or even at 3 year $^{-1}$, indicating that the signal cannot be modeled by a single sinusoidal component. At lower altitudes, the fundamental component becomes negligible as well as the harmonics. Nevertheless, this suggests that CloudSat observations can be interpreted as the superposition of multiple periodic components.




Figure 3. Scatter plot of the altostratus LWC observations.




Figure 4. Empirical covariance: (a) 2-dimensional covariance ($\delta t, \delta z$), (b) slice at $\delta z = 0$ km and (c) slice at $\delta t = 0$ year.

200 4.2 Proposition of an adequate model

The observations Ψ are modeled as the addition of a random process Φ and a random noise process B , such as:

$$\Psi(t, z) = \Phi(t, z) + B(t, z).$$

We dodge the complexity of handling 4 dimensions and choose to do an in-depth analysis of a time-altitude dependent model while omitting the latitude-longitude dimensions. This simplification can be justified when the spatial extension of the dataset is not too important. This is equivalent to assimilate the dataset to a single spatial location by averaging over the geographic components. All random processes are modeled as Gaussian (Rasmussen and Williams, 2005):

$$\begin{aligned} \Phi &\sim \mathcal{N}(m_\Phi, \mathbf{R}_\Phi), \\ B &\sim \mathcal{N}(\mathbf{0}, \mathbf{R}_B), \\ \Psi &\sim \mathcal{N}(m_\Phi, \mathbf{R}_\Psi), \end{aligned} \tag{12}$$




Figure 5. Empirical periodogram at (a) $z = 6.71$ km and (b) $z = 6.00$ km.

where \mathbf{R}_Φ , \mathbf{R}_B and \mathbf{R}_Ψ are the covariance matrices associated to the quantity of interest Φ , the noise B and the observations Ψ . Assuming the noise B independent of the random function Φ , we can write: $\mathbf{R}_\Psi = \mathbf{R}_\Phi + \mathbf{R}_B$, see (8).

205 In order to take into account the periodic temporal trend observed in the observations, we consider a non 1-st order stationary model. Moreover, the mean is chosen not to be a function of the altitude in order to limit the number of parameters. Considering the time variations, we chose a periodic mean with 1 fundamental and 2 harmonics such that:

$$m_\Phi(t) = \beta_0 + \beta_1 \cos\left(\alpha_1 + \frac{2\pi t}{T}\right) \\ + \beta_2 \cos\left(\alpha_2 + \frac{2\pi t}{T/2}\right) + \beta_3 \cos\left(\alpha_3 + \frac{2\pi t}{T/3}\right)$$

210 where β_0 , β_1 , β_2 and β_3 are respectively the amplitude of the continuous component, the fundamental and the 2 harmonics, α_i (with $i = 1, 2, 3$) are the phase parameters of the fundamental and the harmonics and T is the period.

In order to further simplify we resort to the widely used separable model for the covariance (Genton, 2007) so that the spatio-temporal covariance factors into a purely spatial and a purely temporal component. This reduces the number of parameters allowing for computationally efficient estimation and inference, it is then more tractable to run different test on the dataset. Thus, the separable covariance model writes

215 $\mathcal{R}_\Phi(\delta_t, \delta_z) = r_\Phi \mathcal{R}_{\Phi_t}(\delta t) \mathcal{R}_{\Phi_z}(\delta z),$

where r_Φ is the variance of the quantity of interest. We chose an exponential covariance model for the time covariance

$$\mathcal{R}_{\Phi_t}(\delta t) = \exp[-|\delta t|/l_t],$$

where l_t is the correlation time. The choice of this covariance model is motivated by the shape of the empirical covariance described in the previous section (it decreases more rapidly than a Gaussian). The altitude covariance is set to be:

$$220 \quad \mathcal{R}_{\Phi_z}(\delta z) = \exp \left[-(\delta z/l_z)^2 \right],$$

where l_z is the correlation length in altitude. Finally, we use a white noise:

$$\mathcal{R}_B(\delta t, \delta z) = r_B \text{Dirac}(\delta t, \delta z),$$

where r_B is the variance of the noise. Finally, we have a model consisting of 12 parameters, including 4 parameters for covariance, and 8 parameters for the mean

$$225 \quad \boldsymbol{\theta} = [r_\Phi, l_t, l_z, \beta_0, \beta_1, \beta_2, \beta_3, \alpha_1, \alpha_2, \alpha_3, T, r_B].$$

Using this notation we can denote the parametric model for the mean $\mathbf{m}_\Psi(\delta t; \boldsymbol{\theta})$ and for the covariance $\mathbf{R}_\Psi(\delta t, \delta z; \boldsymbol{\theta})$. In the following section, we describe the strategy to estimate this set of parameters.

5 Model parameter estimation, optimization and final kriging equations

In this section, we present the strategy for the estimation of the model parameters $\boldsymbol{\theta}$. We choose the Maximum A Posteriori 230 (hereafter abbreviated MAP) estimate for $\boldsymbol{\theta}$ which is the mode of the *posterior* probability density function (pdf) in Bayesian statistics. The analysis of the estimated parameters and corresponding model is deferred to Sect. 6 for the 1-dimensional case and Sect. 7 for the 2-dimensional case.

5.1 The MAP estimator

In the previous section, we proposed a parametric model for the mean $\mathbf{m}_\Psi(\delta t; \boldsymbol{\theta})$ and for the covariance $\mathbf{R}_\Psi(\delta t, \delta z; \boldsymbol{\theta})$ of the 235 random variables Ψ , which decomposes into a model for the quantity of interest Φ and the noise B . The problem is to find an estimate $\hat{\boldsymbol{\theta}}$ of the true parameters $\boldsymbol{\theta}^*$ from the observations ψ . We introduce the *posterior* pdf:

$$\pi(\boldsymbol{\theta}|\psi) = \frac{f(\psi|\boldsymbol{\theta})\rho(\boldsymbol{\theta})}{f(\psi)},$$

where $f(\psi|\boldsymbol{\theta})$ is the so-called likelihood function, which expresses the probability of the observations given the parameter $\boldsymbol{\theta}$, $\rho(\boldsymbol{\theta})$ is the *prior* distribution for the parameters $\boldsymbol{\theta}$ and $f(\psi)$ is the marginal pdf for the observations. Since we model the 240 observations by a Gaussian pdf conditionally to the parameters $\boldsymbol{\theta}$ (see (12)), the likelihood function $f(\psi|\boldsymbol{\theta})$ writes:

$$f(\psi|\boldsymbol{\theta}) = (2\pi)^{-N/2} (\det \mathbf{R}_\Psi)^{-1/2} \exp \left[-\frac{1}{2} (\psi - \mathbf{m}_\Psi)^t \mathbf{R}_\Psi^{-1} (\psi - \mathbf{m}_\Psi) \right], \quad (13)$$

where we have omitted the dependence in $\boldsymbol{\theta}$, δt and δz for \mathbf{R}_Ψ and \mathbf{m}_Ψ to weigh down the notations. Since the *posterior* pdf $\pi(\boldsymbol{\theta}|\boldsymbol{\psi})$ is by definition positive, we consider the co-log posterior $\text{CLP}(\boldsymbol{\theta}) = -\log \pi(\boldsymbol{\theta}|\boldsymbol{\psi})$, which expands as follows:

$$\begin{aligned} \text{CLP}(\boldsymbol{\theta}) &\# -\log f(\boldsymbol{\psi}|\boldsymbol{\theta}) - \log \rho(\boldsymbol{\theta}) \\ &\# \frac{1}{2} (\boldsymbol{\psi} - \mathbf{m}_\Psi)^t \mathbf{R}_\Psi^{-1} (\boldsymbol{\psi} - \mathbf{m}_\Psi) \\ &\quad + \frac{1}{2} \log \det \mathbf{R}_\Psi - \log \rho(\boldsymbol{\theta}) \end{aligned} \quad (14)$$

where $\#$ stands for “equality up to an additive constant” (not function of $\boldsymbol{\theta}$). The minimum of (14) is the maximum of the *posterior* pdf. Thus, minimizing the CLP gives the MAP estimate. To complete, we consider a uniform *prior* $\rho(\boldsymbol{\theta})$ on a given domain Θ :

245 $\rho(\boldsymbol{\theta}) = \mathcal{U}_\Theta(\boldsymbol{\theta})$

where Θ is a hyper-rectangle in \mathbb{R}^P describing a range of possible values for each component of $\boldsymbol{\theta}$. For our model $P = 12$. Finally, we obtain:

$$\begin{aligned} \text{CLP}(\boldsymbol{\theta}) &\# (\boldsymbol{\psi} - \mathbf{m}_\Psi)^t \mathbf{R}_\Psi^{-1} (\boldsymbol{\psi} - \mathbf{m}_\Psi) \\ &\quad + \log \det \mathbf{R}_\Psi - \log \mathcal{U}_\Theta(\boldsymbol{\theta}) \end{aligned} \quad (15)$$

The function CLP is infinite when $\boldsymbol{\theta} \notin \Theta$ because of the term $-\log \mathcal{U}_\Theta(\boldsymbol{\theta})$. The minimizer $\hat{\boldsymbol{\theta}}$, given by:

$$\hat{\boldsymbol{\theta}} = \arg \min_{\boldsymbol{\theta}} \text{CLP}(\boldsymbol{\theta}), \quad (16)$$

is the MAP estimate of the true parameter $\boldsymbol{\theta}^*$.

5.2 Optimization procedure

250 An optimization procedure is required in order to determine the minimizer $\hat{\boldsymbol{\theta}}$ of the CLP. In this study, we use an Iterative Conditional Mode strategy with a Golden-Section search (ICM-GS) for each component (Kiefer, 1953; Press et al., 1992). We then have an algorithm composed of an outer and an inner loop.

The inner loop optimizes w.r.t. one component at a time, say θ_p , based on a golden-section search. It stops when the variation of θ_p is smaller than a given threshold, say ϵ_p . The outer loop repeats the scan of the P components and stops when the variation 255 of $\boldsymbol{\theta}$ becomes smaller than a second given threshold denotes by η . The values ϵ_p and η have been fixed so as to reach stable results. The specific threshold values used in optimization procedure are reported in Table 1.

Table 1. Threshold values ϵ_p used for each parameters and global threshold value η .

ϵ_{r_Φ}	ϵ_{l_t}	ϵ_{l_z}	ϵ_{r_B}	ϵ_{β_0}	ϵ_{β_1}	ϵ_{β_2}	ϵ_{β_3}	ϵ_{α_1}	ϵ_{α_2}	ϵ_{α_3}	η
$1e-5$	$1e-7$	$1e-1$	$1e-6$	$1e-3$	$1e-2$	$1e-2$	$1e-4$	$1e-2$	$1e-5$	$1e-2$	$1e-3$

5.3 Kriging equation with \hat{m}_Φ and \hat{R}_Ψ

The interpolation/prediction solution given the observations ψ is performed using the kriging estimator developed in Sect. 3. The kriging equations were initially developed assuming a known mean m_Φ and covariance \mathbf{R}_Ψ . In reality, we only have 260 at hand the parametric mean \hat{m}_Φ and covariance \hat{R}_Ψ that depend on the estimated parameter $\hat{\theta}$. Thus, the estimation of the quantity of interest Φ in t_0 is written:

$$\varphi_0 = \hat{r}_{\Psi\Phi_0}^t \hat{R}_\Psi^{-1} (\psi - \hat{m}_\Phi) + \hat{m}_\Phi(t_0), \quad (17)$$

where \hat{m}_Φ is a vector of estimated mean at the positions of the observations. Finally, we can compute the minimum mean square error:

$$265 \quad \mathcal{E} = \hat{r}_\Phi - \hat{r}_{\Psi\Phi_0}^t \hat{R}_\Psi^{-1} \hat{r}_{\Psi\Phi_0}. \quad (18)$$

This last expression is equal to the variance of the estimation error, which we note $\sigma^2 = \text{var}[\hat{\Phi}_0 - \Phi_0]$. It gives an estimation of the forecast error associated with the kriging technique. However, it should be kept in mind that this variance does not account for the uncertainty in the model parameter estimate $\hat{\theta}$. Therefore, this expression tends to underestimate the true estimation variance. This feature has been well documented in (Cressie, 1993; Montero et al., 2015). We leave the assessment of this 270 impact for further development of this model. Note that the kriging stage is essentially a computational step that does not present any major difficulty except for the computational burden when dealing with large dimensions.

6 The 1-dimensional time case

We start our analysis with the 1-dimensional case at a single altitude. We give a detailed analysis of the estimated parameters according to the MAP criterion and the ICM-GS algorithm, both described in the previous section. We concentrate on the 275 estimation over the west of Europe and analyse the impact of reducing the considered geographic area. We then proceed to the kriging of observations in the interpolation and prediction case.

6.1 Estimation over a European area at a single altitude

In order to analyze the results of the previously described parameter estimation procedure, we first consider observations associated to altostratus clouds over an European area extending from 10°W to 10°E in longitude and 30°N to 60°N in latitude 280 (see Fig. 1) and use a subsampled database (at 1/50 rate) at a single altitude of $z \simeq 6$ km. This corresponds to a set of $N_{\text{obs}} = 3653$ observations. The period parameter T has been readily set to 1 year so as to represent the seasonality observed in the data.

Fig. 6 shows the evolution of CLP during the optimization process which, as expected, decreases with the iterations and stabilizes after ~ 5 iterations. The current value of the parameters changes at each iteration and stabilizes as shown in Fig. 7. 285 Even though the decrease of CLP along the iterative optimization process indicates convergence towards a minimum, there is no guarantee that it is a global minimum. This is highlighted in Fig. 8(a) which represents the isocontour plot of CLP as a




Figure 6. Evolution of CLP during the optimization process.




Figure 7. Evolution of each parameters along the optimization process. The x -axes represents the iteration index while the y -axes represent the current value for each parameter. Parameters associated to the covariance of the quantity of interest are represented in red (○); parameters associated to the mean of the quantity of interest are represented in blue (○) and the noise variance is represented in green (○).




Figure 8. (a) Contour plot of CLP as a function of r_Φ and l_t . There is a global minima characterized by $\hat{r}_\Phi \simeq 2.2 \times 10^{-2}$, $\hat{l}_t \simeq 3.6 \times 10^{-6}$ and a corresponding $CLP(\hat{\theta}) = -1.67 \times 10^4$ and a local minima characterized by $\hat{r}_\Phi \simeq 2 \times 10^{-2}$, $\hat{l}_t \simeq 9.1 \times 10^{-4}$ and $CLP(\hat{\theta}) = -1.65 \times 10^4$. (b) CLP as a function of r_Φ while fixing all the other parameters to their initial values. This shows why the first two iterations of the optimization algorithm are moving away from the global minimum shown in the contour plot because the CLP is strictly monotonic inside the considered region.

function of (r_Φ, l_t) while the other parameters are fixed to their estimated values. Additionally, we plot the current values of r_Φ and l_t at each iteration of the ICM-GS algorithm in Fig. 8(a). In Fig. 8(b) we plotted the CLP as a function of r_Φ while fixing all the other parameters to their initial values to highlight the fact that the first two iterations of the optimization algorithm are moving away from the global minimum shown in the contour plot. The contour plot has been represented by varying values of r_Φ and l_t while setting the other parameters to their estimated value at the end of the optimization procedure so that it is not representative of the CLP at the initial states. It clearly shows a global and a local minimum. The global minimum is reached for $(\hat{r}_\Phi, \hat{l}_t) \simeq (2 \times 10^{-3}, 3.75 \times 10^{-6})$. The estimated value $\hat{l}_t = 3.75 \times 10^{-6}$ corresponds to ~ 2 minutes. This correlation time is very small compared to the 9-year period used to train the parameters. It indicates that there is essentially no correlation for observations later than 6 minutes apart. Indeed, the exponential covariance model reaches a 5% correlation at about $3l_t$

(Banerjee et al., 2014). This feature can be either explained by the model, the observations or the phenomenon itself. However, it is then evident that such a model will not be very efficient in the perspective of long-term forecast.

As stated in (Rasmussen and Williams, 2005), each local minimum corresponds to a particular interpretation of the data, and a careful analysis may be required to choose one model instead of the other. In the next section, we will further discuss the 300 presence of two minima of the CLP in relation with two competing models. An exhaustive search for all local minima requires to compute the criterion $CLP(\theta)$ for all possible combinations of parameters in the domain Θ , which is impossible. Instead, we performed a set of 10 optimizations with random initialization to track down additional local minima and converged toward the same minimizer. Hence we conclude there is no other local minima than the one previously pointed out (*cf.* Fig. 8(a)).

6.2 Discussion on the reduction of the geographic area

305 In order to assess the influence of the considered geographic area, the optimization procedure is run by reducing the geographic extent of the dataset. We consider 9 areas that are represented in Fig. 9, and the estimated parameters $\hat{\theta}$ are summarized in Fig. 10.

The estimated values of the covariance parameters (*i.e.*, \hat{r}_Φ , \hat{l}_t and \hat{r}_B) vary in a complementary fashion, with approximately 310 similar values for zones 1 to 3, a leap of several orders of magnitude for zones 4 to 8, followed by a drop for the 9-th zone. These variations can be traced back to the presence of a local minimum in the domain Θ as it is shown in Fig. 11 where the variation of $CLP - \mu_{CLP}$ with respect to l_t has been represented for each geographic area. All CLPs show the presence of a local and global minima on the considered interval (except for the 9-th area). The variations of CLP w.r.t. the correlation time 315 l_t have similar behavior for areas 1 to 3 with locations of the local and global minimum roughly identical. The same is valid for areas 4 and 5. Concerning areas 6 to 8 (*cf.* Fig. 11(c)), they have a more complex shape with a well defined global minimum but less pronounced local minima at lesser correlation time.

It is also worth highlighting the sliding of the global minimum to the local minimum when switching from zones 1, 2 and 3 to zones 4 and 5 (*cf.* Fig. 11 (a) and (b)). The global minimum of zones 1, 2 and 3 is reached for a smaller correlation time 320 $l_t \sim 10^{-6}$ year which means that the aggregation of spatially distant observations is interpreted as a less correlated process than for smaller geographic area (*i.e.*, zones 4 and 5). In this case, the retrieved model tends towards a white Gaussian noise as the covariance function is close to a Dirac. However, the presence of a local minimum of CLP with a higher correlation time indicates that the observations could also be explained by a second model, provided some additional constraint on the correlation time (*e.g.*, a different *prior* distribution for l_t). In particular, we note that the order of magnitude of the l_t values 325 associated with this local minimum is similar to the l_t values obtained for zones 4 to 7. When we reduce the geographic area, the set of observations tends to be more homogeneous, which plays in favor of models with higher correlation times.

The result obtained for zone 9 has a different interpretation. The estimated correlation time is $\hat{l}_t \simeq 7.13 \times 10^{-7}$, which is several orders of magnitude smaller than every other geographic areas. This is in conjunction with a particularly low estimated noise variance $\hat{r}_B \ll \hat{r}_\Phi$ (*cf.* Fig. 10 (c)). For this geographic area, the estimated \hat{r}_B sticks to the minimum bound of the *prior* density so the optimization procedure is stuck for this parameter. Thus, the MAP estimator compensates for the remaining signal variance by increasing \hat{r}_Φ and decreasing the associated \hat{l}_t , so that \mathcal{R}_Φ tends towards a Dirac function, *i.e.*, a covariance




Figure 9. The 9 geographic areas used for the parameter estimation analysis.

330 associated to white noise. In other words, the estimator is no longer able to separate the quantity of interest Φ from the noise with this dataset. To account for computational burden and stability of the estimated model we will pursue our analysis with the dataset corresponding to the 6-th geographic area.

6.3 Discussion on the 1-dimensional model

To conclude, it seems more interesting to carry on with geographic areas associated with higher correlation time, especially 335 those obtained for zones 4 to 7 which corresponds to $l_t \sim 1 - 6$ days, compared to $l_t \sim 2$ minutes for zones 1 to 3. Considering the exponential covariance model reaches 5% of its total variance at $\delta_t \sim 3l_t$ (Banerjee et al., 2014), the latter tends towards its mean m_Φ after ~ 6 minutes. Note that the models corresponding to zones 4 to 7 are associated with variances r_Φ lower than




Figure 10. Comparison of the estimated parameters for 9 geographic area. Parameters associated to the covariance of the quantity of interest are represented in red (●); parameters associated to the mean of the quantity of interest are represented in blue (●) and the noise variance is represented in green (●).

those obtained for the zones 1 to 3 and a greater noise variance r_B . However, the correlation time corresponding to zones 1 to 3 is very small that the corresponding covariance tends toward a diagonal matrix (*i.e.*, a Dirac function).

340 6.4 Results in interpolation

We start with the interpolation of the LWC over the 9-year period. Note that, despite the regular sampling rate of the CPR, actual observations of altostratus in the European area are sparsely distributed. The temporal positions t_0 at which kriging is applied are defined by $t_0 = [0, 1/N_0, \dots, 9-1/N_0, 9]$, with $N_0 = 4001$. The result of this kriging interpolation is represented in Fig. 12. The variability of the estimated $\hat{\varphi}_0$ are smaller than the actual variability of the observations, which is consistent with 345 the estimated model parameters. Indeed, the estimated noise variance $\hat{r}_B = 2.2 \times 10^{-3} \text{ g}^2 \cdot \text{cm}^{-6}$ is greater than the estimated variance of the quantity of interest $\hat{r}_\Phi = 9.8 \times 10^{-4} \text{ g}^2 \cdot \text{cm}^{-6}$. The periodic nature of the estimate $\hat{\varphi}_0$ is strong, especially when there are no observations (for example during the 5-th year of operation of the satellite). In this case, the result of the kriging interpolation consists in the estimated mean \hat{m}_Φ . In the vicinity of observations, the kriging estimates take a more complex form because of the additional information brought by the neighbouring data. The behavior observed over this period of 9 350 years is not surprising in itself because of the relative weakness of the observed correlation time, which is only $\hat{l}_t = 2.7$ days.




Figure 11. $(CLP - \mu_{CLP})$ as a function of the correlation time l_t around $\hat{\theta}$ for the geographic zones 1 to 3 (a), 3 to 4 (b), 6 to 8 (c) and the 9-th (d). The results for the 9 geographic areas have been grouped in 4 sketches according to their similarities.

We perform another kriging interpolation over a period of ~ 36 days centered on the beginning of the satellite's 3-rd year of operation (cf. Fig. 13(a)). In this figure, we observe that some observations are very closed temporally and have significantly different values of LWC which can explain the high estimated variance of the noise with respect to the variance of the quantity of interest. In this case, the kriging gives an average value of the observations. These observations generally correspond to 355 successive profiles in the 1/50 database, which are separated by 8 seconds. It should be noted that between two successive observations, the satellite moves ~ 54 km which can certainly explain a loss of spatial correlation not taken into account in our model. It is therefore likely that the model will explain this loss of spatial correlation by noise. On the other hand, we notice the presence of peaks in the structure of $\hat{\phi}$ that can be explained by the structure of the chosen temporal covariance, which is in fact characterized by an exponential decay corresponding to the decay observed on the curve of $\hat{\phi}$ when moving away from 360 an observation (cf. Fig. 13 (a)).




Figure 12. Kriging based interpolation over a 9 year period. The training set observations are represented as (●), the kriging estimate as (—), and variance of the estimation error (shaded gray).




Figure 13. (a) Kriging based interpolation over a period of 36 days centered at the beginning of 3-rd in the dataset. The mean is represented in as a dashed black line (—). (b) Kriging based prediction over a period of ~ 36 days centered at the beginning of 3-rd in the dataset. The vertical black line (|) represents the correlation time l_t after the last observation. In (a) and (b), the training set observations are represented as (●), the kriging estimate as black line (—), and the variance of the estimation error in shaded gray.

On all kriging results, we represent the 4σ area, where $\sigma^2 = \text{var}[\hat{\Phi}_0 - \Phi_0]$ is the variance of the estimation error of (18). This variance is decomposed into two terms, the variance $\text{var}[\hat{\Phi}_0]$ and a quadratic term in $\hat{r}_{\Psi\Phi_0} (= \hat{r}_{\Phi\Phi_0})$, which represents the covariance between the object of interest and the observations. At observation location (*i.e.*, Φ_0 is chosen collocated with an observation), $\hat{r}_{\Psi\Phi_0}$ reaches its maximum value and so does the quadratic term $\hat{r}_{\Psi\Phi_0} \hat{R}_{\Psi}^{-1} \hat{r}_{\Psi\Phi_0}$. Consequently, at observation 365 location the minimum mean square error given in (18) is minimal.

6.5 Results in prediction

In order to study the result of kriging prediction, we selected a period that includes observations until a certain date and analyzes the behavior of the prediction beyond the last available observation (*cf.* Fig. 13(b)). In order to facilitate the interpretation, we have subtracted the estimated mean \hat{m}_Φ from the observations ψ and the estimates $\hat{\varphi}_0$ and we have centered the x -axis on 370 the last observation available. This clearly shows a decrease in the kriging result after the last observation towards 0, which indicates that the result tends towards the estimated mean. We plot the value of the correlation time l_t as an indication. In the case of the exponential covariance, we get 5% of the variance r_Φ for a time $\sim 3l_t$ which corresponds to the time from which the kriging estimation tends towards the estimated mean. It is therefore a result consistent with the chosen covariance model and the estimated correlation time. This means that the prediction will be different from the average when an observation is 375 available in a period of approximately 8.2 days (because $l_t \simeq 2.7$ days) before or after the position where the estimation is performed.

6.6 Error analysis

In this section, we interpret in more details the kriging results through the analysis of the variance of the estimation error $\text{var}[\hat{\Phi}_0 - \Phi_0]$ and the estimated noise. This is accomplished with a kriging estimation at temporal positions t_0 collocated to the 380 positions of the observed data (*cf.* Fig. 14 (a)). Fig. 14 (b) represents the corresponding histograms for quantities $(\hat{\varphi}_0 - \hat{m}_\Phi)$ and $(\psi - \hat{m}_\Phi)$. Although the model used to describe the observed data is Gaussian, the histogram of the zero-mean quantity, $(\hat{\varphi}_0 - \hat{m}_\Phi)$, is not exactly Gaussian. It is actually skewed towards zero. However, its range of variation is smaller than the corresponding histogram for the observations, $(\psi - \hat{m}_\Phi)$, which is an expected result as some of the observed variability is attributed to the presence of noise.

385 The variance of the estimation error $\text{var}[\hat{\Phi}_0 - \Phi_0]$ gives an approximation of the variability around the estimated value $\hat{\Phi}_0$. Note that it is impossible to compute exactly the difference $(\hat{\Phi}_0 - \Phi_0)$, but we have estimated the variance of the noise knowing the observations, it is then possible to compute the variance $\text{var}[\hat{\Phi}_0 - \Phi_0]$ (*cf.* Sect. 3). This last variance term can be used to compute *posterior* realizations of the quantity of interest Φ . Fig. 14 (c) represents the variance $\text{var}[\Phi_0]$ and the variance of the estimation error $\text{var}[\hat{\Phi}_0 - \Phi_0]$. The latter has a lower magnitude than $\text{var}[\Phi_0]$ which is consistent with (11) as the term 390 $r_{\Psi\Phi_0}^t \mathbf{R}^{-1} r_{\Psi\Phi_0}^t$ is positive.

Finally, we computed the difference $(\psi - \hat{\varphi}_0)$ (*cf.* Fig. 15(a)). According to (2), this distribution must correspond to the noise distribution. We represented the normalized histogram of $(\psi - \hat{\varphi}_0)$ in Fig. 15 (b) as well as the pdf of the noise $B \sim \mathcal{N}(\mathbf{0}, \hat{r}_B)$. We observe that the two distributions have slightly different shapes but similar ranges of variation. The differences observed 395 are mainly due to the fact that the parameters of a Gaussian model are estimated from observations that do not have a strictly Gaussian distribution. Indeed, we notice that the observations have an asymmetric distribution which extends towards higher values of LWC than the Gaussian distribution.




Figure 14. Kriging results collocated to the observations. (a) Observations ψ (●), estimate $\hat{\varphi}_0$ (●), estimated mean \hat{m}_Φ (—) and variance of the estimation error in shaded gray. (b) Histograms of $(\hat{\varphi}_0 - \hat{m}_\Phi)$ and $(\psi - \hat{m}_\Phi)$. (c) Shaded gray representation of the variance of the estimation error and the variance of the quantity of interest Φ .




Figure 15. (a) Differences $(\psi - \hat{\varphi}_0)$ (●) superimposed on $4\sigma_B$ in shaded gray, and (b) normalized histograms of $(\psi - \hat{\varphi}_0)$ and the pdf of $B \sim \mathcal{N}(\mathbf{0}, \hat{\mathbf{R}}_B)$.

7 The 2-dimensional time-altitude case

In a similar fashion as Sect. 6, we examine the 2-dimensional time-altitude case starting from the parameter estimation followed by the kriging results.




Figure 16. Evolution of each parameters during the optimization process in 2-dimensional case (time-altitude). Parameters associated to the covariance of the quantity of interest are represented in red (○); parameters associated to the mean of the quantity of interest are represented in blue (○) and the noise variance is represented in green (○).

400 7.1 Parameter estimation

We are considering here the estimation of model parameters including the altitude dimension. Following the conclusions of the previous section, we use the observations of the 6-th geographic area. However, for computational reasons, the training dataset is restricted to the first 3 years of observations, which consists of $N_{\text{obs}} = 4167$ observations (cf. Fig. 17).

405 Fig. 16 represents the evolution of each parameter during the optimization process. We observe that all parameters have stabilized towards an estimated value. Moreover, none of the parameters have converged towards the bound defined by the prior.

The estimated values in the 1-dimensional (temporal) and 2-dimensional (time-altitude) cases are consistent from one model to the other (cf. Table 2). Overall, the order of magnitude for most quantities are close. However, we point out some noticeable differences:

Table 2. Comparison between estimated parameters for the 1d and the 2d model. Note that we have fixed $T = 1$ in the optimization procedure.

$\hat{\theta}$	1-D Model	2-D Model
\hat{r}_Φ	$\sim 9.84 \times 10^{-4}$	$\sim 5.3 \times 10^{-3}$
\hat{l}_t	$\sim 7.5 \times 10^{-3}$	$\sim 5.6 \times 10^{-3}$
\hat{l}_z	/	~ 865
\hat{r}_B	$\sim 2.2 \times 10^{-3}$	$\sim 1.7 \times 10^{-3}$
$\hat{\beta}_0$	$\sim 6.7 \times 10^{-2}$	$\sim 6.2 \times 10^{-2}$
$\hat{\beta}_1$	$\sim 4.2 \times 10^{-2}$	$\sim 7.9 \times 10^{-3}$
$\hat{\beta}_2$	$\sim 2.9 \times 10^{-3}$	$\sim 4.4 \times 10^{-3}$
$\hat{\beta}_3$	$\sim 9.7 \times 10^{-3}$	$\sim 3.4 \times 10^{-3}$
$\hat{\alpha}_1$	~ 5.4	~ 5.5
$\hat{\alpha}_2$	~ 6.1	~ 1.5
$\hat{\alpha}_3$	~ 6.1	~ 3.6

410 – there is a factor 10 between the estimated variance \hat{r}_Φ of the time model and the time-altitude model;


– for the 1-dimensional case, $\hat{r}_\Phi < \hat{r}_B$ whereas in the 2-dimensional case we have $\hat{r}_\Phi > \hat{r}_B$, which means adding a dimension to the model helps to find some structure in the data that were missing otherwise;

– the amplitudes and phases of the first and second harmonics ($\hat{\beta}_2$, $\hat{\beta}_3$, $\hat{\alpha}_2$ and $\hat{\alpha}_3$) differ substantially.


415 This last point explains the distinct mean behavior for the two models. The mean of altitude-time model has a lower amplitude which can be explained by the fact that the model must accommodate for a greater variability of observations due to an underlying altitude dependence of the dataset. It is an indication that future development should include this altitude dependence.

7.2 Kriging results


In this section, we present the kriging results obtained for the time-altitude model. The parameters corresponding to this model have been estimated in Sect. 7.1. The kriging estimator is applied on a regular grid of 20×500 positions in time and altitude, 420 this corresponds to a total of 10000 positions. We take $z_0 = [1, \dots, 8]$ km and $t_0 = [2, \dots, 5.5]$ years. In addition, we excluded observations made after the 5-th year. Therefore, we used a set of $N_{\text{obs}} = 6112$ observations. Fig. 17(a) outlines the situation we have just defined. The obtained kriging surface represented in Fig. 17(b) visually fits the observations from Fig. 17. Fig. 17(c) represents the 2-dimensional map of the variance of the estimation error. We note that the variance of the estimation error is minimal when there are observations in the neighborhood, whereas it increases as we move away from the observations. This is 425 an expected result that is consistent with what has been observed in the case of time kriging. Since the interpretation of kriging surfaces is complex, we represent these results at constant altitude (*cf.* Fig. 18). On constant height sections, the estimation structure seems more complex than in the 1-dimensional case, it can be explained by the interaction with observations at



(a)



(b)



(c)

Figure 17. CloudSat observations on geographic area 6. The red rectangle (□) indicates the training dataset, the green rectangle (□) indicates observations not used in the kriging estimate and the plain black lines represent the spatialtemporal positions where the kriging estimates are computed (a). Result of the 2-dimensional kriging on zone 6 (b) and its associated 2-dimensional variance of the estimation error (c).

altitudes above and below the considered altitude. Moreover, we note that the model tends to the mean of the model when we move away from observations. This is especially true in the case of long-term prediction around the 5-th year.




Figure 18. Kriging results at constant altitudes, $z \simeq 2.17$ km (a), $z \simeq 3.62$ km (b), $z \simeq 5.08$ km (c), $z \simeq 5.96$ km (d). We represent the observations ψ (●), the estimations $\hat{\varphi}_0$ (●), the mean (—) as well as 2σ around the estimation $\hat{\varphi}_0$.

430 8 Conclusion and perspectives

The interpolation, in space, and prediction, in time, of the cloud microphysics in medium and long term are of major importance in weather and climate analysis. Since a perfect estimation is obviously unattainable, it is an issue of uncertainty quantification.

In this original work, we develop a statistical spatio-temporal kriging-based approach that is able to interpolate/predict from the dataset and provide uncertainties. Beforehand, it requires in particular estimating the covariance model parameters; it is

435 performed in a Bayesian setting, which allows for estimation and uncertainty quantification. The approach is then applied to a subset of the CloudSat dataset which shows promising results, especially in the 2-dimensional case where detailed structure appears in the quantity of interest.

A natural extension to this approach would be to consider the latitude and longitude variables in order to interpolate horizontally the quantity of interest. **Other extensions should consider the joint estimation of Ice Water Content and Liquid Water Content, the estimation of cloud types (cumulus, stratus, *etc.*) or some nonlinear functions of our**

440 **actual quantity of interest (*i.e.*, some parametric model that depends directly on Liquid Water Content or Ice Water Content like optical properties). The treatment of these problems would require to resort to suitable kriging estimators (co-kriging, disjunctive kriging, *etc.*), or, more likely, some adapted versions of them. Finally, the impact of the parameter uncertainty in the kriging results could be rigorously handle by developing a complete Bayesian hierarchical model.**

445 *Data availability.* The level 2B-CWC-RO CloudSat data used in this study have been downloaded from the CloudSat Data Processing Center
(<http://www.cloudsat.cira.colostate.edu/>).

Author contributions. PM and JFG designed the preliminary study, with continuous discussions from all co-authors along the project. JML developed the model code and performed the simulations. JML prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

450 *Acknowledgements.* The authors would like to thank Audrey Giremus for her contribution to the preliminary step of the work and the two anonymous reviewers for their suggestions and comments.

References

Austin, R. T., Heymsfield, A. J., and Stephens, G. L.: Retrieval of ice cloud microphysical parameters using the CloudSat millimeter-wave radar and temperature, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD010049>, 2009.

Banerjee, S., Carlin, B., and Gelfand, A.: *Hierarchical Modeling and Analysis for Spatial Data*, Second Edition, Chapman & Hall/CRC Monographs on Statistics & Applied Probability, Taylor & Francis, 2014.

Belo-Pereira, M., Dutra, E., and Viterbo, P.: Evaluation of global precipitation data sets over the Iberian Peninsula, *Journal of Geophysical Research: Atmospheres*, 116, [https://doi.org/https://doi.org/10.1029/2010JD015481](https://doi.org/10.1029/2010JD015481), 2011.

Bodas-Salcedo, A., Webb, M., Brooks, M., Ringer, M., Williams, K., Milton, S., and Wilson, D.: Evaluating cloud systems in the Met Office global forecast model using simulated CloudSat radar reflectivities, *Journal of Geophysical Research: Atmospheres*, 113, 2008.

Brockwell, P. J. and Davis, R. A.: *Time Series: Theory and Methods*, Springer Series in Statistics, Springer New York, 2009.

Chen, W.-T., Woods, C. P., Li, J.-L. F., Waliser, D. E., Chern, J.-D., Tao, W.-K., Jiang, J. H., and Tompkins, A. M.: Partitioning CloudSat ice water content for comparison with upper tropospheric ice in global atmospheric models, *Journal of Geophysical Research: Atmospheres*, 116, 2011.

Chiles, J.-P. and Delfiner, P.: *Geostatistics: modeling spatial uncertainty*, Wiley, New York, 1999.

Chonavel, T. and Ormrod, J.: *Statistical Signal Processing: Modelling and Estimation*, Advanced Textbooks in Control and Signal Processing, Springer London, 2002.

Cressie, N.: *Statistics for spatial data*, Wiley series in probability and mathematical statistics: Applied probability and statistics, J. Wiley, 1993.

Cressie, N. and Wikle, C.: *Statistics for Spatio-Temporal Data*, Wiley, 2015.

der Meer, F. V.: Remote-sensing image analysis and geostatistics, *International Journal of Remote Sensing*, 33, 5644–5676, <https://doi.org/10.1080/01431161.2012.666363>, 2012.

Didari, S. and Zand-Parsa, S.: Enhancing estimation accuracy of daily maximum, minimum, and mean air temperature using spatio-temporal ground-based and remote-sensing data in southern Iran, *International Journal of Remote Sensing*, 39, 6316–6339, <https://doi.org/10.1080/01431161.2018.1460500>, 2018.

Diggle, P. J., Ribeiro, P. J., and Christensen, O. F.: *An Introduction to Model-Based Geostatistics*, Springer New York, New York, NY, https://doi.org/10.1007/978-0-387-21811-3_2, 2003.

Eriksson, P., Ekström, M., Rydberg, B., Wu, D. L., Austin, R. T., and Murtagh, D. P.: Comparison between early Odin-SMR, Aura MLS and CloudSat retrievals of cloud ice mass in the upper tropical troposphere, *Atmospheric Chemistry and Physics*, 8, 1937–1948, 2008.

Florio, E. N., Lele, S. R., Chang, Y. C., Sterner, R., and Glass, G. E.: Integrating AVHRR satellite data and NOAA ground observations to predict surface air temperature: a statistical approach, *International Journal of Remote Sensing*, 25, 2979–2994, <https://doi.org/10.1080/01431160310001624593>, 2004.

Gaetan, C. and Guyon, X.: *Modélisation et statistique spatiales*, Mathématiques et Applications, Springer Berlin Heidelberg, 2008.

Genton, M. G.: Separable approximations of space-time covariance matrices, *Environmetrics*, 18, 681–695, <https://doi.org/10.1002/env.854>, 2007.

Heuvelink, G., Griffith, D., Hengl, T., and Melles, S.: Sampling Design Optimization for Space-Time Kriging, *Spatio-Temporal Design: Advances in Efficient Data Acquisition*, pp. 207–230, <https://doi.org/10.1002/9781118441862.ch9>, 2012.

Jewell, S. and Gaussiat, N.: An Assessment of Kriging Based Rain-Gauge-Radar Merging Techniques, *Quarterly Journal of the Royal Meteorological Society*, 141, <https://doi.org/10.1002/qj.2522>, 2015.

Khan, M. S., Grabner, M., Muhammad, S. S., Awan, M. S., Leitgeb, E., Kvicer, V., and Nebuloni, R.: Empirical Relations for Optical Attenuation Prediction from Liquid Water Content of Fog., *Radioengineering*, 21, 2012.

490 Kiefer, J.: Sequential minimax search for a maximum, *Proceedings of the American Mathematical Society*, 4, 502–506, 1953.

Krige, D.: A statistical approach to some basic mine valuation problems on the Witwatersrand, *Journal of the Southern African Institute of Mining and Metallurgy*, 52, 119–139, https://doi.org/10.10520/AJA0038223X_4792, 1951.

495 Lyras, N. K., Kourogiorgas, C. I., and Panagopoulos, A. D.: Cloud attenuation statistics prediction from Ka-band to optical frequencies: Integrated liquid water content field synthesizer, *IEEE Transactions on Antennas and Propagation*, 65, 319–328, 2016.

Marshak, A. and Davis, A.: 3D Radiative Transfer in Cloudy Atmospheres, *Physics of Earth and Space Environments*, Springer Berlin Heidelberg, 2005.

500 Matheron, G.: Principles of geostatistics, *Economic Geology*, 58, 1246–1266, <https://doi.org/10.2113/gsecongeo.58.8.1246>, 1963.

Montero, J.-M., Fernández-Avilés, G., and Mateu, J.: Spatial and Spatio-Temporal Geostatistical Modeling and Kriging, Wiley Series in Probability and Statistics, Wiley, 2015.

Nour, M., Smit, D., and Gamal El-Din, M.: Geostatistical mapping of precipitation: implications for rain gauge network design, *Water Science and Technology*, 53, 101–110, <https://doi.org/10.2166/wst.2006.303>, 2006.

Park, N.-W.: Time-Series Mapping of PM10 Concentration Using Multi-Gaussian Space-Time Kriging: A Case Study in the Seoul Metropolitan Area, Korea, *Advances in Meteorology*, 2016, 1–10, 2016.

505 Press, W. H., Teukolsky, S. A., Vetterling, W. T., and Flannery, B. P.: *Numerical Recipes in C* (2nd Ed.): The Art of Scientific Computing, Cambridge University Press, New York, NY, USA, 1992.

Qu, Z., Barker, H. W., Korolev, A. V., Milbrandt, J. A., Heckman, I., Bélair, S., Leroyer, S., Vaillancourt, P. A., Wolde, M., Schwarzenböck, A., et al.: Evaluation of a high-resolution numerical weather prediction model's simulated clouds using observations from CloudSat, GOES-13 and in situ aircraft, *Quarterly Journal of the Royal Meteorological Society*, 144, 1681–1694, 2018.

510 Rasmussen, C. E. and Williams, C. K. I.: *Gaussian Processes for Machine Learning* (Adaptive Computation and Machine Learning), The MIT Press, 2005.

Ripley, B. D.: *Spatial statistics*, Wiley New York, 1981.

Stephens, G. L. and Kummerow, C. D.: The Remote Sensing of Clouds and Precipitation from Space: A Review, *Journal of the Atmospheric Sciences*, 64, 3742–3765, 2007.

515 Stephens, G. L., Vane, D. G., Boain, R. J., Mace, G. G., Sassen, K., Wang, Z., Illingworth, A. J., O'Connor, E. J., Rossow, W. B., Durden, S. L., Miller, S. D., Austin, R. T., Benedetti, A., Mitrescu, C., and Team, C. S.: The CloudSat mission and the A-Train: A new dimension of space-based observations of clouds and precipitation, *Bull. Amer. Meteorol. Soc.*, 83, 1771–1790, <https://doi.org/10.1175/BAMS-83-12-1771>, 2002.

Varouchakis, E. A., Kamińska-Chuchmala, A., Kowalik, G., Spanoudaki, K., and Graña, M.: Combining Geostatistics and Remote Sensing Data to Improve Spatiotemporal Analysis of Precipitation, *Sensors*, 21, <https://doi.org/10.3390/s21093132>, 2021.

520 Verdin, A., Funk, C., Rajagopalan, B., and Kleiber, W.: Kriging and Local Polynomial Methods for Blending Satellite-Derived and Gauge Precipitation Estimates to Support Hydrologic Early Warning Systems, *IEEE Transactions on Geoscience and Remote Sensing*, 54, 2552–2562, <https://doi.org/10.1109/TGRS.2015.2502956>, 2016.

Vivekanandan, J., Zhang, G., and Politovich, M.: An assessment of droplet size and liquid water content derived from dual-wavelength radar
525 measurements to the application of aircraft icing detection, *Journal of atmospheric and oceanic technology*, 18, 1787–1798, 2001.

Wackernagel, H.: *Multivariate Geostatistics: An Introduction with Applications*, Springer Berlin Heidelberg, 2013.

Wu, D. L., Austin, R. T., Deng, M., Durden, S. L., Heymsfield, A. J., Jiang, J. H., Lambert, A., Li, J.-L., Livesey, N. J., McFarquhar, G. M.,
Pittman, J. V., Stephens, G. L., Tanelli, S., Vane, D. G., and Waliser, D. E.: Comparisons of global cloud ice from MLS, CloudSat, and
correlative data sets, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD009946>, 2009.

530 Yang, S. and Zou, X.: Assessments of cloud liquid water contributions to GPS radio occultation refractivity using measurements from
COSMIC and CloudSat, *Journal of Geophysical Research: Atmospheres*, 117, 2012.

Zakeri, F. and Mariethoz, G.: A review of geostatistical simulation models applied to satellite remote sensing: Methods and applications,
Remote Sensing of Environment, 259, 112 381, <https://doi.org/https://doi.org/10.1016/j.rse.2021.112381>, 2021.