# Answers to the Referee 1 about the paper amt-2021-361:

"Towards the use of conservative thermodynamic variables in data assimilation: **preliminary results a case study** using ground-based microwave radiometer measurements"

> by Pascal Marquet, Pauline Martinet, Jean-François Mahfouf, Alina Lavinia Barbu, and Benjamin Ménétrier.

> > January 21, 2022

# 1) Answers to the referee RC1

• The paper demonstrates the analyses of model profiles with the assimilation of ground- based microwave radiometer (MWR) brightness temperatures with a 1D-Var scheme and the use of two conservative thermodynamic variables as control variables in a case study. The presented work shows the potential of using the described conservative ther- modynamic variables for 1D-Var MWR retrievals, and the benefit of the obtained, less weather-dependent, model error covariance matrices. It also hints at the benefit of MWR data for data assimilation in the studied fog case and is therefore a valuable step towards the use of MWR data in operational NWP systems.

• The methodology is well explained, the experiment is carefully analysed and results are clearly presented. The assessment is done both in observation and model space. The manuscript is well written, and the figures are illustrative of the relevant results.

• I support the publication of this article after the authors addressed the comments and suggestions listed below.

We thank you for this review and have taken into account almost all your comments and suggestions, as described below.

Please find, also, a document in "difference" mode where it is possible to isolate all changes in colour mode (from red to blue).

# 2) General comments

## Section 2.1 / The moist-air entropy potential temperature

• The section explains the most important points in the derivation of the moist-air entropy potential temperature and provides an extensive list of references for further information. Although it is beyond the scope of this manuscript to go in full details here, I found the description of the conservation laws (line 74ff) a little short and difficult to follow for people not familiar with these variables. Some sentences are also quite long and complicated. I would very much welcome a reformulation of this paragraph.

Several of the sentences have been rewritten in this paragraph, and some of the sentences in the conclusion have been moved in this paragraph.

# Section 4.3 (Vertical profiles of analysis increments) and Section 5 (Conclusions)

• Do you have the possibility to make a stronger link between the increments and potential improvements in the model profiles?

Two new figures 7 have been introduced and commented in the text at the end of section 4.3.

It is explained that:

"Some radiosoundings (RS) have been launched during the SOFOG3D IOPs. As only one RS profile was launched at 05:21~UTC on the case study presented in the article, no statistical evaluation of the profile increments can be carried out. However, we have conducted an evaluation of the analysis increments obtained at 5 and 6 UTC (the 1D-Var retrievals being performed at a 1h temporal resolution in line with the operational AROME assimilation cycles) around the RS launched time. The AROME temperature background profile extracted at 6 UTC has a vertical structure close to the RS launched at 05:21~UTC.

Figs 7 compare the AROME background profile and 1D-Var analyses performed with the REF and EXP experiments against the RS profile. The temperature increments are a step in the right direction by cooling the AROME background profile in line with the observed RS profile. The two 1D-Var analyses are close to each other, but the EXP analysis produces a temperature profile slightly cooler compared to the REF analysis. In terms of absolute humidity ( $\rho_v = p_v/(R_v T)$ , with  $p_v$  the partial pressure and  $R_v$  the gas constant for water vapour), the background profile already exhibits a similar structure compared to the RS profile. The EXP profile is slightly moister than the REF profile from the surface up to 3500 m which leads to a somewhat better agreement with the RS profile below 1500 m. The 1D-Var increments are thus small and close between the two experiments, although an improvement of the integrated water vapor (IWV) is observed when compared to the background, with a difference against the RS IWV reduced from almost 1 kg m<sup>-2</sup> in the background to less than 0.4 kg m<sup>-2</sup> in the 1D-Var analyses. These analyses confirm the improvement brought to the model profiles by both the REF and EXP analysis increments, with some enhanced improvement for EXP. "

• Are the increments located at the right place and going in the desired direction?

One can notice from results displayed in Figure 7 that, in the boundary layer (below 1000 m), the corrections induced by the assimilation are improving the profiles with respect to radiosounding data (temperature decrease and humidity increase). The analysis profiles from EXP are slightly closer to the RS profiles than those from REF.

• You provide a good indication for temperature increments with respect to MWR measurements in line 271ff. A comparison to soundings, as you mention in line 321, could be especially valuable. Would it be possible to provide a preliminary assessment?

This preliminary assessment is shown by the addition of Figure 7 in the revised paper and the corresponding discussion given section 4.3.

# 3) Specific comments

### Title

• I suggest changing the title to 'Towards the use of conservative thermodynamic variables in data assimilation: a case study using ground-based microwave radiometer measurements' to make the scope of the manuscript clearer.

The title has been modified as suggested.

## Section 2.2

• The 1D-Var formalism: Very clear and concise description of 1D-Var, pleasant to read! It would maybe be good to include a reference to a more detailed description of 1D-Var theory.

Two additional references are given in this section regarding 1D-Var retrieval theory: Rodgers (1976) and Thépaut and Moll (1990).

## Section 2.3

• The conversion operator: To help make it easier to understand, it would be helpful to add in this section that this conversion from  $((\theta_s)_a, q_t)$  to  $(T, q_v, q_l)$  is needed to feed RTTOV after the analysis.

Additional explanation is given regarding the use of the conversion operator in the 1D-Var: "The second term on the right hand side of Eq. (11) and (12) indicates that the conversion operator  $\mathcal{L}$  is needed to compute the brightness temperatures from the observation operator  $\mathcal{H}$ . Indeed RTTOV-gb requires profiles of temperature, specific humidity and liquid water content as input quantities. This space change is required at each step of the minimisation process. For the computation of the gradient of the costfunction  $\nabla_z \mathcal{J}$ , the linearized version (adjoint) of  $\mathcal{L}$  is also necessary. In practice, the operator  $\mathbf{L}^T$ provides the gradient of the brightness temperatures with respect to the conservative variables, knowing the gradient with respect to the classical variables."

## Section 4.1

• The background error cross correlations: I am not familiar with the variable  $(\theta_s)_a$ . Could you make a little clearer why you expect a link between  $(\theta_s)_a$  and  $q_t$  in clear-sky atmosphere (line 234)? If this should be evident from section 2.1 it would maybe be good to highlight it more.

This aspect is explained in the revised version by pointing to Equations (3) and (4) that show the linear relationship between  $(\theta_s)_a$  and  $q_t$ .

# Section 4.2

• 1D-Var analysis fit to observations: In order to better compare the REF and EXP analyses in the regions of concern, it might be interesting to compute and compare RMSE and bias only within the blue rectangular boxes.

The statistics of the fit to the observation have been recomputed in the regions of concern corresponding to the blue rectangular boxes. We included the results in the new Table 2 and commented the results directly in the manuscript with the following sentences:

These statistics have also been calculated by restricting the dataset to the two dashed rectangular boxes represented in Figs 5 (b) and (c). A significant improvement is observed for channels the most sensitive to liquid water in the afternoon with a RMSE decreased from 4.3 K in the background to 0.57 K in REF and 0.37 K in EXP. For all the computed statistics, EXP always provides the best performance in terms of RMSE. Table 2 summarizes the bias and RMSE values obtained for the different samples.

## Conclusions

• Line 300ff Maybe this paragraph would better suit to section 2.1, where the variables are introduced and described.

Indeed, this paragraph is moved into the section 2.1, as suggested.

• Line 326ff Do you expect results of TB to vary a lot after including q i in the conversion operator? Where there ice clouds in the presented case (line 327)?

We have added into the conclusion the following paragraph:

Indeed, inclusion of  $q_i$  in the conversion operator should lead to more realistic retrieved profiles of cloud condensates, and a 1D-var system with only  $q_l$  can create water clouds at locations where ice clouds should be present, as done in our experiment around 400 hPa between 15 and 24 UTC. However, since the frequencies of HATPRO are not sensitive to ice water content, the fit of simulated Tbs to observations could be reduced. In consequence, the synergy with an instrument sensitive to ice water clouds, such as a W-band cloud radar, is necessary to get improved retrievals of both  $q_i$  and  $q_l$ .

# 4) Technical corrections

## 4.1) The main text:

• Line 5: '[...] that are currently highly dependent on weather conditions [...]'  $\rightarrow$  '[...] that are highly

dependent on weather conditions when using classical variables, [...]'

The new text is introduced.

• Line 28: 'scheme'  $\rightarrow$  'schemes'

The name 'schemes' is used.

• Line 38: '[...] that data assimilation systems used to be [...]'  $\rightarrow$  '[...] that most data assimilation systems are [...]'

The new text is introduced.

• Line 67: Use  $s_{d0}(T_0, p_0)$  in the formula, as in line 69.

We have used  $s_{d0}(T_0, p_0)$  in the formula.

• Line 71: State that the explanation for  $\Lambda_r$  follows later in the section.

This sentence is added.

• Line 73: '[...] sublimation.'  $\rightarrow$  '[...] sublimation, respectively.'

The word 'respectively' is used.

• Line 106: Remove the point at the end of the title.

This is done.

• Line 143: 'set-up'  $\rightarrow$  'setup'

This change is made.

• Line 147: '[...] such a HATPRO [...]'  $\rightarrow$  '[...] such as a HATPRO [...]'

This change is introduced, together with a larger change suggested by the RC2.

• Line 155: 'lowest' instead of 'first few'

The word 'lowest' is introduced.

• Line 192: I propose to mention RTTOV-gb here after H because it was already mentioned earlier in the text.

The change is made.

• Line 195: comma after '(LWP)'

The comma is added.

• Line 201: No new paragraph before line 201

This 'new paragraph' is canceled.

• Line 236-237:  $B_z(\theta_s, q_t) \to B_z((\theta_s)_a, q_t)$ ?

This change is made (twice).

• Line 276: 'similarity'  $\rightarrow$  'similarities'

This change is done.

• Lines 284-287: Suggestion: 'Most of the liquid water is created in low clouds. Additionally, increments of  $q_l$  above 600 hPa are larger and more extended vertically and in time in EXP, where condensation occurs over a thicker atmospheric layer between 500 hPa and 300 hPa after 12 UTC. In the REF experiment, the creation of liquid water above 500 hPa only reaches values of 0.3 g/kg sporadically, for example at 21UTC. In the EXP setup, [...]'

We have introduced the proposed sentences.

• Line 289: '[...] keeps unchanged the vertical structure of the  $q_l$  profile [...]'  $\rightarrow$  '[...] keeps the vertical structure of the  $q_l$  profile unchanged [...]'

We have introduced your text.

• Lines 289-290: 'Liquid water is added where it already existed  $[...]' \rightarrow$  'In REF, liquid water is only added where it already exists [...]'

We have introduced your text.

• Line 293: '[...] show similar structures to [...]'  $\rightarrow$  '[...] show structures similar to [...]'

We have introduced your text.

• Line 296: 'value' instead of 'interest' ?

This change is done.

• Line 297: 'for assimilating'  $\rightarrow$  'to assimilate'

This change is done.

• Line 299: 'over South-Western France'  $\rightarrow$  'in South-Western France'

This change is done.

• Line 303: '(in e.g. fronts, [....])'

This change is done.

• Line 305: 'significance' instead of 'interest' ?

This change is done.

• Line 324: 'increment'  $\rightarrow$  'increments'

This change is done.

## 4.2) The Figures:

• Figure 1: Would it be possible to provide this figure with slightly larger font size?

The font of the annotations have been enlarged.

• Figure 2: – Do you mean 2 K instead of 0.2 K in the caption? – 'Relative Humidity' without capital letters – Could you add a description of the arrows (2(b), 2(d)) in the caption?

It is indeed 2 K; we write 'relative humidity'; we add:

"The vertical arrows in (b) and (d) indicate areas where potential temperatures are almost homogeneous or constant along the vertical."

• Figure 5: – Please make the numbering of the channels consistent between figure (1-13) and caption (0-12). Would it maybe be possible to even enumerate the channels in the figure after the MWR channel numbers (1-2,4-14 as channel 3 was removed, if I am correct)? – Provide units to the color bar. – I find it a little difficult to compare the values between panels (a) and (b)/(c) and really point out the magnitude of the reduction of deviations. Would it be possible to use the same range in the color bar of all three panels? Or alternatively at least point out the different scaling in the caption for clarity.

The same range of color bar cannot be used for the three panels because background departures have much larger values (up to 15 K) than analysis residuals (up to 3 K). However, the same scale has been adopted for the analysis residuals of the two 1D-Var systems. In order to avoid any visual misinterpretation of the plots, the color map is now different for the background departures and for the analysis residuals.

• Figure 6: I suggest centering the color map to fit 0 to the change of color between blue and red.

The color maps have been changed and centered to 0 between blue and red.

# Towards the use of conservative thermodynamic variables in data assimilation: preliminary results a case study using ground-based microwave radiometer measurements

Pascal Marquet<sup>1</sup>, Pauline Martinet<sup>1</sup>, Jean-François Mahfouf<sup>1</sup>, Alina Lavinia Barbu<sup>1</sup>, and Benjamin Ménétrier<sup>2</sup>

<sup>1</sup>CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France <sup>2</sup>INP, IRIT, Université de Toulouse, Toulouse, France

**Correspondence:** Pascal Marquet (pascal.marquet@meteo.fr, pascalmarquet@yahoo.com) Pauline Martinet (pauline.martinet@meteo.fr)

Abstract. This study aims at introducing two conservative thermodynamic variables (moist-air entropy potential temperature and total water content) into a one-dimensional variational data assimilation system (1D-Var) to demonstrate the

<sup>5</sup> benefit for future operational assimilation schemes. This system is assessed using microwave brightness temperatures from a ground-based radiometer installed during the field campaign SOFGO3D\_SOFOG3D\_dedicated to fog forecast improvement.

- An underlying objective is to ease the specification of background error covariance matrices that are currently that are highly dependent on weather conditions when using classical variables, making difficult the optimal retrievals of cloud and thermodynamic properties during fog conditions.
- <sup>15</sup> Background error covariance matrices for these new conservative variables have thus been computed by an ensemble approach based on the French convective scale model AROME, for both all-weather and fog conditions. A first result shows that the use of these matrices for the new variables reduces
- <sup>20</sup> some dependencies to the meteorological conditions (diurnal cycle, presence or not of clouds) compared to usual variables (temperature, specific humidity).

Then, two 1D-Var experiments (classical vs. conservative variables) are evaluated over a full diurnal cycle char-<sup>25</sup> acterized by a stratus-evolving radiative fog situation, using hourly brightness temperatures.

Results show, as expected, that analysed brightness temperatures by the 1D-Var are much closer to the observed ones than background values for both variable choices. This is es-

<sup>30</sup> pecially the case for channels sensitive to water vapour and liquid water. On the other hand, analysis increments in model

space (water vapour, liquid water) show significant differences between the two sets of variables.

#### 1 Introduction

Numerical Weather Prediction (NWP) models at convective 35 scale need accurate initial conditions for skillful forecasts of high impact meteorological events taking place at small-scale such as convective storms, wind gusts or fog. Observing systems sampling atmospheric phenomena at small-scale and high temporal frequency are thus necessary for that purpose 40 (Gustafsson et al., 2018). Ground-based remote-sensing instruments (e.g. rain and cloud radars, radiometers, wind profilers) meet such requirements and provide information on wind, temperature, and atmospheric water (vapour and hydrometeors). Moreover, data assimilation systems are evolv- 45 ing towards ensemble approaches where hydrometeors can be initialized together with usual control variables. This is the case for the Météo-France NWP limited area model AROME (Seity et al., 2011; Brousseau et al., 2016) where, on top of wind (U, V), temperature (T) and specific humidity  $q_v$ , sev- 50 eral hydrometeor mass contents can be initialized (cloud liquid water  $q_l$ , cloud ice water  $q_i$ , rain  $q_r$ , snow  $q_s$  and graupel  $q_a$ ) (Destouches et al., 2021). However, these variables are not conserved during adiabatic and reversible vertical motion. 55

The accuracy of the analysed state in variational scheme schemes highly depends on the specification of the so-called background error covariance matrix. Background error variances and cross-correlations between variables are known to be dependent on weather conditions (Montmerle and Berre, 60 2010; Michel et al., 2011). This is particularly the case during fog conditions with much shorter vertical correlation length-scales at the lowest levels and large positive cross-correlations between temperature and specific humid-

- <sup>5</sup> ity (Ménétrier and Montmerle, 2011). In this context, Martinet et al. (2020) have demonstrated that humidity retrievals could be significantly degraded if sub-optimal background error covariances are used during the minimization. New ensemble approaches allow a better approximation of back-
- <sup>10</sup> ground error covariance matrices but rely on the capability of the ensemble data assimilation to correctly represent model errors, which might not always be the case during fog conditions. This is why it would be of interest to examine, in a data assimilation context, the use of variables that are more <sup>15</sup> suitable when water phase changes take place.

It is well-known that that most data assimilation systems used to be were based on the assumptions of homogeneity and isotropy of background error correlations. To release these hypotheses, Desroziers and Lafore (1993)

- <sup>20</sup> and Desroziers (1997) implemented a coordinate change inspired by the semi-geostrophic theory to test flow-dependent analyses with case studies from the Front-87 field campaign (Clough and Testud, 1988), where the local horizontal coordinates were transformed into the semi-geostrophic
- <sup>25</sup> space during the assimilation process. Another kind of flowdependent analyses were made by Cullen (2003) and Wlasak et al. (2006) who proposed a low-order Potential Vorticity (PV) inversion scheme to define a new set of control variables. Similarly, analyses on potential temperature  $\theta$  were

<sup>30</sup> made by Shapiro and Hastings (1973) and Benjamin et al. (1991), and more recently by Benjamin et al. (2004) with moist virtual  $\theta_v$  and moist equivalent  $\theta_e$  potential temperatures.

The aim of the paper is to test a one-dimensional data assimilation method that would be less sensitive to the average vertical gradients of the  $(T, q_v, q_l, q_i)$  variables. To this end, two conservative variables will be proposed, generalizing previous uses of  $\theta$  (as a proxy for the entropy of dry air) to moist-air variables suitable for data assimilation. <sup>40</sup> The new conservative variables are the total water content  $q_t = q_v + q_l + q_i$  and the moist-air entropy potential temperature  $\theta_s$  defined in Marquet (2011), which generalize the two well-known conservative variables  $(q_t, \theta_l)$  of Betts (1973).

- The focus of the study will be on a fog situation from the 45 SOFOG3D field campaign using a one-dimensional variational (1D-Var) system for the assimilation of observed microwave brightness temperatures sensitive to T,  $q_v$  and  $q_l$ from a ground-based radiometer. Short-range forecasts from the convective scale model AROME (Seity et al., 2011) will
- <sup>50</sup> be used as background profiles, the fast radiative transfer model RTTOV-gb (De Angelis et al., 2016; Cimini et al., 2019) will allow to accurately simulate the brightness temperatures, and suitable background error covariance matrices will be derived from an ensemble technique.

Section 2 presents the methodology (conservative variables, 1D-Var, change of variables). Section 3 describes the experimental setting, the meteorological context, the observations and the different components of the 1D-Var system. The results are commented in Section 4. Finally, conclusions and perspectives are given in Section 5.

#### 2 The methods

This section presents the methodology chosen for this study. The definition of the moist-air entropy potential temperature  $\theta_s$  is introduced, as well as the formalism of the 1D-Var assimilation system, before describing the "conservative variable" conversion operator.

#### 2.1 The moist-air entropy potential temperature

The The motivation for using the absolute moist-air entropy potential temperature in atmospheric science was first described by Richardson (1919, 1922), and then fully <sup>70</sup> formalized by Hauf and Höller (1987). The method is to take into account the absolute value for dry air and water vapour and to define a moist-air entropy potential temperature variable called  $\theta_s$  is defined in Marquet (2011) to provide.

However, the version of  $\theta_s$  published in Hauf and Höller (1987) was not really synonymous with the moist-air entropy. This problem has been solved with the version of Marquet (2011) by imposing the same link with the specific entropy of moist air (s) as in the dry-air so formalism of Bauer (1908), leading to:

75

90

$$s = c_{pd} \ln\left(\frac{\theta_s}{T_0}\right) + s_{d0}(\underline{T_0, p_0}), \qquad (1)$$

where  $c_{pd} \approx 1004.7 \,\mathrm{J\,K^{-1}\,kg^{-1}}$  is the specific heat of dry air dry-air specific heat at constant pressure,  $T_0 = 273.15 \,\mathrm{K}$  is a standard temperature and  $s_{d0}(T_0, p_0) \approx 6775 \,\mathrm{J\,K^{-1}\,kg^{-1}}$ the reference dry-air is the reference dry-air entropy at  $T_0$ and at the standard pressure  $p_0 = 1000 \,\mathrm{hPa}$ . Because  $c_{pd}$ ,  $T_0$ , and  $s_{d0}(T_0, p_0)$  are constant terms,  $\theta_s$  defined by (1) is synonymous with, and has the same physical properties as, the moist-air entropy s.

The conservative aspects of this potential temperature  $\theta_s$ and its meteorological properties (in e.g. fronts, convection, cyclones) have been studied in Marquet (2011), Blot (2013) and Marquet and Geleyn (2015). The links with the definition of the Brunt-Väisälä frequency and the potential vorticity are described in Marquet and Geleyn (2013) and Marquet (2014), while the significance of the absolute entropy to describe the thermodynamics of cyclones is shown in Marquet (2017) and Marquet and Dauhut (2018).

Only the first order approximation  $\underbrace{of}_{\mathfrak{S}_{s}}$  noted  $(\theta_s)_1$  in 100 Marquet (2011), will be considered in the following, with that

#### P. Marquet et al.: Data assimilation using conservative thermodynamic variables

writes:

$$\theta_s \approx (\theta_s)_1 = \theta \exp\left(-\frac{L_{vap} q_l + L_{sub} q_i}{c_{pd} T}\right) \exp(\Lambda_r q_t), \quad (2)$$

where  $\theta = T (p/p_0)^{\kappa}$  is the dry-air potential temperature, p the pressure,  $\kappa \approx 0.2857$ ,  $L_{vap}(T)$  and  $L_{sub}(T)$  the la-

 $_{5}$  tent heat of vaporization and sublimation. sublimation, respectively. The explanation for  $\Lambda_{r}$  follows later in the section.

The first term  $\theta$  in on the right-hand side of (2) leads to a first conservation law (invariance) during adiabatic com-<sup>10</sup> pression and expansionand with the joint, with joint and opposite variations of T and p to keep keeping  $\theta$  constant. Here lies the motivation for using  $\theta$  to describe dry-air convective processes, and also in data assimilation systems

- by Shapiro and Hastings (1973) and Benjamin et al. (1991). The first exponential explains another on the right-hand side of (2) explains a second form of conservation law: due to . Indeed, this exponential is constant for reversible and adiabatic phase changesfor which  $e_{pd} dT \approx e_p dT = L_{vap} dq_t + L_{sub} dq_t$ , with  $e_p$  the moist-air
- <sup>20</sup> value, for which  $d(c_{pd}T) \approx d(L_{vap}q_l + L_{sub}q_i)$  due to the approximate conservation of the moist-static energy  $c_{pd}T - L_{vap}q_l - L_{sub}q_i$ , and with therefore joint variations of the numerator and denominator of this first exponential, leaving this exponential invariant, and a constant fraction into
- <sup>25</sup> the first exponential. It should be mentioned that the product of  $\theta$  by the this first exponential forms the Betts (1973) conservative variable  $\theta_l$ , which is presently used together with  $q_t$  to describe the moist-air turbulence in GCM and NWP models.
- <sup>30</sup> While the Betts variable variable  $\theta_l$  was established with the assumption of a constant total water content  $q_t$  in Betts (1973), the second exponential in (2) sheds new light on conservation cases a third and new conservation law, where the entropy of moist air can remain constant despite
- <sup>35</sup> changes in the total water  $q_t$ . This occurs in regions where water vapour turbulence transport takes place, <u>or</u> via the evaporation process over oceans, or at the edges of clouds via entrainment and detrainment processes.

We consider here "open-system" thermodynamic pro-40 cesses, for which the second exponential takes into account the impact on moist-air entropy when the changes in specific content of water vapour is-are balanced, numerically, by opposite changes of that of dry air, namely with  $dq_d = -dq_t \neq$ 0. In this case, as stated in Marquet (2011), the changes in 45 moist-air entropy depend on reference values (with subscript "r") according to  $d[q_d (s_d)_r + q_t (s_v)_r]$ , and thus with  $(s_d)_r$ and  $(s_v)_r$  being constant and with the relation  $q_d = 1 - q_t$ , it leads leading to  $[(s_v)_r - (s_d)_r] dq_t$ .

This explains the new term  $\Lambda_r = [(s_v)_r - (s_d)_r]/c_{pd} \approx$ <sup>50</sup> 5.869 ± 0.003, which depends on the absolute reference entropies for water vapour  $(s_v)_r \approx 12671 \,\mathrm{J \, K^{-1} \, kg^{-1}}$  and dry air  $(s_d)_r \approx 6777 \,\mathrm{J \, K^{-1} \, kg^{-1}}$ . This also explains that these "open-system" thermodynamic effects can be taken into account to highlight regimes where the specific moist-air entropy (s),  $\theta_s$  and  $(\theta_s)_1$  can be constant in spite of variable values of despite changes in  $q_t$ , which may decrease or increase on the vertical (see Marquet, 2011, for such examples).

Although it should be possible to use  $(\theta_s)_1$  as a control variable for assimilation, it appeared desirable to define an additional approximation of this variable for a more "regular" and more "linear" formulation, insofar as tangent-linear and adjoint versions are needed for the 1D-Var system. Considering the approximation  $\exp(x) \approx 1 + x$  for the two exponentials in (2), neglecting the second order terms in  $x^2$ , also neglecting the variations of  $L_v(T)$  with temperature and assuming a no-ice hypothesis ( $q_i = 0$ ), the new variable writes:

$$(\theta_s)_a = \theta \left[ 1 + \Lambda_r q_t - \frac{L_{vap}(T_0) q_l}{c_{pd} T} \right], \tag{3}$$

$$(\theta_s)_a = \frac{1}{c_{pd}} \left(\frac{p_0}{p}\right)^{\kappa} \left[ c_{pd} \left( 1 + \Lambda_r \, q_t \right) T - L_{vap}(T_0) \, q_l \right], \quad (4)$$

where  $L_{vap}(T_0) \approx 2501 \text{ kJ kg}^{-1}$ . This formulation corresponds to  $S_m/c_{pd}$ , where  $S_m$  is the Moist Static Energy defined in Marquet (2011, Eq. 73) and used in the ECMWF<sup>1</sup> NWP global model by Marquet and Bechtold (2020).

The new potential temperature  $(\theta_s)_a$  remains close to  $(\theta_s)_1$  (not shown) and keeps almost the same three conservative properties described for  $(\theta_s)_1$ . This new conservative variable  $(\theta_s)_a$  will be used along with the total water content  $q_t = q_v + q_l$  in the data assimilation experimental context described in the following sections.

#### 2.2 The 1D-Var formalism.

The general framework describing the retrieval of atmospheric profiles from remote-sensing instruments by statistical methods can be found in Rodgers (1976). In the following we present the main equations of the one-dimensional variational formalism. Additional details are given in Thépaut and Moll (1990) who developed the first 1D-Var inversion applied to satellite radiances using the adjoint technique.

The 1D-Var data assimilation system searches for an optimal state (the analysis) as an approximate solution of the  $_{90}$ problem minimizing a cost function  $\mathcal{J}$  defined by:

$$\mathcal{J}(x) = \frac{1}{2} (x - x_b)^T \mathbf{B}_{\mathbf{x}}^{-1} (x - x_b) + \frac{1}{2} [y - \mathcal{H}(x)]^T \mathbf{R}^{-1} [y - \mathcal{H}(x)].$$
(5)

The symbol T represents the transpose of a matrix.

The first (background) term measures the distance in  $_{95}$  model space between a control vector x (in our study, T,

80

<sup>&</sup>lt;sup>1</sup>European Centre for Medium range Weather Forecasts

 $q_v$  and  $q_l$  profiles) and a background vector  $x_b$ , weighted weighted by the inverse of the background error covariance matrix ( $\mathbf{B}_{\mathbf{x}}$ ) associated with the vector x. The second (observation) term measures the distance in the observation space

- <sup>5</sup> between the value simulated from the model variables  $\mathcal{H}(x)$ (in our study, the radiative transfer model RTTOV-gb) and the observation vector y (in our study, a set of microwave brightness temperatures from a ground-based radiometer), weighted weighted by the inverse of the observation error <sup>10</sup> covariance matrix (**R**). The solution is searched iteratively
- by performing several evaluations of  $\mathcal{J}$  and its gradient:

$$\nabla_x \mathcal{J}(x) = \mathbf{B_x}^{-1} (x - x_b) - \mathbf{H}^T \mathbf{R}^{-1} [y - \mathcal{H}(x)], \qquad (6)$$

where **H** is the Jacobian matrix of the observation operator representing the sensitivity of the observation operator to <sup>15</sup> changes in the control vector x (**H**<sup>T</sup> is also called the adjoint

of the observation operator).

#### 2.3 The conversion operator

The 1D-Var assimilation defined previously with the variables  $x = (T, q_v, q_l)$  can be modified to use the conservative variables  $z = ((\theta_s)_a, q_l)$ . A conversion operator that projects the state vector from one space to the other can be written as  $x = \mathcal{L}(z)$ . In the presence of liquid water  $q_l$ , an adjustment to saturation is made to separate its contribution to the total water content  $q_t$  from the water vapour content  $q_v$ . This is <sup>25</sup> equivalent to distinguishing the "unsaturated" case from the

"saturated" one. Therefore, starting from initial conditions  $(T_I, q_I) = (T, q_v)$  and using the conservation of  $(\theta_s)_a$  given by Eq. (4), we look for the variable  $T^*$  such that:

$$T^* + \alpha q_{sat}(T^*) = T_I + \alpha q_I, \tag{7}$$

where 
$$\alpha = \frac{L_{vap}(T_0)}{c_{pd} \left(1 + \Lambda_r q_t\right)}$$
 (8)

and  $q_{sat}(T^*)$  is the specific humidity at saturation.

For the unsaturated case  $(q_v < q_{sat}(T^*))$ , we obtain the variables  $(T, q_v, q_l)$  directly from Eq. (4):

$$q_{l} = 0, \ q_{v} = q_{t} \text{ and } T = (\theta_{s})_{a} \left(\frac{p}{p_{0}}\right)^{\kappa} \frac{1}{1 + \Lambda_{r} q_{t}}.$$
 (9)

For the saturated case  $(q_v \ge q_{sat}(T^*))$ , we write:

$$q_l = q_t - q_{sat}(T^*)$$
 and  $q_v = q_{sat}(T^*)$ . (10)

In this situation, it is necessary to calculate implicitly the temperature  $T^*$ , given by Equation (7). We compute numerically an approximation of  $T^*$  by an iterative Newton's algo-<sup>40</sup> rithm.

Taking into account this change of variables, the costfunction can be written as:

$$\mathcal{J}(z) = \frac{1}{2} (z - z_b)^T \mathbf{B}_{\mathbf{z}}^{-1} (z - z_b) + \frac{1}{2} [y - \mathcal{H}(\mathcal{L}(z))]^T \mathbf{R}^{-1} [y - \mathcal{H}(\mathcal{L}(z))]. \quad (11)$$

Then, its gradient given by Eq. (6) becomes:

$$\nabla_{z} \mathcal{J}(z) = \mathbf{B}_{\mathbf{z}}^{-1} (z - z_{b}) - \mathbf{L}^{T} \mathbf{H}^{T} \mathbf{R}^{-1} [y - \mathcal{H}(\mathcal{L}(z))], \quad (12)$$

45

60

where  $\mathbf{L}^T$  is the adjoint of the conversion operator  $\mathcal{L}$ .

The second term on the right hand side of Eq. (11) and (12) indicates that the conversion operator  $\mathcal{L}$  is needed to compute the brightness temperatures from the observation operator  $\mathcal{H}$ . Indeed RTTOV-gb requires profiles of temperature, specific humidity and liquid water content as input quantities. This space change is required at each step of the minimisation process. For the computation of the gradient of the cost-function  $\nabla_z \mathcal{J}$ , the linearized version (adjoint) of  $\mathcal{L}$  is also necessary. In practice, the operator  $\mathbf{L}^T$ provides the gradient of the brightness temperatures with respect to the conservative variables, knowing the gradient with respect to the classical variables.

#### 3 The experimental set-upsetup

The numerical experiments to be presented afterwards will use measurements made during the SOFOG3D field experiment<sup>2</sup> (SOuth west FOGs 3D experiment for processes study) that took place from 1 October 2019 to 31 March 2020 in southwestern France to advance understanding of <sup>65</sup> small-scale processes and surface heterogeneities leading to fog formation and dissipation.

Many instruments were located at the Saint-Symphorien super-site (Les Landes region), such a HATPRO microwave radiometer (Rose et al., 2005)such as a HATPRO (Humidity and Temperature PROfiler, Rose et al., 2005), a 95 GHz BASTA Doppler cloud radar (Delanoë et al., 2016), a Doppler lidar, an aerosol lidar, a surface weather station and a radiosonde station. One objective of this campaign was to test the contribution of the assimilation of such instrumentation on the forecast of fog events by NWP models.

#### 3.1 The 9 February 2020 situation

This section presents the experimental context of 9 February 2020 at the Saint-Symphorien site characterized by (i) a <sup>80</sup> radiative fog event observed in the morning and (ii) the development of low-level clouds in the afternoon and evening.

Figure 1 shows a time series of cloud radar reflectivity profiles (W-band at 95 GHz) measured by the BASTA instrument (Delanoë et al., 2016) in the first few lowest hundred meters (top panel) between 9 February 2020 at 00 UTC and 10 February 2020 at 00 UTC. The instrument reveals a thickening of the fog between midnight and 04 UTC. The fog layer thickness is located between 90 m and 250 m. After 04 UTC, the fog layer near the ground rises, lifting in a "stratus" type cloud (between 100 and 300 m). After 08 UTC, the

<sup>&</sup>lt;sup>2</sup>https://www.umr-cnrm.fr/spip.php?article1086



**Figure 1.** Reflectivity profiles at 95 GHz (dBZ) measured by the BASTA cloud radar in the first 500 m (top) and up to 12000 m altitude (bottom), with UTC hours in abscissas, for the day of 9 February 2020 at Saint-Symphorien (Les Landes region). From http://basta.projet. latmos.ipsl.fr/?bi=bif

stratus cloud dissipates. In the bottom panel BASTA observations up to 12000 m ( $\approx$  200 hPa) indicate low-level clouds after 14 UTC, generally between 1000 m ( $\approx$  900 hPa) and 2000 m ( $\approx$  780 hPa), with a fairly good agreement with  $_5$  AROME short-range (1 h) forecasts (see Fig. 2 (f)). Optically thin (reflectivity below 0 dBZ) high altitude ice clouds are also captured by the radar.

Figure 2 depicts the diurnal cycle evolution in terms of vertical profiles of: (a) absolute temperature T, (b) dry-air <sup>10</sup> potential temperature  $\theta$ , (c) water vapour specific content  $q_v$ , (d) entropy potential temperature  $(\theta_s)_a$ , (e) cloud liquid water specific content  $q_l$  and (f) relative humidity (*RH*), from

1 h AROME forecasts (background) of 9 February 2020 at Saint-Symphorien. At this stage, it is important to indicate <sup>15</sup> that the AROME model has a 90-level vertical discretisation

from the surface up to 10 hPa, with a high resolution in the Planetary Boundary Layer (PBL) since 20 levels are below 2 km.

Figures 2 (e) and (f), for  $q_l$  and RH, show two main satu-<sup>20</sup> rated layers: a fog layer close to the surface between 00 and 09 UTC with the presence of a thin liquid cloud layer aloft at 850 hPa at 00 UTC, and the presence of a stratocumulus cloud between 14 UTC and midnight at 850 hPa. During the night, near surface layers cool down, with a thermal inversion

- <sup>25</sup> that sets at around 01 UTC and persists until 07 UTC. After the transition period between 06 UTC and 09 UTC, when the dissipation of the fog and stratus takes place, the air warms up and the PBL develops vertically (see the black curves plotted where vertical gradients of  $\theta$  in (b) are large). Towards
- <sup>30</sup> the end of the day, the thickness of the PBL remains important until 24 UTC, probably due to the presence of clouds between 800 and 750 hPa that reduces the radiative cooling (see Figs. 2 (c) and (f) for  $q_v$  and RH).

Figure 2(d) reveals weaker vertical gradients for the  $(\theta_s)_a$ <sup>35</sup> profiles, notably with contour lines often vertical and less numerous than those of the T,  $\theta$  and  $q_v$  profiles in (a), (b) and (c), as also shown by more extensive and more numerous vertical arrows in (d) than in (b). Here we see the impact of the coefficient  $\Lambda_r \approx 5.869$  in Eqs. (3)-(4), which allows the vertical gradients of  $\theta$  in (b) and  $q_v$  in (c) to often compensate each other in the formula for  $(\theta_s)_a$ . This is especially true between 980 hPa and 750 hPa in the morning between 04 and 10 UTC, and also within the dry and moist boundary layers during the day.

Note that the dissipation of the fog is associated with a homogenization of  $(\theta_s)_a$  in (d) from 04 to 05 UTC in the whole layer above, in the same way as the transition from stratocumulus toward cumulus was associated with a cancellation of the vertical gradient of  $(\theta_s)_1$  in the Fig. 6 of Marquet and Geleyn (2015). This phenomenon cannot be easily deduced from the separate analysis of the gradients of  $\theta$  and  $q_v$  in (b) and (c). Therefore, three air mass changes can be clearly distinguished during the day. The vertical gradients of  $(\theta_s)_a$  are stronger during cloudy situations, first (i) at night and early morning before 04 UTC and just above the fog, then (ii) at the end of the day above the top-cloud level at 800 hPa; with (iii) turbulence-related phenomena in between that mix the air mass and  $(\theta_s)_a$ , up to the cloudy layer tops that evolve between 950 and 800 hPa from 13 UTC to 17 UTC.

The observations to be assimilated are presented in  $_{60}$  the following. The HATPRO (Humidity And Temperature PROfiler) The HATPRO MicroWave Radiometer (MWR) measures brightness temperatures (*TB*) at 14 frequencies (Rose et al., 2005) between 22.24 and 58 GHz: 7 are located in the water vapour absorption band-K-band and 7 are located in the oxygen absorption band-K-band (see the Table 1). For our study, one of the channels the third channel (at 23.84 GHz) was eliminated because of a receiver failure identified during the campaign. In this preliminary study, we

Channel numbers	1	2~	$\mathbf{X}_{\sim}$	<u>4</u>	5_	<u>6</u>	.7
K-band Frequencies (GHz)	22.24	23.04	$\mathbf{X}_{\sim}$	25.44	26.24	27.84	31.4
K-band $\sigma_o(K)$	1.34	1.71	$\underset{\sim}{\mathbf{X}}$	1.08	1.25	1.17	1.19
Channel numbers	<u>8</u>	<u>9</u>	10	11	12	13	14
V-band Frequencies (GHz)	51.26	52.28	53.86	54.94	56.66	57.3	<u>58</u>
V-band $\sigma_o(K)$	3.21	3.29	1.30	0.37	0.42	0.42	0.36

 Table 1. Channel numbers, band frequencies (GHz) and observation uncertainties (K) prescribed in the observation-error-covariance matrix (from Martinet et al., 2020)

have only considered the zenith observation geometry of the radiometer for the sake of simplicity.

The radiative transfer model  $\mathcal{H}$  <u>RTTOV-gb</u> needed to simulate the model equivalent of the observations, together with  $_{5}$  the choice of the control vector and the specification of the background and observation error matrices, are presented in the next section.

#### 3.2 The components of the 1D-Var

- In 1D-Var systems, the integrated liquid water content, Liq-<sup>10</sup> uid Water Path  $(LWP)_{x}$  can be included in the control vector x as initially proposed by Deblonde and English (2003) and more recently used by Martinet et al. (2020). A first experimental set-up has been defined where the minimization is performed with the control vector being  $(T, q_v, LWP)$ . It
- <sup>15</sup> will be considered as the reference being named "REF". The 1D-Var system chosen for the present study is the one developed by the EUMETSAT NWP SAF<sup>3</sup>, where the minimisation of the cost-function is solved using an iterative procedure proposed by Rodgers (1976) with a Gauss-Newton <sup>20</sup> descent algorithm.

During the minimization process, only the amount of integrated liquid water is changed. In this approach, the two "moist" variables  $q_v$  and LWP are considered to be independent (no cross-covariances for background errors be-

- <sup>25</sup> tween these variables). The second experimental framework, where the control vector is  $z = ((\theta_s)_a, q_t)$ , corresponding to the conservative variables, is named "EXP". The numerical aspects of the 1D-Var minimisation are kept the same as in "REF".
- <sup>30</sup> Then, a set of reference matrices  $\mathbf{B}_{\mathbf{x}}(T,q_v)$  has been estimated every hour using the Ensemble Data Assimilation (EDA) system of the AROME model on 9 February 2020. These matrices were obtained by computing statistics from a set of 25 members providing 3 h forecasts for a subset of
- <sup>35</sup> 5000 points randomly selected in the AROME domain to obtain a sufficiently large statistical sample. Then, matrices associated with fog areas, and noted  $\mathbf{B}_{\mathbf{x}}(T, q_v)_{fog}$ , were computed every hour by applying a fog mask (defined by areas where  $q_l$  is above  $10^{-6} kg kg^{-1}$  for the three lowest model <sup>40</sup> levels), in order to select only model grid points for which fog is forecast in the majority of the 25 AROME members.

The background error covariance matrices  $\mathbf{B}_{\mathbf{z}}((\theta_s)_a, q_t)$  and  $\mathbf{B}_{\mathbf{z}}((\theta_s)_a, q_t)_{fog}$  were obtained in a similar way.

The observation errors are those proposed by Martinet et al. (2020) with values between 1 and 1.7 K for humidity <sup>45</sup> channels (frequencies between 22 and 31 GHz), values between 1 and 3 K for transparent channels affected by larger uncertainties in the modelling of the oxygen absorption band (frequencies between 51 and 54 GHz) and values below 0.5 K for the most opaque channels (frequencies between 55 and <sup>50</sup> 58 GHz).

The RTTOV<sup>4</sup> radiative transfer model is used to calculate brightness temperatures in different frequency bands from atmospheric temperature, water vapour and hydrometeor profiles together with surface properties (provided by outputs from the AROME model). This radiative transfer model has been adapted to simulate ground-based microwave radiometer observations (RTTOV-gb) by De Angelis et al. (2016).

#### 4 Numerical results

The 1D-Var algorithm was tested on the day of 9 February 2020 with observations from the HATPRO microwave radiometer installed at Saint-Symphorien. This section presents and discusses the results obtained on three aspects: (1) the study of background error cross-correlations; (2) the performance of the 1D-Var assimilation system in observation space by examining the fit of simulated TB with respect to the observed ones; and (3) in model space in terms of analysis increments for temperature, specific humidity and liquid water content.

### 4.1 The background error cross correlations

Figure 3 displays for the selected day at  $\frac{06.06}{0.06}$  UTC the crosscorrelations between T and  $q_v$  (top) and between  $(\theta_s)_a$  and  $q_t$  (bottom), with (right) and without (left) fog mask. For the classical variables the correlations are strongly positive in the saturated boundary layer with the fog mask from levels 75 to 90 (between 1015 and 9501015 and 950 hPa), while with profiles in all-weather conditions the correlations between T and  $q_v$  are very weak in the lowest layers. On the other hand, the atmospheric layers above the fog layer exhibit neg-

70

<sup>&</sup>lt;sup>3</sup>Numerical Weather Prediction Satellite Application Facility

<sup>&</sup>lt;sup>4</sup>Radiative Transfer for the TIROS Operational Vertical Sounder



**Figure 2.** Vertical profiles derived from 1 h forecasts of AROME background for all hours of the day 9 February 2020 at Saint-Symphorien (Les Landes region in France) for: (a) absolute temperature T every 0.22 K; (b) dry-air potential temperature  $\theta$  every 0.2 K, (c) water-vapour specific content  $q_v$  every 1 g/kg, (d) entropy potential temperature ( $\theta_s$ )<sub>a</sub> every 0.2 K, (e) cloud liquid-water specific content  $q_l$  (contoured for 0.00001 g/kg and 0.002 g/kg, then every 0.1 g/kg above 0.1 g/kg) and (f) Relative Humidity relative humidity (RH) every 10 %. The black curves (solid and dashed lines) represent the PBL heights determined from maximum of vertical gradients of  $\theta$ . The vertical arrows in (b) and (d) indicate areas where potential temperatures are almost homogeneous or constant along the vertical.

ative correlations between temperature and specific humidity along the first diagonal.

When considering conservative variables, the correlations along the diagonal show a consistently positive siganal in the troposphere (below level 20 located around 28020 located around 280 hPa). Contrary to the classical variables, which are rather independent in clear-sky atmospheres as previously shown by Ménétrier and Montmerle (2011), the  $\mathbf{B}_{\mathbf{z}}$  matrix reflects the physical link between the two new variables (shown by Eq. (4)) as diagnosed from the 10 AROME model. The correlations are positive with and without fog mask. This result shows that the matrix  $\mathbf{B}_{\mathbf{z}}(\theta_s, q_t)$  $\mathbf{B}_{\mathbf{z}}((\theta_s)_{a}, q_t)$  is less sensitive to fog conditions than the  $\mathbf{B}_{\mathbf{x}}$ matrix. It could therefore be possible to compute a  $\mathbf{B}_{\mathbf{z}}(\theta_s, q_t)$  $\mathbf{B}_{\mathbf{z}}((\theta_s)_{a}, q_t)$  matrix without any profile selection criteria 15 that would be nevertheless suitable for fog situations, resulting in a more robust estimate. This result is key for 1D-Var retrievals which are commonly used in the community of ground-based remote sensing instruments to provide databases of vertical profiles for the scientific community. In 5 fact, the accuracy of 1D-Var retrievals is expected to be more

robust with less flow-dependent B matrices.

It has also been noticed that these background error statistics are less dependent on the diurnal cycle and on the meteorological situation (e.g. in the presence of fog at 06 UTC <sup>10</sup> and low clouds at 21 UTC), contrary to the  $\mathbf{B}_{\mathbf{x}}(T, q_v)$  matrix where there is a reduction in the area of positive correlation in the lowest layers between 06 UTC and 21 UTC (Fig. 4).

The 1D-Var results are now assessed in observation space by examining innovations (differences between observed <sup>15</sup> and simulated brightness temperatures) from AROME background profiles and residuals. In the following, we have only used background error covariance matrices estimated at 06 UTC with fog mask, for a simplified comparison framework of the two 1D-Var systems.

#### **20 4.2 1D-Var analysis fit to observations**

Figure 5 presents both (a) innovations and (b,c) residuals obtained with the two 1D-Var systems (b: REF and c: EXP) for the 13 channels (1,2,4-14) and for each hour of the day. The innovations are generally positive for water vapour <sup>25</sup> sensitive channels during the day, and negative for temperature channels, especially in the morning. The differences are mostly between -2.5 and 5 K. For channels 6,7 and 8, 9 and 10, which are sensitive to liquid water content, the innova-

tions can reach higher values exceeding 10 K (in the after-<sup>30</sup> noon) or being around -5 K (in the morning).

In terms of residuals, as expected from 1D-Var systems, both experiments significantly reduce the deviations of the observed TB from those calculated using the background profiles, especially for the first eight chan- $_{35}$  nels sensitive to water vapour and liquid water. We can

- note that the residuals are not as reduced for channel 9 (52.28 GHz) compared to other channels. Indeed, channels 8 and 9 (51.26 and 52.28 GHz) suffer from larger calibration uncertainties (Maschwitz et al., 2013) and
- <sup>40</sup> larger forward model uncertainties dominated by oxygen line mixing parameters (Cimini et al., 2018) than other temperature sensitive channels. However, by comparing simulated brightness temperatures (TB) with different absorption models (Hewison, 2007), or through a monitoring
- <sup>45</sup> with simulated TB from clear-sky background profiles (Angelis et al., 2017; Martinet et al., 2020), larger biases are generally observed only at 52.28 GHz. Consequently, the higher deviations observed in Figure 5 for channel 9 mostly originate from larger modelling and calibration
- <sup>50</sup> uncertainties, which are taken into account in the assumed instrumental errors (prescribed observation errors of about 3 K for these two channels compared to <1 K for other

# temperature sensitive channels) and also possibly from larger instrumental biases.

The temperature channels used in the zenith mode are <sup>55</sup> modified less or very little, the deviations from the background values being much smaller than for the other channels. During the second half of the day, characterized by the presence of clouds around 800 hPa (see Figs. 2 (e) and (f)), the residual values are largely reduced in the frequency bands sensitive to liquid water for channels 5, 6and 7, 7 and 8, especially for EXP as shown by the comparison of the pixels in the dashed rectangular boxes in Figs. 5 (b) and (c). Residuals are also slightly reduced for EXP in the morning and during the fog and low temperature period for the first five channels (1,2,4-6) between 2 and 8 UTC.

In order to quantify these results on 9 February 2020 dataset (all hours and all channels), the bias and root mean square (RMS) error values are computed for the background and the analyses produced by REF and EXP. The innova-70 tions are characterized by a RMS error of 3.20 K and a bias of 1.32 K. Both assimilation experiments reduce these two quantities by modifying model profiles. The RMS errors are 0.71 K for EXP and 0.72 K for REF and the biases are -0.17 K for EXP and -0.11 K for REF. These 75 statistics have also been calculated by restricting the dataset to the two dashed rectangular boxes represented in Figs. 5 (b) and (c). A significant improvement is observed for the most sensitive channels to liquid water in the afternoon with a RMSE decreased from 4.3 K in the background to 0.57 K  $_{80}$ in REF and 0.37 K in EXP. For all computed statistics, EXP always provides the best performance in terms of RMSE. Table 2 summarizes the bias and RMSE values obtained for the different samples.

#### 4.3 Vertical profiles of analysis increments

After examining the fit of the two experiments to the observed TBs, we assess the corrections made in model space. Figure 6 shows the increments of (a), (b) temperature, (c), (d) specific humidity, and (e), (f) liquid water for the two experiments REF (left panels) and EXP (right panels). In addition, <sup>90</sup> the increments of  $(\theta_s)_a$  are shown in (g)\_(h).

85

The temperature increments are mostly located in the lower troposphere (below 650 hPa) with a dominance of negative values of small amplitude (around 0.5 K). This is consistent with negative innovations observed on temperature <sup>95</sup> channels highlighting a warm bias in the background profiles. The areas of maximum cooling take place in cloud layers (inside the thick fog layer below 900 hPa until 9 UTC and around 700 hPa after 12 UTC). The increments are rather similar between REF and EXP, but the positive increments <sup>100</sup> appear to be larger with EXP (e.g. at 08 and 20 UTC around 800 hPa).

Concerning the profiles associated with moist variables, the structures show similarity similarities between the two experiments but with differences in intensity. During the 105



**Figure 3.** Background error cross-correlation matrices at 06 UTC 9 February 2020 with (right) and without (left) fog mask. Top: between the classical variables  $(T, q_v)$  denoted in the axes by "T" and "Qv", respectively. Bottom: between the new conservative variables  $((\theta_s)_a, q_t)$  denoted in the axes by "Thetas" and "Qtot", respectively. The axes correspond to the levels of the AROME vertical grid (1 at the top and 90 at the bottom). Correlations are between -1 and 1 units.

**Table 2.** Bias / RMSE (K) of the background and analyses produced by EXP and REF against MWR TB observations. Statistics are computed either using all data or restricting to channels number 1 to 5 between 2 and 8 UTC or channels 7 to 9 between 10 and 24 UTC (these two sub-samplings represent the dashed rectangular boxes in Figs. 5 (b) and (c)).

	Background	REF	EXP
All data	1.3/2.2	-0.11/0.72	-0.17/0.71
Channels 1 to 6, 2 to 8 UTC	1.5/2.2	0.11/0.3	0.08/0.3
Channels 6 to 8, 10 to 24 UTC	2.7/4.3	0.16/0.57	-0.12/0.37

night and in the morning, the  $q_v$  increments near the surface are negative. These negative increments are projected into increments having the same sign as T by the strong positive cross-correlations of the  $\mathbf{B}_{fog}$  matrix up to 900 hPa (Fig. 3). Thus, the largest negative temperature and specific humidity  $_{5}$  increments remain confined in the lowest layers.



Figure 4. Same as Fig. 3, but at 21 UTC.

Liquid water is added in both experiments between 03 UTC and 07 UTC, close to the surface, where the Jacobians of the most sensitive channels to  $q_l$  are the largest (not shown(6 to 8) have significant values in the fog layer <sup>5</sup> present in the background (see Figure 2e). After 14 UTC, values of  $q_v$  between 850 and 700 hPa and  $q_l$  around 800 hPa are enhanced in both cases, with larger increments for the REF case, in particular at 20 UTC and around midnight. Most of the liquid water is created in low clouds, while for <sup>10</sup> EXP, Additionally, increments of  $q_l$  aloft are more extended above 600 hPa are larger and more extended vertically and in time in time and in the vertical. For example, at 21 UTC in EXP, where condensation occurs over a thicker atmospheric layer between 500 hPa and 300 hPa after 12 UTC. In the

15 REF experiment, the creation of liquid water reaching above

500 hPa only reaches values of 0.3  $\frac{q}{kq}$  is observed around 400 hPa . For EXP after 12 UTC, condensation occurs over a thicker atmospheric layer between 500 and 300 hPa. g/kg sporadically, for example at 21 UTC. In this experimental set-up, condensed water can be created or removed over the 20 whole column by means of the supersaturation diagnosed at each iteration of the minimisation process (since RTTOV-gb needs  $(T, q_v, q_l)$  profiles for the TB computation.) This is a clear advantage over REF which keeps unchanged of EXP over REF, which keeps the vertical structure of the  $q_l$  pro-25 file unchanged from the background. Liquid water is In REF, liquid water is only added where it already existed exists in the background because once the LWP variable is updated, the analyzed  $q_l$  profile is just modified proportionally to the ratio between the LWP of the analysis and of the back-  $_{30}$ 



**Figure 5.** Differences in observed (channels 0-1, 2 and 4 to 5-7 being located between 22 and 31 GHz and channels 6-8 to 12-14 being located between 51 and 58 GHz, HATPRO radiometer) and simulated (with RTTOV-gb) brightness temperatures (in Kelvin): (a) from AROME background profiles; (b) from 1D-Var analyses from the REF configuration; and (c) from 1D-Var analyses from the EXP configuration for all hours of the day on 9 February 2020 at Saint-Symphorien (Les Landes region). The dashed blue boxes indicate the channels and hours where EXP is improved with respect to REF. Color-bars are in unit of K.

ground, as explained in more details by Deblonde and English (2003).

The profiles of increments for  $(\theta_s)_a$  show similar structures to show structures similar to the increments of  $q_v$  around 800 hPa and to the increments of *T* below, where temperature Jacobians are the largest (not shown). see Figure 7 from De Angelis et al. (2016)). The conversion of *T*,  $q_v$  and  $q_t$  changes obtained with REF into  $(\theta_s)_a$  increments (Fig. 6g) highlights the main differences between the two systems. They take place around 800 hPa with larger increments produced by the new 1D-Var particularly between 11 and 14 UTC.

Some radiosoundings (RS) have been launched during the SOFOG3D IOPs. As only one RS profile was launched at 05:21 UTC on the case study presented in the article, 15 no statistical evaluation of the profile increments can be carried out. However, we have conducted an evaluation of the analysis increments obtained at 5 and 6 UTC (the 1D-Var retrievals being performed at a 1h temporal resolution in line with the operational AROME assimilation cycles) around the 20 RS launched time. As the AROME temperature background profile extracted at 6 UTC was found to have a vertical structure closer to the RS launched at 05:21 UTC, figures 7 compare the AROME background profile and 1D-Var analyses performed with the REF and EXP experiments valid 25 at 6 UTC against the RS profile.

The temperature increments are a step in the right direction by cooling the AROME background profile in line with the observed RS profile. The two 1D-Var analyses are close to each other, but the EXP analysis produces a temperature 30 profile slightly cooler compared to the REF analysis. In terms of absolute humidity  $(\rho_v = p_v/(R_v T))$ , with  $p_v$  the partial pressure and  $R_v$  the gas constant for water vapor), the background profile already exhibits a similar structure compared to the RS profile. The 1D-Var increments are thus 35 small and close between the two experiments. However, we can note that the EXP profile is slightly moister than the REF profile from the surface up to 3500 m which leads to a somewhat better agreement with the RS profile below 1500 m. In terms of integrated water vapor (IWV), a 40 significant improvement of the background IWV with respect to the RS IWV is observed with a difference reduced from almost 1 kg m<sup>-2</sup> in the background to less than 0.4 kg m<sup>-2</sup> in the analyses. These analyses confirm the improvement brought to the model profiles by both the REF and EXP 45 analysis increments, with some enhanced improvement for EXP.

#### 5 Conclusions

The aim of this study was to examine the interest-value of using moist-air entropy potential temperature  $(\theta_s)_a$  and total water content  $q_t$  to study fog initiation and dissipation at small scale. A as new control variables for variational assimilation schemes. In fact, the use of control variables less dependent to vertical gradients of  $(T, q_v, q_l, q_i)$  variables should ease the specification of background error covariance



**Figure 6.** Profiles of analysis increments resulting from two 1D-Var experiments: REF (left) and EXP (right), for: (a)-(b) T in  $\frac{K}{K}$  K; (c)-(d)  $q_v$  in  $\frac{kg/kg}{g}$  g kg<sup>-1</sup>; (e)-(f)  $q_l$  in  $\frac{g/kg}{g}$  g kg<sup>-1</sup> and (g)-(h)  $(\theta_s)_a$  in  $\frac{K}{K}$  K. Color-bars have the same units (K or g kg<sup>-1</sup>) as the variables.



**Figure 7.** Vertical profiles of absolute temperature T (top panel, in K) and absolute humidity  $\rho_v$  (bottom panel, in g m<sup>-3</sup>) for the 9 February 2020 and showing: the RS launched at 05 : 21 UTC (solid black); the AROME background valid at 06 UTC (solid blue); and the 1D-Var retrievals at 06 UTC obtained with REF (dashed blue) and EXP (dotted-dashed blue). Integrated water vapor (IWV) retrievals are also compared to the RS IWV (in the blue box in the bottom panel).

# matrices, which play a key role in the quality of the analysis state in operational assimilation schemes.

To that end, a 1D-Var system has been used for assimilating to assimilate brightness temperature (TB) ob-

servations from the ground-based HATPRO microwave radiometer installed at Saint-Symphorien (Les Landes region over-in South-Western France) during the SOFOG3D measurement campaign (winter 2019-2020).

The motivations for using the entropy of moist air in atmospheric science by taking into account the 10 absolute value for dry air and water vapour were described by Richardson (1922) and formalized by Hauf and Höller (1987), with the moist-air potential temperature variables  $\theta_s$  and  $(\theta_s)_1$  defined by Marquet (2011). The conservative aspects of these variables 15 and their meteorological properties (fronts, convection, cyclones) have been studied in Marquet (2011), Blot (2013)and Marquet and Geleyn (2015). The links with the definition of the Brunt-Väisälä frequency and the potential vorticity are described in Marquet and Geleyn (2013) and 20 Marquet (2014), while the interest of the absolute entropy to describe the thermodynamics of cyclones is shown in Marquet (2017) and Marquet and Dauhut (2018). The use of the  $\theta_s$  variable in data assimilation can thus be considered as an additional and natural step that continues these series of 25 studies.-

The 1D-Var system has been adapted to consider these new quantities as control variables. Since the radiative transfer model needs profiles of temperature, water vapour and cloud liquid water for the simulation of TB, an adjustment process 30 has been defined to obtain these quantities from  $(\theta_s)_a$  and  $q_t$ . The adjoint version of this conversion has been developed for an efficient estimation of the gradient of the cost-function. Dedicated background error covariance matrices have been estimated from the Ensemble Data Assimilation system of AROME. We first demonstrated that the matrices for the new variables are less dependent on the meteorological situation (all-weather conditions vs. fog conditions) and on the time of the day (stable conditions by night vs. unstable conditions during the day) leading to potentially more 40 robust estimates. This is an important result as the optimal estimation of the analysis depends on the good accurate specification of the background error covariance matrix which is known to highly vary with weather conditions when using classical control variables. 45

The comparison of the The new 1D-Var system with a elassical system using temperature, water vapour and liquid water path shows has produced rather similar results in terms of fit of the analysis to observed TB. On the other hand, atmospheric increments on values when compared to the classical one using temperature, water vapour and liquid water ean be locally very different path. Nevertheless, quantitative results reveal smaller biases and *RMS* values with the new system in low cloud and fog areas. It has also been noticed that atmospheric increments are somewhat different in cloudy conditions between the two systems.

The encouraging results obtained from this feasibility study need to be consolidated by complementary studies. The For example, in the stratocumulus layer that formed during the afternoon, the new 1D-Var induces larger temperature increments and reduced liquid water corrections. Moreover, its capacity to generate cloud condensates in clear-sky regions of the background has been demonstrated. As a

- <sup>5</sup> preliminary validation, the retrieved profiles from the 1D-Var should be compared against independent observations (e.g. radiosoundings available from have been compared favourably against an independent observation data set (one radiosounding launched during the SOFOG3D field cam-
- <sup>10</sup> paign). The new 1D-Var leads to profiles of temperature and absolute humidity slightly closer to observations in the planetary boundary layer.

These encouraging results obtained from this feasibility study need to be consolidated by complementary studies.

- <sup>15</sup> Observed brightness temperatures at lower elevation angles should be included in the 1D-Var for a better constraint on temperature profiles within the atmospheric boundary layer. Indeed, larger differences in the temperature increment increments might be obtained between the classical 1D-Var
- 20 system and the 1D-Var system using the new conservative variables when additional elevation angles are included in the observation vector. Other case studies from the field campaign could also be examined to confirm our first conclusions.
- Finally, the conversion operator could be improved by accounting not only for liquid water content  $q_l$  but also for ice water content  $q_i$  (e.g. using a temperature threshold criteria). Indeed, Indeed, inclusion of  $q_i$  in the conversion operator should lead to more realistic retrieved profiles of
- <sup>30</sup> cloud condensates, and a 1D-var system with only  $q_l$  can create water clouds at locations where ice clouds should be present, as done in our experiment around 400 hPa between 15 and 24 UTC. However, since the frequencies of HATPRO are not sensitive to ice water content, the fit of simulated
- <sup>35</sup> *TBs* to observations could be reduced. In consequence, the synergy with an instrument sensitive to ice water clouds, such as a W-band cloud radar, would be necessary for improved retrievals of both  $q_i$  and  $q_l$ . It is worth noticing that the variable  $(\theta_s)_a$  can easily be generalized to the case of the ice
- <sup>40</sup> phase and mixed phases by taking advantage of the general definition of  $\theta_s$  and  $(\theta_s)_1$ , where  $L_{vap} q_l$  is simply replaced by  $L_{vap} q_l + L_{sub} q_i$ .

*Code and data availability.* The numerical code of the radiative transfer model RTTOV-gb together with the associated resources

- <sup>45</sup> (coefficient files) can be downloaded from http://cetemps.aquila. infn.it/rttovgb/rttovgb.html and from https://nwp-saf.eumetsat.int/ site/software/rttov-gb/. The 1D-Var software has been adapted from the NWP SAF 1D-Var provided here: https://nwp-saf. eumetsat.int/site/software/1d-var/ and is available on request to
- <sup>50</sup> pauline.martinet@meteo.fr. The instrumental data are available on the AERIS website dedicated to the SOFOG3D field experiment: https://sofog3d.aeris-data.fr/catalogue/. AROME backgrounds are available on request to pauline.martinet@meteo.fr. Quicklooks from the cloud radar BASTA are available on: http://basta.projet.

latmos.ipsl.fr. The BUMP library to compute background error matrices, developed in the framework of the JEDI project led by the JCSDA (Joint Center for Satellite Data Assimilation, Boulder, Colorado), can be downloaded at https://github.com/JCSDA/saber.

*Author contributions.* PMarq supervised the work of AB, contributed to the implementation of the new conservative variables in <sup>60</sup> the computation of new background error covariance matrices, participated to the scientific analysis and manuscript redaction. JFM developed the conversion operator and adjoint version, participated in the scientific analysis and manuscript redaction. PMart supervised the modification of the 1D-Var algorithm, supported the use <sup>65</sup> of the EDA to compute background error covariance matrices, provided the instrumental data used in the 1D-Var and participated in the manuscript revision. AB adapted the 1D-Var algorithm and processed all the data, prepared the figures and participated in the manuscript revision. BM developed and adapted the BUMP library <sup>70</sup> to compute the background error covariance matrices for the 1D-Var, and participated in the manuscript revision.

*Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. The authors are very grateful to the two 75 anonymous reviewers who suggested substantial improvements to the article. The instrumental data used in this study are part of the SOFOG3D experiment. The SOFOG3D field campaign was supported by METEO-FRANCE and ANR through grant AAPG 2018-CE01-0004. Data are managed by the French national cen-80 ter for Atmospheric data and services AERIS. The MWR network deployment was carried out thanks to support by IfU GmbH, the Köln University, the Met-Office, Laboratoire d'Aérologie, Meteoswiss, ONERA, and Radiometer Physics GmbH. MWR data have been made available, quality controlled and processed in the 85 frame of CPEX-LAB (Cloud and Precipitation Exploration LABoratory, www.cpex-lab.de), a competence center within the Geoverbund ABC/J by acting support of Ulrich Löhnert, Rainer Haseneder-Lind and Arthur Kremer from the University of Cologne. This collaboration is driven by the European COST actions ES1303 90 TOPROF and CA18235 PROBE. Julien Delanoë and Susana Jorquera are thanked for providing the cloud radar quicklooks used in this study for better understanding the meteorological situation. Thibaut Montmerle and Yann Michel are thanked for their support on the use of the AROME EDA to compute background error co- 95 variance matrices. The work of Benjamin Ménétrier is funded by the JCSDA (Joint Center for Satellite Data Assimilation, Boulder, Colorado) UCAR SUBAWD2285.

#### References

Angelis, F. D., Cimini, D., Löhnert, U., Caumont, O., Haefele, A., 100 Pospichal, B., Martinet, P., Navas-Guzmán, F., Klein-Baltink, H., Dupont, J.-C., et al.: Long-term observations minus background monitoring of ground-based brightness temperatures from a microwave radiometer network, Atmospheric Measurement Techniques, 10, 3947–3961, 2017.

- Bauer, L. A.: The relation between "potential temperature" and "entropy", Phys. Rev. (Series I), 26, 177–183, https://doi.org/10.1103/PhysRevSeriesI.26.177, 1908.
- Benjamin, S. G., Brewster, K. A., Brümmer, R., Jewett, B. F., Schlatter, T. W., Smith, T. L., and Stamus, P. A.: An Isentropic Three-Hourly Data Assimilation System Using ACARS Aircraft Observations, Mon.
  Wea. Rev., 119, 888–906, https://doi.org/10.1175/1520-
- 10 Wea. Rev., 119, 888–906, https://doi.org/10.1175/1520-0493(1991)119<0888:AITHDA>2.0.CO;2, 1991.
- Benjamin, S. G., Grell, G. A., Brown, J. M., Smirnova, T. G., and Bleck, R.: Mesoscale Weather Prediction with the RUC Hybrid Isentropic-Terrain-Following Coordinate Model,
- <sup>15</sup> Mon. Wea. Rev., 132, 473–494, https://doi.org/10.1175/1520-0493(2004)132<0473:MWPWTR>2.0.CO;2, 2004.
- Betts, A. K.: Non-precipitating cumulus convection and its parameterization, Quart. J. Roy. Meteor. Soc., 99, 178–196, https://doi.org/10.1002/qj.49709941915, 1973.
- 20 Blot, E.: Etude de l'entropie humide dans un contexte d'analyse et de prévision du temps, Rapport de stage d'approfondissement EIENM3, Ecole Nationale de la Météorologie. Toulouse, 2013.
- Brousseau, P., Seity, Y., Ricard, D., and Léger, J.: Improvement of the forecast of convective activity from the AROME-
- France system, Quart. J. Roy. Meteorol. Soc., 142, 2231–2243, https://doi.org/https://doi.org/10.1002/qj.2822, 2016.
- Cimini, D., Rosenkranz, P. W., Tretyakov, M. Y., Koshelev, M. A., and Romano, F.: Uncertainty of atmospheric microwave absorption model: impact on ground-based radiometer simulations and retrievals, Atmospheric Chemistry and Physics, 18, 15231–
- and retrievals, Atmospheric Chemistry and Physics, 18, 15231– 15259, 2018.
- Cimini, D., Hocking, J., De Angelis, F., Cersosimo, A., Di Paola, F., Gallucci, D., Gentile, S., Geraldi, E., Larosa, S., Nilo, S., Romano, F., Ricciardelli, E., Ripepi, E., Viggiano, M., Luini,
- L., Riva, C., Marzano, F. S., Martinet, P., Song, Y. Y., Ahn, M. H., and Rosenkranz, P. W.: RTTOV-gb v1.0 – updates on sensors, absorption models, uncertainty, and availability, Geosci. Model Dev., 12, 1833–1845, https://doi.org/10.5194/gmd-12-1833-2019, 2019.
- <sup>40</sup> Clough, S. A. and Testud, J.: The Fronts-87 experiment and mesoscale frontal dynamics project, WMO Bulletin, 37, 276– 281, 1988.
- Cullen, M. J. P.: Four-dimensional variational data assimilation: A new formulation of the background-error co-
- 45 variance matrix based on a potential-vorticity representation, Quart. J. Roy. Meteor. Soc., 129, 2777–2796, https://doi.org/https://doi.org/10.1256/qj.02.10, 2003.
- De Angelis, F., Cimini, D., Hocking, J., Martinet, P., and Kneifel, S.: RTTOV-gb – adapting the fast radiative transfer model
- 50 RTTOV for the assimilation of ground-based microwave radiometer observations, Geosci. Model Dev., 9, 2721–2739, https://doi.org/10.5194/gmd-9-2721-2016, 2016.
- Deblonde, G. and English, S.: One-Dimensional Variational Retrievals from SSMIS-Simulated Observations, J. Appl. Me-
- teor. Climatol., 42, 1406–1420, https://doi.org/10.1175/1520-0450(2003)042<1406:OVRFSO>2.0.CO;2, 2003.
- Delanoë, J., Protat, A., Vinson, J.-P., Brett, W., Caudoux, C., Bertrand, F., du Chatelet, J. P., Hallali, R., Barthes, L., Haeffelin, M., and Dupont, J.-C.: BASTA: A 95-GHz

FMCW Doppler Radar for Cloud and Fog Studies, Jour- 60 nal of Atmospheric and Oceanic Technology, 33, 1023–1038, https://doi.org/10.1175/JTECH-D-15-0104.1, 2016.

- Desroziers, G.: A Coordinate Change for Data Assimilation in Spherical Geometry of Frontal Structures, Mon. Wea. Rev., 125, 3030–3038, https://doi.org/10.1175/1520-0493(1997)125<3030:ACCFDA>2.0.CO;2, 1997.
- Desroziers, G. and Lafore, J.-P.: A Coordinate Transformation for Objective Frontal Analysis, Mon. Wea. Rev., 121, 1531–1553, https://doi.org/10.1175/1520-0493(1993)121<1531:ACTFOF>2.0.CO;2, 1993.
- Destouches, M., Montmerle, T., Michel, Y., and Ménétrier, B.: Estimating optimal localization for sampled background-error covariances of hydrometeor variables, Quart. J. Roy. Meteorol. Soc., 147, 74–93, https://doi.org/https://doi.org/10.1002/qj.3906, 2021.
- Gustafsson, N., Janjić, T., Schraff, C., Leuenberger, D., Weissmann, M., Reich, H., Brousseau, P., Montmerle, T., Wattrelot, E., Bučánek, A., Mile, M., Hamdi, R., Lindskog, M., Barkmeijer, J., Dahlbom, M., Macpherson, B., Ballard, S., Inverarity, G., Carley, J., Alexander, C., Dowell, D., Liu, S., <sup>80</sup> Ikuta, Y., and Fujita, T.: Survey of data assimilation methods for convective-scale numerical weather prediction at operational centres, Quart. J. Roy. Meteorol. Soc., 144, 1218–1256, https://doi.org/https://doi.org/10.1002/qj.3179, 2018.
- Hauf, T. and Höller, H.: Entropy and potential temperature, <sup>85</sup> J. Atmos. Sci., 44, 2887–2901, https://doi.org/10.1175/1520-0469(1987)044<2887:EAPT>2.0.CO;2, 1987.
- Hewison, T. J.: 1D-VAR Retrieval of Temperature and Humidity Profiles From a Ground-Based Microwave Radiometer, IEEE Transactions on Geoscience and Remote Sensing, 45, 2163– 90 2168, https://doi.org/10.1109/TGRS.2007.898091, 2007.
- Marquet, P.: Definition of a moist entropy potential temperature: application to FIRE-I data flights, Quart. J. Roy. Meteorol. Soc., 137, 768–791, https://doi.org/10.1002/qj.787, 2011.
- Marquet, P.: On the definition of a moist-air potential vor- 95 ticity., Quart. J. Roy. Meteorol. Soc., 140, 917–929, https://doi.org/10.1002/qj.2182, 2014.
- Marquet, P.: A Third-Law Isentropic Analysis of a Simulated Hurricane, J. Atmos. Sci., 74, 3451–3471, https://doi.org/10.1175/JAS-D-17-0126.1, 2017.
- Marquet, P. and Bechtold, P.: A new Estimated Inversion Strength (EIS) based on the moist-air entropy, Research activities in Earth system modelling. Working Group on Numerical Experimentation. Report No. 50. WCRP (Blue Book) Report No.12/2020. Ed. E. Astakhova, WMO, Geneva, 105 50, 1–2, http://bluebook.meteoinfo.ru/uploads/2020/docs/04\_ Marquet\_Pascal\_NewEntropyEIS.pdf, 2020.
- Marquet, P. and Dauhut, T.: Reply to "Comments on 'A third-law isentropic analysis of a simulated hurricane", J. Atmos. Sci., 75, 3735–3747, https://doi.org/10.1175/JAS-D-18-0126.1, 2018. 110
- Marquet, P. and Geleyn, J.-F.: On a general definition of the squared Brunt-Väisälä frequency associated with the specific moist entropy potential temperature, Quart. J. Roy. Meteorol. Soc., 139, 85–100, https://doi.org/10.1002/qj.1957, 2013.
- Marquet, P. and Geleyn, J.-F.: Formulations of moist thermodynamics for atmospheric modelling, in: Parameterization of Atmospheric Convection. Vol II: Current Issues and New Theories, edited by Plant, R. S. and Yano, J.-

70

75

100

I., *World Scientific*, pp. 221–274, Imperial College Press, https://doi.org/10.1142/9781783266913\_0026, 2015.

- Martinet, P., Cimini, D., Burnet, F., Ménétrier, B., Michel, Y., and Unger, V.: Improvement of numerical weather predic-
- tion model analysis during fog conditions through the assimilation of ground-based microwave radiometer observations: a 1D-Var study, Atmos. Meas. Tech., 13, 6593–6611, https://doi.org/10.5194/amt-13-6593-2020, 2020.
- Maschwitz, G., Löhnert, U., Crewell, S., Rose, T., and Turner,
- D. D.: Investigation of ground-based microwave radiometer calibration techniques at 530 hPa, Atmos. Meas. Tech., 6, 2641– 2658, https://doi.org/10.5194/amt-6-2641-2013, 2013.
- Ménétrier, B. and Montmerle, T.: Heterogeneous backgrounderror covariances for the analysis and forecast of fog
- 5 events, Quart. J. Roy. Meteor. Soc., 137, 2004–2013, https://doi.org/10.1002/qj.802, 2011.
- Michel, Y., Auligné, T., and Montmerle, T.: Heterogeneous Convective-Scale Background Error Covariances with the Inclusion of Hydrometeor Variables, Mon. Wea. Rev., 139, 2994– 3015, https://doi.org/10.1175/2011MWR3632.1, 2011.
- Montmerle, T. and Berre, L.: Diagnosis and formulation of heterogeneous background-error covariances at the mesoscale, Quart. J. Roy. Meteor. Soc., 136, 1408–1420, https://doi.org/10.1002/qj.655, 2010.
- 25 Richardson, L. F.: Atmospheric stirring measured by precipitation, Proc. Roy. Soc. London (A), 96, 9–18, 1919.
- Richardson, L. F.: Weather prediction by numerical process, pp. 1– 229, Cambridge University Press, 1922.
- Rodgers, C. D.: Retrieval of atmospheric temperature and composi-
- tion from remote measurements of thermal radiation, Rev. Geophys., 14, 609–624, https://doi.org/10.1029/RG014i004p00609, 1976.
- Rose, T., Crewell, S., Löhnert, U., and Simmer, C.: A network suitable microwave radiometer for operational monitor-
- <sup>35</sup> ing of the cloudy atmosphere, Atmos. Res., 75, 183–200, https://doi.org/10.1016/j.atmosres.2004.12.005, 2005. Seity, Y., Brousseau, P., Malardel, S., Hello, G., Bénard, P.,
- Bouttier, F., Lac, C., and Masson, V.: The AROME-France convective-scale operational model, Mon. Wea. Rev., 139, 976–991, https://doi.org/10.1175/2010MWR3425.1, 2011.
- Shapiro, M. A. and Hastings, J. T.: Objective cross-section analyses by Hermite polynomial interpolation on isentropic surfaces, J. Appl. Meteorol. Climatol., 12, 753–762, https://doi.org/10.1175/1520-450(1973)012<0753:OCSABH>2.0.CO;2, 1973.
- Thépaut, J.-N. and Moll, P.: Variational inversion of simulated TOVS radiances using the adjoint technique, Quart. J. Roy. Meteor., 116, 1425–1448, 1990.
- Wlasak, M., Nichols, N. K., and Roulstone, I.: Use
- 50 of potential vorticity for incremental data assimilation, Quart. J. Roy. Meteor. Soc., 132, 2867–2886, https://doi.org/https://doi.org/10.1256/qj.06.02, 2006.