

Review: Feofilov et al.

The paper deals with comparing the CALIOP and AEOLUS climatological backscatter data (half year in 2019). While the idea is good and has a high potential, the methodology and the presentation style are poor.

The wavelength difference between the two different lidars in space is only poorly accounted for, using an empirical 50-years old formula. Here, the authors should have used temperature and pressure profiles from NWP (as e.g. provided with Aeolus data) to calculate the molecular backscatter in the UV and visible range (i.e. at 532 nm) to make a real conversion of the scattering ratio and compare apples to apples.

Besides that, I also have the feeling that misinterpretation of Aeolus data is done while not taking into account the high contribution of molecular backscatter to the scattering ratio (see for example specific statement below under 7.).

Thus, the manuscript suffers from a significant methodological weakness and any conclusion drawn from the current applied methodology is very questionable.

Furthermore, the presentation style needs to be improved. The language is hard to read and sometime really not understandable. E.g., already the abstract is hard to understand.

Also, the title does not at all reflect the content of the paper. Furthermore, I could not follow some of the argumentations. Often, statements are made without justification.

Therefore, I recommend the rejection of the manuscript, while at the same time encouraging the authors to re-submit a paper once the methodology and presentation style has been significantly improved.

Some comment general comments below, more detailed comments are given in the commented manuscript.

Major comments:

1. Title:

The title does not reflect the content of the paper. In fact, the authors focus only on the cloud detection capability based on scattering ratios. Furthermore, the whole instruction deals only with clouds and not a single word about scattering ratios is written.


2. Definition of scattering ratio

The scattering ratio which is the essential part of this manuscript has never been properly defined. According to the reference which is given, I assume that, "the ratio between the total backscatter by particles and molecules and the molecular backscatter" (according to Flamantm 2008) is meant, i.e. the ratio between the total backscatter (represented by particles and molecules) **to** the molecular backscatter.

3. Wavelength conversion


The conversion the authors use to account for the different wavelengths of CALIOP and AEOLUS is poor.

For example, I have made a sketch using an arbitrary atmospheric molecular backscatter coefficient profile and a height-constant particle backscatter coefficient (equal at both wavelengths) of $7 \times 10^{-6} \text{ m}^{-1} \text{ sr}^{-1}$ in order to obtain a scattering ratio at 532 nm shortly above 5 as given by the authors as detection threshold for clouds. The input is illustrated below:




As one can see, the molecular backscatter coefficients vary significantly with altitude and between the wavelengths.

As a result, also the scattering ratio varies with altitude even though the particle backscatter coefficient is constant, as shown below:



This figures also shows that the simple empirical conversion applied by the authors is not valid and given the fact that pressure and temperature profiles are provided with Aeolus data, a much better job could have been done to account for the difference wavelengths of Aeolus and CALIOP.

Furthermore, assuming that the given threshold (SNR at 532 nm > 5) is valid for detecting a cloud, it also indicates that a cloud with the same optical properties (same particle backscatter) would be detected differently at different heights. See Figure below (same as above but differ y-axis)



For the case I simulated, CALIOP would “detect” a cloud (orange line) for all heights while with the Aeolus converted SR profiles would not lead to a cloud detection (SNR below 5) for some parts of the height profile.

For the real, physical, conversion this is even true up to 12 km height (blue line).

Thus, using the scattering ratio for cloud detection is not appropriate in my opinion. You could have used the scattering ratio from the satellite data to calculate the real particle backscatter.

I.e. by using the molecular backscatter which you calculate from the meteorological data which is included in the satellite data (AUX-MET in Aeolus data).

Despite all my own doubts concerning this conversion, the authors themselves state: “We would like to stress here that no linear scaling applied uniformly to SRs at all heights could change the ratio of high cloud detection frequency to low cloud detection frequency of ALADIN.”

Therefore, I wonder: Why they are doing so?

4. Scattering ratio threshold:

The choice of this threshold $SR > 5$ is not clear to me and seems very arbitrary and without justification.

What happens if this threshold changes?

5. Different vertical resolution

The different vertical resolution for Aeolus and Calipso is not sufficiently discussed.

6. Language

Language and phrasing need to be improved. It is hardly understandable and not well explained. Please use simple sentences.

Furthermore, “insider information of Aeolus” need to be explained otherwise it is not understandable for non-Aeolus experts.

7. Specific comments in addition to the pdf (examples)

- Some statements are either simply wrong or wrongly phrased, e.g.:
“...is characterized by lower sensitivity **to high clouds** above ~7 km than CALIOP, that we explain by lower SNR for ALADIN at these heights that is due both to physical reasons (smaller backscatter at 355 nm)”
Why should there be a smaller backscatter at 355 nm? This is in absolute contradiction to all my knowledge! The particle backscatter coefficient could be equal in clouds (Angström of 0), but the molecular backscatter coefficient is for sure higher (see plots) and thus the total backscatter is for sure also higher! Could you please comment?
- Abstract: Just one of many examples:
“(b) the cloud detection agreement is better for the lower layers. Above ~7 km, the ALADIN product demonstrates lower sensitivity because of lower backscatter at 355 nm”
I do not understand this statement. First of all: What do you mean? The volume backscatter coefficient, the particle backscatter coefficient, the molecular backscatter coefficient? It is not clear! And I also do not know why any of these should be lower at 355 nm compared to 532 nm (and 1064 nm).
- Abstract last sentence: Is not understandable. What values are this? What is a cloud detection agreement value? Abstracts should be self-explaining and understandable.
- Not all references are in alphabetical order
- Some mistakes in the names of the references, please check



Comparing scattering ratio products retrieved from ALADIN/Aeolus and CALIOP/CALIPSO observations: sensitivity, comparability, and temporal evolution

Artem G Feofilov¹, H el ene Chepfer¹, Vincent Noel², Rodrigo Guzman¹, Cyprien Gindre¹ and Marjolaine Chiriaco³

¹LMD/IPSL, Sorbonne Universit e, UPMC Univ Paris 06, CNRS,  cole polytechnique, Palaiseau, 91128, France

²Laboratoire d'A erologie, CNRS/UPS, Observatoire Midi-Pyr en es, 14 avenue Edouard Belin, Toulouse, France

³LATMOS/IPSL, Univ Versailles Saint-Quentin en Yvelines, Guyancourt, France

Correspondence to: Artem G Feofilov (artem.feofilov@lmd.polytechnique.fr)

10 Abstract.

The spaceborne active sounders have been contributing invaluable vertically resolved information of atmospheric optical properties since the launch of CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation) in 2006. To ensure the continuity of climate studies and monitoring the global changes, one has to understand the differences between lidars operating at different wavelengths, flying at different orbits, and utilizing different observation geometries, receiving paths, and detectors. In this article, we show the results of an intercomparison study of ALADIN (Atmospheric Laser Doppler Instrument) and CALIOP (Cloud Aerosol Lidar with Orthogonal Polarization) lidars using their scattering ratio (SR) products for the period of 28/06/2015–31/12/2019. We suggest an optimal set of collocation criteria ($\Delta\text{dist} < 1^\circ$; $\Delta\text{time} < 6\text{h}$), which would give a representative set of collocated profiles and we show that for such a pair of instruments the theoretically achievable cloud detection agreement for the data collocated with aforementioned criteria is 0.77 ± 0.17 . The analysis of a collocated database consisting of ~78000 pairs of collocated nighttime SR profiles revealed the following: (a) in the cloud-free area, the agreement is good indicating low frequency of false positive cloud detections by both instruments; (b) the cloud detection agreement is better for the lower layers. Above ~7 km, the ALADIN product demonstrates lower sensitivity because of lower backscatter at 355 nm and because of lower signal-to-noise ratio; (c) in 50% of the analyzed cases when ALADIN reported a low cloud not detected by CALIOP, the middle level cloud hindered the observations and perturbed the ALADIN's retrieval indicating the need for quality flag refining for such scenarios; (d) large sensitivity to lower clouds leads to skewing the ALADIN's cloud peaks down by -0.5 ± 0.4 km, but this effect does not alter the polar stratospheric cloud peak heights; (e) temporal evolution of cloud agreement quality does not reveal any anomaly for the considered period, indicating that hot pixels and laser degradation effects in ALADIN have been mitigated at least down to the uncertainties in the following cloud detection agreement values: $61 \pm 16\%$, $34 \pm 18\%$, $24 \pm 10\%$, $26 \pm 10\%$, and $22 \pm 12\%$ at 0.75 km, 2.25 km, 6.75 km, 8.75 km, and 10.25 km, respectively.

Author: Subject: Sticky Note Date: 14 06 2021 09:43:49

Abstract not understandable

Author: Subject: Comment on Text Date: 14 06 2021 09:44:08

poor phrasing

Author: Subject: Comment on Text Date: 14 06 2021 11:26:31

What do these numbers mean? It is not understandable without reading the paper



1 Introduction

Clouds play an important role in the energy budget of our planet: optically thick clouds reflect the incoming solar radiation, leading to cooling of the Earth, while thinner clouds act as “greenhouse films”, preventing escape of the Earth’s long-wave radiation to space. Climate feedback analyses reveal that clouds are a large source of uncertainty for the climate sensitivity of climate models and, therefore, for the predicted climate development scenarios (e.g. Nam et al., 2012; Chepfer et al., 2014; Vaillant de Guélis et al., 2018). Understanding the Earth’s radiative energy budget requires knowing the cloud cover, their geographical and altitudinal distribution, temperature, composition, as well as the optical properties of cloud particles and their concentration.

Satellite observations have been providing a continuous survey of clouds over the whole globe. IR sounders have been observing our planet since 1979: from the TOVS (TIROS Operational Vertical Sounder) instruments (Smith et al., 1979) onboard the NOAA polar satellites to the AIRS (Atmospheric InfraRed Sounder) spectrometer (Chahine et al., 2006) onboard Aqua (since 2002) and to the IASI (Infrared Atmospheric Sounder Interferometer) instrument (Chalon et al., 2001; Hilton et al., 2012) onboard MetOp (since 2006), with increasing spectral resolution. Despite an excellent daily coverage and daytime/nighttime observation capability (Menzel et al., 2016; Stubenrauch et al., 2017), the height uncertainty of the cloud products retrieved from the observations performed by these spaceborne instruments is limited by the width of their channels’ contribution functions, which is on the order of hundreds of meters, and the vertical profile of the cloud cannot be retrieved with accuracy needed for climate feedback analysis. This drawback is eliminated by active sounders, the very nature of which is based on altitude-resolved detection of backscattered radiation, and the vertical profiles of the cloud parameters are available from the CALIOP (Cloud-Aerosol Lidar with Orthogonal Polarization) lidar (Winker et al., 2003) and CloudSat radar (Stephens et al., 2002) since 2006, CATS (Cloud-Aerosol Transport System) lidar on-board ISS provided measurements for over 33 months starting from the beginning of 2015 (McGill et al., 2015). The ALADIN (Atmospheric Laser Doppler Instrument) lidar on-board Aeolus (Krawczyk et al., 1995; Stoffelen et al., 2005; ADM-Aeolus Science report, 2008) has been measuring horizontal winds and aerosols/clouds since September 2018. More lidars are planned – in 2023, the ATLID (ATmospheric LIDar)/EarthCare instrument (Hélière et al., 2012) will be launched and other space-borne lidars are in the development phase. Even though all active instruments share the same measuring principle – a short pulse of laser or radar electromagnetic radiation is sent to the atmosphere and the time-resolved backscatter signal is collected by the telescope and is registered in one or several receiver channels, the wavelength, pulse energy, pulse repetition frequency (PRF), telescope diameter, orbit, detector, and many other parameters are not the same for any given pair of current or future instruments. These differences are responsible for the active instruments’ capability of detecting atmospheric aerosols and/or hydrometeors for given atmospheric scenario and observation conditions (day, night, averaging distance). At the same time, there is an obvious need of ensuring the continuity of global spaceborne measurements and obtaining a seamless transition between the satellite missions (Chepfer et al., 2018).



This work seeks to address this issue using ALADIN/Aeolus spaceborne wind lidar operating at 355 nm and CALIOP/CALIPSO atmospheric lidar operating at 532 nm. Even though the main goal of ALADIN is wind detection (Reitebuch et al., 2020; Straume et al., 2020), the calibration of which does not rely on absolute calibration of the detected radiation, its products include atmospheric optical properties and such a comparison serves the inter-calibration purposes. In addition, the methods developed in the course of this study, and the interpretation of the results will set the stage for the future validation of the ATLID/EarthCare instrument and other spaceborne lidars.

The structure of the article is as follows. In Section 2, we describe the datasets used in this study, explain the collocation criteria, and provide an estimate of the best possible theoretically achievable agreement for two instruments in given configuration. In Section 3, we strive to provide a multifaceted view of the collocated dataset and discuss the observed differences. Section 4 concludes the article.

2 Datasets and methods

We start this section with the description of ALADIN/Aeolus optical properties dataset followed by the description of CALIOP/CALIPSO product and its modification aimed at matching the sampling and averaging of Aeolus product. In the next steps, we define the procedures and criteria for the comparison of these two products.

2.1 AEOLUS

A detailed description of the Aeolus mission and its instrument can be found in (Krawczyk et al., 1995; Stoffelen et al., 2005; ADM-Aeolus Science report, 2008; Flamant et al., 2017) and here we provide only a brief description of the lidar and the details necessary for understanding the key differences between the compared instruments. The Aeolus satellite carries a Doppler wind lidar called ALADIN, which operates at 355 nm wavelength and is composed of a transmitter, a Cassegrain telescope, and a receiver capable of separating the molecular (Rayleigh) and particular (Mie) backscattered photons (HSRL, high spectral resolution lidar). The lidar is aimed 35° from nadir and 90° to the satellite track, its orbit is inclined at 96.97° and the instrument overpasses the equator at 6h and 18h of local solar time (LST), see also Table 1 to compare with CALIOP.

The laser emitter sends 15 ns long pulses of 355 nm radiation down into the atmosphere 50 times per second. The telescope collects the light that is backscattered from air molecules, aerosols and hydrometeors. The received backscatter signal in Mie receiver passes through a Fizeau interferometer, which produces a linear fringe whose position on the ACCD (Accumulation Charge Coupled Device) detector of this channel is linked to the wind velocity. As for the Rayleigh receiver, it uses a dual-filter Fabry-Pérot interferometer, which throws two images on the ACCD detector of this channel, and the wind speed is defined from the ratio of intensity of these two images (Chanin et al., 1989). Besides the winds, the Aeolus processing algorithms retrieve the optical properties of the observed atmospheric layers (Ansmann et al., 2007; Flamant et al., 2017). The vertical resolution of the instrument is adjustable, but the total number of points in a vertical profile is defined by a number of rows of the detector dedicated to this purpose (24). The observation priorities changed throughout the period of the mission.

Author: Subject: Cross-Out Date: 17 05 2021 14:55:03

Author: Subject: Comment on Text Date: 14 06 2021 09:45:08
This statement is very vague

Author: Subject: Sticky Note Date: 17 05 2021 14:56:29
not a single word of scattering ratios so far

Author: Subject: Highlight Date: 17 05 2021 14:59:05



(Bley et al., 2021), and for the majority of the period considered in this work (see below), the vertical sampling of both Mie
95 and Rayleigh channels between 2 km and 22 km was equal to 1 km whereas the sampling below 2 km varied from 0.25 to
1 km. The native horizontal resolution of 140 m of the instrument is sacrificed to achieve a better signal to noise ratio both
onboard by accumulating the detected profiles and on the ground by averaging the downloaded profiles at different steps of
the processing chain (Flamant et al., 2017).

The present study has been done using the pilot L2A dataset from Aeolus, **Prototype v3.10**, which is available for a limited
100 **period of ALADIN's observations, from 28/06/2019 through the 31/12/2019**. According to (Flamant et al., 2017), the L2A
data is produced from the L1B product of this instrument and it contains height profiles of Mie and Rayleigh co-polarized
backscatter and extinction coefficients, **scattering ratios**, and lidar ratios (Flamant et al., 2008; Lolli et al., 2013) along the lidar
line-of-sight. For the end user, the profiles are provided both on observation scale (87 km averages) and on smaller scales after
applying scene classification, but for the purposes of the present work, the scattering ratio on the scale of 87 km is an optimal
105 choice.

In Fig. 1(a-c), we show the observation geometry and sampling of ALADIN's L2A product as well as three variables retrieved
from its observations, namely, the APB (Attenuated Particulate Backscatter), the AMB (Attenuated Molecular Backscatter),
and the ATB (Attenuated Total Backscatter). The white dashed lines in Fig. 1 represent the line of sight of the instrument.
One has to note, however, that in the real life the ALADIN's line of sight is pointed perpendicular to the flight direction; at the
110 same time, **the horizontal variability of the observed scene is nearly the same in latitudinal and longitudinal directions** at
100 km distance, so the **sketch gives an idea of** the comparability of the physical parameters observed by ALADIN (Fig. 1a-
c) and CALIOP (Fig. 1d). The atmospheric scene used in Fig. 1 has been calculated for demonstration purposes for two
wavelengths, 355 nm (Fig. 1a,b,c) and 532 nm (Fig. 1d) from the output of the EAMv1 (Energy Exascale Earth System Model
(E3SM) atmosphere model version 1) atmospheric model (Kasch et al., 2019) **for the conditions of autumn equinox in Northern**
115 **hemisphere**. This data has been obtained with the help of the COSP2 (the Cloud Feedback Model Intercomparison Project
Observational Simulator Package, v2) package, which is capable of simulating the atmospheric observables for spaceborne
instruments (Swales et al., 2018). The CALIOP is built into COSP2 (Chepfer et al., 2008) whereas the ALADIN is not yet a
part of this package, so we used the 355 nm calculations by COSP2 (Reverdy et al., 2015) at fine grid corresponding to
ALADIN's original laser pulse frequency rate and **modified them in accordance with the ALADIN's vertical and horizontal**
120 **averaging**. The **cloud variability along the satellite's track has been estimated from the gridded EAMv1 data using the**
parameterization of (Boutle et al., 2014). **Figure 1 also serves as an illustration to theoretically achievable cloud detection**
agreement discussed below.

For each profile corresponding to an inclined dashed line in Fig. 1, we extracted the corresponding scattering ratio (SR) column
of **sca_optical_properties** group of variables where SCA stands for standard correct algorithm (Flamant et al., 2017). An
125 important companion of such a column is a corresponding quality flag column, which we scanned looking for the points
characterized either **by high Mie signal-to-noise ratio (SNR) or by high Rayleigh SNR**, and by a flag that indicates an absence

- Author: Subject: Highlight Date: 14 06 2021 11:05:32
No reference given. You need to explain what this means and what is the difference to operational data
- Author: Subject: Comment on Text Date: 18 05 2021 09:52:16
the scattering ratio you use is never defined, as it is essential for this work, you should do so
- Author: Subject: Comment on Text Date: 17 05 2021 15:05:39
this is not true and heavily depends on the scene
- Author: Subject: Comment on Text Date: 17 05 2021 15:07:53
it gives a wrong idea, because scenes are usually by far not so homogenous
- Author: Subject: Comment on Text Date: 17 05 2021 15:08:41
for what do you need the autumn equinox in this simulation?
- Author: Subject: Comment on Text Date: 17 05 2021 15:09:42
what does this mean?
- Author: Subject: Comment on Text Date: 14 06 2021 09:41:29
I don't understand, more explanation needed. I do not see any cloud in this figure
- Author: Subject: Comment on Text Date: 14 06 2021 11:07:13
nobody who is not familiar with Aeolus L2A data structure will understand this
- Furthermore, it is not trace-able/understandable for anyone not within an Aeolus Cal/Val team yet. A proper reference should be given or, if not available, a more detailed description of the data set needs to be given here. E.g., what is the difference to other Aeolus data? Why have you used this data set, etc.
- Author: Subject: Comment on Text Date: 14 06 2021 11:06:50
what does this mean: HIGH SNR? >2, >10, >100?



of signal attenuation. Presumably, these flags are necessary and sufficient for a valid SR profile, which can be then compared with that of CALIOP.

2.2 CALIPSO-GOCCP

130 CALIOP, a two-wavelength polarization-sensitive nadir viewing lidar, provides high-resolution vertical profiles of aerosols and clouds. Its 705 km orbit is inclined at 98.05° and it overpasses the equator at 1h30 and 13h30 LST, see also Table 1. It uses three receiver channels: one measuring the 1064 nm backscatter intensity and two channels measuring orthogonally polarized components of the 532 nm backscattered signal. Cloud and aerosol layers are detected by comparing the measured 532 nm signal return with the return expected from a molecular atmosphere.

135 The CALIPSO-GOCCP (GCM Oriented Cloud Calipso Product) was initially designed to evaluate GCM cloudiness (Chepfer et al., 2010). It is derived from CALIPSO L1/NASA products at LMD/IPSL with the support of NASA/CNES, ICARE, and ClimServ and it contains observational cloud diagnostics including the instantaneous scattering ratio (profiles) at the native horizontal resolution of CALIOP (333 m) and at ~0.5 km vertical resolution. This makes it a good reference dataset for ALADIN retrievals because it can be easily recalculated to the latter's horizontal and vertical grids considering the corresponding horizontal averaging. Since the CALIOP is not a HSRL, the detailed information on AMB and APB is not available, and one has to compare the SR products. Correspondingly, we convert the ALADIN's SR retrieved at 355 nm to SR at 532 nm using the following equation:

$$140 \quad SR_{532} = SR_{355} \times 3.3 - 2.3 \quad (1)$$

145 which is derived from (Collis and Russell, 1976) in an assumption that their fitting parameter Λ (see their Section 4.3.1) is equal to 3. The choice of the fitting parameter is not crucial for the purposes of the present work because the conversion described by Eq. 1 is linear and it does not change the altitude distribution of the SR. On the other hand, using the same physical parameter is highly advisable for the comparisons we are intending to perform. Theoretically, one could have validated the parameters of Eq. 1 using the collocated data under consideration, but, looking ahead, one can say that the spread of the values is too large to do it with reasonable uncertainty, so we will stay with Eq. 1 in the framework of this paper and in Appendix A we justify our choice of conversion coefficients using the collocated data.

2.3 Collocation criteria

155 As for any collocation, there is a trade-off between the quality of collocation and the number of collocated pairs of profiles. As we show below, in the case of AEOLUS and CALIPSO, this tradeoff is supplemented with a requirement of a representative geographical coverage, because imposing a strict temporal overlap criterion dramatically changes the latitudinal distribution of the collocated points. Since the horizontal averaging and resolution of the Aeolus Prototype_v3_10 product is 87 km, there is no much sense in collocating the data with the accuracy better than this value. On the other hand, a fractional standard deviation f_c of cloud water content at 1° (~111 km) distance is about 0.5 for a cloud cover of 1 (Boutle et al., 2014), and there

-
- Author: Subject: Comment on Text Date: 14.06.2021 11:07:30
I am not sure if these flags are valid in these kind of data. These data are all preliminary. You should discuss this.
-
- Author: Subject: Comment on Text Date: 18.05.2021 09:56:56
This statement is in contradiction to Figure 1, where you clearly see that it is not nadir but only close to nadir.
-
- Author: Subject: Comment on Text Date: 18.05.2021 09:56:24
I guess you mean the altitude of the orbit, but this is not written here. Please state correctly.
-
- Author: Subject: Comment on Text Date: 18.05.2021 10:26:48
GCM never explained.
-
- Author: Subject: Comment on Text Date: 14.06.2021 09:54:36
This is not state of the art and not acceptable - see plots in my text.
-
- Author: Subject: Comment on Text Date: 14.06.2021 09:55:42
I do not understand this statement. And the justification given in the Appendix is not sufficient in my opinion.



is a risk of comparing incoherent quantities, so we took $\Delta\text{time} = 1^\circ$ as a limit for the collocations and created several subsets based on the Δtime , the absolute value of the difference between two collocated measurements. In Fig. 2, we show three such subsets, and the Table 2 provides the information about the other cases we considered. On the one hand, one can see that a strict collocation criterion of $\Delta\text{time} < 1\text{h}$ (Fig. 2a) provides the information only about two narrow zones in the Southern and Northern polar regions. On the other hand, an excellent geographical coverage shown in Fig. 2c comes at the cost of mixing up the cases, which differ by almost one day that is unacceptable from the point of view of temporal variation. In addition, this case is characterized by unequal distribution of Δtime throughout the globe. Finally, a subset corresponding to $\Delta\text{time} < 6\text{h}$ (Fig. 2b) has been chosen for the analysis. Over the oceans, the diurnal effects in cloud distribution associated with this difference are small (e.g. Noel et al., 2018; Chepfer et al., 2019; Feofilov and Stubenrauch, 2019) and the land represents one third of the analyzed cases. To avoid the risks associated with the solar contamination, we picked up only the **night-time cases**, which yield about 7.8×10^4 pairs of SR profiles. In supplementary materials, we provide the complete collocated database, which corresponds to the last line, 4th column of Table 2 (3.7×10^5 collocations), for further analysis by the interested teams.

2.4 Estimating the theoretically achievable agreement between two collocated datasets

To justify the collocation criteria and to estimate the theoretically possible agreement for the clouds detected by two instruments in a given setup and for the selected Δtime and Δdist values, we have performed a numerical experiment using the same calculated data as we used in Fig. 1. This time, we picked up the “thick curtain” at 532 nm, calculated at the resolution of CALIOP (333m) and created artificial pairs of “collocated” data with the Δdist distribution modulated by that of a real collocated dataset. The “reference”, CALIOP profile has been composed using 2000 individual SR profiles covering 67 km region that is somewhat less than the 87 km covered by ALADIN. This averaging is supposed to catch the mean atmospheric properties and at the same time it is not supposed to go too far from the ALADIN footprint location. The “test” SR profile was created from the SR averages, considering both the ALADIN’s off-nadir pointing and its 87 km averaging. To imitate the diurnal variation, we modulated the SRs using the 6-hour diurnal cycle amplitudes for land and ocean retrieved from active and passive observations (Noel et al., 2018; Chepfer et al., 2019; Feofilov and Stubenrauch, 2019) and added them to the comparison. Besides testing a noise-free simulation, we also checked the effects introduced by instrumental noise for CALIOP. Since ALADIN is not yet part of COSP2, we used the estimates from (Kinsmann et al., 2007). Overall, we considered about 1×10^5 pairs of pseudo-collocated data and we present the results of cloud detection in Fig. 3. We define the cloud detection agreement as follows: for each altitude bin, the cloud detection agreement is a ratio of a number of cases when both instruments have detected a cloud ($\text{SR} > 5$) to a total number of joint observations. For a given altitude bin, the cloud amount is a ratio of number of cases with $\text{SR} > 5$ to a total number of profiles for a single instrument, and the normalized cloud detection agreement is a ratio of the former to the latter. As one can see, the normalized cloud detection quality is mostly defined by a horizontal variability of aerosols/hydrometeors and by differences in viewing geometries of two instruments. Observation noise and diurnal variation play the secondary role, and according to our estimates the saturation effects in 355 nm and 532 nm channels associated with opaque clouds (Guzman et al., 2017) do not add more than 2% to the cloud detection mismatch (not shown in

Author: Subject: Comment on Text Date: 14.06.2021 11:07:47

As Aeolus is flying at a dusk-dawn orbit, how can you select night time cases? Or is this valid for Calipso only? But then you have a bias, right?

Author: Subject: Comment on Text Date: 14.06.2021 09:56:04

I really did not understand what you are doing here. Maybe a sketch or flowchart could help.

Why do you need to imitate diurnal variation? Please explain!

Author: Subject: Comment on Text Date: 14.06.2021 11:09:06

The choice of this threshold is not clear to me and seems very arbitrary. Furthermore, the use of a scattering ratio for cloud detection is questionable to me at all, as the scattering ratio depends on temperature and pressure and thus on height even with uniformly distributed particle load as shown in the attached plots. Using one threshold would mean that you detect a cloud at one height while you may not detect a similar cloud at another height. Thus, using the scattering ratio for cloud detection is not appropriate in my opinion. You could have used the scattering ratio from the satellite data to calculate the real particle backscatter coeff. by using the molecular backscatter which you calculate from the meteorological data which is included in the satellite data as well.

Author: Subject: Comment on Text Date: 18.05.2021 15:10:14

I do not “see” that. At least in Fig. 3 it is not obvious. To what you are referring to? And can you give more explanation?

Author: Subject: Comment on Text Date: 14.06.2021 10:00:39

Any evidence or proof for such a statement? Again to what are you referring?

Author: Subject: Comment on Text Date: 18.05.2021 15:12:08

is this valid for ALADIN as well?



Fig 3 for the sake of clarity) Overall, the theoretically achievable agreement for the collocated data at a given setup can be estimated as 0.77 ± 0.17 for cloud detection

3 Results and discussion

3.1 Zonal averages

195 To give a general overview of the agreement between two products, we have split the database to latitudinal zones: 90S–60S, 60S–30S, 30S–30N, 30N–60N, 60N–90N (Fig 4). As it was stated above, we recalculate the SR_{333} values retrieved from ALADIN observations to SR_{332} using Eq. 1. Even though the zonal mean statistics does not imply using collocated data, we do it to avoid any incoherence in sampling different geographic areas. By using exactly the same number of profiles collocated within 1° , we ensure the same coverage and sampling by both lidars. In the detection efficiency of different cloud types were the same for two instruments, the plots would have been close to each other, because the horizontal variability of clouds would cancel out due to averaging over a large number of profiles within the zone and the diurnal variation is small over oceans, which constitute two thirds of the cases used to build Fig 4 (Neel et al., 2016; Chepfer et al., 2019; Feofilov and Stubenrauch, 2019). Analyzing the Fig 4, one can note the following: (1) the SR/altitude histograms of CALIOP (Fig. 4a-e) are characterized by two distinct peaks corresponding to low-level and high-level clouds; this feature is coherent with other observations, e.g. with GEWEX (Global Energy and Water cycle Experiment) cloud assessment (Stubenrauch et al., 2013); (2) the SR/altitude histograms built for SR_S retrieved from ALADIN's observations (Fig. 4f-j) are characterized by a smoother occurrence frequency plot where the two-peak structure is less pronounced than for CALIOP; (3) even though ALADIN detects some clouds in polar stratosphere (PSCs), its overall sensitivity to high clouds (>7 km) is lower than that of CALIOP; (4) both rows show certain consistency of zone-to-zone change up to ~ 3 km altitude while the behavior above requires a more detailed view.

200 We would like to stress here that no linear scaling applied uniformly to SR s at all heights could change the ratio of high cloud detection frequency to low cloud detection frequency of ALADIN. The same is true for CALIOP. In the next step, we compare the “instantaneous” profiles provided by CALIOP and ALADIN having in mind the peculiarities of cloud detection sensitivity differences observed in Fig 4.

3.2 Comparing pseudo-individual profiles at ALADIN's L2A product resolution

215 To address the high cloud detection sensitivity, we have inspected the 6h nighttime subset of collocated data, looking for the cases, which would satisfy the following criteria: (1) both instruments should have at least one strong SR peak; (2) the vertical position of this peak detected by one instrument should match that of the peak detected by a second instrument within 1 km; (3) the CALIOP SR profile should have a secondary peak at or above 9 km (Fig. 5a-j). For the comparison purposes, the panels in Fig. 5 represent the individual profiles belonging to the same 5 zones as the panels of Fig. 4. For the sake of simplicity, we compare the $SR_{333}(z)$ profiles recalculated to $SR_{332}(z)$, but we also show the source $SR_{333}(z)$ profiles for reference purposes.

220 Regarding the conversion using Eq. 1, the strong peaks selected this way demonstrate a qualitative agreement between the

- Author: Subject: Comment on Text Date: 18 05 2021 15:14:57
which products? SR or cloud product? Unclear
- Author: Subject: Comment on Text Date: 18 05 2021 15:15:04
phrasing
- Author: Subject: Comment on Text Date: 18 05 2021 15:16:04
phrasing
- Author: Subject: Comment on Text Date: 18 05 2021 15:17:14
maybe because your transformation of the SR is incorrect?
- Author: Subject: Comment on Text Date: 18 05 2021 15:18:07
if you know that, why have you done so?
- Author: Subject: Comment on Text Date: 14 06 2021 09:07:46
What is true for CALIOP?
- Author: Subject: Comment on Text Date: 14 06 2021 11:09:47
is the feasible according to the range-bin setting of Aeolus. In principle, Aeolus can also have 2 km thick range bins. Can you comment on this?
- Author: Subject: Comment on Text Date: 14 06 2021 10:08:26
I do not see a reason to use the "sake of simplicity" here. A correct conversion is needed.



Author: Subject: Comment on Text Date: 14 06 2021 10:09:39
This is not convincing

Author: Subject: Comment on Text Date: 14 06 2021 10:10:13
how do you account for the different vertical resolution?

Author: Subject: Comment on Text Date: 14 06 2021 10:11:10
This statement is not clear for me

Author: Subject: Comment on Text Date: 14 06 2021 10:14:44
Phrasing needs to be improved

peak values calculated from SR_{335} and peak retrieved SR_{332} values. In Appendix A, we demonstrate the correlation between individual pairs of CALIOP and ALADIN SR profiles; the conclusion of this exercise is that it matches using Eq. 1, but the uncertainties of the analysis do not allow to refine the conversion coefficients. As for the potential capability of ALADIN to detect high clouds, the subset Fig. 5a-e represents the cases, for which the instrument was capable of retrieving the peak of the same magnitude and height as the peak detected by CALIOP. Even though these cases exist, they are far less frequent than those shown in Fig. 5f-j. We did not detect and correlation between the collocation criteria ($\Delta dist$; $\Delta time$) and the frequency of occurrence of these cases, it's just a statistical observation that both types of cases exist and the former are less frequent than the latter. This observation gives a hint that the instrumental part provides the backscatter information sufficient for some cloud detection up to 20 km, but the detection algorithm suppresses noisy solutions. The PSC detection discussed below (see also Fig. 4f) confirms this assumption because the vertical extent and the composition of these clouds yield a strong signal. Further speculations on this subject are beyond the scope of the present article, but we believe that the high cloud detection agreement might be improved by studying the collocated cases provided in the supplementary materials and by applying different noise filtering techniques in the L0→L1→L2 elements of the ALADIN retrieval chain. Figures 5k-o will be discussed below in the context of low-level cloud observations.

3.3 Cloud detection agreement

To illustrate the peculiarities of zonal and altitudinal behavior of cloud detection agreement between two considered instruments, we have split the collocated data into four groups (Fig. 6). For each altitude/latitude grid point, we have estimated the number of cases when both instruments have detected a cloud ($SR_{332}(z) > \delta$), when neither of instruments has detected a cloud, when only CALIOP has detected a cloud, and when only ALADIN has detected a cloud. For the sake of simplicity, we will call them YES_YES, NO_NO, YES_NO, and NO_YES cases. It is clear that in the ideal experiment the number of mismatched cases (YES_NO and NO_YES) should tend to zero. From the study presented in Section 2.4, we expect that the ratio of $(YES_YES + NO_NO) / (YES_YES + NO_NO + YES_NO + NO_YES)$ should be about 0.77 ± 0.17 if both instruments detect the clouds with the same efficiency. In Fig. 6a we show the ratio of YES_YES cases to the total number of collocated profiles per altitude/latitude bin. This panel resembles a typical cloud amount plot, and this is expected because in the case of an ideal agreement the aforementioned ratio is equivalent to cloud amount definition. Below, we will also discuss the YES_YES statistics normalized to cloud amount, but at this point we also want to study the other cases, which cannot be normalized this way. Even though the distribution in Fig. 6a looks physical, the absolute numbers are somewhat low and this is explained by YES_NO and NO_YES distributions (Fig. 6c and d, respectively). As for NO_NO agreement (Fig. 6b), it is close to 100% in the high-altitude area where there are no clouds. This indicates that the noise-induced false detection rate of both instruments is low, and this is a good sign.

If we consider the mismatch of YES_NO type (Fig. 6c), we will see that the altitudinal/zonal distribution of the mismatch occurrence frequency resembles that of the YES_YES type. A part of mismatch can be explained by theoretically allowed cloud detection disagreement discussed in Section 2.4. However, the occurrence frequency of YES_NO cases above 3 km is



-
- Author: Subject: Comment on Text Date: 14.06.2021 10:17:13
What does it mean? It could be also a cause of your rough conversion of the scattering ratio and/or the range-bin thickness of Aeolus?
-
- Author: Subject: Comment on Text Date: 14.06.2021 10:18:28
Phrasing! Is this really only one cloud?
-
- Author: Subject: Comment on Text Date: 14.06.2021 10:19:55
Can you explain, how these false peaks could develop? It is not clear to me

255 roughly twice that of YES_YES cases, and this indicates the retrieval sensitivity issue of ALADIN. The NO_YES mismatches (Fig. 6d) require specific attention because they are not expected from the methodological point of view: the cloud extinction at 355 nm is larger than at 532 nm and the observation geometry of ALADIN makes the optical paths $1 / \cos(SVA) = 1.22$ times longer than those for CALIOP, where SVA stands for satellite viewing angle of 55° . The typical individual profiles corresponding to NO_YES mismatches are shown in Fig. 5k-o. As one can see, despite the unfavorable observation conditions (e.g. an opaque cloud with peak SR_{352} value of ~ 22 at 9 km in Fig. 5l), ALADIN reports two valid points beneath the cloud, whereas it does not report anything at 9 km height where CALIOP sees a thick cloud. These cases do need our special attention. On the one hand, many cases of this type are over the ocean, so one can rule out the surface echo mixed with atmospheric backscatter and treated like an atmospheric signal. On the other hand, the NO_YES cases are often accompanied by the structures similar to those presented in Fig. 5k,l,n which are probably provoked by a presence of a cloud at these heights. The perturbations to the extinction and backscatter profile caused by these structures might propagate downwards, thus causing the appearance of the false peaks in the lower layers of ALADIN's data. This indicates a need for a quality flag refinement in the lower layers in the presence of a thick cloud above and the improvement of thick cloud detection itself. Apparently, the CALIOP cloud retrievals beneath thick clouds do not suffer from these effects.

To test whether the aforementioned disagreements are at least partially caused by the cloud definition and SR recalculation to another wavelength and whether the agreement could be improved, we varied the SR threshold for ALADIN, assuming the $\pm 50\%$ uncertainty on the parameters forming the coefficients of Eq. 1. However, this exercise yielded no optimum value for SR threshold: its lowering for ALADIN increased the number of YES_YES and reduced the number of YES_NO cases, but at the same time it increased the frequency of NO_YES cases. Correspondingly, increasing the threshold reduced the number of NO_YES cases, but it adversely affected the YES_YES agreement. Summarizing this comparison, one can conclude that

275 (a) a cloud detected by CALIOP is detected by ALADIN in $\sim 50\%$ of cases for clouds below ~ 3 km and in $\sim 30\%$ of cases for higher clouds; (b) in the cloud-free area, the agreement between the datasets is good that indicates a low frequency of false positive detections by both instruments; (c) one half of the cases when ALADIN detects a cloud missed by CALIOP should be attributed to false positive detection of the low cloud in the presence of a higher opaque cloud, which perturbs the retrieval in the lower layers.

280 3.4 Cloud altitude detection sensitivity

Besides marking the profile elements as “cloudy” and “not cloudy” and comparing the cloud detection statistics as we did in the previous section, it would be interesting to obtain cloud peak detection statistics for pairs of collocated profiles like those shown in Fig. 5. This exercise is not aimed at revealing any altitude offset in backscatter signal registration, because this part of experimental setup is robust in both instruments. But, as we saw in Fig. 4 and Fig. 6, the sensitivity of ALADIN to high clouds is lower than to lower clouds and a convolution of sensitivity curve with the backscatter profile can skew the cloud peak position and the average cloud height. To illustrate this effect, we have carried out the following analysis. For each pair

285



of collocated profiles selected for YES_YES plot (Fig. 6a), we scanned through ALADIN profile step by step, looking for a local maximum, which we define as a set of the following conditions:

$$SR(i) > SR_{threshold}; SR(i) > SR(i - 1); SR(i) > SR(i + 1) \quad (2)$$

290 where $SR_{threshold}$ is the cloud detection threshold at 532 nm, which is equal to 5. For each local peak found, we have searched for a peak or for a maximal value of CALIOP's SR profile in the vicinity of ± 3 km from the peak height determined from ALADIN. The choice of a "reference" dataset in this case depends on the detection probability, and if one chooses CALIOP as a reference, the distance to the nearest ALADIN peak might be spoiled by lower probability of cloud detection by ALADIN and the distribution will be skewed. The search limits are arbitrary and they have been chosen from inspecting the collocated
295 profiles taking into account the natural variability of cloud heights at distances of about 100 km, estimated from the analysis of CALIOP data used in this study (~75% of clouds move vertically by less than 1 km, ~8% by 1–2 km, ~5% by 2–3 km, ~4% by 3–4 km, ~3% by 4–5 km and ~5% by more than 5 km). The differences between the ALADIN's and CALIOP's cloud peak heights have been stored and then averaged in the corresponding latitude/altitude bins (Fig. 7). As one can see, the cloud height detection agreement is better than 0.2 km below ~3 km and, surprisingly, for some of high-altitude zones. For the tropical
300 zone, this is probably linked with thick Ci clouds which should be reliably detected by both instruments. For the Southern polar zone, this figure reveals the PSCs, which are barely visible in Fig. 6a, but which can be seen in Fig. 4f for ALADIN. These clouds form at very low temperatures and are composed of ice particles yielding a reflection, which is reliably detected at both wavelengths if the layer is thick (e.g. Adriani et al., 2004; Snels et al., 2021). As for the clouds between ~3 km and ~10 km height, the height sensitivity effects skew the effective cloud height detected by ALADIN downwards by 0.5–1.0 km.
305 This is coherent with Fig. 4, which shows lower frequency of occurrence of high clouds detected by ALADIN. At least a part of the cloud peak shifts in the 3–5 km layer should be attributed to the reasons discussed for NO_YES statistics and these differences should reduce when the aforementioned quality flags for cloud-perturbed retrievals are fixed.

3.5 Temporal evolution of cloud detection agreement

ALADIN is a relatively young instrument, and its calibration/validation activity is still on the way (Baars et al., 2020; Donovan
310 et al., 2020; Kanitz et al., 2020; Reijebuch et al., 2020; Straume et al., 2020). This includes, but is not limited to internal calibration and comparisons with other observations. The Aeolus mission faced a number of technical issues, which hindered obtaining the planned specifications. These issues are related to several factors: (a) laser power degradation (60 mJ/pulse instead of 80 mJ/pulse) and signal losses in the emission and reception paths (33%) that results in lower signal to noise ratio (SNR) than planned, (b) telescope mirror temperature effects biasing the wind detection and calibration of Mie and Rayleigh
315 channels of ALADIN, (c) constantly increasing number of hot pixels of both ACCD detectors (Weiler et al., 2021) leading to errors both in wind speed and in retrieved optical parameters of the atmosphere (the number of hot pixels increased by a factor of 1.4 during the period considered in this work). The Aeolus teams managed to mitigate some of these adverse effects (e.g. Baars et al., 2020; Weiler et al., 2021), and it would be interesting to see whether the pilot L2A dataset, Prototype_v3 10 is



free of cloud detection quality trends. If true, this would indicate a good calibration and consistent processing of Level 0 through Level 1 to Level 2A.

In Fig. 8 and 9 we show the temporal evolution of cloud detection agreement per height bins. The panels of Fig. 8 are consistent with those of Fig. 6, whereas Fig. 9 considers only the evolution of YES_YES statistics, which corresponds to Fig. 6a and Fig. 8a, normalized by cloud amount. Unfortunately, the period available for analysis does not cover the whole year, so the plots can be affected by seasonal variation of cloud distributions. Still, the latitudinal and longitudinal coverage of collocated data does not change throughout the year and a mixture of Northern and Southern hemispheres should partially compensate for seasonal anomalies. The signatures one should be looking for are experimental artefacts linked with laser power degradation, hot pixels appearance, and bias corrections. If these issues are not properly compensated, the “agreement panels” (Fig. 8a,b) should demonstrate a decrease in occurrence frequency with time and the occurrence frequency in “disagreement panels” (Fig. 8c,d) should increase with time. As one can see, this is not the case: visually, all 4 panels of Fig. 8 do not show any anomaly, which would go beyond their noise levels (a special region corresponding to a forced bin size reduction in the period of 28/10/2019–10/11/2019 is marked by white dashed lines in Fig. 8 and should not be considered at heights below 2250m). To quantify the tendencies and to compare them with noise levels, we have normalized Fig. 8a (YES_YES cases) by cloud amount per altitude/time bin. This procedure helps to get rid of seasonal variation of clouds. The results presented in Fig. 9 confirm the previous conclusions regarding the altitude distribution of cloud detection agreement: for the clouds below 3 km it is better than for higher ones (61±16% and 34±18% for 0.75 and 2.25 km, respectively versus 24±10%, 26±10%, and 22±12% for 6.75 km, 8.75 km, and 10.25 km, respectively). As for the tendencies, the low-level clouds demonstrate an improvement towards the end of the year whereas the agreement for 6.75 km and 10.25 km becomes slightly worse by the end of the considered period.

If we compare the hot pixels distribution for Mie and Rayleigh channel ACCD detectors at the beginning and at the end of the time scale of Fig. 8 and 9 (Table 2 of Weiler et al., 2021), we will see 3 and 5 new hot pixels for Mie and Rayleigh matrices, respectively. Even though the Rayleigh matrix pixels are not directly linked to cloud detection, their information is used for the ALADIN SR calculations. For Mie matrix, the lowermost hot pixel, which appeared during the considered period, corresponds to ~15 km height and this cannot affect the tendencies shown in Fig. 9. As for new Rayleigh hot pixels, the lowermost two correspond to 1 km height, the next two – to 5 km height, and the last one – to 18 km. This information does not explain the observed behavior, either. Overall, considering relatively large error bars for all five altitudinal sections presented in Fig. 9b and the variety of the observed slopes, one cannot make a sound conclusion neither regarding the deterioration (or the improvement) of cloud detection agreement nor regarding the link between hot pixels appearance and change of cloud detection quality. A proper conclusion is that one does not detect the tendencies beyond the variability limits of the analyzed parameter and that the hot pixels appearance cannot be tracked from the cloud agreement plot, indicating that compensation for hot pixels effects (Weiler et al., 2021) works properly within the discussed uncertainty limits. The same can be said regarding the other known technical issues: the signal losses in the emission and reception paths do not transform into



a clear signature in cloud detection agreement plots. Moreover, they should have affected the detection of low and high clouds in the same way that is not observed in Fig. 8 and 9.

4. Conclusions

355 The active sounders are advantageous for atmospheric and climate studies because they provide atmospheric parameters at altitude resolved scale with high accuracy. For continuity of climate studies and monitoring the global changes it is essential to understand the differences between spaceborne lidars operating at different wavelengths, flying at different orbits, and utilizing different observation geometries, receiving paths, and detectors. In this article, we addressed an intercomparison of ALADIN and CALIOP lidars using their scattering ratio products (CALIPSO-GOCCP and Aeolus L2A, Prototype_v3_10) for the period of 28/06/2019–31/12/2019.

360 Using the COSP2 lidar simulator coupled with output from the EAMv1 model and a horizontal cloud variability parameterization, we estimated a theoretically achievable agreement in cloud detection of 0.77 ± 0.17 for these two instruments with their orbits, averaging, and observation geometry.

On the one hand, the spatial collocation criterion of 1° chosen in this work is based on averaging distance of Aeolus L2A Prototype_v3_10 data. On the other hand, the temporal collocation criterion of $\Delta\text{time} < 6\text{h}$ is a tradeoff between the geographical coverage of the collocated profiles, their number, and uniformity of Δtime distribution throughout the globe. With the named criteria, we managed to find $\sim 7.8 \times 10^4$ collocated nighttime profiles, which underwent a series of analysis summarized here. For the simplicity of the comparison with CALIOP, we converted SR_{355} of ALADIN to SR_{332} and we discuss the sensitivity of the results to the conversion parameters.

370 Overall, the SR product of ALADIN is characterized by lower sensitivity to high clouds above $\sim 7\text{ km}$ than CALIOP, that we explain by lower SNR for ALADIN at these heights that is due both to physical reasons (smaller backscatter at 355 nm) and technical reasons (hot pixels, lower emission and lower transmissivity of receive path than planned). Large sensitivity to lower clouds leads to prioritizing the lower cloud solutions to higher ones in the case of a continuous cloud or a double layer. This skews the ALADIN's cloud peak height in pairs of ALADIN/CALIOP profiles by $\sim 0.5 \pm 0.4\text{ km}$ downwards. Interestingly, the

375 agreement of PSC peak heights does not suffer from these effects. We explain this by large vertical extent and composition of PSCs, which make them a better target for ALADIN than the tropospheric clouds. In the cloud-free area, the agreement between two instruments is good indicating low rate of noise-induced false detection for both instruments. Last, but not least, the temporal evolution of cloud agreement does not reveal any statistically significant change during the considered period. This indicates that hot pixels and laser energy and receiving path degradation effects in ALADIN have been mitigated at least
380 down to the uncertainties of the following cloud detection agreement values: $61 \pm 16\%$, $34 \pm 18\%$, $24 \pm 10\%$, $26 \pm 10\%$, and $22 \pm 12\%$ estimated at 0.75 km , 2.25 km , 6.75 km , 8.75 km , and 10.25 km , respectively. We believe that the provided collocated dataset will facilitate the further analysis and improvement of ALADIN L2A data.

-
- Author: Subject: Comment on Text Date: 14.06.2021 10:28:59
why should there be a smaller backscatter at 355? This is in absolute contradiction to all my knowledge? Particle backscatter coefficient could be equal (Angström 0). Molecular backscatter coefficient is for sure higher and thus total backscatter is for sure also higher!
-
- Author: Subject: Comment on Text Date: 14.06.2021 10:30:01
phrasing poor



Appendix A

The analysis of the collocated data may enable the researcher not only to validate one dataset against another one, but also to validate a physical concept or to retrieve an important model parameter (e.g. Holl et al., 2010; Feofilov and Petelina, 2010; Feofilov et al., 2012; Virtanen et al., 2018). In this section, we report the results of a validation attempt aimed at the retrieval of the scaling coefficients used in Eq. 1 and through them the model assumptions. To do this, we searched the collocated database for the events which would satisfy the following criteria: (a) the ALADIN SR profile should contain at least one valid point with the corresponding quality flags (see Section 2.2) and with SR higher than halved $SR_{threshold}$; (b) the profiles should fit the selection criteria used for cloud altitude detection sensitivity (Section 3.4); (c) the CALIOP peak should contain more than one point to avoid sampling problems. For these profiles, we picked up not only the major peak values, but also the secondary peak values if the vertical agreement of the profiles was good like in Fig. 5a,c,d,e. The corresponding pairs of SR_{CALIOP} and $SR_{ALADIN333}$ values have been binned using the 0.2×0.07 SR bins, which reflect the differences between SR_{332} and SR_{333} . The corresponding frequency occurrence distribution for this dataset is shown in Fig. A1. Even though the SR pairs exist for opaque domain, the spread increases and the values beyond $SR_{332} = 10$ are neither informative nor suitable for the maximal probability search algorithm (see Dawkins et al., 2018) used for the analysis. Like in Fig. 11 of (Dawkins et al., 2018), the red dots in Fig. A1 represent the centers of Gaussian fit to perpendicular transects. White dashed line shows a linear fit to the dataset represented by these red dots, and the corresponding conversion is given by the following equation:

$$SR_{522} = SR_{355} \times (3.8 \pm 1.0) - (3.3 \pm 1.4) \quad (A1)$$

Even though the coefficients in Eq. A1 differ from those of Eq. 1, the black dashed line in Fig. A1 representing Eq. 1 does not significantly deviate from the white dashed line representing Eq. A1 and both lines fit the maximum probability plot within its uncertainty limits. We conclude that the collocated dataset proves the basic equations used to derive Eq. 1 though its uncertainties do not allow to retrieve the corresponding fitting parameter Λ of (Collis and Russell, 1976) from such a comparison.

405

This page contains no comments



Data availability

The collocated dataset used in this work can be downloaded from ResearchGate repository using the following link
<https://doi.org/10.13140/RG.2.2.11237.12009> (Feofilov et al., 2021)

Author contribution

410 HC, VN, MC, and AF: conceptualization, investigation, methodology, and validation; RG, CG, and AF: data curation and
formal analysis; AF: writing original draft; AF and HC: review and editing

Competing interests

The authors declare that they have no conflict of interest

Disclaimer

415 The presented work includes preliminary data (not fully calibrated/validated and not yet publicly released) of the Aeolus
mission that is part of the European Space Agency (ESA) Earth Explorer Program. This includes aerosol and cloud products,
which have not yet been publicly released. Aerosol and cloud products will become publicly available by spring 2021. The
processor development, improvement and product reprocessing preparation are performed by the Aeolus DISC (Data,
Innovation and Science Cluster), which involves DLR, DoRIT, ECMWF, KNMI, CNRS, S&T, ABB and Serco, in close
420 cooperation with the Aeolus PDGS (Payload Data Ground Segment)

Acknowledgements

This work is supported by the Centre National de la Recherche Scientifique (CNRS) and by the Centre National d'Etudes
Spatiales (CNES) through the Expecting Earth-Care, Learning from A-Train (EECLAT) project. The processor development,
improvement and product reprocessing preparation are performed by the Aeolus DISC (Data, Innovation and Science Cluster),
425 which involves DLR, DoRIT, ECMWF, KNMI, CNRS, S&T, ABB and Serco, in close cooperation with the Aeolus PDGS
(Payload Data Ground Segment). The authors want to thank P.-L. Ma (PNNL) for providing the outputs of the EAMv1
atmospheric model and F. Ehlers (EOP-SMA/ESTEC/ESA), A. Straume (ESTEC/ESA), and O. Reiterbuch (DLR) for their
comments on the preliminary version of the manuscript.



430 References

- ADM-Aeolus Science Report, SP-1311, ISBN 978-92-9221-404-3, ISSN 0379-6566, Coordinated by: P Ingmann, Mission Science Division, Atmospheric Unit, Published by: ESA Communication Production Office, The Netherlands, 121pp, 2008
- Adriani, A , Massoli, P , Di Donfrancesco, G , Cairo, F , Moriconi, M J , Snels M : Climatology of polar stratospheric clouds based on lidar observations from 1993 to 2001 over McMurdo Station, Antarctica, *J Geophys Res* , 109, D24211, doi:10.1029/2004JD004800, 2004
- 435 Ansmann, A , Wandinger, U , Le Rille, O , Lajas, D , and Straume, A G : Particle backscatter and extinction profiling with the spaceborne high-spectral-resolution Doppler lidar ALADIN: methodology and simulations *Appl Optics*, 46 (26), 6606-6622, doi:10.1364/AO.46.006606, 2007
- Baars, H , Geiß, A , Wandinger, U , Herzog, A , Engelmann, R , Bühl, J , Radenz, M , Seifert, P , Ansmann, A , Martin, A ,
440 Leinweber, R , Lehmann, V , Weissmann, M , Cress, A , Filioglou, M , Komppula, M and Reitebuch, O : First Results from the German Cal/Val Activities for Aeolus, *EPJ Web Conf* 237, 01008, doi:10.1051/epjconf/202023701008, 2020
- Bley, S, II Consulting Group B.V., Aeolus Atmospheric Sampling - Range Bin Settings, available at <https://www.aeolus.esa.int/confluence/pages/viewpage.action?spaceKey=CALVAL&title=Vs+002,2021>**
- Boutle, I A , Abel, S J , Hill, P G , Morcrette, C J : Spatial variability of liquid cloud and rain: observations and microphysical effects, *Q J R Meteorol Soc* , 140, 583–594, doi:10.1002/qj.2140, 2014
- 445 Chanin, M L , Garnier, A , Hauchecorne, A and Porteneuve, J : A Doppler lidar for measuring winds in the middle atmosphere, *Geophys Res Lett* 16(11), 1273–1276, doi:10.1029/GL016i011p01273, 1989
- Chahine, M T , and 30 Coauthors: AIRS: Improving weather forecasting and providing new data on green-house gases, *Bull Amer Meteor Soc* , 87(7), 911–926, doi:10.1175/BAMS-87-7-911, 2006
- 450 Chalon G , Cayla F R , Diebel D : IASI: An advance sounder for operational meteorology, *Proc 52nd Congress of IAF, Toulouse France, CNES*, available online at http://smc.cnes.fr/documentation/IASI/Publications/PRESENTATION_IAF_2001.pdf, 2001
- Chepfer H , Bony, S , Winker, D , Chiriaco, M , Dufresne, J -L , Sèze, G : Use of CALIPSO lidar observations to evaluate the cloudiness simulated by a climate model, *Geophys Res Let* , 35, L15704, doi:10.1029/2008GL034207, 2008
- 455 Chepfer H , Bony, S , Winker, D , Cesana, G , Dufresne, J -L , Minnis, P , Stubenrauch, C J , and Zeng, S : The GCM Oriented Calipso Cloud Product (CALIPSO-GOCCP) *J Geophys Res* , 115, D00H16, doi:10.1029/2009JD012251, 2010
- Chepfer, H , Noel, V , Winker, D , and Chiriaco, M : Where and when will we observe cloud changes due to climate warming?, *Geophys Res Lett* , 41, 8387–8395, doi:10.1002/2014GL061792, 2014
- Chepfer H , Noel, V , Chiriaco, M , Wielicki, B , Winker, D , Loeb, N , and Wood, R : The potential of multi-decades space-born lidar to constrain cloud feedbacks, *J Geophys Res Atmos* , DOI:10.1002/2017JD027742, 2018
- 460 Chepfer, H , Brogniez, H , and Noel, V : Diurnal variations of cloud and relative humidity profiles across the tropics, *Sci Rep* , 9, 16045, doi:10.1038/s41598-019-52437-6, 2019



- Collis, R. T. H., Russell, P. B.: Lidar measurement of particles and gases by elastic backscattering and differential absorption, *Laser Monitoring of the Atmosphere, Topics in Applied Physics*, 14, ISBN 978-3-540-07743-5 Springer-Verlag, 71–150, doi:10.1007/3-540-07743-X_18, 1976
- 465 Dawkins, E. C. M., Feofilov, A., Rezac, L., Kutepov, A. A., Janches, D., Höffner, J., X. Chu, X., Lu, X., Mlynarczyk, M. G., and Russell III, J.: Validation of SABER v2 Operational temperature data with ground-based lidars in the mesosphere-lower thermosphere region (75–105 km), *J Geophys Res Atmos*, 123, 9916–9934, doi:10.1029/2018JD028742, 2018
- 470 Donovan, D. P., Marseille, G.-J., de Kloe, J., and Stoffelen, A.: AEOLUS L2 activities at KNMI, *EPJ Web Conf* 237, 01002, doi:10.1051/epjconf/20202370100, 2020
- Feofilov, A. G., Chepfer, H., Noel, V., Guzman, R., Gindre, C., and Chiriaco, M.: Colocated ALADIN/Aeolus and CALIOP/CALIPSO observations for the period of 28/06/2019–31/12/2019, *ResearchGate*, <https://doi.org/10.13140/RG.2.2.11237.12009>, 2021
- Feofilov, A. G. and Petelina, S. V.: Relation between mesospheric ice clouds, temperature, and water vapor determined from 475 Odin/OSIRIS and TIMED/SABER data, *J Geophys Res*, 115, D18305, doi:10.1029/2009JD013619, 2010
- Feofilov, A. G., Kutepov, A. A., She, C.-Y., Smith, A. K., Pesnell, W. D., and Goldberg, R. A.: CO₂(v₂)-O quenching rate coefficient derived from coincidental SABER/TIMED and Fort Collins lidar observations of the mesosphere and lower thermosphere, *Atmos Chem Phys*, 12, 9013–9023, doi:10.5194/acp-12-9013-2012, 2012
- Feofilov, A. G. and Stubenrauch, C. J.: Diurnal variation of high-level clouds from the synergy of AIRS and IASI space-borne 480 infrared sounders, *Atmos Chem Phys*, 19, 13957–13972, doi:10.5194/acp-19-13957-2019, 2019
- Guzman, R., Chepfer, H., Noel, V., Vaillant de Guelis, T., Kay, J. E., Raberanto, P., Cesana, G., Vaughan, M. A., and Winker, D. M.: Direct atmosphere opacity observations from CALIPSO provide new constraints on cloud-radiation interactions, *J Geophys Res Atmos*, 122, 1066–1085, doi:10.1002/2016JD025946, 2017
- Hélière, A., Gelsthorpe, R., Le Hors, L., and Toulemon, Y.: ATLID, the Atmospheric Lidar on board the EarthCARE Satellite, 485 *Proceedings of the ICSSO (International Conference on Space Optics)*, Ajaccio, Corse, France, Oct 9-12, 2012, paper: ICSSO-065, 2012
- Hilton, F., and 42 Coauthors, *Hyperspectral Earth observation from IASI: Five years of accomplishments*, *Bull Am Meteorol Soc*, 93, 347–370, doi:10.1175/BAMS-D-11-00027.1, 2012
- Flamant, P., Cuesta, J., Denneulin, M.-L., Dabas, A., and Hubert, D.: ADM-Aeolus retrieval algorithms for aerosol and cloud 490 products, *Tellus*, 60(2), 273–288, doi:10.1111/j.1600-0870.2007.00287.x, 2008
- Flamant, P. H., Lever, V., Martinet, P., Flament, T., Cuesta, J., Dabas, A., Olivier, M., and Huber, D.: ADM-Aeolus L2A Algorithm Theoretical Baseline Document Particle spin-off products, AE-TN-IPSL-GS-001, V5.5, ESA, available online at <https://earth.esa.int/eogateway/documents/20142/0/Aeolus-L2A-Algorithm-Theoretical-Baseline-Documents/pdf>, 83pp, 2017
- Holl, G., Buehler, S. A., Rydberg, B., and Jiménez, C.: Collocating satellite-based radar and radiometer measurements – 495 methodology and usage examples, *Atmos Meas Tech*, 3, 693–708, doi:10.5194/amt-3-693-2010, 2010



- Kanitz, T., and 24 Coauthors, ESA's Lidar Missions Aeolus and EarthCARE, *EPJ Web Conf* 237, 01006, doi:10.1051/epjconf/202023701006, 2020
- Krawczyk, R., Ghibaudo, J.-B., Labandibar, J.-Y., Willetts, D. V., Vaughan, M., Pearson, G. N., Harris, M. R., Flamant, P. H., Salamitou, P., Dabas, A. M., Charasse, R., Midavaine, T., Royer, M., and Heibel, H.: ALADIN: an atmosphere laser Doppler wind lidar instrument for wind velocity measurements from space, *Proc SPIE* 2581, *Lidar Techniques for Remote Sensing II*, (15 December 1995); doi:10.1117/12.228509, 1995
- Lolli, S., Delaval, A., Loth, C., Garnier, A., and Flamant, P. H.: 0.355-micrometer direct detection wind lidar under testing during a field campaign in consideration of ESA's ADM-Aeolus mission, *Atmos Meas Tech*, 6, 3349–3358, doi:10.5194/amt-6-3349-2013, 2013
- 505 Nam C., Bony, S., Dufresne, J. L., Chepfer, H.: The 'too few, too bright' tropical low-cloud problem in CMIP5 models, *Geophys Res Lett*, 39, 21, doi:10.1029/2012GL053421, 2012
- Noel, V., Chepfer, H., Chiriaco, M., and Yorks, J.: The diurnal cycle of cloud profiles over land and ocean between 51° S and 51° N, seen by the CATS spaceborne lidar from the International Space Station, *Atmos Chem Phys*, 18, 9457–9473, doi:10.5194/acp-18-9457-2018, 2018
- 510 McGill, M. J., Yorks, J. E., Scott, V. S., Kupchok, A. W., and Selmer, P. A.: The Cloud-Aerosol Transport System (CATS): A technology demonstration on the International Space Station, *Proc Spie*, 9612, doi:10.1117/12.2190841, 2015
- Menzel, W. P., Frey, R. A., Borbas, E. E., Baum, B. A., Cureton, G., and Bearson, N.: Reprocessing of HIRS Satellite Measurements from 1980 to 2015: Development towards a consistent decadal cloud record, *J Appl Meteorol Clim*, 55, 2397–2410, doi:10.1175/JAMC-D-16-0129.1, 2016
- 515 Rasch, P., and 41 Coauthors: An Overview of the Atmospheric Component of the Energy Exascale Earth System Model, *J Adv Model Earth Syst*, jame20932, doi:10.1029/2019MS001629, 2019
- Reitebuch, O., and 27 Coauthors: Initial Assessment of the Performance of the First Wind Lidar in Space on Aeolus, *EPJ Web Conf* 237, 01010, doi:10.1051/epjconf/202023701010, 2020
- Reverdy M., Chepfer, H., Donovan, D., Noel, V., Cesana, G., Hoareau, C., Chiriaco, M., Bastin, S.: An EarthCARE/ATLID simulator to evaluate cloud description in climate models, *J Geophys Res Atmos*, 120(21), 11090–11113, doi: 10.1002/2015JD023919, 2015
- Smith, W. L.: The TIROS-N operational vertical sounder, *Bull Am Meteorol Soc*, 60, 1177–1187, 1979
- Snels, M., Colao, F., Cairo, F., Shuli, I., Scoccione, A., De Muro, M., Pitts, M., Poole, L., and Di Liberto, L.: Quasi-coincident observations of polar stratospheric clouds by ground-based lidar and CALIOP at Concordia (Dome C, Antarctica) from 2014 to 2018, *Atmos Chem Phys*, 21, 2165–2178, doi:10.5194/acp-21-2165-2021, 2021
- 525 Stephens, G. L., Hakuba, M. Z., Webb, M. J., Lebsack, M., Yue, Q., Kahn, B. H., Hristova-Veleva, S., Rapp, A. D., Stubenrauch, C. J., Elsaesser, G. S., and Slingo, J.: Regional Intensification of the Tropical Hydro-logical Cycle During ENSO, *Geophys Res Lett*, 45, 4361–4370, doi: 10.1029/2018GL077598, 2018



- Stoffelen, A , Pailleux, J , Källén, E , Vaughan, J M , Isaksen, L , Flamant, P , Wergen, W , Andersson, E , Schyberg, H ,
530 Culoma, A , Meynart, R , Endemann, M , and Ingmann, P : The Atmospheric Dynamics Mission For Global Wind Field
Measurement, *Bull Amer Meteor Soc* , 86, 73–87, doi: 10.1175/BAMS-86-1-73, 2005
- Straume, A G , and 27 Coauthors: ESA's Space-Based Doppler Wind Lidar Mission Aeolus – First Wind and Aerosol Product
Assessment Results, *EPJ Web Conf* 237, 01007, doi:10.1051/epjconf/20202370100, 2020
- Stubenrauch, C J , and 22 Coauthors: Assessment of global cloud datasets from satellites: Project and database initiated by
535 the GEWEX radiation panel, *Bull Am Meteorol Soc* , 94(7), 1031–1049, doi:10.1175/BAMS-D-12-00117.1, 2013
- Stubenrauch, C J , Feofilov, A G , Protospadaki, S E , and Armante, R : Cloud climatologies from the infrared sounders
AIRS and IASI: strengths and applications, *Atmos Chem Phys* , 17, 13625–13644, doi:10.5194/acp-17-13625-
2017, 2017
- Swales, D J , Pincus, R , & Bodas-Salcedo, A : The cloud feedback model intercomparison project observational simulator
540 package: Version 2 Geoscientific Model Development, 11, 77– 81, doi:10.5194/gmd-11-77-2018, 2018
- Vaillant de Guélis T , H Chepfer, Noel, V , Guzman, R , Bonazzola, M , and Winker, D M : Space lidar observations constrain
longwave cloud feedback, *Nature Sci Rep* , 8:16570, doi:10.1038/s41598-018-34943-1, 2018
- Virtanen, T H , Kolmonen, P , Sogacheva, L , Rodriguez, E , Saponaro, G , and de Leeuw, G : Collocation mismatch
uncertainties in satellite aerosol retrieval validation, *Atmos Meas Tech* , 11, 925–938, doi:10.5194/amt-11-925-2018, 2018
- 545 Weiler, F , Kanitz, T , Wernham, D , Rennie, M , Huber, D , Schillinger, M , Saint-Pe, O , Bell, R , Parrinello, T , and
Reitebuch, O : Characterization of dark current signal measurements of the ACCDs used on-board the Aeolus satellite, *Atmos
Meas Tech Discuss* [preprint], doi:10.5194/amt-2020-458, in review, 2020
- Winker, D M , Vaughan, M A , Omar, A H , Hu, Y , Powell, K A , Liu, Z , Hunt, W H , and Young, S A : Overview of the
CALIPSO Mission and CALIOP Data Processing Algorithms, *J Atmos Ocean Tech* , 26, 2310–2323,
550 doi:10.1175/2009JTECHA1281.1, 2009




Instrument	Orbit inclination [deg]	Equator crossing LT [h]	Off-nadir angle [deg]	PRF [Hz]	Native resolution [µs]	L2 resolution resolution [µs]
ALADIN	96.97	6:00 / 18:00	35	50.0	140 (H) x 1000 (V)	87000 (H) x 1000 (V)
CALIOP	98.00	01:30 / 13:30	3	20.1	333 (H) x 60 (V)	333 (H) x 500 (V)

Table 1: Comparison of orbital parameters, viewing geometries, and resolutions of ALADIN and CALIOP instruments

Δtime [h]	Daytime ×1E3	Night-time ×1E3	Total ×1E3	Remarks
< 1	4.3	3.7	8	Narrow polar zone
< 3	13.1	11.2	24.3	Broader polar zone
< 6	91	78	169	All zones covered
< 12	135	116	251	Unequal distribution of Δtime
< 24	176	146	322	Unequal distribution of Δtime

Table 2: Number of collocated cases for Adist < 1° and different Δtime values



560 Figure 1: Observation geometry, averaging, and retrieved parameters for (a-c) ALADIN/Aeolus at its L2A resolution of 87 km and
565 (d) CALIOP/CALIPSO at its native resolution: (a) Attenuated particulate backscatter (APB) at 355 nm; (b) Attenuated molecular
backscatter (AMB) at 355 nm; (c) Attenuated total backscatter (ATB) at 355 nm; (d) Attenuated total backscatter (ATB) at 532 nm.
The scene has been calculated for demonstration purposes using COSP2 simulations with the EAMV1 model data as an input. White
dashed lines stand in (a-c) for ALADIN's observation paths for centers of averaged profiles and in (d) for CALIOP averaged
observation path corresponding to averaged ALADIN on the left and for individual CALIOP profiles on the right (with its 3° off-
nadir viewing angle). ALADIN observes the atmosphere at 35° to the nadir and perpendicular to the flight direction. This inclination
is schematically shown as an inclined line lying in lidar curtain plane whereas the real projection to the same plane should be a
vertical line.

Author: Subject: Sticky Note Date: 17 05 2021 15:06:59
only valid for a cloud free scene IF at all

Author: Subject: Sticky Note Date: 14 06 2021 11:30:37
hard to understand. A clearer description is needed in the caption what the dashed lines mean. It is also not clearly evident while reading the
text. Furthermore, I recommend to use different colours for the dashed line and then clearly describe which dashed line represents what




Figure 2: Geographical distribution of collocated points for (a) $\Delta\text{time} < 1$ h; (b) $\Delta\text{time} < 6$ h; (c) $\Delta\text{time} < 24$ h for $\Delta\text{dist} < 1^\circ$.






Figure 3: Estimating theoretical cloud detection agreement using pseudo-collocated scattering ratio (SR) data calculated using COSP2 lidar simulator coupled with the output of the EAMv1 atmospheric model. For each altitude bin, the agreement is defined as a ratio of number of cases when both CALIOP and ALADIN have detected a cloud to a total number of simulations for a given bin. The cloud amount corresponds to a ratio of number of cases when a reference CALIOP SR value without noise was above the detection threshold to a total number of simulations for a given bin. The normalized cloud detection agreement represents a ratio of the former to the latter.


575




580 Figure 4: Zonal mean comparison for the $\Delta\text{time} < 6\text{h}$, $\Delta\text{dist} < 1^\circ$ collocated nighttime data subset (see Table 2): (a)-(e) CALIOP averages; (f)-(j) ALADIN averages, converted to SR at 532 nm for comparison purposes; (a,f) 90S-60S; (b,g) 60S-30S; (c,h) 30S-30N; (d,i) 30N-60N; (e,j) 60N-90N.



585 **Figure 5: Pseudo-instantaneous comparisons of collocated ALADIN L2A SR profiles and CALIOP SR profiles averaged over 67 km along the track: (a, f, k) 90S-60S; (b, g, l) 60S-30S; (c, h, m) 30S-30N; (d, i, n) 30N-60N; (e, j, o) 60N-90N; (a-e) cases confirming ALADIN's capability to detect high-level clouds; (f-j) cases showing the cases when ALADIN misses a high cloud detected by CALIOP; (k-n) cases explaining the presumably false detection of a low level cloud by ALADIN; (o) a case with a real low cloud detected by both instruments with an extra point near the surface reported by ALADIN.**




590 **Figure 6: Cloud detection agreement: a) both CALIOP and ALADIN have detected a cloud (YES/YES cases); b) neither CALIOP nor ALADIN has detected a cloud (NO/NO cases); c) CALIOP has detected a cloud whereas ALADIN has not detected a cloud (YES/NO cases); d) CALIOP does not detect a cloud whereas ALADIN has detected a cloud (NO/YES cases).**



595 Figure 7: Cloud altitude detection sensitivity represented as a height difference between the CALIOP local peak height and corresponding ALADIN's cloud peak height or maximal SR height found in the ± 3 km vicinity of CALIPSO peak. The subset corresponding to YES_YES selection (Fig. 6a) was used. White dashed isoline corresponds to colored area in Fig. 6a (occurrence frequency of about 5% and higher).


This page contains no comments



600

Figure 8: Temporal evolution of cloud detection agreement for the period of 28/06/2019–31/12/2019. The legend is consistent with that of Fig. 6: a) YES/YES; b) NO/NO; c) YES/NO; d) NO/YES. White vertical dashed lines correspond to the period of Air Motion Vector (AMV) campaign (28/10/2019–10/11/2019), which is characterized by smaller bin sizes and, therefore, larger SNRs for Mie and Rayleigh channels up to the height of 2250m.


This page contains no comments



605

Figure 9: Temporal evolution of normalized cloud detection agreement for the period of 28/06/2019-31/12/2019: a) YES_YES statistics of Fig. 8a normalized by cloud amount; b) the same information presented for 5 heights as linear fits in 2D with error bars. The color scheme for panel (a) is consistent with that of Fig. 8.

This page contains no comments



610

Figure A1: Correlation between individual pairs of CALIOP and ALADIN scattering ratio profiles, for all altitudes. The colors of the bins represent the occurrence frequencies for 0.2×0.07 SR bins, as a function of both CALIOP's SR_{355} and ALADIN's SR_{355} . For each point along the diagonal, a Gaussian was fitted to the data points lying along a perpendicular transect and the central point of the Gaussian is plotted as a red filled circle. The white dashed line represents a linear fit to these points. For comparison, black dashed line shows the fit given by Eq. 1.

615