

A New Machine Learning based Analysis for Improving Satellite Retrieved Atmospheric Composition Data: OMI SO₂ as an Example

Can Li^{1,2}, Joanna Joiner², Fei Liu^{2,3}, Nickolay A. Krotkov², Vitali Fioletov⁴, Chris McLinden⁴

¹Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD 20740, USA

²Atmospheric Chemistry and Dynamics Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

³Goddard Earth Sciences Technology and Research (GESTAR) II, Morgan State University, Baltimore, MD 21251, USA

⁴Environment and Climate Change Canada, Toronto, Ontario, Canada

Correspondence to: Can Li (can.li@nasa.gov)

Abstract. Despite recent progress, satellite retrievals of anthropogenic SO₂ still suffer from relatively low signal-to-noise ratios. In this study, we demonstrate a new machine learning data analysis method to improve the quality of satellite SO₂ products. In the absence of large ground truth datasets for SO₂, we start from SO₂ slant column densities (SCDs) retrieved from the Ozone Monitoring Instrument (OMI) using a data-drive, physically based algorithm and calculate the ratio between the SCD and the root mean square (RMS) of the fitting residuals for each pixel. To build the training data, we select presumably clean pixels with small SCD/RMS ratios (SRRs) and set their target SCDs to zero. For polluted pixels with relatively large SRRs, we set the target to the original retrieved SCDs. We then train neural networks (NNs) to reproduce the target SCDs using predictors including SRRs for individual pixels, solar zenith, viewing zenith and phase angles, scene reflectivity and O₃ column amounts, as well as the monthly mean SRRs. For data analysis, we employ two NNs: 1) one trained daily to produce analysed SO₂ SCDs for polluted pixels each day and 2) the other trained once every month to produce analysed SCDs for less polluted pixels for the entire month. Test results for 2005 show that our method can significantly reduce noise and artifacts over background regions. Over polluted areas, the monthly mean NN analysed and original SCDs generally agree to within ±15%, indicating that our method can retain SO₂ signals in the original retrievals except for large volcanic eruptions. This is further confirmed by running both the NN analysed and the original SCDs through a top-down emission algorithm to estimate the annual SO₂ emissions for ~500 anthropogenic sources, with the two datasets yielding similar results. We also explore two alternative approaches to the NN-based analysis method. In one, we employ a simple linear interpolation model to analyse the original SCD retrievals. In the other, we develop a PCA-NN algorithm that uses OMI measured radiances, transformed and dimension-reduced with a principal component analysis (PCA) technique, as inputs to NNs for SO₂ SCD retrievals. While the linear model and the PCA-NN algorithm can reduce retrieval noise, they both underestimate SO₂ over polluted areas. Overall, the results presented here demonstrate that our new data analysis method can significantly improve the quality of existing OMI SO₂ retrievals. The method can potentially be adapted for other sensors and/or species and enhance the value of satellite data in air quality research and applications.

1 Introduction

Sulfur dioxide (SO₂) and its oxidation product in the atmosphere, sulfate aerosols, have significant impacts on air quality, visibility, ecosystems, and the weather and climate. For over two decades, spaceborne hyperspectral ultraviolet (UV) instruments (e.g., Eisinger and Burrows, 1998; Krotkov et al., 2006; Nowlan et al., 2011; Theys et al., 2017) have been providing global observations of anthropogenic SO₂ sources such as coal-fired power plants, metal smelters, and the oil and gas industry (e.g., Fioletov et al., 2016; McLinden et al., 2016; Zhang et al., 2019). More recently, the quality of satellite SO₂ data products has substantially improved thanks to the development of data driven retrieval techniques. In particular, the principal component analysis (PCA) based algorithm (Li et al., 2013; 2017a; 2020) and the covariance-based retrieval algorithm (COBRA, Theys et al., 2021) have helped to reduce the noise and artifacts of SO₂ retrievals from several sensors including the Ozone Monitoring Instrument (OMI), Ozone Mapping and Profiler Suite (OMPS), and TROPOspheric Monitoring Instrument (TROPOMI), enabling the detection and quantification of relatively small point sources (e.g., Fioletov et al., 2015, Theys et al., 2021).

Despite these progresses, satellite remote sensing of anthropogenic SO₂ remains challenging. The signal of anthropogenic SO₂ is relatively weak as compared with volcanic sources. With an atmospheric lifetime of ~1 day (e.g., Lee et al., 2011), SO₂ emitted from human activities is also more concentrated in the boundary layer, where the sensitivity of satellite instruments is limited by the low surface albedo, strong Rayleigh scattering, and interferences from O₃ absorption in the UV (e.g., Krotkov et al., 2008). As a result, the noise in satellite SO₂ retrievals is relatively large even for data driven algorithms. For example, the standard deviation (1- σ noise) of OMI PCA SO₂ slant column densities (SCDs) over the remote Pacific is ~0.2-0.3 DU (Dobson Unit, 1 DU = 2.69×10^{16} molecules/cm², Li et al., 2020), far greater than the typical SCDs retrieved outside of the most polluted areas (e.g., Persian Gulf, eastern China, and Norilsk, Russia). The retrieval noise can be reduced by spatially and temporally averaging the data (Krotkov et al., 2008). However, relatively small but noticeable artifacts still exist in monthly or annual mean OMI SO₂ (e.g., negative values over arid and semi-arid areas), indicating systematic biases that cannot be averaged out. While there was little drift in the mean OMI SO₂ SCDs over remote regions from 2005 to 2019, the retrieval noise grew by ~10% during the same period (Li et al., 2020), presumably due to instrument degradation. With the recent large decreases in SO₂ emissions and signals in many regions (e.g., Krotkov et al., 2016; Li et al., 2017b), the increase in retrieval noise makes the analyses and applications of satellite SO₂ data even more challenging, especially for long-term monitoring. It is thus imperative to further enhance the quality of satellite SO₂ data products.

In recent years, machine learning has emerged as a powerful tool in satellite remote sensing of atmospheric composition. Capable of incorporating large, diverse datasets and modelling complex, nonlinear functions, techniques such as neural networks (NN) and random forests (RF) have been utilized to solve various problems. For instance, a number of studies trained NN or RF models to infer surface concentrations of pollutants from satellite observations, including particulate matter (e.g., Huang et al., 2021; Liu et al., 2019; Zheng et al., 2021), NO₂ (e.g., Chan et al., 2021), and SO₂ (e.g., Zhang et al. 2022). NNs have also been used to speed up radiative transfer calculations (e.g., Castellanos and da Silva, 2019; Nada et al., 2019)

Moved (insertion) [1]

Deleted: These issues pos

Deleted: e challenges for

Moved up [1]: the recent large decreases in SO₂ emissions in many regions (e.g., Krotkov et al., 2016; Li et al., 2017b)

Deleted: in light of

and to retrieve O₃ profiles (e.g., Muller et al., 2003; Xu et al., 2017) and total columns (Muller et al., 2004), isoprene amounts (Wells et al., 2020), and aerosol layer height (Chimot et al., 2017). For SO₂, De Santis et al. (2021) demonstrated a NN retrieval algorithm using the operational TROPOMI product for training in their case study of Mt. Etna. Piscini et al. (2014) attempted NN-based SO₂ and volcanic ash retrievals using thermal infrared measurements from MODIS (the Moderate Resolution Imaging Spectroradiometer). Hedelt et al. (2019) also developed near-real-time volcanic SO₂ height retrievals using the Full-Physics Inverse Learning Machine (FP-ILM) method, a technique later adapted for OMI by Fedkin et al. (2021). While these studies have demonstrated the potential of machine learning for SO₂ retrievals, they all focus on volcanic SO₂. To our knowledge, so far there have been no published studies demonstrating the use of machine learning techniques for anthropogenic SO₂ retrievals.

A major obstacle in developing machine learning retrieval algorithms for anthropogenic SO₂ is the lack of high-quality, ground-truth training data. As mentioned above, existing satellite SO₂ products provide global coverage, but the signal-to-noise ratios are typically small for anthropogenic sources. Ground air quality monitors generally offer good data quality and long-term measurements, but they do not represent the entire atmospheric column. Aircraft measurements and surface based remote sensing instruments (e.g., MAX-DOAS) have been used to evaluate satellite retrievals, but they are quite sparse. The FP-ILM method circumvents this data availability issue by using a large set of model-simulated synthetic radiance spectra in training. However, the models may not fully represent the various geophysical processes and instrument characteristics that affect satellite measurements. This can lead to substantial errors, and FP-ILM retrievals [of volcanic SO₂ height](#) are currently limited to satellite pixels with sizable SO₂ amounts (> 20 DU).

Here, we introduce a new data analysis method to further improve the quality of satellite retrieved SO₂. In the absence of sufficient ground truth data, we compile our training data by analysing existing OMI SO₂ SCD retrievals and the associated fitting errors, assuming that retrievals with greater SCDs and smaller fitting errors can be trusted more than those with smaller SCDs and larger errors. This allows us to train NNs to reduce noise and artifacts in the original retrievals, meanwhile retaining SO₂ signals over major emission source areas. The rest of the paper is organized as follows: Section 2 describes our methodology and setups for NN training. Section 3 provides some example results. This is followed by a more detailed discussion on the NN analysed SCDs in Section 4 and conclusions in Section 5.

2 Data and methodology

The flowchart in Fig. 1a presents an overview of our data analysis method. We start from existing OMI PCA SO₂ retrievals (Section 2.1) and calculate the ratio between the SCD and the root mean square (RMS) of the fitting residuals (SCD/RMS ratio, SRR) for each pixel, as well as the statistics of the SRRs for the entire month. This provides input to a data classification scheme (Fig. 1b, Section 2.2) that assigns OMI pixels from each day into different groups ("clean", "polluted", "in-between" and "high-SRR"). The pixels within each group are then either processed with one of the two neural networks (pre-trained NN1 for clean and in-between pixels, daily trained NN2 for polluted pixels, Fig. 1c, Section 2.3) or retain their original

Formatted: Subscript

retrieved SCDs (for high-SRR pixels). In the end, the OMI pixels from different groups are merged into the final analysed SCD dataset.

2.1 Analysis of OMI SO₂ data

To demonstrate our methodology, we use data from OMI, a Dutch/Finnish UV/Visible spectrometer that has been flying on the National Aeronautics and Space Administration (NASA)'s Aura spacecraft in a Sun synchronous orbit since 2004 (Levelt et al., 2018). OMI measures backscattered solar radiation between 270 and 500 nm in the local afternoon (local equator crossing time: ~13:45) at a relatively high spatial ($13 \times 24 \text{ km}^2$ at nadir) and spectral (~0.5 nm) resolution. We focus on the year 2005, when all cross-track positions (rows) of OMI's 2-dimensional detectors were taking nominal measurements, providing daily global coverage.

For SO₂ data, we use SCDs retrieved from NASA version 2 OMI standard SO₂ algorithm based on the PCA spectral fitting technique. The algorithm has been described in detail elsewhere (Li et al., 2020) and is only briefly introduced here. The algorithm uses OMI-measured Sun-normalized Earthshine radiances within the spectral range of 310.5-340 nm and processes each row of individual OMI orbits separately. The ~1600 OMI pixels from a given row in a given orbit are first filtered to exclude those with large solar zenith angles ($\text{SZA} > 75^\circ$) or potentially strong SO₂ signals (e.g., volcanic plumes, [by examining the ozone residuals at 313/314 and 314/315 nm wavelength pairs, see Li et al., 2017a for details](#)). Next, the spectra of the remaining pixels are analysed utilizing a PCA technique to extract spectral features (principal components, PCs). The leading PCs that account for the most spectral variances are typically associated with geophysical (e.g., O₃ absorption and rotational Raman scattering, RRS) or instrumental (e.g., dark current, wavelength shift) factors that interfere with SO₂ retrievals. For each pixel, up to 30 leading PCs, along with the SO₂ cross sections, are fit to the measured radiances to estimate the SO₂ SCD while minimizing the interferences. This multi-step (data filtering, PCA analysis, and spectral fitting) procedure is iterated a few times. To avoid collinearity in fitting, the PCs are also examined to exclude those potentially containing SO₂ spectral signatures. For this study, the standard algorithm has been modified to use the new collection 4 OMI level 1B (L1B) radiance and irradiance data, instead of the collection 3 data for the current standard OMI SO₂ product. No obvious differences were found between the SCDs retrieved from the two collections. In addition, the RMS of the fitting residuals (i.e., the differences between the measured and the fit normalized radiance spectra) for each pixel has been added to the output.

In order to compare the SO₂ signal vs. the fitting error, we calculate the SCD/RMS ratio (SRR) for each pixel. The pixel-level SRRs are also gridded into monthly mean (SRR_m) at $0.25^\circ \times 0.25^\circ$ resolution (Fig. 2). At middle and low latitudes, the overall spatial distribution of SRR_m (Fig. 2b) is quite similar to that of the monthly mean SCDs (Fig. 2a). On the other hand, the bias in SRR_m is smaller at high latitudes due to generally greater fitting errors at larger SZAs, allowing us to better distinguish polluted areas from background regions. In the following steps (Sections 2.2 and 2.3), SRR_m is utilized as an indicator of the likelihood of an OMI SO₂ retrieval over a certain area to represent a positive SO₂ value. For each day of the month, we also calculate the mean and standard deviation of SRRs for 3° latitude bands, using all pixels within each band after

Deleted:)

removing outliers (SRRs outside of $\pm 5\sigma$ from the mean). The monthly medians of the daily mean (\overline{SRR}) and standard deviation
135 (σ_{SRR}) are then taken from each latitude band as inputs to the OMI pixel classification scheme (Section 2.2)

2.2 Classification of OMI pixels

The purpose of the pixel classification scheme (Fig. 1b) is to compile a training dataset by selecting pixels in two categories:
1) the first for clean pixels in which the retrieved SCDs are relatively small while the fitting errors are relatively large (i.e.,
negative or small positive SRRs) so that they can be considered largely SO₂-free and 2) the second in which the retrieved SCDs
140 are large while the fitting errors are relatively small (i.e., large SRRs). In this category for polluted pixels, the retrieved SCDs
are assumed to be close to the truth. There are two additional categories. The third is for pixels that fall in-between the clean
and polluted categories. For these pixels, an unambiguous classification cannot be made and they are excluded from the training
dataset. The fourth category (high SRR) is for pixels that have very large SRRs (> 300). Such pixels are few but are also
excluded from the training, as they tend to have a disproportionately large influence on the trained NNs.

145 In addition to the SRRs of individual OMI pixels, the classification scheme also takes into account the location
(latitude/longitude) of the pixels, as well as the general performance of the PCA algorithm for the latitude bands in which they
are located. A pixel with a specific SCD/RMS ratio of SRR_i is considered to be polluted, if:

$$SRR_i > \overline{SRR} + a_1 \sigma_{SRR} . \quad (1)$$

The pixel would be considered to be clean, if:

150 $SRR_i < \overline{SRR} + a_2 \sigma_{SRR} , \quad (2)$

where \overline{SRR} and σ_{SRR} are the monthly medians of the daily mean and standard deviation of SRRs for the corresponding latitude
band, respectively. a_1 and a_2 are scaling factors (see Fig. 1b for values) that have been adjusted through trial and error in order
to 1) minimize the artifacts in NN analysed SCDs over background areas and 2) maximize the retained original SO₂ signals
over polluted areas. Both factors depend on the location of the pixels and the monthly mean SRRs (SRR_m). As shown in Fig.
155 1b, a_1 and a_2 are large, if the pixel is located in an area with a small SRR_m (< 3). In this case, the area is generally unpolluted
and the likelihood of a pixel containing a positive SO₂ value is low. Thus, more pixels are classified as clean. On the other
hand, for polluted areas with large SRR_m (> 5), both a_1 and a_2 are kept small so that more pixels would be classified as polluted.
For areas that are moderately polluted (i.e., $3 < SRR_m < 5$), a_1 and a_2 are linearly interpolated based on the SRR_m . One may
also notice that a_1 and a_2 are smaller for low (30°S-30°N) and middle (30°S-60°S and 30°N-60°N) latitudes than for high
160 latitudes. This helps to reduce the positive bias in the original SCDs near the northern edge of the domain (Fig. 2a). [We also
tested a simple classification scheme with constant \$a_1\$ and \$a_2\$ everywhere, and found that it produces relatively large positive
biases over high latitudes and negative biases over low latitude source areas, as compared with the more complicated scheme
described above.](#) It should also be pointed out that the areas affected by the south Atlantic anomaly (SAA) are not subject to
classification and excluded from the training dataset.

165 [Using the classification scheme, one can also develop a simple method to reduce retrieval artifacts, by assuming that
clean pixels should have zero SCDs while polluted pixels should retain their original SCDs and by estimating SCDs for pixels](#)

that fall in-between through a linear interpolation (between zero and the original SCDs). As will be demonstrated in Section 4.4, such an approach produces negative biases over polluted areas. It is thus advantageous to employ the more complex NN-based method for the present study.

170 2.3 Training of neural networks

For training data, we use the OMI pixels identified as either clean or polluted by the classification scheme. For a typical day, approximately ~800000 out of ~1 million OMI pixels are classified as clean, and ~10000 (~1%) as polluted. Given the scarcity of ground truth SO₂ data, we set the training target (SCD_{target}) to zero for the clean pixels and to the original SCDs for the polluted pixels. Note that unlike the PCA spectral fitting algorithm, data from all 60 rows are pooled together in the training

175 [so that a relatively large sample of polluted pixels is available](#). We also include several candidate predictors in the training data, including SCD/RMS ratios for the individual pixels (SRR_i), the monthly mean SCD/RMS ratios (SRR_m) where the pixels are located, the cosines of solar zenith angles (SZA, θ_0), viewing zenith angles (θ) and phase angles (ϕ), the O₃ column amounts from the OMI total O₃ product (OMTO3, Bhartia, 2005), and the scene reflectivity (R) at 354 nm from the OMI Raman cloud product (OMCLDRR, Joiner and Vasilkov, 2006). The function of a neural network (NN) is then to use the input predictors

$$180 \text{ or features to predict the output } \text{SCD}_{\text{target}}: \\ \text{SCD}_{\text{target}} = f_{\text{NN}}(\text{SRR}_i, \text{SRR}_m, \theta_0, \theta, \phi, R, O_3). \quad (3)$$

To optimize the set of predictors, we carried out a number of tests using different combinations (See Table 1 for example results). Among the predictors, SRR_i is well correlated with SCD_{target} and has the largest impact on the performance of the NNs. Indeed, the NN without SRR_i produces the lowest correlation coefficient (r) and the largest root mean square error (RMSE) between the analysed SCDs (SCD_{NN}) and SCD_{target} (Table 1). SRR_m provides geospatial context for the NNs so that 185 higher SCDs tend to be assigned to polluted areas. In the particular example in Table 1, a simple NN using just SRR_i and SRR_m as predictors produces SCD_{NN} that agrees reasonably well with SCD_{target} ($r = 0.958$, RMSE = 0.0517 DU). The angles, O₃ column amounts, and scene reflectivity all affect the signal-to-noise ratio of OMI measurements and the quality of SO₂ retrievals (Li et al., 2020). Adding them as predictors generally leads to small but noticeable improvements in the performance 190 of the NNs (Table 1). While the NN with all seven predictors has slightly worse performance than the NN without SRR_m for this case, including SRR_m as a predictor helps to retain signals over SO₂ source areas. We also tested additional predictors (e.g., the terrain pressure and the scene pressure) and found no discernible improvements in the overall performance of the NNs. Hereafter we use all seven predictors as specified in Eq. 3 in the NNs.

The architecture of the NNs in this study (Fig. 1c) is similar to that employed by Joiner et al. (2021, 2022) for 195 reconstruction of RGB images from hyperspectral radiances. [A similar architecture has also been used to capture changes in gross primary production \(GPP\) from satellite reflectance data \(Joiner and Yoshida, 2020\)](#). Briefly, the artificial feedforward NNs are implemented in IDL (Interactive Data Language) and contain two hidden layers, each with 14 nodes (twice the number of predictors), and an output layer with one node. [Experiments using more \(up to 30\) nodes in each hidden layer yield little difference in the performance of the NNs](#). The activation functions are a soft sign for the first hidden layer, a logistic (sigmoid)

Deleted:

for the second hidden layer, and a bent identity for the output layer. [If we replace the activation functions in both hidden layers with ReLU \(Rectified Linear Unit\), the NNs converge faster in training but increase the SCDs over background areas by ~0.01-0.02 DU \(Fig. S1\).](#) An adaptive moment estimation (Adam) optimizer (Kingma and Ba, 2014) with a learning rate of 0.1 is used to minimize the error. Inputs and outputs are normalized so that they each have a mean of zero and a unit standard deviation.

For each month, we train a neural network (NN1, Fig. 1) utilizing data from 5 days (the 5th, 10th, 15th, 20th and 25th days of the month). Half of the clean and polluted pixels are used in the training and the rest for evaluation. We notice that NN1 well reproduces SCD_{target} for clean pixels and also for polluted pixels that have SCDs up to ~4-5 DU, but it produces a low bias for larger SCDs. This is likely due to the imbalance between the clean and polluted categories in the training data. To mitigate this issue, we use the pre-trained NN1 only for clean and in-between pixels (Fig. 1a) and a separate neural network (NN2) for polluted pixels from each day (Fig. 1a). NN2 has the same architecture as NN1 but is trained daily with half of the polluted pixels. Alternatively, we can also train NN2 using data from multiple days and apply the pre-trained multi-day model to the entire month. As compared with the daily trained NN2, SCD_{NN} produced by the multi-day model is similar but slightly lower over some polluted areas (e.g., eastern China). To maximize the retained SO_2 signals over those regions, we use daily trained NN2 in the present study.

In the final step (Fig. 1a), the SCD_{NN} outputs from NN1 and NN2 are merged with the original SCDs for high-SRR pixels to produce the final NN analysed SCDs.

3 Results

3.1 Daily comparisons of SO_2 SCDs

In Fig. 3, we compare the NN analysed SO_2 SCDs (SCD_{NN}) and the target SCDs (SCD_{target}) from the 16th of January, April, July and October 2005, for independent pixels that are not part of the training. There is generally good agreement between SCD_{NN} and SCD_{target} , with $r > 0.93$ and RMSE at ~0.02-0.03 DU for all four days. The vast majority of clean pixels as identified by the classification scheme have SCD_{NN} between -0.1 and 0.1 DU, indicating substantial reduction in the retrieval noise as compared with the original retrievals ($1-\sigma$ noise of ~0.2-0.3 DU), although a small fraction of the clean pixels still have SCD_{NN} as large as ± 0.5 DU. The slopes from the linear regression analysis are between 0.95 and 0.98, suggesting slight underestimates in SCD_{NN} . There is also some scatter for the polluted pixels particularly at higher SCDs (> 2 DU). The number of pixels having large SCD_{target} are relatively small and this limit in the training data may affect the performance of NNs under high SCD conditions (such as for volcanic plumes). We repeated the analysis for the whole year and found similar results for most days. On average, the correlation coefficient from the daily comparisons is 0.948 ± 0.0309 (hereafter results are shown as mean \pm standard deviation), the RMSE is 0.0343 ± 0.0194 DU, while the slope is 0.966 ± 0.0409 . There are four days with RMSE > 0.1 DU (April 6, June 11, July 13, and August 14). All four have relatively large errors over areas affected by volcanic plumes,

again suggesting that the NN performance may deteriorate at high SCDs. Overall, the comparisons here point to quite good performance of the NNs in reproducing the target SCDs.

As compared with the original SCDs, the NN analysed SCDs have much reduced noise and artifacts over background areas and largely retain SO₂ signals over polluted regions. This is evident from Fig. 4 which shows the original SO₂ SCDs, the NN analysed SCDs, their differences, and the mean SCDs for different latitude bands over generally clean areas (monthly mean SRR < 3) for April 16, 2005 as an example. As can be seen from the figure, the NN analysed SCDs show little variation with latitudes as compared with the original PCA retrievals (Fig. 4d). The differences between the two (Fig. 4c) are similar to the original SCDs (Fig. 4a) over most background areas, as ~80% of the pixels are identified as clean and have SCD_{NN} within ±0.1 DU. The differences are quite small over polluted regions (e.g., eastern China, Sichuan Basin, Norilsk), as pixels over those areas tend to be classified as polluted and have SCD_{NN} close to their original retrievals. It is worth mentioning that even though the SAA affected areas are excluded from training, the analysed SCDs over those areas still show smaller noise than the original ones. One potential reason is that retrievals over the SAA areas tend to have relatively large RMS, and the use of SRRs partially cancels out the relatively noisy SCDs.

3.2 Comparisons of monthly SO₂ SCDs

The monthly maps in Fig. 5 for March 2005 show consistent results with the daily comparisons in Section 3.1. While the monthly mean SCDs from the original PCA retrievals (Fig. 5a) are close to zero for most background areas, biases are evident over certain regions. For example, there are patches of negative SCDs (approximately -0.1 DU) at ~40-60°N and over the oceans near the equator. Another noticeable feature is the negative bias over the relatively bright arid and semi-arid land surfaces such as the Sahara desert, the Arabian peninsula, and the Taklimakan and Gobi deserts. It is possible that the retained PCs (derived from hundreds of pixels from each OMI row) do not fully capture certain interfering factors for those areas. The exact reasons for these artifacts are unknown and beyond the scope of the present study. In any case, they are largely removed through our NN-based analysis (Fig. 5b). Meanwhile, there is no obvious difference between the original and analysed SCDs over major SO₂ source areas (Fig. 5c), evidence that the NNs have learned to preserve the SO₂ signals over those areas.

One may notice that outside of the source regions, the difference map in Fig. 5c is not identical to the original SCD map in Fig. 5a. For example, the differences are slightly more negative than the original SCDs over parts of Canada, Mongolia, and Russia. Most pixels have SCD_{NN} near zero, but some pixels with noisy, positive original SCDs could be misclassified as polluted, resulting in a small positive bias in SCD_{NN} for these areas. This is also noticeable in Fig. 5d that shows the mean original and NN analysed SCDs within 1° latitude bands over clean areas. The NN SCDs have generally less structure, indicating reduced artifacts, but a positive bias of ~0.02 DU can be found north of 60°N. Mean SCD maps for other months (January, April, July, October 2005, see Fig. S2 in the supplemental information) show quite similar results. For areas/periods strongly influenced by relatively large volcanic eruptions (e.g., Sierra Negra (Galapagos Islands) eruption in October 2005), the NNs have difficulty completely reproducing the strong SO₂ signals. This again points to the slightly deteriorated performance of NNs under high SO₂ conditions, as already discussed.

Deleted: and

Deleted:

Deleted: t

Deleted: c

Deleted: l

270 A close-up look at the NN-analysed SCDs and their differences from the original SCDs over eastern China is given
in Fig. 6. For polluted areas (analysed SCDs > 0.15 DU), the relative differences are typically within $\pm 20\%$, with a mean of
4% (with the original SCDs being greater). For background areas, the relative differences are close to $\pm 100\%$ as expected for
clean pixels. Comparisons for other major anthropogenic source areas including India, the Middle East, South Africa, the
275 eastern U.S., and Norilsk, Russia yield similar results (see Figs. S3-S7 in the supplemental information). The mean relative
differences for polluted areas in these regions are all within $\pm 15\%$, ranging between -11% for the eastern U.S. and 14% for the
Middle East. In comparison, the relative differences for areas affected by large volcanic plumes are greater, for example
reaching 20% on average over part of the southeast Pacific during the October 2005 Sierra Negra eruption (see Fig. S8 in the
supplemental information).

Deleted: 2

Deleted: 6

Deleted: 7

4 Discussion

280 4.1 Original and analysed SO₂ SCDs as a function of SRRs

The results presented in Section 3 demonstrate that our NN-based analysis can reduce noise and artifacts for clean pixels,
meanwhile largely retaining the original SCDs for polluted pixels. However, some key questions remain unanswered. Namely,
given the somewhat subjective criteria used in the pixel classification scheme (Section 2.2) to build the training data, do we
risk removing real SO₂ signals as noise (i.e., over-correction) and/or keeping noise/artifacts as signals (i.e., under-correction)?
285 Another related question is: how do the NNs treat pixels that are not in the training data (i.e., the pixels that fall in-between
the clean and polluted categories)? To shed light on these issues, we calculate the monthly mean SO₂ SCDs as a function of
pixel-level SCD/RMS ratios (SRR_{*i*}) from the original retrievals (Panel a of Figs. S9-S13 in the supplemental information) and
the analysed data (Panel b of Figs. S9-S13 in the supplemental information), as well as their differences (Fig. 7, left) for March
2005. In addition, we also calculate the mean original and NN analysed SCDs as a function of latitude for different ranges of
290 SRR_{*i*} (Fig. 7, right).

Deleted: 7

Deleted: , left

Deleted: Fig. 7, center

Deleted: right

For pixels having $SRR_i < \overline{SRR}$ (see Section 2.1 for definitions of \overline{SRR} and σ_{SRR}), the original SCD map (Fig. S9a)
shows no obvious hotspots even over the major SO₂ source areas. All such pixels would be classified as clean and indeed the
mean NN analysed SCDs (Fig. S9b) from these pixels are zero everywhere. The mean analysed SCDs (Fig. 7b) are also near
295 zero at all latitudes.

Deleted: 7

Deleted: 7

The next group of pixels have $\overline{SRR} < SRR_i < \overline{SRR} + \sigma_{SRR}$ (Fig. 7, second row). Most pixels in this group, except for
those near large SO₂ sources at low latitudes (30°S-30°N), would also be classified as clean. Similar to the first group, there
are no obvious SO₂ hotspots in the original SCD map (Fig. S10a). The analysed SCDs (Fig. S10b) are similarly near zero
almost everywhere, with notable exceptions over some degassing volcanoes (Anatahan, Nyiragongo, and Vanuatu) and heavily
polluted areas (Sichuan Basin and Norilsk). The case of Norilsk is particularly interesting. Given the thresholds for high
300 latitudes (Fig. 1b), all pixels in this group over Norilsk would be classified as clean, but the NNs seem to be able to override

Deleted: 7

Deleted: d

Deleted: 7

Deleted: e

the classification based on factors other than SRRs. [The mean analysed SCDs are around zero for all latitude bands, smaller than the original SCDs \(Fig. 7d\).](#)

For the group of pixels having intermediate SRRs ($\overline{SRR} + \sigma_{SRR} < SRR_i < \overline{SRR} + 2\sigma_{SRR}$, Fig. 7, third row), the original SCD map (Fig. [S11a](#)) contains enhanced SO₂ signals over source areas but also artifacts over background regions. The pixels in this group would be classified as clean, polluted, or in-between depending on their SRR_i and locations. In general, the NNs are able to largely eliminate the artifacts and retain signals over SO₂ source areas for this group (Fig. [7c](#)), although there are remaining small positive biases [both](#) near the northern edge of the domain and at [lower latitudes \(e.g., around 30°S\) as shown in Fig. 7f](#).

For the following group ($\overline{SRR} + 2\sigma_{SRR} < SRR_i < \overline{SRR} + 3\sigma_{SRR}$, Fig. 7, fourth row), almost all pixels would have a classification of either polluted or in-between. NNs reduce the retrieval artifacts in this group particularly at middle to high latitudes (Figs. [7g and 7h](#)). The relatively small changes at low latitudes can probably be attributed to the more relaxed thresholds for pixels to be classified as polluted and in-between (Section 2.2). Using more stringent thresholds may further reduce the artifacts in the tropics, but this may also lead to low bias over pollution sources ([See Fig. 8b for example](#)).

For the final group ($SRR_i > \overline{SRR} + 3\sigma_{SRR}$, Fig. 7, fifth row), almost all pixels are identified as polluted. As a result, the differences between the original and the analysed SCDs are quite small except over the SAA affected areas ([Figs. 7i and 7j at around 30°S](#)) where the pixels are not part of the training data and the noise is reduced by the NNs. Overall, it is encouraging that the NN analysed SCDs show improvements over the original ones for all ranges of the SRRs.

[4.2 Sensitivity of NN analysed SCDs to the pixel classification scheme](#)

[We further test the sensitivity of NN analysed SCDs to the settings of the pixel classification scheme, by altering the a₁ and a₂ parameters \(Eq. 1 and 2\). In one experiment, we scale both parameters by 90% \(i.e., reduced by 10% from the baseline values as specified in Fig. 1b\). This leads to more pixels being classified as polluted and greater monthly mean SCDs \(Fig. 8a\). The increase in the SCDs is ~0.01-0.02 DU on average over relatively clean areas \(Fig. 8c\) and slightly larger over some source areas \(e.g., eastern China\) but typically less than 0.1 DU. In another experiment, both a₁ and a₂ are scaled by 110% \(i.e., increased by 10% from the baseline values\), resulting in SCD reductions of ~0.01 DU over clean areas \(Fig. 8c\). Some source areas \(e.g., Norilsk\) show slightly more reductions \(Fig. 8b\) that are still typically less than 0.1 DU. Overall, the tests here point to moderate sensitivity of the NN-based analysis to the settings of the pixel classification scheme. An overly stringent scheme may lead to over-correction and low biases over source areas, whereas an overly relaxed scheme may result in positive biases. For our particular study, any over- or under-correction appears to be minor for major source areas, given the relatively small differences between the analysed and the original SCDs \(see Sect. 3.2\). But if one is to apply the technique to other datasets \(e.g., different instruments and/or species\), the pixel classification scheme will need to be tested and optimized. For long-term analysis of a dataset from a single instrument \(e.g., OMI SO₂ for the entire mission\), the scheme will need to be verified using data from different years, although we envision that a constant set of a₁ and a₂ parameters over time will probably](#)

Deleted: 7g

Deleted: 7h

Deleted: and Fig. 7i

Deleted: 2

Deleted: k

Deleted: and Fig. 7l

Deleted: ,

be more suitable to avoid artificial trends introduced by time-dependent parameters. For future studies, we plan to develop a more systematic way for pixel classification, for example, by using more objective metrics.

355 4.3 SO₂ emission estimates using the original and NN analysed SCDs

Deleted: 2

Another test involves running both the original and NN analysed SCDs through a top-down emission estimation algorithm to derive annual SO₂ emissions from large point sources. Here we focus on anthropogenic sources, given the low bias in the NN analysed SCDs for large volcanic plumes. We infer SO₂ emissions by fitting oversampled and smoothed OMI vertical column densities (VCDs) to a 3-parameter (i.e., total mass, lifetime and plume spread) function of horizontal coordinates and wind speeds (Fioletov et al., 2015). To convert SCDs to VCDs, we use the same air mass factors (AMFs, $VCD = SCD/AMF$) as in Fioletov et al. (2016). For wind fields, we use the average winds between the surface and ~1 km from GEOS-5 Forward Processing for Instrument Teams (FP-IT) assimilated products that have been co-located with OMI (OMUFPITMET; available at https://disc.gsfc.nasa.gov/datasets/OMUFPITMET_003/summary). The OMI pixels are then rotated around known source locations according to wind directions such that all observations are aligned in the upwind-downwind direction (Fioletov et al., 2015). Following Fioletov et al. (2016), we prescribe the SO₂ lifetime (6 h) and the parameter describing the spread of the emitted plume (20 km) to obtain more robust fitting results. Only one parameter, the total SO₂ mass, is estimated from the fit. We further derive SO₂ emissions by dividing the fitted total SO₂ mass by the prescribed lifetime. For fitting uncertainty, we calculate the one standard deviation error in the fitted parameter by taking the square root of the diagonal elements of the covariance matrix of the parameter.

370 As shown in Fig. 9a, the two sets of emission estimates agree quite well ($r > 0.99$, slope > 0.96), suggesting the NN analysis has largely preserved SO₂ signals in the original retrievals. In general, the estimated emissions using the NN analysed SCDs are slightly smaller than those based on the original retrievals, particularly for relatively small sources (< 20 kt, 10^3 tonnes, per year). While on the surface this may suggest loss of some real SO₂ signals in our analysis for relatively small sources, the emission uncertainties (Fig. 9b) for those sources also become much smaller when using the NN analysed data.

Deleted: 8

375 This leads to greater emission/uncertainty ratios (Fig. 9c) for those sources, implying that the reduced noise/artifacts in the analysed data may facilitate SO₂ source detection and quantification. We note that the results here should be interpreted with caution, given that OMI sensitivity to sources < 30 kt/year is quite limited (Fioletov et al., 2015).

Deleted: 8

Deleted: 8

4.4 Can a simple linear interpolation model reproduce NN analysed SCDs?

Deleted: 3

380 Given the seemingly simple assumptions made about the clean and polluted pixels during the training process (Section 2.3), one may also ask whether there is any advantage to using the NN-based data analysis approach. To test this, we apply the same pixel classification scheme as described in Section 2.2 and build a simple model by assigning zero SCDs to the clean pixels, the original SCDs to the polluted pixels, and by linearly interpolating between zero and the original SCDs for pixels that fall in-between (based on the SRRs for those pixels and the corresponding thresholds as defined in Eq. 1 and Eq. 2).

390 The mean SCDs for March 2005 (Fig. 10a), produced with this simple linear interpolation model, appear to be quite
similar to those produced with the NN-based analysis (Fig. 5b). This is not surprising since the majority of pixels are classified
as clean, and NN analysed SCDs for those pixels are also close to zero. The plot of mean SCDs as a function of latitude (Fig.
395 10c) also indicates overall comparable results for relatively clean areas between the two methods, although the linear model
has more obvious step changes at 30°N and 30°S probably related to the pixel classification scheme. Over pollution source
areas (e.g., eastern China), on the other hand, the linear model has a substantial negative bias as compared with the NN-based
400 approach (see the SCD difference map in Fig. 10b). Additionally, the noise is also larger over the SAA areas for the linear
model. This comparison demonstrates some advantages in the NN-based approach, particularly for preserving SO₂ signals
over source areas. It should be mentioned that the simple linear model tested here can be potentially improved by including
more predictors such as those used in the NNs (e.g., monthly SRRs, the Sun-satellite geometry, and O₃). But such a multi-
regression model may need to be optimized locally for different regions and can be more challenging to implement, as
compared with the NNs.

Deleted: 9

Deleted: 4

Deleted: 9

4.5 Implementation of a PCA-NN SO₂ fitting algorithm

Deleted: 4

So far, we have relied on the output from the existing PCA SO₂ algorithm as input to the NNs; therefore, our method can be
viewed as an additional data processing step following the spectral fit. For a potential alternative to this approach, we also
attempt to build an NN-based SO₂ SCD fitting algorithm that uses the measured radiances as inputs and the NN analysed SCDs
405 for training targets. As with the PCA SO₂ algorithm, the NN fitting algorithm uses the logarithm of Sun-normalized Earthshine
radiances at 310.5-340 nm and processes each OMI row separately with individually trained NNs. We pool the data from 12
days in 2005 (the 10th day of each month), generating a training dataset that contains about 200000 pixels for each row. To
reduce the data dimension of the inputs, a PCA technique is combined with the NNs in this PCA-NN fitting algorithm as in
Joiner et al. (2022). We conduct PCA on the radiance spectra and include the coefficients of the first 50 leading PCs as
410 predictors in the NNs. Experiments using fewer (as few as 20) or more (up to 100) PCs generally result in larger errors in the
retrieved SCDs. In addition to the PC coefficients, the NNs also use four other parameters (solar zenith angles, O₃ column
amounts, scene reflectivity, and monthly mean SRR ratios) as predictors. Viewing zenith angles are not included since the
training is carried out separately for each row. We also exclude the phase angles, given that adding them as a predictor leads
to no discernible improvements in the algorithm performance. The SRRs for individual pixels are also excluded, as the PCA-
415 NN algorithm is designed to run independently from the PCA SO₂ algorithm after the training phase. While the monthly mean
SRRs also originate from the PCA retrievals, they essentially provide geospatial context on the spatial distribution of SO₂ and
can be potentially replaced with other datasets such as SO₂ emission inventories or model simulated SO₂.

For the PCA-NN algorithm, we utilize a NN architecture similar to that in Fig. 1c, with the only difference being that
the number of nodes in each hidden layer is now 108 (twice the number of the predictors). For each row, we train an NN using
420 half of the pixels and the rest for evaluation. The pre-trained NNs are then applied to SO₂ SCD retrievals for April 16, 2005, a
day not used in the training.

The results shown in Fig. 14 indicate that the PCA-NN algorithm can reduce the retrieval noise over background areas as compared with the original PCA SO₂ algorithm. However, over polluted areas and degassing volcanoes, the PCA-NN retrieved SO₂ is biased low (Fig. 14c). This suggests that the PCA-NN algorithm, with its present implementation, cannot yet achieve the same level of performance as our NN-based data analysis on the original PCA retrievals. It is possible that due to the much smaller number of polluted pixels as compared with the clean ones, some spectral signatures of SO₂ are lost in the first 50 or even 100 PCs, leading to the low bias over polluted areas. The NNs may need to include more PCs as predictors or directly use radiances without the transformation. A separate set of NNs trained on a refined dataset that contains more polluted pixels may also help to mitigate the bias. But applying these NNs to retrievals would require some prior knowledge about the status of the pixels (whether they are polluted or clean). Also, the PCA-NN retrievals show some striping features, probably reflecting the different performance of the NNs for different rows despite the use of the same architecture. The reason for this row-to-row change in performance is not yet understood. Nonetheless, the PCA-NN algorithm shows promises and will be the subject of more in-depth studies in the future. For example, the training performance may improve if the architecture is optimized for each row.

Deleted: 0

Deleted: 0

5 Conclusions

We have developed a new machine learning based method to analyse satellite retrieved atmospheric composition data, with the aim to reduce the noise and artifacts while retaining the signals in the original retrievals. To demonstrate this approach, we use OMI SO₂ SCDs retrieved with the PCA-based spectral fitting algorithm as an example. A key parameter in the analysis method is the SRR, the ratio between the retrieved SCD and the RMS of the fitting residuals. Based on prior knowledge about the global distribution of SO₂ pollution (from existing in situ measurements and model simulations), we assume that a given pixel with a small (large) SRR is likely clean (polluted) and its real SCD should be close to zero (the original retrieved SCD). This allows us to overcome the lack of ground truth data and build a training dataset for SO₂ by selecting clean and polluted pixels from the original retrievals.

We then train neural networks (NNs) using the compiled dataset. The NNs contain two hidden layers with 14 nodes each and one node in the output layer for the analysed SCDs. The predictors for the NNs include SRRs for individual pixels, solar zenith, viewing zenith and phase angles, scene reflectivity, and O₃ column amounts, as well as the monthly mean SRRs. The latter provide context for the spatial distribution of SO₂, whereas the other predictors (angles, O₃ and reflectivity) affect the quality of the original SCDs. The function of the NNs is to connect these predictors to the target SCDs (zero for clean pixels, the original SCDs for polluted pixels in the training data). For data analysis, we employ a hybrid model (Fig. 1) that includes two NNs: 1) an NN pre-trained using 5 days of data from each month to produce analysed SO₂ SCDs for pixels that are clean or moderately polluted (i.e., those with SRRs in between clean and polluted pixels) for the entire month and 2) an NN trained daily to produce analysed SCDs for the polluted pixels each day. This hybrid model helps to maximize the retained SO₂ signals over source areas.

460 Results for 2005 show that the NNs can well reproduce the target SCDs and largely reduce noise and artifacts in the original retrievals. For polluted areas, the monthly mean SCDs from the analysis are mostly within $\pm 15\%$ from the original retrievals, indicating that the NNs are able to preserve SO₂ signals. This is confirmed by another experiment in which the NN analysed and original SCDs are used to estimate the SO₂ emissions for ~500 anthropogenic sources in 2005, with both datasets yielding largely similar results. For relatively small sources (< 20 kt/year), the emission estimates based on the analysed SCDs
465 are generally smaller, but the uncertainties for those sources are reduced even more, although OMI has quite limited sensitivity to such small sources. One remaining issue is that the NNs perform slightly worse for high SO₂ conditions such as plumes from large volcanic eruptions (e.g., the 2005 Sierra Negra eruption). This will be the focus of future studies to further improve the method. [Also the NNs analysed SCDs show moderate sensitivity to the settings of the pixel classification scheme. Therefore the scheme needs to be tested, especially for different instruments and/or species, to minimize over- or under-correction.](#)
470 Overall, it is quite encouraging that the NNs seem to have improved the quality of SCDs for pixels from different ranges of SRRs.

We also compare two alternative approaches with the NN-based analysis method. In one test, we employ a simple linear interpolation model to analyse the original retrievals. The linear model can largely match the performance of NNs over background areas, but underestimates SO₂ over polluted regions. In another test, we develop a PCA-NN algorithm that first
475 transforms OMI measured radiances using a PCA technique and then uses the resulting PC coefficients as predictors in NNs (trained with NN analysed SCDs) for SO₂ retrievals. Again, the PCA-NN algorithm can reduce retrieval noise but also has a low bias over SO₂ source areas. One advantage of the PCA-NN algorithm is its computation speed (approximately a factor of two faster than the original PCA algorithm in our limited tests) that can make it useful for high resolution instruments such as TROPOMI or TEMPO (Tropospheric Emissions: Monitoring of Pollution). Further improvement in the PCA-NN SO₂
480 algorithm may be possible through, for example, refinement of the training data and will be the subject for follow-up studies. [The lack of the high-quality training data has been a major obstacle for training NNs to conduct retrievals using radiances \(or PCA transformed radiances\). Our analysis method can contribute to such efforts by providing training data with improved quality as compared with the original retrievals.](#)

In summary, our new machine learning based data analysis method shows promises in further improving satellite
485 retrievals of atmospheric composition. [In a way, our analysis method can be viewed as a more advanced version of the Pacific sector correction \(PSC\), a quite common and well-established practice to reduce retrieval artifacts for species such as SO₂ \(e.g., Theys et al., 2017\).](#) While we focus on OMI SO₂ in this study, the method can also be potentially applied to other instruments (e.g., TROPOMI) and/or species (e.g., HCHO). The improved data quality [from such analyses](#) will likely enhance the value of satellite data in air quality research and applications such as reducing the uncertainty in top-down emission
490 estimates.

Code and data availability

Collection 4 OMI L1B radiance and irradiance data are available, free of charge, at Goddard Earth Sciences Data and Information Services Center (https://disc.gsfc.nasa.gov/datasets/OML1BRUG_004/summary). The experimental OMI PCA SO₂ SCDs and NN analysed SCDs are available upon request from the corresponding author. Code used to analyse data and produce figures in this paper is also available upon request from the corresponding author.

Author contribution

CL and JJ designed the NN-based analysis method. CL implemented the method, performed tests, and prepared the manuscript. JJ provided the code and NN architecture used in the study. FL conducted top-down emission estimates. VF and CM designed and provided the emission algorithm. All authors commented on the manuscript.

500 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We would like to thank Dr. Arlindo da Silva of NASA Goddard Space Flight Center for comments and suggestions on the interpretation of the NN analysed results. We also thank the NASA Earth Science Division (ESD) Aura Science Team program for funding of the OMI SO₂ product development and analysis. The Ozone Monitoring Instrument (OMI) is a Dutch/Finnish instrument flying aboard the NASA Earth Observing System Aura spacecraft. The OMI project is managed by the Royal Meteorological Institute of the Netherlands (KNMI) and the Netherlands Space Agency (NSO).

References

- Bhartia, P. K.: OMI/Aura Ozone (O₃) Total Column 1-Orbit L2 Swath 13x24 km V003, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: March 30, 2022, doi:10.5067/Aura/OMI/DATA2024, 2005.
- Castellanos, P. and da Silva, A.: A neural network correction to the scalar approximation in radiative transfer, *J. Atmos. Ocean. Technol.*, 36, 819-832, <https://doi.org/10.1175/JTECH-D-18-0003.1>, 2019.
- Chan, K. L., Khorsandi, E., Liu, S., Baier, F., and Valks, P.: Estimation of surface NO₂ concentrations over Germany from TROPOMI satellite observations using a machine learning method, *Remote Sens.*, 13, 969. <https://doi.org/10.3390/rs13050969>, 2021.

- Chimot, J., Veefkind, J. P., Vlemmix, T., de Haan, J. F., Amiridis, V., Proestakis, E., Marinou, E., and Levelt, P. F.: An exploratory study on the aerosol height retrieval from OMI measurements of the 477 nm O₂-O₂ spectral band using a neural network approach, *Atmos. Meas. Tech.*, 10, 783–809, <https://doi.org/10.5194/amt-10-783-2017>, 2017.
- 520 Eisinger, M. and Burrows, J. P.: Tropospheric sulfur dioxide observed by the ERS-2 GOME instrument, *Geophys. Res. Lett.*, 25(22), 4177–4180, doi:10.1029/1998GL900128, 1998.
- De Santis, D., Petracca, I., Corradini, S., Guerrieri, L., Picchiani, M., Merucci, L., Stelitano, D., Del Frate, F., Prata, F., and Schiavon, G.: Volcanic SO₂ near-real time retrieval using TROPOMI data and neural networks: The December 2018 Etna test case, 2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS, 8480-8483, doi:10.1109/IGARSS47720.2021.9554915, 2021.
- 525 Fedkin, N. M., Li, C., Krotkov, N. A., Hedelt, P., Loyola, D. G., Dickerson, R. R., and Spurr, R.: Volcanic SO₂ effective layer height retrieval for the Ozone Monitoring Instrument (OMI) using a machine-learning approach, *Atmos. Meas. Tech.*, 14, 3673–3691, <https://doi.org/10.5194/amt-14-3673-2021>, 2021.
- Fioletov, V., McLinden, C., Krotkov, N. A., and Li, C.: Lifetimes and emissions of SO₂ from point sources estimated from OMI, *Geophys. Res. Lett.*, 42, doi:10.1002/2015GL063148, 2015.
- 530 Fioletov, V. E., McLinden, C. A., Krotkov, N., Li, C., Joiner, J., Theys, N., Carn, S., and Moran, M. D.: A global catalogue of large SO₂ sources and emissions derived from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 16, 11497-11519, doi:10.5194/acp-16-11497-2016, 2016.
- Hedelt, P., Efremenko, D. S., Loyola, D. G., Spurr, R., and Clarisse, L.: Sulfur dioxide layer height retrieval from Sentinel-5 Precursor/TROPOMI using FP_ILM, *Atmos. Meas. Tech.*, 12, 5503–5517, <https://doi.org/10.5194/amt-12-5503-2019>, 2019.
- 535 Huang, C., Hu, J., Xue, T., Xu, H., and Wang, M.: High-resolution spatiotemporal modeling for ambient PM_{2.5} exposure assessments in China from 2013 to 2019, *Environ. Sci. Technol.*, 55, 2152–2162, <https://dx.doi.org/10.1021/acs.est.0c05815>, 2021.
- Joiner, J. and Vasilkov, A. P.: First results from the OMI rotational-Raman scattering cloud pressure algorithm, *IEEE Trans. Geophys. Remote Sens.*, 44, 1272–1282, 2006.
- 540 [Joiner, J., and Yoshida, Y.: Satellite-based reflectances capture large fraction of variability in global gross primary production \(GPP\) at weekly time scales, *Agric. For. Meteorology* 291, 108092. doi:10.1016/j.agrformet.2020.108092, 2020.](#)
- [Joiner, J., Fasnacht, Z., Gao, B.-C., and Qin, W.: Use of multi-spectral visible and near-infrared satellite data for timely estimates of the Earth's surface reflectance in cloudy and aerosol loaded conditions: Part 2 - image restoration with HICO satellite data in overcast conditions, *Front. Remote Sens.*, 2. doi:10.3389/frsen.2021.721957, 2021.](#)
- 545 [Joiner, J., Fasnacht, Z., Qin, W., Yoshida, Y., Vasilkov, A. P., Li, C., Lamsal, L., and Krotkov, N.: Use of hyper-spectral visible and near-infrared satellite data for timely estimates of the Earth's surface reflectance in cloudy and aerosol loaded conditions: Part 1 - application to RGB image restoration over land with GOME-2, *Front. Remote Sens.*, 2, 716430, doi:10.3389/frsen.2021.716430, 2022.](#)
- 550 Kingma, D. P. and Ba, J.: Adam: A method for stochastic optimization, arXiv preprint arXiv:1412.6980, 2014.

- Krotkov, N. A., Carn, S. A., Krueger, A. J., Bhartia, P. K., and Yang, K.: Band residual difference algorithm for retrieval of SO₂ from the Aura Ozone Monitoring Instrument (OMI), *IEEE Trans. Geosci. Remote Sensing*, 44, 1259–1266, doi:10.1109/TGRS.2005.861932, 2006.
- 555 Krotkov, N. A., McClure, B., Dickerson, R. R., Carn, S. A., Li, C., Bhartia, P. K., Yang, K., Krueger, A. J., Li, Z., Levelt, P. F., Chen, H., Wang, P., and Lu, D.: Validation of SO₂ retrievals from the Ozone Monitoring Instrument over NE China, *J. Geophys. Res.-Atmos.*, 113, D16S40, <https://doi.org/10.1029/2007JD008818>, 2008.
- Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V., Swartz, W. H., Bucsela, E. J., Joiner, J., Duncan, B. N., Boersma, K. F., Veefkind, J. P., Levelt, P. F., Fioletov, V. E., Dickerson, R. R., He, H., Lu, Z. and Streets, D. G.: Aura OMI observations of regional SO₂ and NO₂ pollution changes from 2005 to 2015, *Atmos. Chem. Phys.*, 560 16, 4605–4629, <https://doi.org/10.5194/acp-16-4605-2016>, 2016.
- Lee, C., Martin, R. V., Van Donkelaar, A., Lee, H., Dickerson R. R., Krotkov, N., Richter, A., Vinnikov, K., and Schwab, J. J.: SO₂ emissions and lifetimes: Estimates from inverse modeling using in situ and global, space-based (SCIAMACHY and OMI) observations, *J. Geophys. Res.-Atmos.*, 116, D06304, <https://doi.org/10.1029/2010JD014758>, 2011.
- Levelt, P. F., Joiner, J., Tamminen, J., Veefkind, J. P., Bhartia, P. K., Stein Zweers, D. C., Duncan, B. N., Streets, D. G., Eskes, 575 H., van der A, R., McLinden, C., Fioletov, V., Carn, S., de Laat, J., DeLand, M., Marchenko, S., McPeters, R., Ziemke, J., Fu, D., Liu, X., Pickering, K., Apituley, A., González Abad, G., Arola, A., Boersma, F., Chan Miller, C., Chance, K., de Graaf, M., Hakkarainen, J., Hassinen, S., Ialongo, I., Kleipool, Q., Krotkov, N., Li, C., Lamsal, L., Newman, P., Nowlan, C., Suleiman, R., Tilstra, L. G., Torres, O., Wang, H., and Wargan, K.: The Ozone Monitoring Instrument: overview of 14 years in space, *Atmos. Chem. Phys.*, 18, 5699–5745, <https://doi.org/10.5194/acp-18-5699-2018>, 2018.
- 570 Li, C., Joiner, J., Krotkov, N. A. and Bhartia, P. K.: A fast and sensitive new satellite SO₂ retrieval algorithm based on principal component analysis: application to the ozone monitoring instrument, *Geophys. Res. Lett.*, 40, 6314–6318, doi:10.1002/2013GL058134, 2013.
- Li, C., Krotkov, N. A., Carn, S., Zhang, Y., Spurr, R. J. D., and Joiner, J.: New-generation NASA Aura Ozone Monitoring Instrument (OMI) volcanic SO₂ dataset: algorithm description, initial results, and continuation with the Suomi-NPP Ozone 575 Mapping and Profiler Suite (OMPS), *Atmos. Meas. Tech.*, 10, 445–458, doi:10.5194/amt-10-445-2017, 2017a.
- Li, C., McLinden, C., Fioletov, V., Krotkov, N., Carn, S., Joiner, J., Streets, D., He, H., Ren, X., Li, Z., and Dickerson, R. R.: India is overtaking China as the world's largest emitter of anthropogenic sulfur dioxide, *Sci. Rep.*, 7, 14304, doi: 10.1038/s41598-017-14639-8, 2017b.
- Li, C., Krotkov, N. A., Leonard, P. J. T., Carn, S., Joiner, J., Spurr, R. J. D., and Vasilkov, A.: Version 2 Ozone Monitoring 580 Instrument SO₂ product (OMSO2 V2): new anthropogenic SO₂ vertical column density dataset, *Atmos. Meas. Tech.*, 13, 6175–6191, <https://doi.org/10.5194/amt-13-6175-2020>, 2020.
- Liu, J., Weng, F., and Li, Z.: Satellite-based PM_{2.5} estimation directly from reflectance at the top of the atmosphere using a machine learning algorithm, *Atmos. Environ.*, 208, 113-122, <https://doi.org/10.1016/j.atmosenv.2019.04.002>, 2019.

McLinden, C. Fioletov, V., Shephard, M., Krotkov, N., Li, C., Martin, R. V., Moran, M. D. and Joiner, J.: Space-based
585 detection of missing sulfur dioxide sources of global air pollution, *Nature Geosci.*, DOI: 10.1038/NGEO2724, 2016.

Müller, M. D., Kaifel, A. K., Weber, M., Tellmann, S., Burrows, J. P., and Loyola, D.: Ozone profile retrieval from Global
Ozone Monitoring Experiment (GOME) data using a neural network approach (Neural Network Ozone Retrieval System
(NNORSY)), *J. Geophys. Res.*, 108, 4497, doi:10.1029/2002JD002784, 2003.

Müller, M. D., Kaifel, A., Weber, M., and Burrows, J. P.: Neural network scheme for the retrieval of total ozone from Global
590 Ozone Monitoring Experiment data, *Applied Optics*, 41, 5051-5058, 2004.

Nanda, S., de Graaf, M., Veefkind, J. P., ter Linden, M., Sneep, M., de Haan, J., and Levelt, P. F.: A neural network radiative
transfer model approach applied to the Tropospheric Monitoring Instrument aerosol height algorithm, *Atmos. Meas. Tech.*,
12, 6619–6634, <https://doi.org/10.5194/amt-12-6619-2019>, 2019.

Nowlan, C. R., Liu, X., Chance, K., Cai, Z., Kurosu, T. P., Lee, C., and Martin, R. V.: Retrievals of sulfur dioxide from the
595 Global Ozone Monitoring Experiment 2 (GOME-2) using an optimal estimation approach: Algorithm and initial validation, *J.*
Geophys. Res., 116, D18301, doi:10.1029/2011JD015808, 2011.

Piscini, A., Picchiani, M., Chini, M., Corradini, S., Merucci, L., Del Frate, F., and Stramondo, S.: A neural network approach
for the simultaneous retrieval of volcanic ash parameters and SO₂ using MODIS data, *Atmos. Meas. Tech.*, 7, 4023–4047,
<https://doi.org/10.5194/amt-7-4023-2014>, 2014.

600 Theys, N., De Smedt, I., Yu, H., Danckaert, T., van Gent, J., Hörmann, C., Wagner, T., Hedelt, P., Bauer, H., Romahn, F.,
Pedernghana, M., Loyola, D., and Van Roozendael, M.: Sulfur dioxide retrievals from TROPOMI onboard Sentinel-5 Precursor:
algorithm theoretical basis, *Atmos. Meas. Tech.*, 10, 119–153, <https://doi.org/10.5194/amt-10-119-2017>, 2017.

Theys, N., Fioletov, V., Li, C., De Smedt, I., Lerot, C., McLinden, C., Krotkov, N., Griffin, D., Clarisse, L., Hedelt, P., Loyola,
D., Wagner, T., Kumar, V., Innes, A., Ribas, R., Hendrick, F., Vlietinck, J., Brenot, H., and Van Roozendael, M.: A sulfur
605 dioxide Covariance-Based Retrieval Algorithm (COBRA): application to TROPOMI reveals new emission sources, *Atmos.*
Chem. Phys., 21, 16727–16744, <https://doi.org/10.5194/acp-21-16727-2021>, 2021.

Wells, K. C., Millet, D. B., Payne, V. H., Deventer, M. J., Bates, K. H., de Gouw, J. A., Graus, M., Warneke, C., Wisthaler,
A., and Fuentes, J. D.: Satellite isoprene retrievals constrain emissions and atmospheric oxidation, *Nature*, 585, 225–233,
<https://doi.org/10.1038/s41586-020-2664-3>, 2020.

610 Xu, J., Schüssler, O., Loyola R., D., Romahn, F., and Doicu, A.: A novel ozone profile shape retrieval using Full-Physics
Inverse Learning Machine (FP_ILM), *IEEE J. Sel. Top. Appl.*, 10, 5442–5457,
<https://doi.org/10.1109/JSTARS.2017.2740168>, 2017.

Zhang, Y., Gautam, R., Zavala-Araiza, D., Jacob, D. J., Zhang, R., Zhu, L., Sheng, J., and Scarpelli, T.: Satellite-observed
changes in Mexico's offshore gas flaring activity linked to oil/gas regulations, *Geophys. Res. Lett.*, 3, 1879–1888,
615 10.1029/2018gl081145, 2019.

Zhang, S., Mi, T., Wu, Q., Luo, Y., Grieneisen, M. L., Shi, G., Yang, F., Zhan, Y.: A data-augmentation approach to deriving long-term surface SO₂ across Northern China: Implications for interpretable machine learning, *Science of the Total Environment*, 827, 154278, <https://doi.org/10.1016/j.scitotenv.2022.154278>, 2022.


620 Zheng, T., Bergin, M., Wang, G., and Carlson, D.: Local PM_{2.5} hotspot detector at 300 m resolution: A random forest-convolutional neural network joint model jointly trained on satellite images and meteorology, *Remote Sens.*, 13, 1356, <https://doi.org/10.3390/rs13071356>, 2021.

Table 1. The correlation coefficient (r) and root mean square error (RMSE) between the NN analysed OMI SO₂ SCDs (SCD_{NN}) using different predictors and the target SCDs (SCD_{target}). The NNs are trained using data from July 5, 10, 15, 20 and 25, 2005.

625 The comparisons shown here are for pixels from the same days but not included in the training.


Predictors	r	RMSE (DU)
SRR_t	0.642*	0.180*
$SRR_t + SRR_m$	0.958	0.0517
$SRR_t + SRR_m + \theta$	0.962	0.0491
$SRR_t + SRR_m + R + \theta$	0.968	0.0451
$SRR_t + SRR_m + R + \theta_0 + \theta$	0.976	0.0393
$SRR_t + SRR_m + R + \theta_0 + \theta + \phi$	0.976	0.0392
$SRR_m + R + \theta_0 + \theta + \phi + O_3$	0.793	0.111
$SRR_t + R + \theta_0 + \theta + \phi + O_3$	0.978	0.0374
$SRR_t + SRR_m + R + \theta_0 + \theta + \phi + O_3$	0.976	0.0388

*Results shown are from a simple linear regression analysis.



630 **Figure 1: (a) Flow chart of the SO₂ analysis method. (b) Scheme for classification of OMI pixels as "clean", "polluted", "in-between" and "high-SRR". (c) Setups of the neural networks for SO₂ SCD analysis.**

(a) SO₂ SCDs



(b) SO₂ SCD/RMS




Figure 2: (a) Monthly mean OMI SO₂ SCDs for March 2005 showing enhanced SO₂ signals over major anthropogenic source areas (e.g., China, the eastern U.S., India, and South America) as well as degassing volcanoes. Note the positive bias at northern high latitudes. (b) Monthly mean SCD/RMS ratio (SRR) from the same sample of OMI pixels as in (a). The SRR map also shows major SO₂ sources but has reduced bias at high latitudes as compared with the SCD map.

635





Figure 3: Scatter plots between the NN analysed SO₂ SCDs and the target SO₂ SCDs for clean and polluted OMI pixels from the 16th day of (a) March, (b) April, (c) July, and (d) October 2005. Only pixels not used in the training of the neural networks are shown. Colours represent the number of data points within each 0.1 DU (in NN SCDs) by 0.1 DU (in target SCDs) bin. The solid line in each panel represents the best fit through the data from the simple linear regression analysis between NN and target SCDs. The slope and intercept from the regression are given in each panel, along with the correlation coefficient (r), root mean square error (RMSE), and number of pixels (N).

640



645

Figure 4: The (a) original, (b) NN analysed OMI SO₂ SCDs and (c) their differences for April 16, 2005, (d) Mean SO₂ SCDs for 1° latitude bands over generally clean areas (monthly mean SRR < 3), calculated from (red) the original and (blue) NN analysed SCDs for the same day.

Deleted: and
Deleted: and (c) their differences




Figure 5: Similar to Figure 4 but showing monthly means for March 2005.






Figure 6: (a) The NN analysed SO₂ SCDs and (b) their relative differences from the original SCDs over eastern China for March 2005.



660 Figure 7: Left: the differences between the original and the NN analysed OMI SO₂ SCDs for March 2005. Right: the mean (red) original and (blue) NN analysed SCDs for 3° latitude bands for the same month. Different rows show results from pixels that have SCD/RMS ratios (SRR) within different ranges based on the monthly medians of the daily mean (\overline{SRR}) and standard deviation (σ_{SRR}) of SRRs for their corresponding latitude bands: (a-j) $SRR_i < \overline{SRR}$, (c-d) $\overline{SRR} < SRR_i < \overline{SRR} + \sigma_{SRR}$, (e-f) $\overline{SRR} + \sigma_{SRR} < SRR_i < \overline{SRR} + 2\sigma_{SRR}$, (g-h) $\overline{SRR} + 2\sigma_{SRR} < SRR_i < \overline{SRR} + 3\sigma_{SRR}$ and (i-j) $SRR_i > \overline{SRR} + 3\sigma_{SRR}$.

Deleted: (First column
 Deleted:)
 Deleted: T
 Deleted: original, (second column)
 Deleted: and (third
 Deleted: column
 Deleted:)
 Deleted: their
 Deleted: differences
 Deleted: for March 2005
 Deleted: -
 Deleted: c
 Deleted: d
 Deleted: f
 Deleted: g
 Deleted: i
 Deleted: j
 Deleted: l
 Deleted: p
 Deleted: r



685 **Figure 8:** (a) Differences in the analysed OMI SO₂ SCDs for March 2005 between NNs trained using pixels classified with a₁ and a₂
 (see Eqs. 1 and 2) scaled to 90% of the baseline values in the classification scheme and those trained with the baseline scheme. (b)
 690 Same as (a) but for a₁ and a₂ scaled to 110% of the baseline values. (c) Mean SCDs for 1° latitude bands over relatively clean areas
 (monthly mean SRR < 3) using NNs trained with pixels from different classification schemes: (black) the baseline a₁ and a₂, (blue)
 a₁ and a₂ scaled to 90% and (red) a₁ and a₂ scaled to 110% of the baseline values.






Figure 9: Scatter plots comparing (a) the annual emission estimates for 485 large point sources for 2005, (b) the uncertainties in the emission estimates, and (c) the ratios between the emission estimates and the uncertainties using the NN analysed vs. the original SCDs. All sources shown here are anthropogenic and have emission estimates at least twice the uncertainties for both datasets.

Deleted: 8

(a) SCDs, Linear Interpolation



(b) SCD Diff., Linear - NN



(c) Mean SCDs





Figure 10: (a) Monthly mean OMI SO₂ SCDs for March 2005, analysed using a simple linear interpolation model. (b) The differences in the analysed SCDs between the linear model and the neural networks. (c) Mean SCDs for 1° latitude bands over generally clean areas (monthly mean SRR < 3), calculated from the SCDs from (red) the linear interpolation model and (blue) the NNs.

Deleted: 9



705 **Figure 11:** OMI SO₂ SCDs for April 16, 2005 retrieved using (a) the original PCA algorithm and (b) a PCA-NN algorithm, (c) the differences between the two retrievals, and (d) mean SO₂ SCDs for 1° latitude bands over relatively clean areas (monthly mean SRR < 3), calculated from (red) the original and (blue) PCA-NN retrievals.

Deleted: 0

Deleted: and