

1 Earth Observations from the Moon surface: dependence on lunar libration

2 Nick Gorkavyi¹, -Nickolay Krotkov², Alexander Marshak²

3 ¹ Science Systems and Applications, Inc., Lanham, MD, USA

4 ² National Aeronautics and Space Administration (NASA), Goddard Space Flight Center (GSFC),
5 Greenbelt, MD, USA

6

7 **Correspondence:** Nick Gorkavyi (nick.gorkavyi@ssaihq.com)

8 **Abstract.** Observing the Earth from the Moon surface has important scientific advantages. The
9 angular diameter of the Earth as seen from the Moon surface is 1.8° -2.0° 1.9° ± 0.1° (the angular
10 size varies due to the change in the Earth-Moon distance-between the Earth and the Moon). The
11 libration of the Moon in latitude reaches an amplitude of 6.68° and has α-the main period of 27.21
12 days (or 653.1 hours). The libration of the Moon in longitude, reaching an amplitude of 7.9°, has a
13 period of 27.55 days (or 661.3 hours). This causes the center of the Earth to move in the Moon's sky
14 in a rectangle measuring 13.4° × 15.8°. The trajectory of the Earth's motion in this rectangle
15 changes its shape with a period of 6 years. This apparent librational movement of the Earth in the
16 Moon's sky complicates observations of the Earth. The paper proposes to turn this disadvantage
17 into an advantage and place a multi-slit spectrometer on the Moon surface on a fixed platform. The
18 libration motion and the daily rotation of the Earth will act as a natural replacement for the
19 scanning mechanism.

20

21 1 Introduction

22

23 The scientific benefits of observations from the Moon for the Earth, exoplanet and astrophysics
24 studies are discussed in several recent papers (Marshak et al., 2020; Gorkavyi et al., 2021; Boyd et
25 al., 2022). Although current Earth-observing satellites can produce high-resolution images, Low
26 Earth Orbit (LEO) sensors can only scan a small portion of the globe at a given time, while
27 Geosynchronous Equatorial Orbit (GEO) sensors can provide temporally continuous, though lower-
28 resolution observations of a significant, but fixed, portion of the Earth's disk. The Earth
29 Polychromatic Imaging Camera (EPIC) on the Deep Space Climate Observatory (DSCOVR) clearly
30 stands apart, observing the entire Sun-illuminated Earth from the L1 Sun-Earth Lagrange point
31 (Marshak et al., 2018). The L1 location, however, limits phase angles to a nearly backscattering
32 direction (a phase angle is between 2° and 12°). The phase angle interval from 2° to 12° is
33 determined by the trajectory of the DSCOVR space observatory, which does not rest at the Lagrange
34 point, but moves around it. It is too 'noisy' to transmit data directly from the Earth-Sun line; thus,
35 the phase angle is bigger than 2°. A compact, lightweight, autonomous camera and spectrometer on
36 the Moon's surface offers a unique opportunity to complement these observations and image the full
37 range of Earth phases, potentially advancing Earth science in many ways (Marshak et al., 2020;
38 Gorkavyi et al., 2021):


39

- 40 1. observing ocean/cloud glints reflection-forat different phase angles;
- 41 2. comprehensive whole-globe monitoring of transient volcanic and aerosol clouds, including the
42 strategically important (for climate studies) polar regions not covered by GEO;
- 43 3. detecting of polar mesospheric and stratospheric clouds;
- 44 4. estimating the bidirectional surface reflectance factor (BRF) and full phase-angle integrated
albedo;

- 45 | 5. monitoring and quantifying changes in vegetationed land;
46 | 6. simultaneous imaging of the day and night parts (i.e., the twilight zone) during crescent phases of
47 | the Earth and shadowed parts illuminated by the Moon.

48 | The first telescopic image of the Earth from the Moon was obtained during the expedition
49 | Apollo 16 in 1972 using an ultraviolet telescope (Carruthers and Page, 1972) – see Fig-Figure 1. In
50 | later years, prospects for lunar observations of the Earth have been discussed in many papers (e.g.,
51 | Foing, 1996; Moccia and Renga, 2010). Observations of the Earth with instruments mounted on the
52 | Moon have been actively discussed in recent years (Hamill, 2016). Impressive prospects for
53 | observing the Earth from the Moon in the visible spectrum are demonstrated in the pictures taken by
54 | the Lunar Reconnaissance Orbiter (LRO) in 2015 (Fig-Figure 2). The difference in distances Earth -
55 | Moon and Earth – L1 leads to the fact that a telescope with the same field of view sees the Earth
56 | with different angular sizes and resolutions (Figure 2).

57 |



58 |


59 | **Figure 1. Left:** A Far-Ultraviolet Camera/Spectrograph was operated on the lunar surface during the
60 | Apollo 16 mission, April 1972 (Credits: NASA/Apollo 16). **Right:** The Earth, photographed in far-
61 | ultraviolet light (1304 angstroms) by astronaut John W. Young. Credits: G. Carruthers (NRL) et
62 | al./Far UV Camera/NASA/Apollo 16; based on the image AS16-123-19657 (Mason, 2019).

63 | One of the objectives of the Chinese space program is Moon-based observation of the Earth
64 | (Li et al., 2019; Guo et al, 2019). A lunar lander Chang’e-3 (landed on the Moon in 2014 and is still
65 | working) is equipped with a 5-cm ultraviolet telescope and extreme UV camera and studied changes
66 | in the Earth's plasmasphere in the UV range (He et al., 2016).

67 | One of the tasks facing the US Artemis program is: “Use the Moon as a platform for Earth-
68 | observing studies... The observations from the Moon will have higher resolution than would similar
69 | observations made from L1. Myriad science investigations targeting topics such as lightning, Earth’s
70 | albedo, atmosphere, and exosphere..., the oceans, infrared emission, and radar interferometry may
71 | be accomplished from the surface of the Moon. The Moon also offers a unique vantage point for
72 | full-disk observations...” (Artemis III Science, 2020).

73 | Because of tidal locking, the Moon's rotation around its axis is synchronized with its orbital
74 | rotation around the Earth. Therefore, the Moon always faces the Earth on one side, and the task of
75 | observing the Earth from the Moon seems simple: the Earth must hang motionless in the lunar sky,
76 | rotating around its axis. In reality, the Earth moves along a complex trajectory in the sky of the
77 | Moon due to lunar librations in latitude and longitude. On one hand, the librations of the Moon

78 cause the Earth to shift from the field of view of the lunar telescope, which forces one to turn the
79 telescope to track the Earth observable movement in the Moon sky (guiding); on other hand, for a
80 fixed slit spectrometer, the librations of the Moon can be useful, because they make the Earth move
81 through the fixed field of view of the instrument (across the slit). These librations can serve as a
82 natural mechanism for scanning the Earth when observed from the Moon, which allows the use of
83 slit spectrometers and telescopes on a fixed platform. This paper takes into account lunar librations
84 and analyzes the conditions for observing the Earth from the Moon.



85

86

87 **Figure 2.** A unique view of Earth from the LRO's vantage point in orbit around the Moon (October
88 12, 2015). LRO was about 134 km above the Moon's far_side crater Compton (55°N, 104°E). The
89 photograph is a combination of images in seven color bands from a wide-angle camera (WAC) and
90 black and white images from two narrow angle linear pushbroom cameras (NACs) with atthe linear
91 (one-dimensional) array ofrom 5064 elements. Each NAC camera has a field of view of 2.86°.
92 Image Credit: NASA/GSFC/Arizona State University ([https://www.nasa.gov/image-](https://www.nasa.gov/image-feature/goddard/lro-earthrise-2015)
93 [feature/goddard/lro-earthrise-2015](https://www.nasa.gov/image-feature/goddard/lro-earthrise-2015)). LRO data (Burns et al., 2012; Keller et al., 2016) can be of
94 great help in planning Earth observations from the lunar surface. Insert shows an image of the same
95 part of the Earth taken by DSCOVR/EPIC on the same day (October 12, 2015) as it would be seen
96 by the LRO from the Lagrange point.

97

98


99

2 Librations of the Moon

100

101 Observations of the Earth from the Moon require accounting for the geometry of the relative
 102 position of the centers of the Earth and the Moon, the inclinations of their axes, and [the](#) libration
 103 effects (Meeus, 1991,2000; Guo et al., 2018; Xu and Chen, 2019; Huang et al., 2020).

104
 105 **Libration of the Moon in latitude.** The angle between Moon's axis of rotation (NS [or North-South](#))
 106 and the normal to the plane of its orbit around Earth (PP') is 6.68° ([Fig.Figure 3](#)). This causes the
 107 libration of the Moon in latitude with the same amplitude and with a period of draconic month
 108 $T_D=27.21222$ days or 653.0933 h (the interval between consecutive passages of the Moon through
 109 the same node of the orbit; an orbital node is either of the two points where a lunar orbit intersects
 110 an ecliptic plane to which it is inclined) – see, for example, Meeus (1991, 2000). As a result of this
 111 inclination, parts of the Moon polar regions are accessible for observations from the Earth.



112
 113 **Figure 3.** Libration in latitude results from an inclination of 6.68° between the Moon's axis of
 114 rotation (NS) and the normal to the plane of its orbit around Earth (PP'). The hemisphere of the
 115 Moon that is visible from Earth at point D is marked in yellow; black is invisible hemisphere at
 116 point B. Additional areas of the lunar surface that become available for observation at points A and
 117 C are marked in orange. View from the point close to the ecliptic plane.


118
 119 **Libration of the Moon in longitude.** The Moon moves around the Earth in an elliptical orbit with
 120 an average eccentricity (or deviation of an orbit from circularity) $e = 0.055$ (it varies between
 121 0.0255 [and](#) $\div 0.0775$) and a period of anomalistic month $T_A=27.55455$ days or 661.3092 h - the
 122 interval between consecutive passages of the Moon through the perigee $r_{min} = a(1 - e)$ or the
 123 apogee $r_{max} = a(1 + e)$ of its orbit, where a is the semi-major axis $a=384,399$ km. ([Fig.Figure 4](#)).
 124 This causes libration in longitude with an amplitude of 7.9° (see, for example, Meeus (1991, 2000)).

125 The longitudinal libration consists of two components:

126 1. An ellipse with a small eccentricity (in the first approximation) can be described as a
 127 circle around the Earth, which is shifted from the center of the circle (point O) by the distance
 128 $OE = ea$ (see [Figure- 4](#)). This displacement leads to the fact that the observer from the Earth begins
 129 to see part of the lateral surfaces of the Moon ([Fig.Figure 4](#)).

130 2. If the Moon was moving along an orbit with a uniform speed, then its visible part would
 131 always be directed to the center of the orbit (point O in [Fig.Figure 4](#)). But the speed of the moon
 132 changes due to the ellipticity of the orbit. If at perigee the Moon is turned to the Earth as in
 133 [Fig.Figure 4](#) at point C, then in a quarter of the anomalistic month $T_A/4$, it should turn 90°
 134 counterclockwise at point D (see the red solid arrow at point D). But due to the high velocity along
 135 the orbit segment CD, the Moon arrives at point D faster than $T_A/4$, so the Moon does not have time
 136 to turn 90° (see the red dashed arrow at point D). This further increases the area of the Moon's
 137 surface visible from Earth. On segment DA, the Moon's orbital speed slows down, and the Moon has
 138 time to turn 180° at point A. With the slow motion of the Moon along segment AB, the Moon has
 139 time to rotate around its axis by more than 90° , which again increases the surface area available for


140 observations from the Earth. On segment BC, the Moon's orbital velocity increases: the Moon
 141 passes through the BC segment faster than $T_A/4$, so it does not have time to turn 90 degrees and, as a
 142 result of this lag, the Moon returns to its initial position at point C.



143
 144 **Figure 4.** Libration in longitude results from the eccentricity of the Moon's orbit. It can reach 7.9° in
 145 amplitude. The point A is the apogee of the lunar orbit ($AE = r_{max}$); C – the perigee ($EC = r_{min}$);
 146 $AO = a$; $OE = ae$; B and D are co-vertices; the semi-minor axis $OB = OD = a\sqrt{1 - e^2}$. Additional
 147 areas of the lunar surface that become available for observation are marked in orange. View from the
 148 North Ppole.

149
 150
 151 Figure 5 shows the librations of the Moon in longitude and latitude for the 2022 (Espanak, 2021).
 152 The selenographic coordinate system repeats the Earth's, therefore, the selenographic center of the
 153 Moon's disk is the intersection point of the lunar equator and the lunar prime meridian. The
 154 selenographic zero corresponds to the average position of the center of the visible disk of the Moon.
 155 At any particular moment in time, the center of the visible disk of the Moon can shift from the
 156 selenographic zero due to libration. If the apparent center shifts along the lunar equator, then we call
 157 this shift the relative longitude of the libration; if it shifts along the meridian, then we call this shift
 158 the relative latitude of the libration. In other words, latitude, and longitude libration (or relative
 159 libration) is the visual displacement of the selenographic center of the Moon's disk (0° lunar latitude
 160 and 0° lunar longitude) relative to the center of the visible disk of the Moon. Libration in longitude
 161 correlates with variations in the Earth-Moon distance and changes more strongly with time than
 162 libration in latitude.

163 The libration of the Moon discussed above (Figures 3 and 4) is called optical libration. The tidal
 164 action of the Earth causes physical libration associated with a change in the period of the Moon's
 165 own rotation. Physical libration is only 2 arc minutes, that is, much less than optical libration. In the
 166 calculations (Figure 5 and below), physical libration is taken into account along with optical.



167

168 **Figure 5.** Variability of the Moon orientation and orbit during the 2022 starting from January 1,
 169 2022 (Espenak, 2021). The distance between the Earth and the Moon is measured between the
 170 centers of the bodies, so it does not depend on libration, which is measured in angles relative to the
 171 centers of the bodies.
 172 **(a)** longitude libration (red) and latitude libration (black). The straight dashed line corresponds to the
 173 case of zero libration, that is, when the center of the visible lunar disk coincides with the zero point
 174 of selenographic longitude and latitude; **(b)** the Earth-Moon distance. The time-averaged distance
 175 between the centers of Earth and the Moon is 385,000 km; the minimal distance is 356,500 km and
 176 the maximal distance 406,700 km.




Figure 6. The relationship between the Earth-Moon distance and lunar librations in latitude (left) and longitude (right) during the 2022–2027 starting from January 1, 2022 (Espenak, 2021). The straight dashed line corresponds to the case of zero libration. The change in the distance between the Earth and the Moon depends on the eccentricity of the lunar orbit, so it does not depend on the degree of libration of the Moon.


3 Visual librations of the Earth

Obviously, the discussed librations of the Moon are directly related to the observation point on the Moon surface: if the observation point on the Moon changes its angle relative to the Earth-Moon line, then the Earth also changes its position in the lunar sky by the same amount, but different sign (in an approximation where the size of the Moon can be neglected compared to the Moon-Earth distance). In other words, if the lunar telescope raises its line of sight up by +5 degrees above the line connecting the centers of the Earth and the Moon, then the Earth goes down from the telescope's line of sight by -5 degrees.

An observer can see the Earth from any point in the Moon's visible hemisphere (Fig-Figure 67). The location of the observer will affect the i) position of the zero point of Earth libration in latitude and longitude and ii) orientation of the trajectory of the apparent libration of the Earth in the sky of the Moon.

From the point of view of an Earth observer, lunar librations in latitude and longitude are measured as a relative displacement from the lunar zero longitude and longitude - that is, from the point of the lunar disk taken as the zero point and located in the center of the visible disk of the Moon, near the crater Möstig A. From the point of view of a lunar observer located near this crater at the intersection of the lunar equator and the lunar zero meridian (see point O in Fig-Figure 67), the Earth hangs above a given point on the lunar surface (at the zenith). Therefore, if the observer moves away from this point along the lunar meridian, for example, to the North pole (point N in Fig-Figure 67), then the apparent position of the center of the Earth will also shift, moving to the horizon. When the observer is at the lunar pole, the Earth will hang on the horizon. If the observer goes again to the

207 equator, but not along the zero meridian, but along the meridian with a longitude of 90° (for
 208 | example, to West, see points NW and W in Fig.Figure 67), that is, along the border between the
 209 visible and invisible hemispheres of the Moon, then the Earth will remain hanging above the lunar
 210 horizon, but will change the apparent tilt of its axis of rotation. For an observer at the Moon's
 211 | equator (points W, O, E in Fig.Figure 76), the Earth's axis will tilt 90° , that is, the Earth will "lay on
 212 its side."
 213




214
 215

216 | **Figure 67.** Observers in different points of the Moon's visible hemisphere. Image of the Moon –
 217 LRO (NASA/GSFC/Arizona State University) [https://www.nasa.gov/feature/goddard/2020/moon-](https://www.nasa.gov/feature/goddard/2020/moon-more-metallic-than-thought)
 218 [more-metallic-than-thought](https://www.nasa.gov/feature/goddard/2020/moon-more-metallic-than-thought). Observer: photo of John W. Young, commander of the Apollo 16 lunar
 219 landing mission (NASA). It is shown how, from the point of view of different observers on the
 220 Moon, the crescent of the Earth is oriented, indicating the Earth's poles. The yellow squares (with
 221 blue top and violet bottom lines) show the astronaut's vertically oriented field of view (FOV) and the
 222 crescent of the Earth that he sees in this FOV (or in frame of the camera).
 223

224 | When the observer reaches the South Pole (point S in Fig.Figure 76), the Earth will be turned
 225 180° relative to it. Similar changes will occur with the trajectories of the Earth in the sky of the
 226 Moon. Libration of the Moon sets the trajectory of the Earth in the lunar sky described by relative
 227 | latitude and longitude (relative to the point of zero libration, marked with a black dot in Figure 78).
 228 The shape of this trajectory (see the red trajectories in Figure 87) is strictly defined and does not
 229 depend on the position of the observer on the Moon's surface. But the height of the point of zero
 230 libration above the horizon depends on the position of the lunar observer, as well as the orientation
 231 of the libration trajectory, that is, the rotation of the visible libration trajectory around this point of
 232 zero libration. An analogy is a picture hanging on the wall of a room. The pattern in the picture does
 233 not depend on the position of the observer, but he can stand on his head and completely change the
 234 orientation of the pattern relative to his field of vision.

235 We can take the latitude and longitude of the libration of the Moon (Giesen, 2018; Espenak,
 236 2021) (Fig.Figure 5) and plot the positions of the Earth in the sky of the Moon for each day
 237 (Fig.Figure 87). Each dot in the Figures 87 abc represents the latitude and longitude for a particular
 238 day.



239

240 **Figure 78.** Visual librations of Earth in the Moon sky during (a) first 6 months 2022; (b) first 6
 241 months 2023; (c) July-September 2024; (d) 2022-2024 (the dots are deleted). The latitude and
 242 longitude of the libration of the Moon (Giesen, 2018; Espenak, 2021) were converted to the relative
 243 Earth libration angles using a sign change. The positions of the Earth in the sky of the Moon are plot
 244 in increments of a day. If there were no lunar libration, then the Earth would be at the black dot in
 245 the center of the figures.

246

247 Figure 78a shows the visual position of the Earth for the first half of 2022. Figure 87b shows
 248 the apparent libration of the Earth for the first half of 2023, and Fig.Figure 87c - for 4 months (July-
 249 September) of 2024. Figure 87d shows the trajectory of the Earth in the sky of the Moon for three
 250 years (2022-2024). The orientation of the libration pattern in Fig.Figure 87 corresponds to the
 251 position of the observer on the line of the zero meridian S-O. At point O, the zero libration point is
 252 above the observer's head, and when the observer moves to the South Pole (point S), the zero
 253 libration point shifts to the horizon.

254 It can be seen that the shapes of the curves along which the Earth moves in the sky of the
 255 Moon change noticeably during 3 years.

256 The movement of the Moon around the Earth can be characterized by three periods:

- 257 • Draconic month $T_D=27.21222$ days or 653.0933 h (the period of movement relative to the
 258 stars ascending or descending node)
- 259 • Anomalistic month $T_A=27.55455$ days or 661.3092 h (the period of movement relative to the
 260 perigee)

261 • Sidereal month $T_S=27.32166$ days or 655.7198 h (the period of movement relative to the
 262 ascending nodes)
 263 A beat is an interference pattern between two slightly different frequencies, perceived as a
 264 periodic variation in amplitude whose rate is the difference of the two frequencies. As a result of 3
 265 slightly different lunar periods we have 3 different beats or precession frequencies.

266 The apsidal precession period is $T_{SA} = 8.85$ years and is found by the formula

$$267 \quad \frac{1}{T_{SA}} = \frac{1}{T_S} - \frac{1}{T_A} \quad (1)$$

268 The nodal precession period is $T_{DS} = 18.6$ years and is found by the formula

$$269 \quad \frac{1}{T_{DS}} = \frac{1}{T_D} - \frac{1}{T_S} \quad (2)$$

270 The librations of the Moon in latitude and longitude follow a six-year cycle, when the
 271 major axis of the lunar orbit has performed one complete revolution with respect to the line of nodes
 272 (Meeus, 1991, 2000; Giesen, 2018)


$$273 \quad \frac{1}{T_{DA}} = \frac{1}{T_D} - \frac{1}{T_A} \quad (3)$$

274 with a period of $T_{DA} = 2190.34$ days or 6 (more precisely, 5.99667) anomalistic years (365.259636
 275 days each). All three periods of precessions are connected:

$$276 \quad \frac{1}{6.00} = \frac{1}{18.6} + \frac{1}{8.85} \quad (4)$$

277 Figure 89a shows the position of the center of the Earth in the lunar sky for six years (2022-2027).
 278 Figure 98b shows the statistics of the distribution of the 2191 positions of the center of the Earth for
 279 this period. The average distribution density of the center of the Earth in squares $1^\circ \times 1^\circ$ (or the
 280 number of entries of the center of the Earth into this square for 6 years) is $N = 2191$ positions/255
 281 pixels = 8.6; in reality N ranges from 0 to 34.

282



283

284 **Figure 98.** Visual libration of Earth during 2022-2027 (6 years, 2191 days or positions): (a)
 285 Trajectory of the center of the Earth in the sky of the Moon; (b) Statistics of the distribution of the
 286 2191 positions of the center of the Earth for 6 years. N - the number of entries of the center of the
 287 Earth into each $1^\circ \times 1^\circ$ grid' cell' for this period.

288 4 Multi-slit spectrometer on a fixed platform

289

290 Spectrometric observations through the slit are a common practice for many satellite observations of
291 the Earth. Scanning of the Earth's surface is usually carried out by movement of the low orbit
292 satellites. For observation from the Moon, it is logical to consider the option when scanning occurs
293 due to the libration and diurnal motion of the Earth. The scientific goals for the slit observation are
294 close in both cases.


295 The angular velocity of a point on the Earth's surface in the field of view of the sensor is caused by
296 two comparable factors: the rotation of the Earth around its axis and lunar libration, which causes a
297 shift in the center of the Earth. The rotation of the Earth around its axis is a well-studied process, but
298 librations of the center of the Earth in the lunar sky are poorly understood and raise many questions.
299 When observing the Earth through the slit of the spectrometer, it will be necessary to take into
300 account both the displacement of the center of the Earth and the Earth rotation.

301 The librational apparent motion of the Earth must be taken into account when observing from the
302 Moon and can also become a natural substitute for scanning (Figures: ~~9+0-11~~). It is proposed to
303 install two fixed mount instruments on the Moon surface, directed towards the Earth:

304 1. A hyperspectral sensor (UV, Vis, NIR, IR) will observe Earth passing through fixed vertical
305 slits. The librations of the Moon and the daily rotation of the Earth will serve as a natural
306 scanning mechanism for this spectrometer. This multi-slit spectrometer can be similar to the
307 six-slit hyperspectral Limb Profiler (LP) on the OMPS aboard Suomi National Polar
308 Partnership (S-NPP) LEO satellite, as well as the single-slit hyperspectral Ozone Monitoring
309 Instrument (OMI) on the NASA Earth Observing System (EOS) Aura satellite. Each LP slit
310 uses approximately 1/6 of the detector matrix. A multi-slit spectrometer for observing the
311 Earth from the surface of the Moon can have 6-8 slits, which field of views are shifted by
312 2.5°. Since the maximum angular size of the Earth is 2°~~about two degrees~~, the angular
313 distance between the lines of sight of neighboring slits must be greater than the angular
314 diameter of the Earth, so that light from the Earth does not hit two slits at the same time.
315 Each slit can use the entire matrix because they scan the Earth at different times in turn and
316 do not interfere with each other.


317 2. A wide field-of-view (WFOV) ~18°-20° camera will continuously image the Earth in any
318 points of trajectory, including a part of the lunar surface with a true-color calibration target.
319 The camera can be hyperspectral, with the inclusion of wavelengths that EPIC uses.

320



321
322
323
324
325
326

Figure 910. Visual positions of the Earth's center during 2026 (green line) from the South Pole of the Moon (point S in Fig. Figure 67) and possible positions of slits of the spectrometer (violet). Blue square is $\sim 18^\circ$ - 20° FOV of fixed mount camera. The calibration bar is used to calibrate color images from a wide-angle camera. Lunar surface is from the photo taken by NASA/Apollo.




327
328
329
330
331
332
333
334
335

Figure 11. Left: Position of the Earth center during July-September 2024 (green line) for an observer located on the edge of the visible hemisphere of the Moon in the region of middle $\sim 45^\circ$ N latitudes (point NW in Fig.7). Violet lines are positions of slits of the spectrometer. **Right:** View of Moon limb with Earth on the horizon, Mare Smythii region (3° N, 85° E), July 20, 1969. NASA/JSC/Apollo 11, AS11-44-6551 (<https://eol.jsc.nasa.gov/SearchPhotos/photo.pl?mission=AS11&roll=44&frame=6551>). Lunar surface is from the photo taken by NASA/Apollo

336
337

It should be noted that the longitude of the observation point on the Moon affects the orientation of the visible trajectory of the Earth in the sky of the Moon. For example, the diagonally elongated

338 trajectory of the Earth for July-October 2024 ([Fig-Figure 87c](#), for the case of lunar longitudes near
 339 0° , for observer in the point S in [Fig-Figure 76](#)) will have a different orientation when observed from
 340 the zone of lunar longitudes of about 45°N for observer in the point NW in [Fig-Figure 67](#) ([Fig. 11](#),
 341 [left](#)). [The orientation of the Earth from a point located near the equator of the Moon is shown in Fig.](#)
 342 [11 \(right\)](#). The orientation of the Earth's libration trajectories in the sky of the Moon will change
 343 accordingly (see [Fig-Figure 102](#)). The sun in the region of the lunar poles moves almost parallel to
 344 the horizon, and in the region of the lunar equator it passes through the zenith, descending vertically
 345 to the horizon or rising from it.
 346



347

348

349

Figure 102. [Positions of the Earth for different Moon observers.](#)

350

351 (a) Position of the Earth center and Earth' phases during June 15-29, 2022 (15 points, green line)
 352 for an observer located on the lunar South Pole (the point S in [Figure 6-7](#)). The Earth is almost
 353 completely illuminated on June 29, 2022, while its axis is tilted to the Sun at an almost maximum
 354 angle (summer in the Northern Hemisphere), which creates good conditions for observing the
 355 Earth's North Pole (marked with a cross and the letter N).

356

357 (b) Earth' phases and visible trajectory for same period for an observer on the equator near
 358 longitude 90°W (the point W in [Figure 6-7](#)).

359

360 (c) Position of the Earth center and Earth' phases during June 15-July 12, 2022 (28 points, green
 361 line) for an observer located on the lunar North Pole (the point N in [Figure 6-7](#)).

362


363 (d) Earth' phases and visible trajectory for same period for an observer on the equator near
 364 longitude 90°E (the point E in [Figure 6-7](#)).

362 | Figure 67 shows the orientation of the Earth's crescent from the point of view of different
363 | observers on the Moon and helps interpret the orientation of the libration pattern in Figure- 102.
364 |

365 | The portion of the illuminated Earth is not changing with the position of the Moon observer, but
366 | it's changing during lunar month (Figure- 120, based on data by Espenak, 2021).

367 | These arguments must be taken into account when planning observations of the Earth from the
368 | Moon (or when communicating between the Moon and the Earth). Satellites located at the Earth-
369 | Moon Lagrange points will move along similar trajectories in the sky of the Moon.

370 | The principal design of the multislit spectrometer is shown in Fig-Figure 113. Its main feature is
371 | that it uses only one matrix detector for many slits. This is due to the fact that such a local object as
372 | the Earth can pass only one slit at a given moment. Therefore, it is possible to image the light from
373 | all slits onto a single matrix without compromising observations, although the problem of scattered
374 | light may exist and should be studied in the development of a specific instrument. Each slit is
375 | directed to a unique position of the Earth in the Moon sky, but the spectral rangedispersion of all
376 | slits is the same. If the spectra are taken with a slit that occupies a length of 4000 pixels on the
377 | detector matrix, then the spectra will be determined from the part of the Earth with a size of ~30 km
378 | along the slit. The effective width of this pixel across the slit (i.e. spatial resolution) will depend on
379 | the frequency of observations, the width of the slit, and the velocity of the Earth moving across the
380 | slit.



381 |
382 | **Figure 131.** The principal design of a spectrometer that has multiple slits (A), a collimator (B), a
383 | prism (C) and a single 2-dimensional (~~2D~~)-detector array (matrix D). The spectrometer merges the
384 | light from all the slits together, but since the Earth is always occupies only one slit, the signals from
385 | different slits do not interfere with each other.
386 |

387 | The angular diameter of the Earth in the sky of the Moon is about 1.8°-2.0°9-degrees. The
388 | typical rate of displacement of the center of the Earth is 1°-2° degrees per day (see Figures- 78, 102).
389 | Therefore, the passage of the Earth through each individual slit of the spectrometer will take 1-2
390 | days. During this time, the Earth makes 1-2 rotations around its axis, which will allow each slit to
391 | receive at least one scan of the entire Earth's surface in one pass. The potential scan frequency
392 | depends on the field of view of the device and on the detector matrix used, so that the spatial pixel
393 | across the slit is comparable to the size of the spatial pixel along the slit. For a detector matrix with a
394 | size of 1000-4000 pixels and a field of view of 5 to 15 degrees, the scan frequency should be 10-100
395 | seconds.

396 | Important Earth science goals for such spectrometer are to complement and improve the
397 | current DSCOVR/EPIC whole-Earth imaging (Gorkavyi et al., 2021). The acquired data will enable

398 estimating aerosol and cloud scattering phase functions, amount of trace gases and surface
399 Bidirectional Reflectance Factor (BRDF).

400

401

402 **5 Conclusion**

403 1. Due to lunar libration, the center of the Earth for an observer on the Moon moves in a rectangular
404 area with dimensions of $13.4^\circ \times 15.8^\circ$. The density of the location of the Earth in this rectangular
405 area is an average of 8.6 per square degree over 6 years (2191 days). The density for different parts
406 of the area varies from 1 to 20.

407 2. The movement of the Earth in the sky of the Moon is characterized by quasi-periodicity with
408 frequencies of ~27 days and 6 years. The rates of displacement of the Earth in the Moon sky reach
409 two degrees per day. The shape of the Earth's trajectory changes from a circle to a straight line (see
410 Figure 7).

411 3. Lunar libration must be taken into account when observing the Earth from the surface of the
412 Moon and during Moon-Earth communications.

413 This paper discusses Earth observations from the Moon surface, both spectroscopically and in the
414 imaging mode. The librations of the Moon in the range of 13° - 16° and the daily rotation of the Earth
415 serve as a natural scanning (or guiding) mechanism for a spectrometer with vertical slits. This
416 greatly simplifies the design of the spectrometer. We suggest ~~that proposed~~ lightweight EPIC-
417 Moon instrument on a fixed platform ~~to serve as a will provide the~~ proof of concept for Earth
418 observations, as well as the whole Earth true-color imagery to the public.

419 The lunar environment may have serious problems for the operation of sensors due to dust settling
420 and impact on moving parts, and due to the influence of high-energy particles and meteoroids. The
421 discussed design of instruments on a fixed mount with no movement of the external parts of the
422 instruments, significantly reduces the dependence of observations on lunar dust. During the use of
423 such instruments on the lunar surface, the rate of dust settling and the degree of degradation of
424 instruments due to radiation will be clarified, which will make it possible to optimize the design of
425 future instruments and protect them as much as possible from a hostile environment.

426

427 Observations of the lunar dust environment during total solar/lunar eclipses (it happens ~2 times a
428 year when the Earth blocks the Sun) has a particular interest: the lunar dust glows from scattered
429 sunlight under forward scattering conditions without saturation of the detector.

430

431 The proximity to the Earth (versus the L1 point) and wide variations in phase angle accessible by a
432 Moon-based camera offer unique advantages for observations of the bidirectional land surface
433 reflectance; ocean/cloud glint reflection; whole-globe monitoring of transient volcanic/aerosol
434 clouds, polar mesospheric and stratospheric clouds; vegetation; the twilight zone and shadowed
435 parts of the Earth illuminated by the Moon.

436

437 **Data availability**

438 A Far-Ultraviolet Camera/Spectrograph data are available at [https://gold.cs.ucf.edu/earths-shining-](https://gold.cs.ucf.edu/earths-shining-upper-atmosphere-from-the-apollo-era-to-the-present/)
439 [upper-atmosphere-from-the-apollo-era-to-the-present/](https://gold.cs.ucf.edu/earths-shining-upper-atmosphere-from-the-apollo-era-to-the-present/). The LRO data are available at

440 (<https://www.nasa.gov/image-feature/goddard/lro-earthrise-2015> and at
441 (<https://www.nasa.gov/feature/goddard/2020/moon-more-metallic-than-thought>). The Apollo data
442 are available at <https://moon.nasa.gov/news/38/nasa-mourns-the-passing-of-astronaut-john-young/>
443 and at https://www.nasa.gov/mission_pages/apollo/missions/index.html. Planetary Ephemeris Data
444 Courtesy of Fred Espenak. Data are available at www.Astropixels.com.

445
446 **Author contributions.** NG developed computer codes and algorithms, analyzed the results, and
447 wrote the manuscript. NK, AM participated in [the](#) algorithm development, analyzing the results and
448 writing the manuscript.

449
450 **Competing interests.** The contact author has declared that neither [they](#) [he](#) nor [their](#) [his](#) co-authors
451 have any competing interests.

452
453 **Acknowledgements.** The authors thank the Apollo, LRO and EPIC/DSCOVR teams for providing
454 the data presented. The authors are grateful to Fred Espenak for useful Planetary Ephemeris data.
455 We also thank Padi Boyd and our colleagues from Goddard Space Flight Center interested in the
456 Earth observations from the Moon surface.

457
458 **Financial support.** NK and AM were supported by the NASA DSCOVR project managed by
459 Richard Eckman. AM was supported by the Goddard Artemis project managed by Michele Gates.
460 NG was partially supported by the NASA Aura project (OMI core team) managed by Ken Jucks.

461
462 **References:**

- 463 Artemis III Science Definition Team Report, NASA/SP-20205009602,
464 [https://www.nasa.gov/sites/default/files/atoms/files/artemis-iii-science-definition-report-](https://www.nasa.gov/sites/default/files/atoms/files/artemis-iii-science-definition-report-12042020c.pdf)
465 [12042020c.pdf](https://www.nasa.gov/sites/default/files/atoms/files/artemis-iii-science-definition-report-12042020c.pdf), 2020.
- 466 Burns, K.N., Speyerer, E.J., Robinson, M.S., Tran, T., Rosiek, M.R., Archinal, B. A.,
467 Howington-Kraus, E. and the LROC Science Team: Digital elevation models and derived products
468 from LROC NAC stereo observations, International Archives of the Photogrammetry, Remote
469 Sensing and Spatial Information Sciences, Volume XXXIX-B4, 2012, XXII ISPRS Congress, 25
470 August – 01 September 2012, Melbourne, Australia, [https://doi.org/10.5194/isprsarchives-XXXIX-](https://doi.org/10.5194/isprsarchives-XXXIX-B4-483-2012)
471 [B4-483-2012](https://doi.org/10.5194/isprsarchives-XXXIX-B4-483-2012), 2012
- 472 Boyd, P.T. et al.: EarthShine: Observing our world as an exoplanet from the surface of the
473 Moon, Journal of Astronomical Telescopes, Instruments, and Systems, 8(1), 014003.
474 <https://doi.org/10.1117/1.JATIS.8.1.014003>, 2022.
- 475 Carruthers, G.R. and Page, T.: Apollo 16 Far-Ultraviolet Camera/Spectrograph: Earth
476 Observations, Science, 177, 788-791, <https://doi.org/10.1126/science.177.4051.788>, 1972.
- 477 Espenak, F.: Planetary Ephemeris Data, 2021,
478 <http://www.astropixels.com/ephemeris/ephemeris.html>
- 479 Foing, B.H.: The Moon as a platform for astronomy and space science. Adv. Space Res., 18,
480 1117-1123, 1996.
- 481 Giesen, J., Moon Libration Applet, <http://www.jgiesen.de/moonlibration/>, 2018
- 482 Gorkavyi, N., Krotkov, N., Gorkavyi, N., Marchenko, S., Vasilkov, A., Knyazikhin, Y.,
483 Kowalewski, M., Torres, O., DeLand, M., Ramsey, M., Christensen, P., and Realmuto, V.: Earth
484 Imaging From the Surface of the Moon With a DSCOVR/EPIC-Type Camera, Front. Remote Sens.
485 2:724074. <https://doi.org/10.3389/frsen.2021.724074>, 2021.
- 486 Guo, H., Fu, W., and Liu, G.: Scientific Satellite and Moon-Based Earth Observation for Global
487 Change, Springer, Singapore, <https://doi.org/10.1007/978-981-13-8031-0>, 2019.

488 Guo, H., Liu, G., and Ding, Y.: Moon-based Earth observation: scientific concept and potential
489 applications, *International Journal of Digital Earth*, 11:6, 546-557,
490 <https://doi.org/10.1080/17538947.2017.1356879>, 2018.

491 Hamill, P.: Atmospheric observations from the moon: A lunar earth-observatory, IGARSS 2016
492 - 2016 IEEE International Geoscience and Remote Sensing Symposium,
493 <https://doi.org/10.1109/IGARSS.2016.7729964>, 2016.

494 He, H., Shen, C., Wang, H. et al.: Response of plasmaspheric configuration to substorms
495 revealed by Chang'e 3", *Sci. Rep.*, 6, 32362, <https://doi.org/10.1038/srep32362>, 2016.

496 Huang, J., Guo, H., Liu, G., Shen, G., Ye, H., Deng, Y., and Dong, R.: Spatio-Temporal
497 Characteristics for Moon-Based Earth Observations, *Remote Sens.*, 12, 2848.
498 <https://doi.org/10.3390/rs12172848>, 2020.

499 Keller, J., Petro, N., Vondrak, R. and the LRO team: The Lunar Reconnaissance Orbiter Mission
500 – Six years of science and exploration at the Moon. *Icarus*, **273**, 2-2.
501 <https://doi.org/10.1016/j.icarus.2015.11.024>, 2016.

502 Li, Ch., Wang, Ch., Weiland, Y., and Lin, Y.: China's present and future lunar exploration
503 program, *Science*, 365, 238-239, <https://doi.org/10.1126/science.aax9908>, 2019.

504 Marshak, A., J. Herman, A. Szabo, K. Blank, S. Carn, A. Cede, I. Geogdzhaev, D. Huang, L.-K.
505 Huang, Y. Knyazikhin, M. Kowalewski, N. Krotkov, A. Lyapustin, R. McPeters, K. Meyer, O.
506 Torres and Y. Yang. Earth Observations from DSCOVR/EPIC Instrument. *Bulletin Amer. Meteor.*
507 *Soc. (BAMS)*, 9, 1829-1850, <https://doi.org/10.1175/BAMS-D-17-0223.1>, 2018.

508 Marshak, A., Krotkov, N., Gorkavyi, N., Marchenko, S., Vasilkov, A., Knyazikhin, Y.,
509 Kowalewski, M., Torres, O., DeLand, M., Ramsey, M., Christensen, P., and Realmuto, V.: Whole
510 Earth imaging from the Moon South Pole (EPIC-Moon), WhitePaper # 2054 in Artemis III Science
511 Def. Team Report, 2020.

512 Mason, T., *Earth's Shining Upper Atmosphere — From the Apollo Era to the Present*, 2019;
513 <https://gold.cs.ucf.edu/earths-shining-upper-atmosphere-from-the-apollo-era-to-the-present/> Meeus,
514 J.: *Astronomical Algorithms*, Willmann-Bell, Richmond, Virginia, 1st English Edition 1991.

515 Meeus, J.: *Mathematical Astronomy Morsels*, Willmann-Bell, Richmond, Virginia, 2nd Printing
516 2000.

517 Moccia, A., Renga, A.: Synthetic Aperture Radar for Earth Observation from a Lunar Base:
518 Performance and Potential Applications, *IEEE Trans. on Aerospace and Electr. Systems*, 46, 1034-
519 1051, <https://doi.org/10.1109/TAES.2010.5545172>, 2010.

520 Xu, Z., and Chen, K.-Sh.: Effects of the Earth's Curvature and Lunar Revolution on the Imaging
521 Performance of the Moon-Based Synthetic Aperture Radar, *IEEE Transactions on Geoscience and*
522 *Remote Sensing*, 57, 5868-5882, <https://doi.org/10.1109/TGRS.2019.2902842>, 2019.

523