

Monitoring greenhouse gases (GHGs) in China: status and perspective

Youwen Sun^{1, 2}, Hao Yin^{1, 2}, Wei Wang^{1, 2}, Changgong Shan^{1, 2}, Justus Notholt³,
Mathias Palm³, Ke Liu⁴, Zhenyi Chen^{5*}, and Cheng Liu^{1,2,6,7,8*}

1. Key Laboratory of Environmental Optics and Technology, Anhui Institute of
Optics and Fine Mechanics, HFIPS, Chinese Academy of Sciences, Hefei
230031, China

2. Department of Precision Machinery and Precision Instrumentation, University
of Science and Technology of China, Hefei, 230026, China

3. University of Bremen, Institute of Environmental Physics, P. O. Box 330440,
28334 Bremen, Germany

4. Kaihang Institute of Measurement and Control Technology, 710065, Xi'an,
China

5. School of Ecology and Environment, Beijing Technology and Business
University, 100048, Beijing, China

6. Center for Excellence in Regional Atmospheric Environment, Institute of
Urban Environment, Chinese Academy of Sciences, Xiamen, 361021, China

7. Key Laboratory of Precision Scientific Instrumentation of Anhui Higher
Education Institutes, University of Science and Technology of China, Hefei,
230026, China

8. Anhui Province Key Laboratory of Polar Environment and Global Change,
University of Science and Technology of China, Hefei, 230026, China

*Correspondence: Cheng Liu (chliu81@ustc.edu.cn) and Zhenyi Chen
(zychen@btbu.edu.cn)

Abstract

In order to establish a creditable greenhouse gases (GHGs) monitoring network to support the goals of carbon peak/neutrality, it is necessary to know what we have done and what we have to do in the future. In this study, we summarize an overview to the status and perspective of GHGs monitoring in China. With decades of effort, China has made a great breakthrough in GHGs monitoring capacity and steadily improved the performance of homemade GHGs monitoring instruments. However, most GHGs monitoring studies are hitherto research-oriented, temporal, sparse, and uncoordinated.

1 It is suggested to take full advantages of various monitoring technologies, monitoring
2 platforms, numerical simulations, and inventory compilation techniques to form a
3 creditable GHGs stereoscopic monitoring and assessment system at an operational
4 level. We envisage that this system can routinely quantify GHGs on national,
5 provincial, regional and even individual scales with high spatiotemporal resolution and
6 wide coverage to support low-carbon policy in China.

7 **Keywords:**

8 Carbon peak; Carbon neutrality; Greenhouse gases; Carbon monitoring

9 **1. Introduction**

10 Climate change is one of the great challenges facing humankind around the globe
11 (Tian et al., 2017; Sun et al., 2021; Liu et al., 2022). According to the United Nations
12 Intergovernmental Panel on Climate Change (IPCC) estimates, in order to achieve the
13 1.5° target of the Paris Agreement, the integrated Earth system must achieve net zero
14 carbon dioxide (CO₂) emissions (also known as "neutrality") by 2050, i.e., the annual
15 CO₂ emissions are equal to the amount of CO₂ reductions through strategies that can
16 either increase carbon sinks or reduce carbon sources, e.g., decarbonization, carbon
17 offset, **energy consumption reduction**, tree-planting, carbon capture and storage (CCS),
18 and carbon sequestration (Zheng et al., 2020a). Carbon neutrality is an important
19 strategy to tackle global climate change (IPCC, 2019). Currently, there are 137
20 countries around the globe have proposed carbon neutrality deadlines through policy
21 announcements or legislation. Most of these countries such as the ones from the
22 European Union, the United States, Japan, the Great Britain, Australia, Canada, New
23 Zealand, South Africa, have committed to achieve carbon neutrality by 2050 (IPCC,
24 2019). A few countries such as Germany **have** brought forward **their** carbon neutrality
25 deadline to 2045 (IPCC, 2019). Since most developed **countries** have already achieved
26 carbon peak, they only need to continue their previous greenhouse gases (GHGs)
27 reduction strategies for achieving their carbon neutrality goals, and thus their carbon
28 reduction tasks are relatively easy to realize (IPCC, 2019). Although the total carbon
29 emissions are still increasing, China is committed to **achieving** the goals of carbon peak
30 by 2030 and carbon neutrality by 2060 (Liu et al., 2022). Considering much of China's
31 economic growths in current stage still rely on the high-carbon energy previously
32 implemented by the developed countries, the next 30 years will be a critical period for

1 China to balance its economic development with industrial transformation. During the
2 14th Five-Year Plan stage, China's ecological civilization construction will step into a
3 critical stage for upgrading its ecological environment quality (Zhao et al., 2021). In
4 addition to **continuing** current pollution control policies, this stage will promulgate a
5 series of carbon reduction measures to achieve an initial low-carbon transformation for
6 economic and social development (Yang et al., 2021a).

7 To tackle climate change, it is critical to have creditable information on GHGs with
8 respect to who, which emission sector and how much quantity are responsible for the
9 emissions (Boesch et al., 2021). This information allows assessment of the effectiveness
10 of GHGs mitigation initiatives, strategies, and policies. Especially it allows assessment
11 of how much GHGs reductions are being met at global, national, sector, or even
12 individual point source. It also allows GHGs trading schemes to be functional since
13 such schemes would have no integrity without credible trading units. This credibility
14 determines if buyers and sellers will have confidence in such trading schemes (Boesch
15 et al., 2021). Accurate GHGs information is also crucial for investigating the
16 relationship between global warming and GHGs (Wunch et al., 2010; Wunch et al.,
17 2011). By accurately capturing the diurnal, monthly, seasonal, and inter-annual
18 variabilities of key GHGs, we can speculate their sources and sinks, reveal the physical
19 and chemical mechanisms that drive their variabilities, predict their future trends, and
20 understand how GHGs emissions interact with the atmosphere and how the climate
21 responds to both natural and anthropogenic GHGs emissions (Wunch et al., 2011b). In
22 addition, a creditable GHGs monitoring system could not only promote the
23 investigation of carbon cycle, but also support the development of chemical transport
24 models (CTMs) and emission inventory compilation technology around the globe
25 (MacFaul, 2007; Yang et al., 2020).

26 In order to establish a creditable GHGs monitoring network to support the goals
27 of carbon peak/neutrality in China, it is necessary to know what we have done and what
28 we have to do in the future. In this study, we summarize an overview to the status and
29 perspective of GHGs monitoring in China. There is a very large number of topics and
30 literature related to GHGs, this work cannot be capable to summarize all of them, but
31 will attempt to condense the major information in the field of GHGs monitoring
32 capacity in China. In section 2, we briefly introduce the history of GHGs monitoring
33 around the globe. Section 3 and section 4 summarize the status and typical advances of
34 GHGs monitoring in China. Section 5 discusses main challenges that need to be

addressed for developing creditable GHGs stereoscopic monitoring network in China. In section 6, we present a perspective for future development of GHGs monitoring in China. Section 7 gives the conclusions.

2. History of GHGs monitoring around the globe

The first continuous monitoring of atmospheric GHGs was started at Mauna Loa (155.6°W, 19.5°N) in Hawaii in 1957 (MacFaul, 2007). In situ GHGs measurements at this station were based on nondispersive infrared (NDIR) spectroscopic technology, and were conducted by the National Oceanic and Atmospheric Administration (NOAA), the United States. Subsequently, NOAA expanded such continuous routine monitoring of atmospheric GHGs to Barrow (156.6°W, 71.3°N), American Samoa (170.6°W, 14.2°S), and the South Pole (59.0°E, 90.0°S) (MacFaul, 2007). Long-term time series of GHGs measurements at the four observatories show that global atmospheric CO₂ concentration increased year by year in the past 50 years (Fig. 1). Furthermore, the Global Atmosphere Watch (GAW) network organized by the World Meteorological Organization (WMO) measures atmospheric GHGs from several ground-based and tower-based stations around the globe (Fig. 2). Currently, the GAW only operates one global station (Mt. Waliguan (100.9°E, 36.3°N)) and three regional stations (Lin'an (119.7°E, 30.3°N), Longfengshan (127.6°E, 44.7°N), and Shangdianzi (117.2°E, 40.7°N)) within China (Fang et al., 2014; Fang et al., 2015a; Fang et al., 2015b; Fang et al., 2016). Most GAW ground-based and tower-based stations use commercially available cavity ring-down spectroscopic (CRDS) instruments to achieve high-precision measurements of GHGs (Gomez-Pelaez et al., 2019). Furthermore, GAW also operates many airborne in situ monitoring instrumentations for atmospheric GHGs monitoring around the globe (Fig. 2). These airborne measurement campaigns include the Intercontinental Chemical Transport Experiment–North America campaign (INTEX-NA) and the CO₂ Budget and Rectification Airborne – Maine experiment (COBRA-ME) over the United States during 2004–2005 (Gerbig et al., 2003; Lin et al., 2006; Singh et al., 2006); the Tropical Warm Pool International Cloud Experiment (TWP-ICE) over Australia in 2006; the HIAPER aircraft campaign following the START-08 and HIPPO campaigns in 2008 and 2009; the Beechcraft King Air aircraft campaign over Tsukuba, Japan in 2009, and Learjet overflights over Lamont, the United States in 2009 (Wunch et al., 2010).

Optical remote sensing techniques sampling the total atmospheric column have been developed throughout the last two decades and have been found to be very useful for monitoring atmospheric GHGs (Wunch et al., 2011b). A series of state-of-the-art satellites with different spatiotemporal resolutions, including SCIAMACHY (Schneising et al., 2012; Dils et al., 2014; Houweling et al., 2014; Buchwitz et al., 2015; Heymann et al., 2015; Kulawik et al., 2016) and TROPOMI (Butz et al., 2012; Veeffkind et al., 2012; Pandey et al., 2019; Wang et al., 2020a; Zhang et al., 2020b; Barre et al., 2021; Pandey et al., 2021; Park et al., 2021; Qu et al., 2021; Sha et al., 2021; Shen et al., 2021) by European Space Agency (ESA), GOSAT and GOSAT-2 by Japan (Butz et al., 2011; Morino et al., 2011; Cogan et al., 2012; Yoshida et al., 2013; Deng et al., 2014; Parker et al., 2020; Boesch et al., 2021), OCO-2 and OCO-3 by the United States (Thompson et al., 2012; Frankenberg et al., 2015; Eldering et al., 2017; Nassar et al., 2017; Patra et al., 2017; Wunch et al., 2017; Wang et al., 2020a; Zheng et al., 2020a; Zheng et al., 2020b; Hu and Shi, 2021; Kiel et al., 2021), TanSat (Liu et al., 2013; Liu et al., 2014; Liu et al., 2018; Yang et al., 2018a; Yang et al., 2018b, c; Zhang et al., 2019; Yang et al., 2020d; Bao et al., 2020; Wang et al., 2020c; Yang et al., 2020a; Yang et al., 2021a; Yang et al., 2021b) and Gaofen-5 (GF-5) series satellites by China (Li et al., 2016; Wu et al., 2018; Zhang et al., 2020a; Zhao et al., 2021), GHGSat by Canada (Varon et al., 2019; Jervis et al., 2021), etc., have been launched to derive the global distributions of GHGs. These satellites mainly measure total columns of GHGs by means of infrared grating or Fourier transform infrared (FTIR) spectrometers through atmospheric limb or nadir observations. **SCIAMACHY, GOSAT, OCO-2, and TanSat have XCO₂ precisions of 2.5ppmv, 1–2 ppmv, ~1 ppmv, and 1–4 ppmv, respectively (Reuter et al., 2011; Nassar et al., 2017; Boesch et al., 2021; Yang et al., 2020a).** Studies with satellite data have yielded anthropogenic CO₂ flux estimates at the scale of megacities or larger regions (Eldering et al., 2017), and recently have extended CO₂ emissions estimate at the scale of an individual facility, such as a single power plant (Nassar et al., 2017; Zheng et al., 2020a). **Jacob et al. (2022) have summarized the capability of current and scheduled satellite observations of atmospheric methane in the shortwave infrared (SWIR) to quantify CH₄ emissions from the global scale down to point sources, where XCH₄ precisions of various satellites were presented.**

Ground-based high resolution FTIR spectrometers are powerful tools for deriving total columns and profiles of GHGs (Wunch et al., 2011a). Both the Total Carbon Column Observing Network (TCCON) and the Network for Detection of Atmospheric

Composition Change-Infrared working group (NDACC-IRWG) use high resolution FTIR spectrometers (mainly IFS120HR/IFS125HR series spectrometers manufactured by Bruker, Germany) to observe total columns and profiles of GHGs and atmospheric pollutants (Chevallier et al., 2011; Messerschmidt et al., 2011; Saito et al., 2012; Kuai et al., 2012; Connor et al., 2016; Kiel et al., 2016; Belikov et al., 2017). The TCCON/NDACC-IRWG networks were operating since 2004/1992 and provided time series of many atmospheric constituents, including GHGs such as H₂O, HDO, CO₂, CH₄, CH₃D, N₂O, SF₆, O₃, C₂H₆, CCl₃F, CCl₂F₂, and CHClF₂. For solar zenith angles (SZAs) of less than 80°, the total errors of X_{CO₂}, X_{CH₄}, and X_{N₂O} are less than 0.25% (~1ppmv), 0.5% (~5ppbv), and 1% (~3 ppbv), respectively (Wunch et al., 2011b). These observations have been extensively used in investigations of carbon cycle, carbon source and transport, satellite validation, development of remote sensing algorithm, and evaluation of atmospheric CTMs. Currently, there are only ~30 TCCON/NDACC-IRWG joint stations around the globe, most of them distributed in Europe and Northern America, and the number of stations in other parts of the globe is sparse (Fig. 3). Currently, only two TCCON stations have been set up in China, the Hefei station (117.2°E, 32.0°N) and the Xianghe station (116.96°E, 39.75°N) (Tian et al., 2017; Wang et al., 2017; Yang et al., 2020c).

Despite their outstanding capabilities such as high precision and stability, the high-resolution IFS120/125HR FTIR spectrometers also have their limitations. They are expensive and ponderous spectrometers; their operation relies on a large number of infrastructure and the maintenance for their optical alignments are difficult and time consuming. In order to address these issues, the usage of cheaper, smaller, and more transportable FTIR spectrometers has been investigated in recent few years. These FTIR spectrometers including EM27/SUN and VERTEX-80/SUN manufactured by Bruker, Germany have been verified to have comparable capacity as the IFS125HR with respect to GHGs monitoring. The transportability of the EM27/SUN and VERTEX-80/SUN spectrometers favor campaign use, and many successful campaigns were conducted by various scientists (Hase et al., 2015; Hedelius et al., 2016; Frey et al., 2019; Vogel et al., 2019; Ars et al., 2020; Jacobs et al., 2020; Tu et al., 2020; Frey et al., 2021; Mermigkas et al., 2021). Generally, scientists first use high accuracy of high resolution FTIR dataset to calibrate the EM27/SUN and VERTEX-80/SUN spectrometers, and then use the transportable spectrometers to derive the emission rate of a city, an industrial facility or a landfill. With the transportable EM27/SUN

spectrometers, the Collaborative Carbon Column Observing Network (COCCON) has been built to derive column-averaged abundances of GHGs over the world (Hase et al., 2015; Frey et al., 2019). The EM27/SUN and VERTEX-80/SUN observations can complement the high-resolution FTIR observations around the globe (Table S2).

In addition to the network-based routine observations, there are also many research-oriented GHGs campaigns around the globe, and these uncoordinated behaviors are too numerous to count accurately (Gerbig et al., 2003; Lin et al., 2006; Zellweger et al., 2016; Gomez-Pelaez et al., 2019; Liu et al., 2021a; Liu et al., 2021b). These GHGs measurements are proceeded using by either CRDS, NDIR spectroscopy, off-axis integrated cavity output spectroscopy (OA-ICOS), gas chromatography with flame ionisation detection (GC/FID) technique, FTIR spectroscopy or differential absorption LIDAR (DIAL) implemented on different platforms (Krings et al., 2011; Krings et al., 2013; Zellweger et al., 2016; Krautwurst et al., 2017; Krings et al., 2018; Krautwurst et al., 2021). For example, the airborne MAMAP (Methane Airborne Mapper) spectrometer developed by University of Bremen can be used to derive point source rates of CH₄ and CO₂ (Krings et al., 2011; Krings et al., 2013; Krings et al., 2018). Japanese scientists have developed a grating based optical spectrum analyzer (OSA) and an optical fiber Fabry-Perot interferometer (FFPI) to measure atmospheric CO₂ and CH₄ total columns (Kobayashi et al., 2010).

Overall, the international community has established a series of monitoring networks to measure GHGs on different spatiotemporal scales. Taking the advantage that all GHGs have spectral absorptions in the infrared waveband, most of these networks are established by means of various spectroscopic instruments. These stereoscopic monitoring networks combining the emission inventory compilation and CTMs have formed the state-of-the-art GHGs monitoring and assessment system (Zellweger et al., 2016; Krautwurst et al., 2021), which is extensively used by the United Nations Framework Convention on Climate Change (UNFCCC) to assess GHGs emissions on global, national and regional scales, and identify who, which emission sector and how much quantity are responsible for respective GHGs emissions (IPCC, 2019).

3. Status of GHGs monitoring in China

The global energy consumption data discloses that China has overtaken the United

States in 2006 as the world's top CO₂ producer, i.e., the biggest anthropogenic contributor to global warming (IPCC, 2019). The severity, extension, complexity, and the need-to-cut scale of GHGs emissions in China are unrivaled compared to other countries (Liu et al., 2022). Facing one of the most serious climate change problems around the globe, China has to address a series of scientific, technical, and management issues to achieve the goals of carbon peak/neutrality. As China pays more and more attention to climate change, Chinese government has put a large effort into the development of GHGs monitoring capacity. Although efforts to monitor GHGs in China have hitherto been largely uncoordinated with the established international networks, the GHGs monitoring capacity has been steadily improved. Chinese scientists have conducted many GHGs monitoring studies in urban agglomerations or typical industrial parks in different city clusters such as the Yangtze River Delta (YRD), North China Plain (NCP), Pearl River Delta (PRD) (Tian et al., 2017; Tian et al., 2018; Wang et al., 2019; Li et al., 2021; Liu et al., 2021b), in background areas such as Waliguan in Qinghai Province and Longfengshan in Heilongjiang Province (Fang et al., 2014; Fang et al., 2015a; Fang et al., 2016), and in offshore areas such as the South China Sea, Yellow Sea, and Bohai Bay (Gerbig et al., 2003; Liu et al., 2021a; Liu et al., 2021b). Research institutions, monitoring technologies, monitoring platforms, monitoring scales, monitoring applications, and typical advances can be summarized in Table 1. We elaborate this as follows.

1. **Monitoring technologies include a variety of active and passive measurement technologies**, which mainly include Electrochemical (EC) sensing technology, Tunable Diode Laser Absorption Spectroscopy (TDLAS), Differential Optical Absorption Spectroscopy (DOAS), FTIR, NDIR, GC/FID, LIDAR, CRDS, OA-ICOS, Photoacoustic Spectroscopy (PAS), etc. All these techniques can be classified as spectroscopic technique except EC technology, which uses capacitive readout cantilevers to detect an absorbed signal (Zellweger et al., 2016; Liu et al., 2022). **In most cases, the EC, TDLAS, NDIR, GC/FID, LIDAR, CRDS, OA-ICOS, PAS techniques are commonly used for active measurement platforms, while DOAS and FTIR techniques are extensively used for both active and passive measurement platforms.** Fig.4 illustrates the principles of spectroscopic techniques for GHGs monitoring. The active measurement techniques (a, f) use artificial light source and the passive measurement techniques (b, c, d, e) use natural light sources such as **sunlight** to monitor GHGs. For both active and passive measurement techniques, the absorbed

1 signals can be detected from direct transmission (a, b), surface reflection (e) or
2 atmospheric scattering (c, d, f). Currently, spectroscopic technique is the only
3 technology that can be used to observe global GHGs from space.

4 2. Monitoring platforms include manual sampling analysis, surface in-situ
5 measurement sites (e.g., laboratory measurement or surface monitoring network),
6 ground-based remote sensing platforms (such as ground-based FTIR, LIDAR, and
7 DOAS observatories), tower-based, airborne, space-based (e.g., GF-5 series, TanSat
8 satellites, and space-borne LIDAR), ship-borne, vehicle-borne, unmanned aerial
9 vehicle (UAV), balloon, tethered balloon and other monitoring platforms. Since any of
10 a single monitoring platform cannot fully meet the requirements of stereoscopic
11 monitoring of GHGs emissions due to its limited coverage or spatial resolution,
12 scientists usually integrate a suite of observation platforms to form a stereoscopic
13 monitoring system (Fig. 5). However, most stereoscopic GHGs monitoring activities in
14 China are hitherto research-oriented, temporal, sparse, and uncoordinated with the
15 established international networks (Fang et al., 2014; Tian et al., 2017; Wang et al.,
16 2017; Tian et al., 2018; Yang et al., 2020c; Liu et al., 2021a; Liu et al., 2021b; Sun et
17 al., 2021).

18 3. Monitoring spatial scale ranges from single point source, single constituent,
19 small scale, regional scale to multi-point sources clustered, multi-constituent, large
20 scale, global scale (Zellweger et al., 2016). Depending on monitoring technologies,
21 constituents and platforms, monitoring temporal resolution ranges from second, minute,
22 hour to day levels. Monitoring spatial resolution ranges from meter, dozens of meters,
23 kilometer to dozens of kilometers. Monitoring accuracy ranges from thousandth level
24 to percent level, and monitoring sensitivity ranges from ppbv to ppmv level. Usually
25 the more abundant GHGs tend to produce stronger spectroscopic absorptions, which
26 makes them easier to be separated from background, and thus can be monitored with
27 high sensitivity. Although traditional EC or manual sampling analysis techniques are
28 capable to measure many GHGs with satisfactory accuracy, they usually have limited
29 coverage and can only measure one constituent at a time. In comparison, spectroscopic
30 technology can have a larger coverage, wider monitoring range, more sensitivity, and
31 can continuous real-time monitoring of multi-constituents at a time. Particularly, a
32 single spectroscopic instrument can simultaneously monitor several GHGs without
33 disturbing the samples, i.e., the monitoring process can be completely unattended. As
34 long as an appropriate waveband is selected, the volume mixing ratio (VMR)

concentrations of some GHGs can be measured with a sensitivity of less than 1 ppmv. The coverage can be extended from several meters to several kilometers without multi-point sampling.

4. Monitoring targets include H₂O, HDO, CO₂, CH₄, CH₃D, N₂O, SF₆, O₃, C₂H₆, CCl₃F, CCl₂F₂, and CHClF₂ (Sun et al., 2018b). Monitoring regions include typical industrial zones, industrial stack emissions, urban atmosphere, ambient atmosphere, remote background regions, offshore regions, wetlands, etc. These GHGs measurements with different spatiotemporal **scales** have been extensively used in investigations of global carbon cycle, GHGs trends, regional GHGs sources and transport, ecological GHGs flux estimate, urban or industrial GHGs emissions estimates, validations of CTMs and emission inventory, multi-platform cross calibration, and algorithm improvement, etc (De Maziere et al., 2018; Tian et al., 2018; Sun et al., 2021).

4. Advances in GHGs monitoring in China

With decades of effort, China has made a great breakthrough in GHGs monitoring capacity and steadily improved the performance of homemade GHGs monitoring instruments. Typical advances in GHGs monitoring in China include, but are not limited to, the following aspects.

1. The TanSat and GF-5 series GHGs satellite payloads developed by China have successfully obtained high precision of global CO₂ distributions (Li et al., 2016; Wu et al., 2018; Zhang et al., 2020a; Zhao et al., 2021; **Cai et al., 2022**). The comparisons with the TCCON, GOSAT, and OCO-2 data show that some key performance indicators such as accuracy, precision, and spatiotemporal resolution of these Chinese GHGs satellites have reached the envisaged requirements (Liu et al., 2013; Liu et al., 2014; Cai et al., 2014; Du et al., 2018; Liu et al., 2018; Yang et al., 2018a; Yang et al., 2018b, c; Li et al., 2019; Zhang et al., 2019; Zhao et al., 2019; Bao et al., 2020; Wang et al., 2020c; Yang et al., 2020a, b; Yang et al., 2020d; Boesch et al., 2021; Yang et al., 2021a; Yang et al., 2021b). Both TanSat and GF-5 series GHGs payloads use passive remote sensing technology to derive global CO₂ distributions from scattered sunlight. As a result, they can only work in the daytime and are also seriously influenced by clouds and aerosols. The first Chinese space-borne CO₂ LIDAR onboard the atmospheric environment monitoring satellite launched on April 16, 2022 **used** active remote sensing technology

1 to derive global CO₂ distributions. Its operation does not rely on sunlight and is less
2 influenced by clouds and aerosols, which will greatly improve the global CO₂ mapping
3 capacity.

4 2. A series of in situ online, ground-based, and airborne instruments have been
5 developed by Chinese scientists to investigate the diurnal, monthly, seasonal, and inter-
6 annual variabilities and spatial distributions of key GHGs (Tang et al., 2006), speculate
7 their sources and sinks, and reveal the physical and chemical mechanisms that drive
8 their variabilities. For example, Chinese scientists have developed a suite of in situ
9 spectroscopic instruments to measure surface VMRs and isotope ratios of GHGs in
10 background atmosphere, sea-air CO₂ flux in coastal ocean boundary layer, and soil-air
11 CO₂ flux in farmland (Gerbig et al., 2003; Liu et al., 2021a; Liu et al., 2021b). They
12 have also developed a suite of ground-based spectroscopic instruments for measuring
13 total columns of GHGs (Tian et al., 2018), vehicle-based spectroscopic instruments for
14 industrial GHGs emissions, and airborne spectroscopic instruments for deriving the
15 spatial distributions of CO₂ in the North China Plain (Wang et al., 2019; Shi et al.,
16 2021).

17 3. The ground-based high-resolution FTIR observatory at Hefei has continuously
18 observed the total columns or profiles of H₂O, HDO, CO₂, CH₄, CH₃D, N₂O, SF₆, O₃,
19 C₂H₆, CCl₃F, CCl₂F₂, and CHClF₂ in eastern China since 2014, and has become a
20 national infrastructure for ground-based validation of GF-5 series GHGs satellites and
21 other space-borne instruments (Sun et al., 2018a; Sun et al., 2018b). The ground-based
22 FTIR measurements at the Hefei observatory meet the TCCON quality requirements
23 and this station has been formally accepted as a TCCON site in 2018. Ground-based
24 FTIR CO₂ measurements at the Hefei observatory showed an increasing change rate of
25 (2.71 ± 0.32) % per year between 2015 and 2019 (Fig. 6). A similar ground-based high-
26 resolution FTIR observatory at Xianghe has also passed the TCCON quality inspection
27 and joined the TCCON network in 2021 (Yang et al., 2020c). Furthermore, a few
28 affiliations started to operate the portable EM27/SUN FTIR spectrometers and became
29 COCCON members in recent three years (Frey et al., 2019; Liu et al., 2022; Che et al.,
30 2022a; Che et al., 2022b; Cai et al., 2021).

31 4. Some Chinese scientists have used the commercial in situ instruments such as
32 Picarro or Licor series GHGs analyzers to investigate the spatiotemporal variabilities
33 and emission flux of GHGs in different regions of China (Lin et al., 2006; Tang et al.,
34 2006; Fang et al., 2016; Tian et al., 2018; Yi et al., 2019; Li et al., 2021; Liu et al.,

2021b). With the public accessible OCO-2 satellite data, Chinese scientists have estimated CO₂ anthropogenic emissions of cities and industrial regions in China. The satellite-based CO₂ emissions are generally in good agreement with the MEIC emission inventory values but are more different from the global gridded EDGAR and ODIAC emission datasets (Zheng et al., 2020a; Zheng et al., 2020b). **Most recently, Chinese scientists have used the public accessible OCO-2 satellite observations to quantify CO₂ emissions down to individual point sources such as middle- to large-size coal power plants over China** (Zheng et al., 2020a; Hu and Shi, 2021).

5. Challenges

Governments around the globe are committed to **providing** credible data to support global carbon budget, which promoted the emergence of the state-of-the-art GHGs monitoring technology in developed countries. As the world's top CO₂ producer, China faces both challenges and opportunities. One of the major challenges **is how** we can accurately monitor GHGs emissions under the complex carbon emission scenarios over China. **GHGs emissions in China are complex and diverse (Liu et al., 2022). GHGs concentrations measured at a specific place include both local generation and long-range transport, which occurs not only near the surface but also in upper atmosphere. In addition, China has a complex ecological environment characterized as high aerosol levels, high variability, and compound pollution mixed with many constituents, which poses unprecedented challenges (i.e., increase monitoring uncertainty) to the establishment of GHGs stereoscopic monitoring network in China.** Fortunately, China can learn experience from other countries (IPCC, 2019). Through in-depth cooperation with international community, China is possible to establish a reliable GHGs monitoring network with international credibility.

Accurate knowledge of regional GHGs emissions requires accurate measurements of GHGs variabilities on different spatial scales, including the in-situ "point" concentration reflecting small-scale level, the "column" concentration reflecting mesoscale level, and more importantly, the "profile" concentration reflecting vertical distribution of GHGs. In order to develop creditable GHGs stereoscopic monitoring network in China, some key technical questions need to be solved, which are summarized as follows.

1. In terms of specific in situ monitoring and remote sensing technologies, how can

different monitoring technologies and monitoring platforms learn from and complement each other by means of intensive comparison, verification and optimization?

2. In terms of organization and implementation, how to take full advantages of various monitoring platforms and technologies, make full use of their strengths and avoid their weaknesses, and make concerted efforts to achieve stereoscopic GHGs monitoring for specific carbon source and carbon sink scenarios?

3. In terms of data fusion, how to assimilate multiple datasets collected from different platforms and technologies to generate a new uniform dataset that have better coverage than the original dataset without reducing their accuracy, which would improve our understanding for carbon cycle mechanism and promote the development of GHGs forecasting model?

6. Future perspectives

Although much of China's economic growths in current stage still rely on the high-carbon energy previously implemented by the developed countries, China is committed to **achieving** the goals of carbon peak by 2030 and carbon neutrality by 2060. In order to support the formulation of low-carbon policies for achieving the goals of carbon peak/neutrality, China should improve its GHGs monitoring capability as soon as possible. **It is suggested to take full advantages of various monitoring technologies, monitoring platforms, numerical simulations, and inventory compilation techniques to form a creditable GHGs stereoscopic monitoring and assessment (M&A) system. Implementation of this M&A system should be coordinated with the established international networks, and routinely quantify GHGs on global, national, provincial, regional, and individual point scales with high spatiotemporal resolution and wide coverage. Improved knowledge of carbon emissions on different scales is very useful for adjustment of low-carbon policy in China.** In view of status, advances, and challenges of China's GHGs monitoring, future developments are expected to focus on the following aspects.

1. The development of high-end GHGs monitoring technology, instrument and core components should be strengthened to improve GHGs monitoring capacity in China. Development priorities include intelligent and miniaturized instruments dedicated for profile and flux of GHGs within multi-sphere ecological environment,

1 and key optical components such as high-resolution spectrometer, light source, solid
2 laser, high-reflectivity mirror, narrow-band filter, detector, etc. It is expected that
3 homemade GHGs monitoring instruments can meet routine GHGs monitoring demands
4 in China in near future.

5 2. It is suggested to routinely monitor GHGs over typical GHGs sources and
6 atmospheric background regions, which favors the verification of GHGs emission
7 inventory and the implement of nationwide carbon trading. At present, the number of
8 GHGs monitoring sites in China remains sparse. The rural regions are rarely covered
9 and there are only few monitoring stations located in western China. GHGs are not
10 included in the atmospheric constituents routinely monitored by the China National
11 Environmental Monitoring Center (CNEMC) network. It is suggested to include key
12 GHGs into China's surface environmental quality monitoring network, which will
13 improve China's GHGs monitoring capacity.

14 3. Since roughly 70% of the Earth is shrouded by clouds at any given moment and
15 GHGs monitoring from space is prone to cloud interference, a single satellite can only
16 provide a small number of observations per year suitable for emission estimates for any
17 given GHGs source. It is necessary to routinely monitor GHGs over China with satellite
18 constellation, which can offer better spatiotemporal resolution and coverage compared
19 to a single satellite alone. Only with high spatiotemporal resolution and coverage, we
20 can routinely quantify GHGs on global, national, provincial, regional, and individual
21 point scales to support adjustment of low-carbon policy.

22 4. Monitoring data quality control, multi-source data fusion, and data sharing
23 platforms should be systematized and standardized. By standardizing data quality
24 control and data fusion technology of multi-source GHGs metadata and establishing a
25 systematic data sharing mechanism, the metadata can be eventually applied in carbon
26 reduction governance and decision-making by management departments.

27 5. It is suggested to establish an inter-departmental management agency for GHGs
28 monitoring in China, where the government serves as the leader, and the technology
29 holders and expert communities are participating. Furthermore, it is of great
30 significance to unite environmental protection industry association of China, Chinese
31 association of environmental science, research institutions, universities, enterprises and
32 other professional communities to build a uniform verification standard for GHGs
33 monitoring, which should be in cooperation with international networks/partners to
34 achieve common global criteria. This verification standard can not only standardize

1 GHGs monitoring technology but also disclose its verification criteria and process,
2 which can promote the **implementation** of new GHGs monitoring technology in China.

3 **7. Conclusions**

4 GHGs monitoring capability in China has achieved rapid improvement in recent
5 years. Relying on homemade technologies and instruments, combined with public
6 accessible space-borne observation instruments and open source remote sensing
7 algorithms, China has conducted a suite of GHGs stereoscopic monitoring studies in
8 different regions of China, but most of them are hitherto research-oriented, temporal,
9 sparse, and uncoordinated with the established international networks. Some key
10 technical indicators such as spatiotemporal resolution, coverage and accuracy need to
11 be further improved. Furthermore, monitoring data quality control, multi-source data
12 fusion, and data sharing platform have not been standardized.

13 In order to support the formulation of green economic policies for achieving the
14 goals of carbon peak/neutrality, China should improve its GHGs monitoring capability
15 as soon. It is suggested to take full advantages of various monitoring technologies,
16 monitoring platforms, numerical simulations, and inventory compilation techniques to
17 form a creditable GHGs stereoscopic monitoring and assessment (M&A) system.
18 Implementation of this M&A system should be coordinated with the established
19 international networks, and routinely quantify GHGs on global, national, provincial,
20 regional, and individual point scales with high spatiotemporal resolution and wide
21 coverage. Improved knowledge of carbon emissions on different scales is very useful
22 for adjustment of low-carbon policy in China.

23 **Data availability.** The geolocations of all sites coordinated by NOAA and WMO, and
24 all FTIR sites coordinated by TCCON, NDACC-IRWG, and COCCON are
25 summarized in the supplement. All other data are available on request of YS
26 (ywsun@aiofm.ac.cn).

27 **Author contributions.** CL designed the study, YS wrote the paper, HY, WW, CS, JN,
28 MP and ZC improved the content.

29 **Competing interests.** None.

30 **Acknowledgements.** This work is supported by the Youth Innovation Promotion

1 Association, CAS (No.2019434).

2 **References**

- 3 Ars, S., Vogel, F., Arrowsmith, C., Heerah, S., Knuckey, E., Lavoie, J., Lee, C., Pak, N.
4 M., Phillips, J. L., and Wunch, D.: Investigation of the Spatial Distribution of
5 Methane Sources in the Greater Toronto Area Using Mobile Gas Monitoring
6 Systems. *Environment Science & Technology*, 54(24): 15671-15679, 2020.
- 7 Bao, Z. Y., Zhang, X. Y., Yue, T. X., Zhang, L. L., Wang, Z., Jiao, Y. M., Bai, W. G.,
8 Meng, X. Y.: Retrieval and Validation of XCO₂ from TanSat Target Mode
9 Observations in Beijing. *Remote Sensing*, 12(18): 3063, 2020.
- 10 Barre, J., Aben, I., Agusti-Panareda, A., Balsamo, G., Bousserez, N., Dueben, P.,
11 Engelen, R., Inness, A., Lorente, A., McNorton, J., Peuch, V. H., Radnoti, G.,
12 Ribas, R.: Systematic detection of local CH₄ anomalies by combining satellite
13 measurements with high-resolution forecasts, *Atmospheric Chemistry and*
14 *Physics*, 21(6): 5117-5136, 2021.
- 15 Belikov, D. A., Maksyutov, S., Ganshin, A., Zhuravlev, R., Deutscher, N. M., Wunch,
16 D., Feist, D. G., Morino, I., Parker, R. J., Strong, K., Yoshida, Y., Bril, A.,
17 Oshchepkov, S., Boesch, H., Dubey, M. K., Griffith, D., Hewson, W., Kivi, R.,
18 Mendonca, J., Notholt, J., Schneider, M., Sussmann, R., Velazco, V. A., Aoki, S.:
19 Study of the footprints of short-term variation in XCO₂ observed by TCCON sites
20 using NIES and FLEXPART atmospheric transport models. *Atmospheric*
21 *Chemistry and Physics*, 17(1): 143-157, 2017.
- 22 Boesch, H., Liu, Y., Tamminen, J., Yang, D. X., Palmer, P. I., Lindqvist, H., Cai, Z. N.,
23 Che, K., Di Noia, A., Feng, L., Hakkarainen, J., Ialongo, I., Kalaitzi, N.,
24 Karppinen, T., Kivi, R., Kivimäki, E., Parker, R. J., Preval, S., Wang, J., Webb, A.
25 J., Yao, L., Chen, H. L.: Monitoring Greenhouse Gases from Space. *Remote*
26 *Sensing*, 13(14): 2700, 2021.
- 27 Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I.,
28 Armante, R., Bergamaschi, P., Blumenstock, T., Bovensmann, H., Brunner, D.,
29 Buchmann, B., Burrows, J. P., Butz, A., Chedin, A., Chevallier, F., Crevoisier, C.
30 D., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J.,
31 Kaminski, T., Laeng, A., Lichtenberg, G., De Maziere, M., Noel, S., Notholt, J.,
32 Orphal, J., Popp, C., Parker, R., Scholze, M., Sussmann, R., Stiller, G. P., Warneke,

1 T., Zehner, C., Bril, A., Crisp, D., Griffith, D. W. T., Kuze, A., O'Dell, C.,
2 Oshchepkov, S., Sherlock, V., Suto, H., Wennberg, P., Wunch, D., Yokota, T.,
3 Yoshida, Y.: The Greenhouse Gas Climate Change Initiative (GHG-CCI):
4 Comparison and quality assessment of near-surface-sensitive satellite-derived
5 CO₂ and CH₄ global data sets. *Remote Sensing of Environment*, 162: 344-362,
6 2015.

7 Butz, A., Galli, A., Hasekamp, O., Landgraf, J., Tol, P., Aben, I.: TROPOMI aboard
8 Sentinel-5 Precursor: Prospective performance of CH₄ retrievals for aerosol and
9 cirrus loaded atmospheres. *Remote Sensing of Environment*, 120: 267-276, 2012.

10 Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C.,
11 Hartmann, J. M., Tran, H., Kuze, A., Keppel-Aleks, G., Toon, G., Wunch, D.,
12 Wennberg, P., Deutscher, N., Griffith, D., Macatangay, R., Messerschmidt, J.,
13 Notholt, J., and Warneke, T.: Toward accurate CO₂ and CH₄ observations from
14 GOSAT. *Geophysical Research Letters*, 38: L14812, 2011.

15 Cai, Z. N., Liu, Y., and Yang, D. X.: Analysis of XCO₂ retrieval sensitivity using
16 simulated Chinese Carbon Satellite (TanSat) measurements. *Science China Earth*
17 *Science*, 57(8): 1919-1928, 2014.

18 Liu, C., Sun, Y. W., Shan, C. G., Wang, W., Notholt, J., Palm, M., Yin, H., Tian, Y.,
19 Gao, J. X., Mao, H. Q.: Long-term observations of atmospheric constituents at the
20 first ground-based high resolution Fourier transform spectrometry observation
21 station in China. *Engineering*, In Press, 10.1016/j.eng.2021.11.022, 2022.

22 Chevallier, F., Deutscher, N. M., Conway, T. J., Ciais, P., Ciattaglia, L., Dohe, S.,
23 Frohlich, M., Gomez-Pelaez, A. J., Griffith, D., Hase, F., Haszpra, L., Krummel,
24 P., Kyro, E., Labuschagne, C., Langenfelds, R., Machida, T., Maignan, F.,
25 Matsueda, H., Morino, I., Notholt, J., Ramonet, M., Sawa, Y., Schmidt, M.,
26 Sherlock, V., Steele, P., Strong, K., Susmann, R., Wennberg, P., Wofsy, S., Worthy,
27 D., Wunch, D., and Zimnoch, M.: Global CO₂ fluxes inferred from surface air-
28 sample measurements and from TCCON retrievals of the CO₂ total column.
29 *Geophysical Research Letters*, 38: L24810, 2011.

30 Cogan, A. J., Boesch, H., Parker, R. J., Feng, L., Palmer, P. I., Blavier, J. F. L.,
31 Deutscher, N. M., Macatangay, R., Notholt, J., Roehl, C., Warneke, T., and Wunch,
32 D.: Atmospheric carbon dioxide retrieved from the Greenhouse gases Observing
33 SATellite (GOSAT): Comparison with ground-based TCCON observations and
34 GEOS-Chem model calculations. *Journal of Geophysical Research-Atmosphere*,

117: D21301, 2012.

Connor, B. J., Sherlock, V., Toon, G., Wunch, D., Wennberg, P. O.: GFIT2: an experimental algorithm for vertical profile retrieval from near-IR spectra. *Atmospheric Measurement Techniques*, 9(8): 3513-3525, 2016.

Cai, Z.; Sun, K.; Yang, D.; Liu, Y.; Yao, L.; Lin, C.; Liu, X. On-Orbit Characterization of TanSat Instrument Line Shape Using Observed Solar Spectra. *Remote Sens.* 2022, 14, 3334. <https://doi.org/10.3390/rs14143334>.

Cai, Z., Che, K., Liu, Y., Yang, D., Liu, C., Yue, X. Decreased Anthropogenic CO₂ Emissions during the COVID-19 Pandemic Estimated from FTS and MAX-DOAS Measurements at Urban Beijing. *Remote Sens.* 2021, 13, 517. <https://doi.org/10.3390/rs13030517>.

Che, K., Liu, Y., Cai, Z. N., Yang, D. X., Wang, H. B., Ji, D. H., Yang, Y., and Wang, P. C.: Characterization of regional combustion efficiency using $\Delta XCO:\Delta XCO_2$ observed by a portable Fourier Transform spectrometer at an urban site in Beijing, 5, doi: 10.1007/s00376-022-1247-7, 2022.

Che K., Cai Z., Liu Y., Wu L., Yang D., Chen Y., Meng X., Zhou M., Wang J., Yao L. and Wang P.. Lagrangian inversion of anthropogenic CO₂ emissions from Beijing using differential column measurements, *Environmental Research Letters*, 17(7): 075001, <https://doi.org/10.1088/1748-9326/ac7477>, 2022.

De Maziere, M., Thompson, A. M., Kurylo, M. J., Wild, J. D., Bernhard, G., Blumenstock, T., Braathen, G. O., Hannigan, J. W., Lambert, J. C., Leblanc, T., Mcgee, T. J., Nedoluha, G., Petropavlovskikh, I., Seckmeyer, G., Simon, P. C., Steinbrecht, W., Strahan, S. E.: The Network for the Detection of Atmospheric Composition Change (NDACC): history, status and perspectives. *Atmospheric Chemistry and Physics*, 18(7): 4935-4964, 2018.

Deng, F., Jones, D. B. A., Henze, D. K., Bousserez, N., Bowman, K.W., Fisher, J. B., Nassar, R., O'Dell, C., Wunch, D., Wennberg, P. O., Kort, E. A., Wofsy, S. C., Blumenstock, T., Deutscher, N. M., Griffith, D. W. T., Hase, F., Heikkinen, P., Sherlock, V., Strong, K., Sussmann, R., Warneke, T.: Inferring regional sources and sinks of atmospheric CO₂ from GOSAT XCO₂ data, *Atmospheric Chemistry and Physics*, 14(7): 3703-3727, 2014.

Dils, B., Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Parker, R., Guerlet, S., Aben, I., Blumenstock, T., Burrows, J. P., Butz, A., Deutscher, N. M., Frankenberg,

C., Hase, F., Hasekamp, O. P., Heymann, J., De Maziere, M., Notholt, J., Sussmann, R., Warneke, T., Griffith, D., Sherlock, V., Wunch, D.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON, *Atmospheric Measurement Techniques*, 7(6): 1723-1744, 2014.

Du, S. S., Liu, L. Y., Liu, X. J., Zhang, X., Zhang, X. Y., Bi, Y. M., and Zhang, L. C.: Retrieval of global terrestrial solar-induced chlorophyll fluorescence from TanSat satellite. *Science Bulletin*, 63(22): 1502-1512, 2018.

Eldering, A., Wennberg, P. O., Crisp, D., Schimel, D. S., Gunson, M. R., Chatterjee, A., Liu, J., Schwandner, F. M., Sun, Y., O'Dell, C. W., Frankenberg, C., Taylor, T., Fisher, B., Osterman, G. B., Wunch, D., Hakkarainen, J., Tamminen, J., Weir, B.: The Orbiting Carbon Observatory-2 early science investigations of regional carbon dioxide fluxes. *Science*, 358(6360): eaam5745, 2017.

Fang, S. X., Luan, T., Zhang, G., Wu, Y. L., and Yu, D. J.: The determination of regional CO₂ mole fractions at the Longfengshan WMO/GAW station: A comparison of four data filtering approaches. *Atmospheric Environment*, 116:36-43, 2015a.

Fang, S. X., Tans, P. P., Steinbacher, M., Zhou, L. X., Luan, T.: Comparison of the regional CO₂ mole fraction filtering approaches at a WMO/GAW regional station in China. *Atmospheric Measurement Techniques*, 8(12): 5301-5313, 2015b.

Fang, S. X., Tans, P. P., Steinbacher, M., Zhou, L. X., Luan, T., and Li, Z.: Observation of atmospheric CO₂ and CO at Shangri-La station: results from the only regional station located at southwestern China, *Tellus B.*, 68(1): 28506, 2016.

Fang, S. X., Zhou, L. X., Tans, P. P., Ciais, P., Steinbacher, M., Xu, L., Luan, T.: In situ measurement of atmospheric CO₂ at the four WMO/GAW stations in China. *Atmospheric Chemistry and Physics*, 14(5): 2541-2554, 2014.

Frankenberg, C., Pollock, R., Lee, R. A. M., Rosenberg, R., Blavier, J. F., Crisp, D., O'Dell, C. W., Osterman, G. B., Roehl, C., Wennberg, P. O., and Wunch, D.: The Orbiting Carbon Observatory (OCO-2): spectrometer performance evaluation using pre-launch direct sun measurements, *Atmospheric Measurement Techniques* 8(1): 301-313, 2015.

Frey, M., Sha, M. K., Hase, F., Kiel, M., Blumenstock, T., Harig, R., Surawicz, G., Deutscher, N. M., Shiomi, K., Franklin, J. E., Bosch, H., Chen, J., Grutter, M., Ohyama, H., Sun, Y. W., Butz, A., Tsidu, G. M., Ene, D., Wunch, D., Cao, Z. S.,

- Garcia, O., Ramonet, M., Vogel, F., and Orphal, J.: Building the COllaborative Carbon Column Observing Network (COCCON): long-term stability and ensemble performance of the EM27/SUN Fourier transform spectrometer. *Atmospheric Measurement Techniques*, 12(3): 1513-1530, 2019.
- Frey, M. M., Hase, F., Blumenstock, T., Dubravica, D., Gross, J., Gottsche, F., Handjaba, M., Amadhila, P., Mushi, R., Morino, I., Shiomi, K., Sha, M. K., De Maziere, M., and Pollard, D. F.: Long-term column-averaged greenhouse gas observations using a COCCON spectrometer at the high-surface-albedo site in Gobabeb, Namibia, *Atmospheric Measurement Techniques*, 14(9): 5887-5911, 2011.
- Gerbig, C., Lin, J. C., Wofsy, S. C., Daube, B. C., Andrews, A. E., Stephens, B. B., Bakwin, P. S., Grainger, C. A.: Toward constraining regional-scale fluxes of CO₂ with atmospheric observations over a continent: 2. Analysis of COBRA data using a receptor-oriented framework, *Journal of Geophysical Research-Atmosphere*, 108(D24): 4756, 2003.
- Gomez-Pelaez, A. J., Ramos, R., Cuevas, E., Gomez-Trueba, V., Reyes, E.: Atmospheric CO₂, CH₄, and CO with the CRDS technique at the Izana Global GAW station: instrumental tests, developments, and first measurement results, *Atmospheric Measurement Techniques*, 12(4): 2043-2066, 2019.
- Hase, F., Frey, M., Blumenstock, T., Gross, J., Kiel, M., Kohlhepp, R., Tsidu, G. M., Schafer, K., Sha, M. K., Orphal, J.: Application of portable FTIR spectrometers for detecting greenhouse gas emissions of the major city Berlin, *Atmospheric Measurement Techniques*, 8(7): 3059-3068, 2015.
- Hedelius, J. K., Viatte, C., Wunch, D., Roehl, C. M., Toon, G. C., Chen, J., Jones, T., Wofsy, S. C., Franklin, J. E., Parker, H., Dubey, M. K., Wennberg, P. O.: Assessment of errors and biases in retrievals of XCO₂, XCH₄, XCO., and XN₂O from a 0.5 cm⁻¹ resolution solar-viewing spectrometer, *Atmospheric Measurement Techniques*, 9(8): 3527-3546, 2016.
- Heymann, J., Reuter, M., Hilker, M., Buchwitz, M., Schneising, O., Bovensmann, H., Burrows, J. P., Kuze, A., Suto, H., Deutscher, N. M., Dubey, M. K., Griffith, D. W. T., Hase, F., Kawakami, S., Kivi, R., Morino, I., Petri, C., Roehl, C., Schneider, M., Sherlock, V., Sussmann, R., Velasco, V. A., Warneke, T., and Wunch, D.: Consistent satellite XCO₂ retrievals from SCIAMACHY and GOSAT using the BESD algorithm, *Atmospheric Measurement Techniques*, 8(7): 2961-2980, 2015.

- 1 Houweling, S., Krol, M., Bergamaschi, P., Frankenberg, C., Dlugokencky, E. J.,
2 Morino, I., Notholt, J., Sherlock, V., Wunch, D., Beck, V., Gerbig, C., Chen, H.,
3 Kort, E. A., Rockmann, T., and Aben I.: A multi-year methane inversion using
4 SCIAMACHY, accounting for systematic errors using TCCON measurements.
5 *Atmospheric Chemistry and Physics*, 14(8): 3991-4012, 2014.
- 6 Hu, Y. Q. and Shi, Y. S.: Estimating CO₂ Emissions from Large Scale Coal-Fired Power
7 Plants Using OCO-2 Observations and Emission Inventories. *Atmosphere*, 12(7):
8 811, 2021.
- 9 IPCC (Intergovernmental Panel on Climate Change): 2019 Refinement to the 2006
10 IPCC Guidelines for National Greenhouse Gas Inventories, 2019.
- 11 Jacobs, N., Simpson, W. R., Wunch, D., O'Dell, C. W., Osterman, G. B., Hase, F.,
12 Blumenstock, T., Tu, Q. S., Frey, M., Dubey, M. K., Parker, H. A., Kivi, R.,
13 Heikkinen, P.: Quality controls, bias, and seasonality of CO₂ columns in the boreal
14 forest with Orbiting Carbon Observatory-2, Total Carbon Column Observing
15 Network and EM27/SUN measurements, *Atmospheric Measurement Techniques*,
16 13(9): 5033-5063, 2020.
- 17 Jacob, D. J., Varon, D. J., Cusworth, D. H., Dennison, P. E., Frankenberg, C., Gautam,
18 R., Guanter, L., Kelley, J., McKeever, J., Ott, L. E., Poulter, B., Qu, Z., Thorpe, A.
19 K., Worden, J. R., and Duren, R. M.: Quantifying methane emissions from the
20 global scale down to point sources using satellite observations of atmospheric
21 methane, *Atmos. Chem. Phys.*, 22, 9617–9646, [https://doi.org/10.5194/acp-22-](https://doi.org/10.5194/acp-22-9617-2022)
22 [9617-2022](https://doi.org/10.5194/acp-22-9617-2022), 2022.
- 23 Jervis, D., McKeever, J., Durak, B. O. A., Sloan, J. J., Gains, D., Varon, D. J., Ramier,
24 A., Strupler, M., and Tarrant, E.: The GHGSat-D imaging spectrometer.
25 *Atmospheric Measurement Techniques*, 14(3): 2127-2140, 2021.
- 26 Kiel, M., Eldering, A., Roten, D. D., Lin, J. C., Feng, S., Lei, R. X., Lauvaux, T., Oda,
27 T., Roehl, C. M., Blavier, J. F., Iraci, L. T.: Urban-focused satellite CO₂
28 observations from the Orbiting Carbon Observatory-3: A first look at the Los
29 Angeles megacity, *Remote Sensing of Environment*, 258: 112314, 2021.
- 30 Kiel, M., Wunch, D., Wennberg, P. O., Toon, G. C., Hase, F., Blumenstock, T.:
31 Improved retrieval of gas abundances from near-infrared solar FTIR spectra
32 measured at the Karlsruhe TCCON station, *Atmospheric Measurement*
33 *Techniques*, 9(2): 669-682, 2016.
- 34 Kobayashi, N., Inoue, G., Kawasaki, M., Yoshioka, H., Minomura, M., Murata, I.,

- 1 Nagahama, T., Matsumi, Y., Tanaka, T., Morino, I., and Ibuki, T.: Remotely
2 operable compact instruments for measuring atmospheric CO₂ and CH₄ column
3 densities at surface monitoring sites, *Atmospheric Measurement Techniques*, 3(4):
4 1103-1112, 2010.
- 5 Krautwurst, S., Gerilowski, K., Borchardt, J., Wildmann, N., Galkowski, M., Swolkien,
6 J., Marshall, J., Fiehn, A., Roiger, A., Ruhtz, T., Gerbig, C., Necki, J., Burrows, J.
7 P., Fix, A., Bovensmann, H.: Quantification of CH₄ coal mining emissions in
8 Upper Silesia by passive airborne remote sensing observations with the Methane
9 Airborne MAPper (MAMAP) instrument during the CO₂ and Methane (CoMet)
10 campaign, *Atmospheric Chemistry and Physics*, 21(23): 17345-17371, 2021.
- 11 Krautwurst, S., Gerilowski, K., Jonsson, H. H., Thompson, D. R., Kolyer, R. W., Iraci,
12 L. T., Thorpe, A. K., Horstjann, M., Eastwood, M., Leifer, I., Vigil, S. A., Krings,
13 T., Borchardt, J., Buchwitz, M., Fladeland, M. M., Burrows, J. P., and
14 Bovensmann, H.: Methane emissions from a Californian landfill, determined from
15 airborne remote sensing and in situ measurements, *Atmospheric Measurement*
16 *Techniques*, 10(9): 3429-3452, 2017.
- 17 Krings, T., Gerilowski, K., Buchwitz, M., Hartmann, J., Sachs, T., Erzinger, J.,
18 Burrows, J. P., Bovensmann, H.: Quantification of methane emission rates from
19 coal mine ventilation shafts using airborne remote sensing data, *Atmospheric*
20 *Measurement Techniques*, 6(1): 151-166, 2013.
- 21 Krings, T., Gerilowski, K., Buchwitz, M., Reuter, M., Tretner, A., Erzinger, J., Heinze,
22 D., Pfluger, U., Burrows, J. P., and Bovensmann, H.: MAMAP - a new
23 spectrometer system for column-averaged methane and carbon dioxide
24 observations from aircraft: retrieval algorithm and first inversions for point source
25 emission rates, *Atmospheric Measurement Techniques*, 4(9): 1735-1758, 2011.
- 26 Krings, T., Neininger, B., Gerilowski, K., Krautwurst, S., Buchwitz, M., Burrows, J. P.,
27 Lindemann, C., Ruhtz, T., Schuttemeyer, D., and Bovensmann, H.: Airborne
28 remote sensing and in situ measurements of atmospheric CO₂ to quantify point
29 source emissions, *Atmospheric Measurement Techniques*, 11(2): 721-739, 2018.
- 30 Kuai, L., Wunch, D., Shia, R. L., Connor, B., Miller, C., and Yung, Y.: Vertically
31 constrained CO₂ retrievals from TCCON measurements, *Journal of Quantitative*
32 *Spectroscopy & Radiative Transfer*, 113(14): 1753-1761, 2012.
- 33 Kulawik, S., Wunch, D., O'Dell, C., Frankenberg, C., Reuter, M., Oda, T., Chevallier,
34 F., Sherlock, V., Buchwitz, M., Osterman, G., Miller, C. E., Wennberg, P. O.,

- 1 Griffith, D., Morino, I., Dubey, M. K., Deutscher, N. M., Notholt, J., Hase, F.,
- 2 Warneke, T., Sussmann, R., Robinson, J., Strong, K., Schneider, M., De Maziere,
- 3 M., Shiomi, K., Feist, D. G., Iraci, L. T., Wolf, J. : Consistent evaluation of ACOS-
- 4 GOSAT, BESD-SCIAMACHY, CarbonTracker, and MACC through comparisons
- 5 to TCCON Atmospheric Measurement Techniques, 9(2): 683-709, 2016.
- 6 Li, M., Feng, K. S., Zheng, H. R., Pan, C., Meng, J., Li, J. S., Guan, D. B., Li, Y.: Supply
- 7 chain effects of China's fast growing marine economy on greenhouse gas
- 8 emissions, Environment Research Letter, 16: 054061, 2021.
- 9 Li, S., Li, C. L., Xu, S. Y., Zhang, H., and Zheng, Y. Q.: Preflight radiometric calibration
- 10 of a carbon dioxide spectrometer. Measurement Science Technology, 30(5):
- 11 055401, 2019.
- 12 Li, Y. F., Zhang, C. M., Liu, D. D., Chen, J., Rong, P., Zhang, X. Y., Wang, S. P.: CO₂
- 13 retrieval model and analysis in short-wave infrared spectrum. Optik, 127(10):
- 14 4422-4425, 2016.
- 15 Lin, J. C., Gerbig, C., Wofsy, S. C., Daube, B. C., Matross, D. M., Chow, V. Y., Gottlieb,
- 16 E., Andrews, A. E., Pathmathevan, M., Munger, J. W.: What have we learned from
- 17 intensive atmospheric sampling field programmes of CO₂? Tellus B., 58:331-343,
- 18 2006.
- 19 Liu, S., Fang, S. X., Liu, P., Liang, M., Guo, M. R., and Feng, Z. Z.: Measurement
- 20 report: Changing characteristics of atmospheric CH₄ in the Tibetan Plateau:
- 21 records from 1994 to 2019 at the Mount Waliguan station, Atmospheric Chemistry
- 22 and Physics, 21(1): 393-413, 2021a.
- 23 Liu S., Feng Z. Z., Lin H. W., Liu P., Liang M., Qing X. M., Xiong H. Y., Qiu S. S., Li,
- 24 J. X., Jiang K., Hong H. X., and Fang S. X.: Changes of Atmospheric CO₂ in the
- 25 Tibetan Plateau From 1994 to 2019, Journal of Geophysical Research-
- 26 Atmosphere, 126(20): e2021JD035299, 2021b.
- 27 Liu, Y., Cai, Z. N., Yang, D. X., Zheng, Y. Q., Duan, M. Z., and Lu, D. R.: Effects of
- 28 spectral sampling rate and range of CO₂ absorption bands on XCO₂ retrieval from
- 29 TanSat hyperspectral spectrometer. Chinese Science Bulletin, 59(14): 1485-1491,
- 30 2014a.
- 31 Liu, Y., Wang, J., Yao, L., Chen, X., Cai, Z. N., Yang, D. X., Yin, Z. S., Gu, S. Y., Tian,
- 32 L. F., Lu, N. M., Lyu, D. R.: The TanSat mission: preliminary global observations,
- 33 Science Bulletin, 63(18): 1200-1207, 2018.
- 34 Liu, Y., Yang, D. X., and Cai, Z. N.: A retrieval algorithm for TanSat XCO₂ observation:

1 Retrieval experiments using GOSAT data, Chinese Science Bulletin, 58(13): 1520-
2 1523, 2013.

3 MacFaul, L.: Monitoring greenhouse gases, Bulletin of America Meteorological
4 Society, 88(8): 1171-1171, 2007.

5 Mermigkas, M., Topaloglou, C., Balis, D., Koukouli, M. E., Hase, F., Dubravica, D.,
6 Borsdorff, T., and Lorente, A.: FTIR Measurements of Greenhouse Gases over
7 Thessaloniki, Greece in the Framework of COCCON and Comparison with
8 S5P/TROPOMI Observations, Remote Sensing, 13(17): 3395, 2021.

9 Messerschmidt, J., Geibel, M. C., Blumenstock, T., Chen, H., Deutscher, N. M., Engel,
10 A., Feist, D. G., Gerbig, C., Gisi, M., Hase, F., Katrynski, K., Kolle, O., Lavric, J.
11 V., Notholt, J., Palm, M., Ramonet, M., Rettinger, M., Schmidt, M., Sussmann, R.,
12 Toon, G. C., Truong, F., Warneke, T., Wennberg, P. O., Wunch, D., and Xueref-
13 Remy, I.: Calibration of TCCON column-averaged CO₂: the first aircraft campaign
14 over European TCCON sites. Atmospheric Chemistry and Physics, 11(21): 10765-
15 10777, 2011.

16 Morino, I. , Uchino, O., Inoue, M., Yoshida, Y., Yokota, T., Wennberg, P. O., Toon, G.
17 C., Wunch, D., Roehl, C. M., Notholt, J., Warneke, T., Messerschmidt, J., Griffith,
18 D. W. T., Deutscher, N. M., Sherlock, V., Connor, B., Robinson, J., Sussmann, R.,
19 and Rettinger, M.: Preliminary validation of column-averaged volume mixing
20 ratios of carbon dioxide and methane retrieved from GOSAT short-wavelength
21 infrared spectra, Atmospheric Measurement Techniques, 4(6): 1061-1076, 2011.

22 Nassar, R., Hill, T. G., McLinden, C. A., Wunch, D., Jones, D. B. A., and Crisp, D.:
23 Quantifying CO₂ Emissions From Individual Power Plants From Space.
24 Geophysical Research Letters, 44(19): 10045-10053, 2017.

25 Pandey, S., Gautam, R., Houweling, S., van der Gon, H. D., Sadavarte, P., Borsdorff,
26 T., Hasekamp, O., Landgraf, J., Tol, P., van Kempen, T., Hoogeveen, R., van Hees,
27 R., Hamburg, S. P., Maasakkers, J. D., and Aben, I.: Satellite observations reveal
28 extreme methane leakage from a natural gas well blowout. Proceedings of the
29 National Academy of Sciences, 116(52): 26376-26381, 2019.

30 Pandey, S., Houweling, S., Lorente, A., Borsdorff, T., Tsvilidou, M., Bloom, A. A.,
31 Poulter, B., Zhang, Z., and Aben, I.: Using satellite data to identify the methane
32 emission controls of South Sudan's wetlands. Biogeosciences, 18(2): 557-572,
33 2021.

- 1 Park, H., Jeong, S., Park, H., Labzovskii, L. D., and Bowman, K. W.: An assessment of
2 emission characteristics of Northern Hemisphere cities using spaceborne
3 observations of CO₂, CO., and NO₂. *Remote Sensing of Environment*, 254:
4 112246, 2021.
- 5 Parker, R. J., Webb, A., Boesch, H., Somkuti, P., Guillo, R. B., Di Noia, A., Kalaitzi,
6 N., Anand, J., S., Bergamaschi, P., Chevallier, F., Palmer, P. I., Feng, L., Deutscher,
7 N. M., Feist, D. G., Griffith, D. W. T., Hase, F., Kivi, R., Morino, I., Notholt, J.,
8 Oh, Y. S., Ohyama, H., Petri, C., Pollard, D. F., Roehl, C., Sha, M. K., Shiomi, K.,
9 Strong, K., Sussmann, R., Te, Y., Velazco, V. A., Warneke, T., Wennberg, P. O.,
10 and Wunch, D.: A decade of GOSAT Proxy satellite CH₄ observations. *Earth*
11 *System Science Data*, 12(4): 3383-3412, 2020.
- 12 Patra, P. K., Crisp, D., Kaiser, J. W., Wunch, D., Sacki, T., Ichii, K., Sekiya, T.,
13 Wennberg, P. O., Feist, D. G., Pollard, D. F., Griffith, D. W. T., Velazco, V. A., De
14 Maziere, M., Sha, M. K., Roehl, C., Chatterjee, A., and Ishijima, K.: The Orbiting
15 Carbon Observatory (OCO-2) tracks 2-3 peta-gram increase in carbon release to
16 the atmosphere during the 2014-2016 El Nino. *Scientific Reports*, 7(1): 13567, 2017.
- 17 Qu, Z., Jacob, D. J., Shen, L., Lu, X., Zhang, Y. Z., Scarpelli, T R., Nesser, H., Sulprizio,
18 M. P., Maasakkers, J. D., Bloom, A. A., Worden, J. R., Parker, R. J., and Delgado,
19 A. L.: Global distribution of methane emissions: a comparative inverse analysis of
20 observations from the TROPOMI and GOSAT satellite instruments. *Atmospheric*
21 *Chemistry and Physics*, 21(18): 14159-14175, 2021.
- 22 Reuter, M., Bovensmann, H., Buchwitz, M., Burrows, J.P., Connor, B.J., Deutscher,
23 N.M., Griffith, D.W., Heymann, J., Keppel-Aleks, G., Messerschmidt, J., Notholt,
24 J., Petri, C., Robinson, J., Schneising, O., Sherlock, V., Velazco, V.A., Warneke,
25 T., Wennberg, P.O., and Wunch, D.: Retrieval of atmospheric CO₂ with enhanced
26 accuracy and precision from SCIAMACHY: validation with FTS measurements
27 and comparison with model results. *Journal of Geophysical Research*, 116, 1,
28 2011.
- 29 Saito, R., Patra, P. K., Deutscher N., Wunch, D., Ishijima, K., Sherlock, V.,
30 Blumenstock, T., Dohe S., Griffith, D., Hase, F., Heikkinen, P., Kyro, E.,
31 Macatangay, R., Mendonca, J., Messerschmidt, J., Morino, I., Notholt, J.,
32 Rettinger, M., Strong, K., Sussmann, R., and Warneke, T.: Technical Note:
33 Latitude-time variations of atmospheric column-average dry air mole fractions of
34 CO₂, CH₄ and N₂O. *Atmospheric Chemistry and Physics*, 12(16): 7767-7777,

1 2012.

2 Schneising, O., Bergamaschi, P., Bovensmann H., Buchwitz M., Burrows J. P.,
3 Deutscher N. M., Griffith, D. W. T., Heymann J., Macatangay R., Messerschmidt,
4 J., Notholt, J., Rettinger M., Reuter M., Sussmann R., Velazco, V. A., Warneke T.,
5 Wennberg P. O., Wunch, D.: Atmospheric greenhouse gases retrieved from
6 SCIAMACHY: comparison to ground-based FTS measurements and model
7 results. *Atmospheric Chemistry and Physics*, 12(3): 1527-1540, 2012.

8 Sha, M. K., Langerock, B., Blavier J. F. L., Blumenstock, T., Borsdorff, T., Buschmann,
9 M., Dehn A., De Maziere, M., Deutscher, N. M., Feist, D. G., Garcia, O. E.,
10 Griffith, D. W. T., Grutter, M., Hannigan, J. W., Hase F., Heikkinen, P., Hermans,
11 C., Iraci, L. T., Jeseck, P., Jones, N., Kivi, R., Kumps, N., Landgraf, J., Lorente,
12 A., Mahieu, E., Makarova, M. V., Mellqvist, J., Metzger, J. M., Morino, I,
13 Nagahama, T., Notholt, J., Ohyama, H., Ortega, I, Palm M., Petri, C., Pollard, D.
14 F., Rettinger, M., Robinson, J., Roche, S., Roehl, C. M., Rohling, A. N.,
15 Rousogonous, C., Schneider, M., Shiomi, K., Smale, D., Stremme, W., Strong, K.,
16 Sussmann, R., Te Y., Uchino, O., Velazco, V. A., Vigouroux, C., Vrekoussis, M.,
17 Wang, P. C., Warneke, T., Wizenberg, T., Wunch, D., Yamanouchi, S., Yang Y.,
18 Zhou, M. Q.: Validation of methane and carbon monoxide from Sentinel-5
19 Precursor using TCCON and NDACC-IRWG stations. *Atmospheric Measurement*
20 *Techniques*, 14(9): 6249-6304, 2021.

21 Shen, L., Zavala-Araiza, D., Gautam, R., Omara, M., Scarpelli, T., Sheng, J. X.,
22 Sulprizio, M. P., Zhuang, J. W., Zhang, Y. Z., Qu, Z., Lu, X., Hamburg, S. P., Jacob,
23 D. J.: Unravelling a large methane emission discrepancy in Mexico using satellite
24 observations. *Remote Sensing of Environment*, 260: 112461, 2021.

25 Singh, H. B., Brune, W. H., Crawford, J. H., Jacob, D. J., Russell, P. B.: Overview of
26 the summer 2004 intercontinental chemical transport experiment - North America
27 (INTEX-A). *Journal of Geophysical Research-Atmosphere*, 111(D24): D24S01,
28 2006.

29 Sun, Y. W., Liu, C., Chan, K., Wang, W., Shan, C. G., Hu, Q., Liu, J. G.: The Influence
30 of Instrumental Line Shape Degradation on the Partial Columns of O₃, CO₂, CH₄
31 and N₂O Derived from High-Resolution FTIR Spectrometry. *Remote Sensing*,
32 10(12): 2041, 2018a.

33 Sun, Y. W., Palm, M., Liu, C., Hase, F., Griffith, D., Weinzierl, C., Petri, C., Wang, W.,
34 Notholt, J.: The influence of instrumental line shape degradation on NDACC gas

1 retrievals: total column and profile. *Atmospheric Measurement Techniques*, 11(5):
2 2879-2896, 2018b.

3 Sun, Y. W., Yin, H., Liu, C., Mahieu, E., Notholt, J., Te, Y., Lu, X., Palm, M., Wang,
4 W., Shan, C. G., Hu, Q. H., Qin, M., Tian, Y., Zheng, B.: The reduction in C₂H₆
5 from 2015 to 2020 over Hefei, eastern China, points to air quality improvement in
6 China. *Atmospheric Chemistry and Physics*, 21(15): 11759-11779, 2021.

7 Shi, T Q., Han, Z. Y., Gong, W., Ma, X., and Han, G.: High-precision methodology for
8 quantifying gas point source emission. *Journal of Cleaner Production*, 320,
9 128672, 2021.

10 Tang, X. L., Liu, S. G., Zhou, G. Y., Zhang, D. Q., Zhou, C. Y.: Soil-atmospheric
11 exchange of CO₂, CH₄, and N₂O in three subtropical forest ecosystems in southern
12 China. *Global Change Biology*, 12(3): 546-560, 2006.

13 Thompson, D. R., Benner, D. C., Brown, L. R., Crisp, D., Devi, V. M., Jiang, Y. B.,
14 Natraj, V., Oyafuso, F., Sung, K., Wunch, D., Castano, R., Miller, C. E.:
15 Atmospheric validation of high accuracy CO₂ absorption coefficients for the
16 OCO-2 mission. *Journal of Quantitative Spectroscopy & Radiative Transfer*,
17 113(17): 2265-2276, 2012.

18 Tian, Y., Sun, Y., Liu, C., Wang, W., Shan, C., Xu, X., Hu, Q.: Characterisation of
19 methane variability and trends from near-infrared solar spectra over Hefei, China.
20 *Atmospheric Environment*, 173: 198-209, 2017.

21 Tian, Y., Sun, Y. W., Liu, C., Xie, P. H., Chan, K., Xu, J., Wang, W., and Liu, J. G:
22 Characterization of urban CO₂ column abundance with a portable low resolution
23 spectrometer (PLRS): Comparisons with GOSAT and GEOS-Chem model data.
24 *Science of the Total Environment*, 612: 1593-1609, 2018.

25 Tu, Q. S., Hase, F., Blumenstock, T., Kivi, R., Heikkinen, P., Sha, M. K., Raffalski, U.,
26 Landgraf, J., Lorente, A., Borsdorff, T., Chen, H. L., Dietrich, F., Chen, J.:
27 Intercomparison of atmospheric CO₂ and CH₄ abundances on regional scales in
28 boreal areas using Copernicus Atmosphere Monitoring Service (CAMS) analysis,
29 COllaborative Carbon Column Observing Network (COCCON) spectrometers,
30 and Sentinel-5 Precursor satellite observations. *Atmospheric Measurement*
31 *Techniques*, 13(9): 4751-4771, 2020.

32 Varon, D. J., Mckeever, J., Jervis, D., Maasakkers, J. D., Pandey, S., Houweling, S.,
33 Aben, I., Scarpelli, T., Jacob, D. J.: Satellite Discovery of Anomalously Large
34 Methane Point Sources From Oil/Gas Production. *Geophysical Research Letters*,

1 46(22): 13507-13516, 2019.

2 Veefkind, J. P., Aben, I., McMullan, K., Forster, H., De Vries, J., Otter, G., Claas, J.,
3 Eskes, H. J., De Haan, J. F., Kleipool, Q., Van Weele, M., Hasekamp, O.,
4 Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga,
5 B., Vink, R., Visser, H., Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor:
6 A GMES mission for global observations of the atmospheric composition for
7 climate, air quality and ozone layer applications. *Remote Sensing of Environment*,
8 120: 70-83, 2012.

9 Vogel, F. R., Frey, M., Staufer, J., Hase, F., Broquet, G., Xueref-Remy, I., Chevallier,
10 F., Ciais, P., Sha, M. K., Chelin, P., Jeseck, P., Janssen, C., Te, Y., Gross, J.,
11 Blumenstock, T., Tu, Q. S., Orphal, J.: XCO₂ in an emission hot-spot region: the
12 COCCON Paris campaign 2015. *Atmospheric Chemistry and Physics*, 19(5):
13 3271-3285, 2019.

14 Wang, C., Guan, K. Y., Peng, B., Chen, M., Jiang, C. Y., Zeng, Y. L., Wu, G H., Wang,
15 S., Wu, J., Yang, X., Frankenberg, C., Kohler, P., Berry, J., Bernacchi, C., Zhu, K.,
16 Alden, C., Miao, G F.: Satellite footprint data from OCO-2 and TROPOMI reveal
17 significant spatio-temporal and inter-vegetation type variabilities of solar-induced
18 fluorescence yield in the US Midwest. *Remote Sensing of Environment*, 241:
19 111728, 2020a.

20 Wang, K., Wang, Y. L., Wang, X. H., He, Y., Li, X. Y., Keeling, R. F., Ciais P., Heimann
21 M., Peng, S. S., Chevallier, F., Friedlingstein, P., Sitch, S., Buermann, W., Arora,
22 V. K., Haverd, V., Jain, A. K., Kato, E., Lienert, S., Lombardozzi, D., Nabel, J. E.
23 M. S., Poulter, B., Vuichard, N., Wiltshire, A., Zeng, N., Zhu, D., Piao, S. L.:
24 Causes of slowing-down seasonal CO₂ amplitude at Mauna Loa. *Global Change*
25 *Biology*, 26(8): 4462-4477, 2020b.

26 Wang, R. W., Xie, P. H., Xu, J., Li, A., Sun, Y. W.: Observation of CO₂ Regional
27 Distribution Using an Airborne Infrared Remote Sensing Spectrometer (Air-IRSS)
28 in the North China Plain. *Remote Sensing*, 11(2): 123, 2019.

29 Wang, S. P., van der, A. R. J., Stammes, P., Wang, W. H., Zhang, P., Lu, N. M., Fang,
30 L.: Carbon Dioxide Retrieval from TanSat Observations and Validation with
31 TCCON Measurements. *Remote Sens*, 12(21), 3626, 2020c.

32 Wang, W., Tian, Y., Liu, C., Sun Y., Liu W., Xie, P., Liu, J., Xu, J., Morino, I., Velazco,
33 V. A., Griffith, D. W. T., Notholt, J., Warneke, T.: Investigating the performance of
34 a greenhouse gas observatory in Hefei, China. *Atmospheric Measurement*

1 Techniques, 10(7): 2627-2643, 2017.

2 Wei, W. D., Li, J. S., Chen B., Wang, M., Zhang, P. F., Guan, D. B., Meng, J., Qian, H.

3 Q., Cheng, Y. H., Kang, C. Q., Feng, K. S., Yang, Q., Zhang, N., Liang, X., and

4 Xue J. J.: Embodied greenhouse gas emissions from building China's large-scale

5 power transmission infrastructure. *Nature Sustainability*, 4(8): 739-747, 2021.

6 Wu, H., Wang, X. H., Ye, H. H., Jiang, Y., and Duan, F. H.: Error analysis of the

7 greenhouse-gases monitor instrument short wave infrared XCO₂ retrieval

8 algorithm. *Journal of Applied Remote Sensing*, 12(1): 016015, 2018.

9 Wunch, D., Toon G C., Blavier, J. F. L., Washenfelder, R. A., Notholt, J., Connor, B. J.,

10 Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column

11 Observing Network. *Philosophical Transactions of the Royal Society A-*

12 *Mathematical Physical and Engineering Sciences*, 369(1943): 2087-2112, 2011.

13 Wunch, D., Toon, G C., Wennberg, P. O., Wofsy, S. C., Stephens B. B., Fischer, M. L.,

14 Uchino, O., Abshire J. B., Bernath, P., Biraud, S. C., Blavier, J. F. L., Boone, C.,

15 Bowman, K. P., Browell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher,

16 N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst,

17 D. F., Jimenez, R., Keppel-Aleks, G., Kort, E. A., Macatangay, R., Machida, T.,

18 Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl C. M., Sawa,

19 Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. A.: Calibration of the

20 Total Carbon Column Observing Network, using aircraft profile data. *Atmospheric*

21 *Measurement Techniques*, 3(5): 1351-1362, 2010.

22 Wunch, D., Wennberg, P. O., Osterman, G., Fisher, B., Naylor, B., Roehl, C. M., O'dell,

23 C., Mandrake L., Viatte C., Kiel M., Griffith, D. W. T., Deutscher, N. M., Velazco,

24 V. A., Notholt, J., Warneke, T., Petri, C., De Maziere, M., Sha, M. K., Sussmann,

25 R., Rettinger, M., Pollard D., Robinson, J., Morino, I., Uchino, O., Hase, F.,

26 Blumenstock, T., Feist, D. G., Arnold, S. G., Strong, K., Mendonca, J., Kivi, R.,

27 Heikkinen, P., Iraci, L., Podolske, J., Hillyard, P., Kawakami, S., Dubey, M. K.,

28 Parker, H. A., Sepulveda, E., Garcia, O. E., Te, Y., Jeseck, P., Gunson, M. R.,

29 Crisp, D., and Eldering, A.: Comparisons of the Orbiting Carbon Observatory-2

30 (OCO-2) XCO₂ measurements with TCCON *Atmospheric Measurement*

31 *Techniques*, 10(6): 2209-2238, 2017.

32 Yang, D., Boesch, H., Liu, Y., Somkuti, P., Cai, Z., Chen, X., Di Noia, A., Lin, C., Lu,

33 N., Lyu, D., Parker, R. J., Tian, L., Wang, M., Webb, A., Yao, L., Yin, Z., Zheng,

34 Y., Deutscher, N. M., Griffith, D. W. T., Hase, F., Kivi, R., Morino, I., Notholt, J.,

1 Ohyama, H., Pollard, D. F., Shiomi, K., Sussmann, R., Te, Y., Velazco, V. A.,
2 Warneke, T., and Wunch, D.: Toward High Precision XCO₂ Retrievals From
3 TanSat Observations: Retrieval Improvement and Validation Against TCCON
4 Measurements. *Journal of Geophysical Research- Atmosphere*, 125(22):
5 e2020JD032794, 2020a.

6 Yang, D. X., Liu, Y., Boesch, H., Yao, L., Di Noia, A., Cai, Z. N., Lu, N. M., Lyu, D.
7 R., Wang, M. H., Wang, J., Yin, Z. S., and Zheng, Y. Q.: A. New TanSat XCO₂
8 Global Product towards Climate Studies, *Advances in Atmospheric Sciences*,
9 38(1): 8-11, 2021a.

10 Yang, D. X., Liu, Y., Cai, Z. N., Chen, X., Yao, L., and Lu, D. R.: First Global Carbon
11 Dioxide Maps Produced from TanSat Measurements, *Advances in Atmospheric*
12 *Sciences*, 35(6): 621-623, 2018a.

13 Yang, D. X., Liu, Y., Feng, L., Wang, J., Yao, L., Cai, Z. N., Zhu, S. H., Lu, N. M., and
14 Lyu, D. R.: The First Global Carbon Dioxide Flux Map Derived from TanSat
15 Measurements, *Advances in Atmospheric Sciences*, 38(9): 1433-1443, 2021b.

16 Yang, Y., Zhou, M. Q., Langerock, B., Sha, M. K., Hermans, C., Wang, T., Ji, D. H.,
17 Vigouroux, C., Kumps, N., Wang, G. C., De Maziere, M., and Wang, P. C.: New
18 ground-based Fourier-transform near-infrared solar absorption measurements of
19 XCO₂, XCH₄ and XCO at Xianghe, China, *Earth System Science Data*, 12(3):
20 1679-1696, 2020c.

21 Yang, Z. D., Bi, Y. M., Wang, Q., Liu, C. B., Gu, S. Y., Zheng, Y. Q., Lin, C., Yin, Z.
22 S., and Tian, L. F.: Inflight Performance of the TanSat Atmospheric Carbon
23 Dioxide Grating Spectrometer, *IEEE Transactions on Geoscience and Remote*
24 *Sensing*, 58(7): 4691-4703, 2020.

25 Yang, Z. D., Zhen, Y. Q., Yin, Z. S., Lin, C., Bi, Y. M., Liu, W., Wang, Q., Wang, L.,
26 Gu, S. Y., and Tian, L. F.: Laboratory spectral calibration of the TanSat
27 atmospheric carbon dioxide grating spectrometer, *Geoscientific Instrumentation*
28 *Methods and Data Systems*, 7(3): 245-252, 2018.

29 Yang, Z. D., Zhen, Y. Q., Yin, Z. S., Lin, C., Bi, Y. M., Liu, W., Wang, Q., Wang, L.,
30 Gu, S. Y., and Tian, L. F.: Prelaunch Radiometric Calibration of the TanSat
31 Atmospheric Carbon Dioxide Grating Spectrometer, *IEEE Transactions on*
32 *Geoscience and Remote Sensing*, 56(7): 4225-4233, 2018c.

33 Yi Y., Liu Y., Cai Z., Fang S., Yang D., Wang Y., Liang, M., Yao, B., Ma Q., Wang M.:

Measuring and comparing in-situ CO₂ and CO profiles with satellite observations and model data. *Atmospheric and Oceanic Science Letters*, DOI: 10.1080/16742834.2019.1649974, 2019.

Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyro, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data. *Atmospheric Measurement Techniques*, 6(6): 1533-1547, 2013.

Zellweger, C., Emmenegger, L., Firdaus, M., Hatakka, J., Heimann, M., Kozlova, E., Spain, T. G., Steinbacher, M., Van Der Schoot, M. V., and Buchmann, B.: Assessment of recent advances in measurement techniques for atmospheric carbon dioxide and methane observations, *Atmospheric Measurement Techniques*, 9(9): 4737-4757, 2016.

Zhang, H., Zheng, Y. Q., Li, S., Lin, C., Li, C. L., Yuan, J. Z., and Li, Y.: Geometric correction for TanSat atmospheric carbon dioxide grating spectrometer, *Sensor Actuator A-Physical*, 293:62-69, 2019.

Zhang, X. Y., Wang, F., Wang, W. H., Huang, F. X., Chen, B. L., Gao, L., Wang, S. P., Yan, H. H., Ye, H. H., Si, F. Q., Hong, J., Li, X. Y., Cao, Q., Che, H. Z., and Li, Z. Q.: The development and application of satellite remote sensing for atmospheric compositions in China, *Atmospheric Research*, 245, 105056, 2020.

Zhang, Y. Z., Gautam, R., Pandey, S., Omara, M., Maasakkers, J. D., Sadavarte, P., Lyon, D., Nesser, H., Sulprizio, M. P., Varon, D. J., Zhang, R. X., Houweling, S., Zavala-Araiza, D., Alvarez, R. A., Lorente, A., Hamburg, S. P., Aben, I., and Jacob, D. J.: Quantifying methane emissions from the largest oil-producing basin in the United States from space, *Science Advance*, 6(17): eaaz5120, 2020b.

Zhao, M. J., Si, F. Q., Zhou, H. J., Jiang, Y., Ji, C. Y., Wang, S. M., Zhan, K., Liu, W. Q.: Pre-Launch Radiometric Characterization of EMI-2 on the GaoFen-5 Series of Satellites, *Remote Sensing*, 13(14), 2843, 2021.

Zheng, B., Chevallier, F., Ciais, P., Broquet, G., Wang, Y. L., Lian, J. H., and Zhao, Y. H.: Observing carbon dioxide emissions over China's cities and industrial areas with the Orbiting Carbon Observatory-2, *Atmospheric Chemistry and Physics*, 20(14): 8501-8510, 2020a.

1 Zheng, B., Geng, G. N., Ciais, P., Davis, S. J., Martin, R. V., Meng, J., Wu, N. N.,
2 Chevallier, F., Broquet, G., Boersma, F., Van Der Ronald, A., Lin, J. T., Guan, D.
3 B., Lei, Y., He, K. B., and Zhang, Q.: Satellite-based estimates of decline and
4 rebound in China's CO₂ emissions during COVID-19 pandemic. *Science*
5 *Advances*, 6(49): eabd4998, 2020b.
6

1 Figures




Fig. 1 Long-term time series of CO₂ at the Mauna Loa, Barrow, American Samoa, and South Pole observatories (adapted from <https://gml.noaa.gov/>)




Fig. 2 Global GHGs monitoring network coordinated by NOAA and WMO. Geolocations of all sites are listed in Table S1. Base map of this figure is from the Basemap package of Python.




Fig. 3 Global FTIR observation networks, including TCCON, NDACC-IRWG, and COCCON networks. Geolocations of all sites are listed in Table S2. Base map of this figure is from the Basemap package of Python.




Fig. 4 Principles of spectroscopic techniques for GHGs monitoring. Active measurement techniques (a, f) use artificial light source and passive measurement techniques (b, c, d, e) use natural light sources such as sun to monitor GHGs.




Fig. 5 Stereoscopic GHGs monitoring network to support carbon peak/neutrality in China




Fig. 6 Ground-based FTIR CO₂ measurements at the Hefei observatory (Shan et al., 2021). The seasonality and interannual variability are represented by black dashed curve and black dashed line, respectively, which are fitted by using a bootstrap resampling model with a 3rd Fourier series plus a linear function.

1 Tables

2

Table 1 Status of GHGs monitoring

Institutions	institutes of Chinese academy of sciences, universities, business units, and enterprises
Technologies	EC, DOAS, TDLAS, FTIR, NDIR, GC/FID, LIDAR, CRDS, OA-ICOS, PAS
Platforms	manual sampling, surface in-situ, ground-based, tower-based, airborne, space-based, ship-borne, vehicle-borne, UAV, balloon, and tethered balloon
Working mode	active, passive, single constituent, and multi-constituent
Target	H ₂ O, HDO, CO ₂ , CH ₄ , CH ₃ D, N ₂ O, SF ₆ , O ₃ , C ₂ H ₆ , CCl ₃ F, CCl ₂ F ₂ , and CHClF ₂
Coverage	single point, small scale, regional scale, large scale, and global scale
Sensitivity	ppbv to ppmv level
Accuracy	thousandth to percent level
Temporal resolution	second, minute, hour to day levels
Spatial resolution	meter, dozens of meters, kilometer to dozens of kilometers
Monitoring regions	typical industrial zones, industrial stack emissions, urban atmosphere, ambient atmosphere, remote background regions, offshore regions, wetlands
Applications	investigations of global carbon cycle, GHGs evolution trends, regional GHGs sources and transport, ecological GHGs flux estimate, urban or industrial GHGs emissions estimates, validations of CTMs and emission inventory, cross calibration, and algorithm improvement
Advances	TanSat and GF-5 series GHGs satellites, space-borne CO ₂ LIDAR, ground-based high resolution remote sensing, the performance for homemade GHGs monitoring instruments have been steadily improved

3