

Mass spectrometry-based *aerosolomics*: a new approach to resolve sources, composition, and partitioning of secondary organic aerosol

Markus Thoma¹, Franziska Bachmeier¹, Felix Leonard Gottwald¹, Mario Simon¹, and Alexander Lucas Vogel¹

¹Institute for Atmospheric and Environmental Sciences, Goethe-University Frankfurt, 60438 Frankfurt am Main, Germany

Correspondence: Alexander L. Vogel (vogel@iau.uni-frankfurt.de), Markus Thoma (thoma@iau.uni-frankfurt.de)

Abstract. Particulate matter (PM) largely consists of secondary organic aerosol (SOA) that is formed via oxidation of biogenic and anthropogenic volatile organic compounds (VOCs). Unambiguous identification of SOA molecules and their assignment to their precursor vapors is a challenge that has so far only succeeded for a few SOA marker compounds, which are now well characterized and (partly) available as authentic standards. In this work, we resolve the complex composition of SOA by a top-down approach based on a newly created *aerosolomics* database, which is fed by non-target analysis results of filter samples from oxidation flow reactor experiments. We investigated the oxidation products from the five biogenic VOCs α -pinene, β -pinene, limonene, 3-carene, and *trans*-caryophyllene and from the four anthropogenic VOCs toluene, *o*-xylene, 1,2,4-trimethylbenzene, and naphthalene. Using ultra-high performance liquid chromatography coupled to a high-resolution (Orbitrap) mass spectrometer, we determine the molecular formula of 596 chromatographically separated compounds based on exact mass and isotopic pattern. We utilize retention time and fragmentation mass spectra as a basis for unambiguous attribution of the oxidation products to their parent VOCs. Based on the molecular-resolved application of the database, we are able to assign roughly half of the total signal of oxygenated hydrocarbons in ambient suburban PM_{2.5} to one of the nine studied VOCs. The application of the database enabled us to interpret the appearance of diurnal compound clusters that are formed by different oxidation processes. Furthermore, by performing a hierarchical cluster analysis (HCA) on the same set of filter samples we identified compound clusters that depend on sulfur dioxide mixing ratio and temperature. This study demonstrates how *aerosolomics* tools (database and HCA) applied on PM filter samples can improve our understanding of SOA sources, their formation pathways, and temperature-driven partitioning of SOA compounds.

1 Introduction

Secondary organic aerosol (SOA) is a complex mixture forming through the oxidation of biogenic (BVOCs) and anthropogenic volatile organic compounds (AVOCs) in the atmosphere. Aerosol particles influence Earth's climate as well as human health (Hallquist et al., 2009; Shrivastava et al., 2017). Earlier work has shown that SOA makes up a large fraction of fine particulate matter (PM_{2.5}, particles with an aerodynamic diameter less than 2.5 μm) (Jimenez et al., 2009; Huang et al., 2014; McDonald et al., 2018). Globally, the emissions of BVOCs are considerably higher than those of AVOCs, with 760–1150 TgC y⁻¹ compared to 140 TgC y⁻¹, respectively (Kari et al., 2019; Shrivastava et al., 2017; Sindelarova et al., 2014). BVOC emissions

25 can mainly be distributed among isoprene (70 %), monoterpenes (11 %), methanol (6 %), and others (13 %) (Sindelarova et al., 2014). AVOCs and BVOCs are not only emitted from different sources, but they also have different SOA yields, and result in different products with distinct different properties. Furthermore, it is known that both organic and inorganic anthropogenic emissions can affect SOA formation from BVOCs (Kari et al., 2019; Xu et al., 2021). Still major knowledge gaps exist on the sources and formation pathways of SOA, its transformation and lifetime in the atmosphere, and its underlying effect on Earth's
30 climate and human health. Furthermore, emissions of anthropogenic and biogenic precursors as well as their atmospheric fate are uncertain, resulting in a discrepancy between measured and modeled SOA (Fuzzi et al., 2015). Improved chemical characterization of ambient SOA can help understanding of sources, formation pathways, and effects on both climate and human health.

Many controlled laboratory studies have increased our mechanistic understanding of the oxidation of volatile organic compounds (VOCs) (Burkholder et al., 2017). However, the ambient atmosphere is usually more complex than chamber experiments, and unaccounted chemical interactions can therefore alter SOA yields that are derived from simple laboratory systems (McFiggans et al., 2019). Furthermore, it is likely that many VOCs are understudied that are relevant for SOA formation. Therefore, a comprehensive top-down investigation of SOA can enable identification of missing important precursor gases or relevant formation pathways. Numerous previous investigations (Glasius et al., 2000; Kristensen et al., 2016; Nozière et al.,
40 2015; Surratt et al., 2007) clearly highlight the advantages of offline measurement techniques, which apply separation techniques like gas chromatography or (ultra-high performance) liquid chromatography (UHPLC) coupled to (high-resolution) mass spectrometry (HRMS), because with these techniques the unambiguous identification of different compounds becomes possible. In recent years, non-target analysis (NTA) of UHPLC-HRMS measurements has become a powerful tool that builds peak lists of all detected compounds in complex samples, and determines the molecular formula based on the exact mass and
45 isotopic pattern. Furthermore, MS²-spectra can be compared to fragmentation libraries and enable database-assisted identification of compounds (Ditto et al., 2018; Ma et al., 2022; Pereira et al., 2021; Pleil et al., 2018). However, there are currently no established databases of atmospheric SOA tracers, which can be applied on measurements of ambient PM_{2.5} filter samples.

Inspired by metabolomics, a tool widely used in the life sciences to identify metabolites, metabolic pathways, and biomarkers (Fiehn, 2002; Johnson et al., 2016), we created an *aerosolomics* database for database-assisted identification of marker
50 compounds (without having the need for authentic standards), and hence enables the investigation of atmospheric transformation pathways of VOCs under different oxidation conditions. The database enables compound matching based on filters from potential aerosol mass (PAM) oxidation flow reactor (OFR) experiments of nine biogenic and anthropogenic VOCs. We applied the database to ambient air filter samples collected in summer 2018 near Vienna (Austria). Figure 1 shows the principal steps of the new *aerosolomics* approach that is based on the comparison between filter samples from OFR experiments and
55 from the ambient. Additionally, a hierarchical cluster analysis (HCA) was performed in order to reduce the complexity of the ambient dataset and to assign compounds to certain formation processes or emission sources. Both strategies combined allow the identification of oxidation products from either biogenic or anthropogenic VOCs, and enable a better understanding of the oxidation conditions and of temperature-driven gas-to-particle partitioning.




Figure 1. Establishment and application of the *aerosolomics* database: The database is filled with the results of several PAM-OFR experiments with different precursors and reactants and compared to the molecular fingerprints of ambient PM_{2.5} filter samples. Matching compounds can be assigned to the corresponding VOC precursor. Knowledge about not-matching compounds, due to missing primary sources or SOA from unaccounted precursors/formation mechanisms, can guide further experiments.

2 Experimental

60 2.1 Oxidation flow reactor experiments

We used a PAM-OFR (Aerodyne Research Inc.) (Kang et al., 2007; Lambe et al., 2011) for laboratory SOA formation of biogenic VOCs (-)- α -pinene (98 %, Alfa Aesar, CAS: 7785-26-4), (+)- β -pinene (98 %, Sigma-Aldrich, CAS: 19902-08-0), (S)-(-)-limonene (96 %, Sigma-Aldrich, CAS: 5989-54-8), (+)-3-carene (≥ 98.5 %, Fluka Analytical, CAS: 498-15-7), and (-)-*trans*-caryophyllene (≥ 98 %, Sigma-Aldrich, CAS: 87-44-5), as well as the anthropogenic VOCs toluene (>99.5 %, Fluka Analytical, CAS: 108-88-3), *o*-xylene (>99.0 %, Fluka Analytical, CAS: 95-47-6), 1,2,4-trimethylbenzene (98 %, Sigma-Aldrich, CAS: 95-63-6), and naphthalene (>99 %, Merck KGaA, CAS: 91-20-3). In dark ozonolysis experiments, all BVOCs were oxidized with externally produced ozone (O₃). Hydroxyl radicals (OH) formed during ozonolysis of VOCs were not scavenged. Under daytime simulations, OH was generated by UV light ($\lambda = 254$ nm) inside the OFR.

We evaporated the VOCs in a heated glass flask purged continuously with nitrogen (N₂, 6.0 purity, Nippon Gases). The aerosol mass concentration was measured with a scanning mobility particle sizer spectrometer (SMPS; consisting of an elec-

trostatic classifier 3082, a differential mobility analyzer 3081A, and an ultrafine condensation particle counter 3776, TSI Inc.). Changing the precursor concentration by varying the temperature inside the flask resulted in aerosol mass concentrations between 20 and 184 $\mu\text{g m}^{-3}$. The individual settings and the resulting mass concentrations are given in Table S1.

In all experiments the nitrogen flow into the reactor was 4.8 L min^{-1} , the oxygen (O_2 , 5.0 purity, Nippon Gases) flow was 1.2 L min^{-1} resulting in a mean residence time of 2.4 minutes. The relative humidity was 55 %. The O_3 concentration was ~ 1 ppm, decreasing to 0.8 ppm under OH conditions. Based on these experimental conditions we calculated the external OH reactivity (Eq. S1) and the OH exposure using the OFR exposure estimator (Peng et al., 2015, 2016). The resulting OH exposures (Table S1) correspond to approximately 0.1–6 days of equivalent atmospheric OH exposure, based on the assumption of an averaged tropospheric OH concentration of 1.09×10^6 molecules cm^{-3} (Li et al., 2018).

The aged air leaving the OFR passed through two 50 cm denuders, packed with charcoal (IAC-402, Infiltec GmbH) and potassium permanganate (KMnO_4 , IAC-630, Infiltec GmbH) in order to remove (reactive) gas-phase compounds. Glass fiber filters (47 mm, Pallflex Emfab Filters, Pall) sampled the formed SOA particles with a flow of 3 L min^{-1} for a duration of 90 minutes. Until sample preparation and analysis, filter samples were packed in aluminium foil and stored at -18°C .

2.2 Ambient air filter sampling campaign

$\text{PM}_{2.5}$ filter samples were collected in August 2018 during a field campaign (48.127°N , 16.534°E), at a suburban background station between the Vienna International Airport in the east and the Schwechat Industrial Park and Vienna city center in the north-west. 52 glass fiber filters (150 mm, Ahlstrom-Munksjö) were sampled for 12 hours starting at 05:00 (UTC) or 17:00 (UTC) respectively, using a high volume sampler (DHA-80, Digital Elektronik AG) at a flow rate of 30 $\text{m}^3 \text{h}^{-1}$. The meteorological parameters (i.e., wind direction, wind speed, and temperature), the trace gas concentration (i.e., nitrogen oxide (NO), nitrogen dioxide (NO_2), sulfur dioxide (SO_2), and carbon monoxide (CO)), as well as the $\text{PM}_{2.5}$ mass concentration were monitored continuously.

2.3 Sample preparation

From each ambient filter sample, one punch (25 mm diameter) was cut in small pieces and extracted in a glass vial using 200 μL of ultrapure water (Milli-Q Reference A+, Merck KGaA) and methanol (Optima LC/MS Grade, Thermo Fisher Scientific Inc.) (90/10, v/v) for 20 minutes on an orbital shaker with 300 rpm. Afterwards, the solvent was drawn up with a syringe (Injekt-F, Braun Melsungen AG) and filtered through a 0.2 μm syringe filter (non-sterile PTFE Syringe Filter, Thermo Fisher Scientific Inc.). In a second step, 100 μL of the solvent mixture was added and the procedure was repeated. 50 μL of the extracted sample was mixed with 5 μL of an internal standard containing isotopically labeled benzoic acid ($\text{C}_6\text{H}_5^{13}\text{CO}_2\text{H}$, 99 atom % ^{13}C , Sigma-Aldrich, $c = 0.1 \text{ mg mL}^{-1}$).

Half of each filter from the OFR experiments was cut in small pieces and extracted analogously to the ambient air filter samples with an adjustment in the eluent volume: 180 μL in the first and 80 μL in the second step was used. Finally 100 μL of the extracted sample was mixed with 10 μL of the internal standard.

2.4 Standard mixture for non-target analysis validation

105 A solution of 13 analytical standards was used to validate UHPLC-HRMS measurements and the NTA workflow with primary attention toward automated compound identification, but also toward fragmentation and adduct formation, which can result in false positives. To cover a variety of atmospherically relevant compounds, the mixture consists of carboxylic acids, organosulfates and -phosphates, as well as nitrogen containing compounds. The injection volume for the analysis was 1 μL . A detailed overview of the substances used and their concentrations in the mixture are given in Table S2.

2.5 UHPLC-(–)HESI-HRMS measurements

110 The extracts of the ambient PM samples were separated by ultra-high performance liquid chromatography (Vanquish Flex, Thermo Fisher Scientific Inc.) on a reversed phase column (Accucore C₁₈, 2.6 μm , 150 \times 2.1 mm, Thermo Fisher Scientific Inc.), ionized in the negative mode using a heated electrospray ionization source (HESI-II Probe, Thermo Fisher Scientific Inc.) and detected with a high-resolution hybrid quadrupole-Orbitrap mass spectrometer (Q Exactive Focus, Thermo Fisher Scientific Inc.). Eluents were ultrapure water (eluent A) and methanol (eluent B), both mixed with 0.1 % formic acid (98 %, 115 Merck KGaA). The injection volume was 5 μL , the flowrate was 400 $\mu\text{L min}^{-1}$, and the temperature was 40 $^{\circ}\text{C}$. The gradient started with 1 % eluent B (0–0.5 min), increased linearly to 99 % B (0.5–14 min), stayed at 99 % B (14–16 min), backflushed in one minute and equilibrated in three minutes resulting in a total run time of 20 minutes. The ion source settings were –3.5 kV spray voltage, 40 psi sheath gas, 8 arbitrary units auxiliary gas, and 262.5 $^{\circ}\text{C}$ capillary temperature. The spectra were recorded in full scan MS with data dependent tandem mass spectrometry (ddMS²) using a higher-energy collisional dissociation (HCD) 120 cell with stepped collision energies of 15, 30 and 45 eV. The scan range in full MS was m/z 50–750 with a resolution of 70 000 at m/z 200. For ddMS² the resolution was 17 500.

A representative selection of 10 ambient PM samples was measured a second time. The selection was based on external influences like wind direction, temperature, time of day and trace gas concentrations. In one sequence, together with filter samples from laboratory OFR experiments, we applied and improved gradient on another reversed phase column (Cortecs 125 Solid Core T3, 2.7 μm , 150 \times 3 mm, with the corresponding VanGuard Cartridge, Waters Corp.). The gradient also started with 1 % B for half a minute, increased linearly to 99 % B in 15 minutes, and held it for 2 minutes. Afterwards, the column was backflushed in 90 seconds and equilibrated in two and a half minutes resulting in a total run time of 21.5 minutes. This dataset is basis for the application of the database on ambient samples.

As a quality control routine, we extracted one filter three times to estimate the reproducibility of the extraction procedure. 130 In addition, we determined the instrument performance by a triplicate measurement of one filter extract. We calculated the relative standard deviation (RSD) for 7 compounds (m/z 115–357 and signal intensities of 3×10^5 – 5×10^8 counts). Averaged over all seven compounds, we determined a mean RSD of 6.7 % for the reproducibility of the extraction procedure and 2.1 % for the instrumental performance.

2.6 Non-target analysis, MS² libraries, hierarchical cluster analysis, and volatility estimation

135 We used Compound Discoverer 3.2 (Thermo Fisher Scientific Inc.) for the NTA of the UHPLC-HRMS raw files. Chromato-
graphic peaks of interest were aligned with a maximum shift of 0.1 minutes in retention time and a mass tolerance of ± 1 ppm.
Ions were detected if the peak intensity was at least 5×10^5 counts for one of the following ions: $[M - H]^-$, $[M - CO_2 - H]^-$,
140 $[M - H_2O - H]^-$, and $[2M - H]^-$. In addition to the mass-to-charge ratio of the detected ion, at least one corresponding iso-
topologue has to be measured. The tolerance between measured and calculated intensity of the isotopologue has to be less
than 30 %. Unknown compounds were then grouped with a retention time tolerance of 0.1 minute and those of them with a
sample-to-blank ratio smaller than five were marked as background. A predicted composition was calculated within ± 1 ppm,
allowing the elements carbon (C), hydrogen (H), bromine (Br), chlorine (Cl), nitrogen (N), oxygen (O), and sulfur (S). Com-
pounds were grouped together as CHO, CHNO, CHOS, CHNOS, and “other” if the elemental composition contained other
heteroatoms. For unidentified compounds the software does not predict a composition under the given conditions. The detailed
145 workflow is given in Table S3.

To be clear on the degree of certainty regarding compound identification, we used the confidence levels from Schymanski
et al. (2014). Probably and tentatively labeled compounds correspond to level 2 and level 3, respectively. We used the mzCloud
database (HighChem LLC, 2013-2021) for comparing MS² spectra of commercial chemicals. Unambiguously identified com-
pounds (reference standard, MS² spectrum) correspond to confidence level 1.

150 Based on the Compound Discoverer results from the OFR experiments, we created a library for every examined chemi-
cal system (e.g., limonene and ozone) using mzVault 2.3 (Thermo Fisher Scientific Inc.), resulting in total 14 libraries of the
aerosolomics database. Every entry in a library contains the exact mass-to-charge ratio, the retention time, the MS² spectrum,
and the relative abundance to the major product of the respective system if the relative abundance is higher than 1 %. These
libraries were implemented in Compound Discoverer and aligned with the identified compounds from the representative se-
lection of the field campaign. An entry in the library was dedicated to a compound in the ambient air filter samples if the
155 difference in the retention times was smaller than 0.2 minutes, the measured mass-to-charge ratios of the ddMS² scans were
within a window of 10 ppm, and the match factor indicating the similarity of the MS² spectra was bigger than 50 %. Detailed
settings of this node are given in Table S3. If a compound appears in several libraries, the match factor was crucial for an
assignment. If match factors were equal, the system in which the compound has the highest relative abundance was chosen.

160 We calculated the effective saturation mass concentration ($\log_{10} C^*$) for each compound with a predicted composition
including at least the elements C, H, and O as well as N and S, based on Li et al. (2016). However, we like to point out
that this parameterization comprises a large molecular corridor and thus leads to a wide range of $\log_{10} C^*$. A bias has been
reported for nitrogen containing compounds (Isaacman-VanWertz and Aumont, 2021), but also for CHO compounds it appears
to be biased. For example, $\log_{10} C^*$ of the atmospheric tracer 3-methyl-1,2,3-butanetricarboxylic acid (C₈H₁₂O₆, MBTCA,
165 Szmigielski et al. (2007)) results in $1.97 \mu\text{g m}^{-3}$, while with SIMPOL.1 (Pankow and Asher, 2008) we find $\log_{10} C^*$ at 298 K
being $-2.2 \mu\text{g m}^{-3}$. However, this difference of four orders of magnitude is certainly an extreme case, as all oxygen atoms of
MBTCA are a part of a carboxylic acid functional group.

We performed a HCA with MATLAB R2020a (The MathWorks) based on the complete data set from the Vienna field campaign including the blank-corrected integrated sample peak areas. After z -transformation, we used an Euclidean distance metric and the Ward algorithm for computing the distance between the clusters. For the compound clusters of the HCA, an intensity-weighted mean of $\log_{10} C^*$ was calculated. The borders of the different volatility classes are given accordingly to Schervish and Donahue (2020). Based on this, the organic compounds can be classified as volatile (VOC), intermediate volatile (IVOC), semi-volatile (SVOC), low volatile (LVOC), extremely-low volatile (ELVOC), and ultra-low volatile (ULVOC).

3 Results and discussion

3.1 Oxidation flow reactor

OFR experiments provided SOA from several individually studied VOCs under different oxidation conditions. NTA of UHPLC-HRMS measurements of the SOA extracts enabled us to populate our *aerosolomics* database with individual oxidation products of the studied VOCs in a qualitative way. We investigated five BVOCs and four AVOCs, and identified 481 and 115 oxidation products, respectively. Each of these 596 oxidation products is listed in the database with the information on precursor, oxidation condition, exact mass-to-charge ratio, retention time, MS^2 -spectrum, and relative abundance in the respective system. Although, we are not able to determine the individual chemical structure of the different SOA compounds, their individual structure leads to compound-specific retention times. Using all these parameters in the presented database allows unambiguous attribution of SOA compounds in ambient samples to their major parent VOC.

3.1.1 SOA compounds from oxidation of biogenic VOCs

We investigated the composition of SOA from the atmospherically most abundant biogenic monoterpenes ($C_{10}H_{16}$) α -pinene, β -pinene, limonene, and 3-carene. The results are shown as mirror spectra in Fig. 2. The upper half of each subplot shows the ozonolysis products under dark conditions, while the lower spectra show the products from OH oxidation (254 nm UV). The most abundant compounds are labeled with the predicted formula and their retention time, however, the database contains these entries of all compounds down to 1 % relative peak intensity.

Panel (a) shows the results of the α -pinene oxidation experiments. Monomers produced during ozonolysis are mainly in the mass range between 140 and 210 Da, dimers are in the range between 300 and 400 Da. The major products during ozonolysis are pinic acid ($C_9H_{14}O_4$ at 8.79 min, Yu et al. (1999), level 1), terpenylic acid ($C_8H_{12}O_4$ at 6.67 min, Claeys et al. (2009), level 2), pinyndiaterpenyl ester ($C_{17}H_{26}O_8$ at 11.28 min, Kahnt et al. (2018), Yasmeen et al. (2010), level 2), $C_8H_{14}O_5$ (at 5.84 min), and $C_8H_{14}O_6$ (at 6.56 min). Oxidation by OH reduces the absolute signal intensity of most oxidation products (see Fig. S1). Furthermore, this oxidation environment prevents the production of dimers and certain monomers, and changes the relative abundance of several monomers. For example, the relative abundance of pinic acid, which is the major compound of the ozonolysis, decreases by 30 % under OH conditions. In contrast, the relative abundance of several other compounds increase, which indicates a higher relevance in the OH system, like terpenylic acid increasing to 100 % relative abundance,




Figure 2. Mass spectra of the detected products from the OFR experiments of five biogenic precursors (a) α -pinene, (b) β -pinene, (c) limonene, (d) 3-carene, (e) and *trans*-caryophyllene. The intensity is normalized to the highest signal of each chemical system. The most abundant compounds of each experiment are labeled with their predicted composition and the according retention time.

$C_{10}H_{16}O_5$ (at 9.28 min) increasing to 88% relative abundance, $C_8H_{14}O_5$ (at 5.84 min) increasing from 28% to 73%, and MBTCA increasing from 4% to 46%. Panel (b) shows the results of the β -pinene oxidation experiments. Here, many compounds are similar to the α -pinene oxidation products, with the exception of dimer formation during OH conditions. For both ozonolysis and OH oxidation, $C_9H_{14}O_4$ is the compound with the highest relative abundance, although the chromatography resolves different isomers: in the upper spectrum (O₃) pinic acid has a relative abundance of 100%, whereas homoterpenylic acid ($C_9H_{14}O_4$ at 7.58 min, Yasmeen et al. (2010), level 2) has a relative abundance of 17%. In the lower spectrum (OH), the relative abundances are reversed, with 100% at 7.57 minutes and 65% at 8.78 minutes. This indicates that different oxidation conditions of the same precursor result in different isomers of $C_9H_{14}O_4$, which can only be resolved with chromatographic separation. Furthermore, homoterpenylic acid does not appear in any other experiment in higher amounts, for which reason it can be used as a specific β -pinene tracer. While the most prominent dimer (pinyl-diterpenyl ester) appears analogously to the α -pinene-system during ozonolysis, here, under OH conditions, β -pinene oxidation results in dimer oxidation products like $C_{19}H_{32}O_9$ (at 13.23 min), in contrast to no dimers in α -pinene-system.

Panel (c) shows the results of the limonene oxidation experiments. In contrast to the other three monoterpenes no dimers were formed, which is in general agreement with Hammes et al. (2019). The ozonolysis shows three major products, hydroxyterpenylic acid ($C_8H_{12}O_5$ at 5.57 min, Kahnt et al. (2014), level 2), ketolimononic acid ($C_9H_{14}O_4$ at 6.44 min, Yasmeen et al. (2011), level 2), and $C_{10}H_{16}O_5$ (at 6.85 min). In the OH system ketolimononic acid becomes the major compound whereas the

215 intensity of hydroxyterpenylic acid increases clearly. Analogous to the β -pinene oxidation, the $C_9H_{14}O_4$ isomer ketolimononic acid can be used as specific limonene tracer due to the missing appearance of this isomer in other experiments.

Panel (d) shows the results of the 3-carene oxidation experiments. Three monomers are the most prominent products in both systems: caric acid ($C_9H_{14}O_4$ at 9.62 min, Yasmeen et al. (2011), level 2), hydroxy-3-caric acid ($C_{10}H_{16}O_4$ at 8.72 min, Larsen et al. (2001), level 3), and 3-caronic acid ($C_{10}H_{16}O_3$ at 10.27 min, Larsen et al. (2001), level 3). The four dimers
220 $C_{19}H_{30}O_5$ (at 12.76 min), $C_{19}H_{28}O_7$ (at 12.77 min), $C_{20}H_{30}O_5$ (at 13.74 min), and $C_{19}H_{28}O_9$ (at 11.40 min) appear during ozonolysis from which three are also reported by Thomsen et al. (2021), tentatively identified as dimers from 3-carene. Under OH conditions the dimers disappear with the exception of $C_{19}H_{30}O_5$.

In addition to the four monoterpenes, we investigated the composition of sesquiterpene-SOA from *trans*-caryophyllene ($C_{15}H_{24}$). During ozonolysis we find one major and four minor products in the mass range between 198 and 302 Da (Fig. 2e).
225 The major compound is tentatively identified as β -norcaryophyllonic acid ($C_{14}H_{22}O_4$ at 11.82 min, van Eijck et al. (2013), Jaoui et al. (2003), level 3). In contrast, the reaction with OH leads to one major and seven minor products in a range from 118 Da up to 382 Da. The major compound $C_{17}H_{26}O_4$ (at 14.03 min) also appears during ozonolysis but only with a relative abundance of 8 %.

Considering BVOC oxidation in general, it is worth mentioning that different isomers of $C_9H_{14}O_4$ are clearly separated by
230 the chromatographic system and we can use them as specific tracers for different BVOCs in the *aerosolomics* database. These and even more isomers are present in ambient filter samples (Fig. S2), which demonstrates the necessity of chromatographic separation if an unambiguous assignment is desired. Furthermore, ion source dimerization is a known phenomenon that hinders the unambiguous identification of atmospheric dimers, or leads to misinterpretation of results from direct-injection HESI. Based on the knowledge of the exact m/z and the mass dependence of the retention time, we can assign ion source related dimers to
235 the associated atmospheric monomer. This allows us an unambiguous distinction between monomers and covalently bonded “real” dimers (Fig. S3).

3.1.2 SOA compounds from oxidation of anthropogenic VOCs

We investigated the composition of SOA from the anthropogenic VOCs 1,2,4-trimethylbenzene, toluene, *o*-xylene, and naphthalene. We carried out only OH-oxidation of AVOCs, because oxidation of aromatic compounds by O_3 is negligible. The
240 filter criteria were similar to the experiments with BVOCs and the resulting spectra are shown in Fig. 3. All experiments show a noticeably lower number of oxidation products compared to biogenic precursors. We observe dimers only in the 1,2,4-trimethylbenzene and the *o*-xylene system.

Panel (a) shows the results of the 1,2,4-trimethylbenzene oxidation. The two most prominent compounds are $C_9H_8O_4$ (at 9.04 min) and $C_5H_6O_4$ (at 4.29 min). The remaining compounds play a minor role due to relative abundances less than
245 10 %.

Panel (b) shows the results from the oxidation of toluene. The five most prominent compounds show a higher relative abundance than 40 % and no compound has a lower abundance than 15 %. Most of these products are highly oxygenated with more than four oxygen atoms with the exception of the highest signal corresponding to $C_5H_6O_3$. All the small ($C \leq 5$) highly




Figure 3. Mass spectra of detected products from OFR experiments of four anthropogenic precursors (a) 1,2,4-trimethylbenzene, (b) toluene, (c) *o*-xylene, (d) and naphthalene. The intensity is normalized to the highest signal of each chemical system. The most abundant compounds of each experiment are labeled with their predicted composition and the according retention time.

oxygenated molecules exhibit also a high polarity ($t_R < 2$ min) compared to the oxidation products of other anthropogenic precursors.

Panel (c) shows the results of the *o*-xylene oxidation, the anthropogenic precursor with the largest number of detected oxidation products ($n = 52$) of the four investigated AVOCs. While no composition could be assigned by the NTA software for the highest signal, due to an invalid isotopic pattern, the most abundant product in panel (d) appears on the same mass trace and the same retention time. This peak was identified as phthalic acid ($C_8H_6O_4$, level 1), which is described as naphthalene SOA tracer by Al-Naiema et al. (2020).

Panel (d) shows the oxidation of naphthalene resulting in the main oxidation product phthalic acid. All other compounds have a relative abundance smaller than 20%. It is worth mentioning that two isomers of $C_8H_6O_3$ appear with a similar relative abundance, but with two distinguishable retention times.




Figure 4. Molecular fingerprint of the representative selection of the field campaign samples. The area of the scatters represents the qualitative mean signal intensity on a linear scale. (a) Retention time vs. molecular mass. (b) Kroll diagram. (c) Van Krevelen diagram. (d) Mass difference between calculated and measured molecular mass (Δ mass) vs. molecular mass. (e) Kendrick mass defect plot. Panels (b), (c), and (d) only include compounds with a predicted composition.

3.2 Aerosolomics-database application on ambient samples

260 3.2.1 Fingerprint

The NTA of the representative selection of the Vienna field campaign extracts results in 1312 compounds shown in panel (a) of Fig. 4 as retention time as a function of molecular mass. The scatter size represents the mean signal intensity of the 10 measurements on a linear scale. The majority are CHO compounds (54.7%), followed by CHOS (17.3%), CHNOS (13.1%), and unidentified (12.3%) compounds (Fig. 5) spanning the range from 100 Da to 350 Da. CHNO and other compounds, especially
 265 halogenated hydrocarbons, play a minor role with proportions of 2.2% and 0.4%, respectively. The entire range of retention times indicates a broad mixture of polarities. Among the most common signals are terpene-derived nitrooxy organosulfates like $C_{10}H_{17}NO_7S$ (Wang et al., 2021), isoprene-derived CHOS like $C_2H_4O_6S$ (Claeys and Maenhaut, 2021) as well as monoterpene oxidation products like MBTCA, terpenylic acid, and pinic acid.

Beside the two series of isoprene-derived CHNOS isomers highlighted in Sect. 3.3.1, three additional series of isomers can be found in the molecular fingerprint: $C_{10}H_{17}NO_7S$ (295 Da), $C_{10}H_{17}NO_9S$ (327 Da), and $C_{10}H_{17}NO_{10}S$ (343 Da) are all described as monoterpene derived SOA (Surratt et al., 2008). Especially $C_{10}H_{17}NO_7S$ plays a prominent role, due to the overall maximum intensity, and it illustrates the anthropogenic influence on the oxidation products from BVOCs in the presence of NO_x and SO_2 . The appearance of the ions NO_3^- and HSO_4^- in the MS^2 spectra prove that these compounds are nitrooxy organosulfates. Overall, 86 % of the CHNOS compounds show both ions in their MS^2 spectra and can also be attributed to this group.

The average carbon oxidation state (\overline{OS}_C) according to Kroll et al. (2011) is in the range between -1.5 and 3 , with the exception of two small CHOS compounds with an \overline{OS}_C of up to 4 (Fig. 4b). The majority of the CHO compounds consist of equal to or less than ten carbon atoms, while the majority of the CHOS compounds consist of equal to or less than five carbon atoms. This pattern indicates the importance of monoterpenes and isoprene as SOA precursors.

The Van Krevelen diagram (Fig. 4c) clarifies the influence of biogenic precursors. On the one hand the majority of the CHO compounds appear in the H/C range between 1.2 and 1.8 , described as a surrogate for biogenic SOA (Daellenbach et al., 2019; Kourtev et al., 2015). On the other hand, compounds with a $H/C < 1.2$, indicating aromatic character, are rare despite proximity to an airport and a large refinery. Even though, several CHNO tracers for biomass burning could be identified, they only play a minor role with regard to noticeably lower signal intensities: $C_7H_5NO_5$ (nitrosalicylic acid, level 3), $C_7H_7NO_4$ (methylnitrocatechol, level 3), $C_7H_6N_2O_5$ (methyl dinitrophenol/dinitrocresol, level 2), $C_7H_6N_2O_6$, $C_6H_5NO_4$ (nitrocatechol, level 2), $C_6H_4N_2O_5$ (dinitrophenol, level 2), $C_8H_7NO_4$ (methylnitrobenzoic acid, level 3), and $C_8H_9NO_5$ (Kitanovski et al., 2012; Salvador et al., 2021).

In order to evaluate optimal settings for chemical composition prediction during NTA and to avoid false predictions, the mass difference between calculated and measured molecular mass (Δ_{mass}) has to be considered. Figure 4d shows that over the entire mass range only a slight shift to negative values (median = -0.29 ppm) can be observed within the space between -1 and 1 ppm deviation. Within the set of allowed elemental compositions an enlargement of the range (± 1 ppm) would increase false predictions due to the increased mathematical possibility of other elemental combinations. The Kendrick mass defect (KMD) plot (Fig. 4e) allows the identification of homologous series. Members of such series have the same Kendrick mass defect (Kendrick, 1963) like $C_nH_{2n-2}O_3$ with $n = 5-15$ (KMD = 0.082), $C_nH_{2n-4}O_4$ with $n = 4-16$ (KMD = 0.119), or $C_nH_{2n-2}O_5$ with $n = 3-14$ (KMD = 0.128). Also CHOS compounds form homologous series like $C_nH_{2n}O_5S$ with $n = 2-8$ (KMD = 0.178), or $C_nH_{2n}O_6S$ with $n = 2-10$ (KMD = 0.201). The appearance of these homologous series can be interpreted as oxygenated aliphatic hydrocarbons of fossil origin, as homologous series of (sulfur-containing) aliphatics can be detected in crude oil.

3.2.2 Aerosolomics database assignment

The results from the NTA of the ambient samples were compared with the outcome from the PAM-OFR experiments in order to estimate the contribution of oxidation products formed by several precursors to ambient $PM_{2.5}$. Out of a total of 580 detected CHO compounds, we can assign 108 CHO compounds to biogenic precursors and 24 CHO compounds to anthro-




Figure 5. Contribution of different molecular formula groupings to the mean signal intensity of the molecular fingerprint from the representative selection shown in Fig. 4. Compounds of the CHO group were compared to the libraries of the individual OFR experiments presented in Sect. 3.1 and dedicated to a plausible SOA precursor and oxidation condition shown in the bar.

305 pogenic VOCs using our *aerosolomics* database. The bar plot in Fig. 5 divides the CHO fraction into the different contributions examined. 40.8 % of the mean signal intensity can be attributed to SOA originated from biogenic precursors (gradations of green and yellow). Out of these, 26.5 % account for experiments with OH oxidation and 14.3 % for ozonolysis experiments. The two major biogenic precursors are α - and β -pinene with a respective share of 17.6 and 13.2 %. The remaining shares are distributed among *trans*-caryophyllene products (4.2 %), among limonene products (3.6 %), and among 3-carene products (2.2 %). Anthropogenic precursors (gradations of magenta) play a minor role with a total contribution of 5.7 %. From that 2 % can be assigned to *o*-xylene, 1.9 % to naphthalene, 1.2 % to 1,2,4-trimethylbenzene, and 0.6 % to toluene. 53.5 % of the mean 310 signal intensity could not be assigned (diagonally hatched).

The majority of the assigned compounds from the representative selection of the Vienna field campaign have molecular masses smaller than 250 Da and a retention time lower than 10 minutes (Fig. S4a). The mean number of carbon atoms is 9 and the mean \overline{OS}_C is -0.4 (Fig. S4b). SOA originated from biogenic precursors is located in the H/C area between 1.2 and 1.8 (Fig. S4c), while compounds with $H/C < 1.2$ can be of aromatic character. The observation that 19 % of the CHO compounds 315 (number-wise) in the database are responsible for nearly 50 % of the mean signal intensity demonstrates the high relevance of the investigated VOCs in SOA formation. Nevertheless, a few compounds with high signal intensities remain unassigned, such as $C_8H_{14}O_5$, $C_7H_6O_2$, $C_9H_{16}O_4$ or the tentatively isoprene derived $C_4H_6O_5$ and $C_4H_8O_4$ (Claeys and Maenhaut, 2021; Krechmer et al., 2015). Considering the retention behaviour and the molecular mass of these unassigned compounds (Fig. S4a) we expect isoprene as well as other monoterpenes to be promising candidates closing this gap. In addition, the precursors 320 already used should be investigated under varying chemical conditions, like further oxidants or more complex mixtures of VOCs. A comprehensive study of isoprene oxidation is planned and the outcome will be uploaded to the *aerosolomics* database in the near future.




Figure 6. Standardized matrix of the detected compounds from the NTA of the filter samples from the Vienna field campaign. Parts of the horizontal dendrogram of the filter samples are shaded in grey for night cluster, yellow for day cluster and blue for clustered samples at decreasing temperatures. The vertical compound clusters are separated into four subclusters according to day- and night-time chemistry, organosulfates, and temperature dependency. For a better differentiation the dendrograms are coloured randomly.

Beside CHO compounds, CHOS and CHNOS compounds play an important role in the overall composition of suburban SOA. Based on these outcome, further experiments with various VOCs and complex mixtures including inorganic trace gases need to be performed and the results have to be added to the *aerosolomics* database.

3.3 Hierarchical cluster analysis

In Fig. 6 we show the results of the HCA (as a heatmap with dendrograms), with the ambient $PM_{2.5}$ filter samples from the Vienna field campaign on the horizontal axis, and the detected compounds of the NTA on the vertical axis. The color code of the heatmap represents the standardized values of the integrated peak intensities after background correction. We find that the driving factors for the clustering of the filter samples are the wind direction overlaid by the diurnal cycle, as well as the influence of decreasing temperatures, as explained in the following section. Clusters that contain mostly night-time samples are shaded by a grey background. The ones that contain mainly daytime samples are shaded by a yellow background, while samples during a low-temperature period are shaded by a blue background. Over the entire period the mean $PM_{2.5}$ mass concentration was $8.7 \pm 4.4 \mu\text{g m}^{-3}$, the mean NO_x concentration was $15.4 \pm 16.7 \mu\text{g m}^{-3}$, the mean SO_2 concentration was $1.5 \pm 1.2 \mu\text{g m}^{-3}$, and the mean CO concentration was $0.17 \pm 0.03 \text{ mg m}^{-3}$. A detailed time series of meteorological data, $PM_{2.5}$, and trace gas concentrations are shown in Fig. S5. Additionally, Fig. S6 shows the distribution of the wind direction of sample clusters a to i. With exception of sample clusters b and f, all clusters show a predominant wind direction. The overall mean $PM_{2.5}$ mass concentration was higher during collection in sample cluster I ($10.3 \pm 4.1 \mu\text{g m}^{-3}$) compared to sample cluster II ($5.0 \pm 2.4 \mu\text{g m}^{-3}$).

340 3.3.1 Compound cluster I – daytime chemistry

Figure 7a–c illustrates the molecular fingerprint of the 373 compounds occurring in compound cluster I. 79 % of the mean signal intensity is caused by CHO compounds, 16 % is unidentified, and 4 % is caused by CHOS compounds. The molecular masses of the compounds are in the range between 100 and 350 Da. CHO compounds have mainly masses smaller than 250 Da and a mean bulk composition of $C_{8.5}H_{12.5}O_5$ which is in good agreement with the appearance of monomers from monoterpene oxidation during daytime. The number of carbon atoms ranges mainly between 4 and 10 (Fig. 7b), indicating biogenic VOCs, like monoterpenes, as potential precursors. The compounds cover a large range in volatility, with $\log_{10} C^*$ values between -4.0 and $6.5 \mu\text{g m}^{-3}$ (Fig. 8a), corresponding mainly to LVOC–IVOC with an intensity weighted $\log_{10} C^*$ mean value of $2.3 \pm 1.6 \mu\text{g m}^{-3}$. Only a small fraction of the detected compounds have a H/C smaller 1.2, indicating an aromatic character. The five most abundant compounds with a predicted composition are MBTCA (*aerosolomics*-database library: α -pinene + OH), hydroxyterpenylic acid, $C_5H_6O_7$, $C_4H_6O_5$, and $C_7H_{10}O_5$. All five compounds are characterized as biogenic SOA-compounds derived from isoprene or monoterpenes (Chen et al., 2018, 2020; Ehn et al., 2012; Müller et al., 2012; Qi et al., 2020).

CHOS compounds appear less important in this compound cluster, based on a 5 % contribution to the mean signal intensity. Nevertheless, about 45 % of the mean CHOS signal intensity can be attributed to monoterpene and isoprene derived SOA (Brüggemann et al., 2020).

It is also remarkable that compounds appearing in this cluster show low standardized values in sample cluster II. Low standardized values indicate lower signal intensities of the MS measurements. However, it needs to be considered that signal intensities are not directly quantitative to its concentration. It is well known, that the ionization efficiency of HESI varies greatly for several compounds as well as compound classes (Kenseth et al., 2020; Ma et al., 2022). Nevertheless, variation in the intensity of a single compound or class of compounds can be qualitatively interpreted as variation of its concentration. Compared to a mean temperature of 25.5 ± 4.8 °C during sample cluster I, the mean temperature of sample cluster II is noticeably lower with 19.3 ± 4.3 °C. Especially the subcluster including the last six samples of the entire field campaign shows the lowest standardized values and the lowest mean temperature of 16.1 ± 3.3 °C. The correlation of lower temperatures and low standardized values can be explained due to the temperature dependency of terpene emissions from plants (Holzke et al., 2006) resulting in a lower biogenic SOA burden in the atmosphere.

365 3.3.2 Compound cluster II – night-time chemistry

The mean signal intensity of the 134 compounds appearing in compound cluster II (Fig. 7d–f) is mainly caused by CHO (78 %) and CHNO (14 %) compounds. The molecular masses reach up to 440 Da and the mean bulk composition is $C_{11.9}H_{18.2}O_{5.1}$. The \overline{OS}_C of these CHO compounds are in the range between -1 and 0.5 (Fig. 7e). Analogously to compound cluster I, the H/C is in the range between 1.2 and 1.8 (Fig. 7f), indicating biogenic SOA. $\log_{10} C^*$ values range from -7.1 to $6.7 \mu\text{g m}^{-3}$ (Fig. 8b) corresponding mainly to LVOC–IVOC with an intensity weighted mean of $2.3 \pm 2.7 \mu\text{g m}^{-3}$. The most prominent compound is pinic acid (*aerosolomics*-database library: α -pinene/ β -pinene + O_3) with a contribution of about 24 % to the

mean signal intensity. It is well described as a α - and β -pinene ozonolysis product (Christoffersen et al., 1998; Glasius et al., 2000).

In contrast to sample cluster I, compounds from nighttime chemistry form a higher ratio of dimers with oxidation products up to 20 carbon atoms (Fig. 7e). 14 of 23 CHO dimers occurring in compound cluster II are also reported from OFR experiments performed by Kristensen et al. (2016) and agree with our findings from the OFR experiments described in Sect. 3.1. These 14 dimers are responsible for 72 % of the mean dimer signal intensity. Among them are pinyl-diaterpenyl ester $C_{17}H_{26}O_8$ (*aerosolomics*-database library: α -pinene/ β -pinene + O_3) and $C_{19}H_{28}O_7$ (*aerosolomics*-database library: α -pinene/3-carene + O_3), both of which are described as esters between pinic acid and terpenylic acid (Gao et al., 2010) or pinic acid and hydroxypinonic acid (Müller et al., 2008), respectively. Furthermore, Yasmeeen et al. (2010) reported pinyl-diaterpenyl ester in nighttime ambient aerosol.

While during daytime the main oxidant of VOCs is OH, at night-time O_3 and NO_3 dominate the oxidation process, resulting in the formation of organonitrates (Kiendler-Scharr et al., 2016). In accordance, we observed several nitrogen containing monomers and dimers including NO_3^- in the corresponding MS^2 spectra. Hence, about 87 % of the CHNO compounds at night are tentatively identified as organonitrates ($R-ONO_2$).

3.3.3 Compound cluster III – organosulfates (CHOS)

Compound cluster III includes 279 compounds (Fig. 7g–i). The mean signal intensity of compound cluster III is mainly caused by CHOS compounds (60 %), followed by unidentified (29 %) and CHNOS compounds (7 %). The molecular masses of the compounds reaches from 120 to 350 Da, those with higher mean signal intensities show only a very limited retention (< 0.9 min) due to high polarity and represent about 72 % of the mean signal intensity. These compounds have ≤ 5 carbon atoms and an \overline{OS}_C between 0 and 4 (Fig. 7h). The generally higher H/C compared to other compound clusters, shown in panel (i), indicate a predominantly aliphatic character of the detected compounds. The saturation vapor pressure ($\log_{10} C^*$) ranges between -8.8 and $5.0 \mu\text{g m}^{-3}$ (Fig. 8c), which is noticeably lower compared to compound clusters I and II. The intensity-weighted mean of $\log_{10} C^*$ of this cluster yields $-0.9 \pm 2.0 \mu\text{g m}^{-3}$.

Several of the chemical formulas can be described as isoprene-derived SOA (Brüggemann et al., 2020; Chen et al., 2018; Nestorowicz et al., 2018; Riva et al., 2016; Surratt et al., 2007). From 174 CHOS compounds in compound cluster III, 163 MS^2 spectra were recorded. From those the vast majority (98 %) shows the fragment at m/z 96.9601 (HSO_4^-), indicating an organosulfate functional group ($R-OSO_3$). The remaining 2 % only show m/z 79.9573 ($SO_3^{\bullet-}$). While this sulfur trioxide radical anion can indeed occur in an organosulfate fragment spectra (Wang et al., 2019), it could also be originated from organosulfonates ($R-SO_3$) (Liang et al., 2020; Liu et al., 2015). Beside one high signal intensity of a CHNOS compound at 295 Da and 9.7 minutes ($C_{10}H_{17}NO_7S$), three further series of isomers are appearing in the CHOS-cluster: We identified six isomers of $C_5H_{10}N_2O_{11}S$, four isomers of $C_5H_{11}NO_9S$, and three isomers of $C_5H_9NO_8S$. All three chemical formulas can be described as isoprene-derived SOA (Nestorowicz et al., 2018; Surratt et al., 2007, 2008). Furthermore, three isomers of $C_5H_9NO_7S$ also appear in this compound cluster, but has not yet been described as isoprene-derived SOA.




Figure 7. Molecular fingerprint of compounds that appear in compound cluster I (a–c), II (d–f), III (g–i), IV (j–l), illustrated as retention time vs. molecular mass, Kroll diagram, and Van Krevelen diagram. The area of the scatters represent the mean signal intensity on a linear scale. The pie chart shows the proportions of the different molecular formula groupings.

405 The distinct increased standardized values of several Cluster-III-compounds in the heatmap between 8. August 2018, 17:00 (UTC, AS-17) and 10. August 2018, 05:00 (UTC, AS-19) can be explained by high SO_2 concentrations up to $18 \mu\text{g m}^{-3}$ around noon on 8 August 2018 (Fig. S5). The nearby airport as a main source can be excluded due to similar high SO_2 concentrations at the Stixneusiedl monitoring station (Umweltbundesamt GmbH, 2021) located 12 km southeast of the airport and thus in the upwind direction of the airport. It is also noticeable, that the standardized values of these Cluster-III-compounds are very low
 410 during northwest trajectories, analogously to compound cluster I (Sec. 3.3.1). Therefore, it is likely that this cluster is mainly linked to long-range transport of pollution from the south-east.

3.3.4 Compound cluster IV – decreasing temperature

Compounds occurring in compound cluster IV (Fig. 7j–l) show clearly increased standardized values during the last six filter samples of the field campaign and are mainly CHO (73 %) and unidentified (17 %) compounds. The majority of the Cluster-
 415 IV-compounds have molecular masses $< 200 \text{ Da}$. The $\overline{\text{OS}}_{\text{C}}$ of those compounds ranges between -1.5 and 1 while the numbers of carbon is less than 10 (Fig. 7k). The majority of the 174 compounds have a non-aromatic character, illustrated in




Figure 8. Calculated effective saturation mass concentrations of compounds appearing in compound cluster I-IV (a-d) with the intensity weighted mean and standard deviation.

panel (l). Dicarboxylic acids, like phthalic acid ($C_8H_6O_4$, level 1), succinic acid ($C_4H_6O_4$, level 3), or maleic/fumaric acid ($C_4H_4O_4$, level 3) are reported as tracers for emissions from biomass burning, vehicular exhaust and fossil fuel combustion (Zhao et al., 2018). Furthermore, this compound cluster contains the homologous series $C_{4-9}H_{6-16}O_4$, $C_{5-10}H_{8-18}O_3$, and
 420 $C_{5-9}H_{10-18}O_3$, which can be interpreted as oxygenated aliphatics of fossil origin.

Compound cluster IV includes $\log_{10} C^*$ -values between -8.6 and $6.5 \mu\text{g m}^{-3}$ (Fig. 8d) corresponding predominantly to SVOC-IVOC with an intensity weighted mean of $3.9 \pm 1.9 \mu\text{g m}^{-3}$. The high standardized values in the dendrogram of these compounds in sample subcluster i can be attributed by their intermediate volatility, which will only occur in the condensed particle phase at low ambient temperature. During sample cluster I, which is characterized by a higher mean temperature, those
 425 compounds would likely partition predominantly into the gas phase.

3.4 Challenges with measurements in HESI positive ionization mode

All results in this paper are based on negative ionization mode measurements. For a comprehensive assessment of the chemical composition of the investigated $PM_{2.5}$ measurements in positive ionization mode ((+)HESI) are essential with respect to relevant compound classes that are considerably better ionized in (+)HESI (e.g., organophosphates, phthalates, pesticides, and
 430 others). Issues due to strong fragmentation or ion-source adduct formation, which are explained in detail in the supplement,

highlight the need of further optimization of the NTA workflow to prevent false identifications and thus misinterpretations of the results.

4 Conclusions

A large fraction of ambient PM_{2.5} consists of anthropogenic and biogenic SOA. The chemical composition of this mixture is highly complex, which hinders the identification and attribution of single molecules to its precursors, potential sources, and formation pathways. In this study, we present two complementary approaches that enable complexity reduction, and identification of precursors, formation pathways, as well as partitioning effects of various IVOCs.

The introduced *aerosolomics* database for compound matching and precursor identification is based on PAM-OFR experiments with five biogenic and four anthropogenic precursors, as well as different oxidizing regimes (OH, O₃). In order to validate the method on ambient samples, we applied the database on a set of PM_{2.5} filter samples. Based on the average composition of these samples, we find that CHO compounds account for the largest proportion with about 55 % of the mean signal intensity. Approximately 45 % of the CHO compounds can be attributed to one of the investigated VOC precursors, which we confirmed by the compound matching procedure. Hence, the compounds are identified based on retention time, exact mass-to-charge ratio, isotopic pattern, and the MS² fragmentation spectrum.

On a one-month set of 12-hour filter samples, we performed a HCA to reduce the complexity due to the large number of compounds detected. The compounds were clustered when their intensities show similar behavior over time, which in turn indicates similar sources or (trans-)formation pathways. The clustering of the various PM_{2.5} filter samples was primarily driven by wind direction, as well as by the diurnal cycle (day/night) and temperature-driven partitioning changes. Known proxies for monoterpene ozonolysis, like pinic acid or α -pinene-derived dimers, were identified in the night-time compound cluster. The SOA aging tracer MBTCA appeared in the daytime compound cluster. A large number of sulfur-containing compounds were clustered together, and this cluster is clearly elevated during south-easterly wind direction. Small IVOCs were clustered and show a high intensity during a cold period. These observations are a proof-of-principle that (1) the presented *aerosolomics* database enables to identify tracers from the oxidation of different VOC precursors, and in combination with HCA we can (2) attribute different oxidation products to either night-time or daytime chemistry, (3) identify periods of multiphase-chemistry processes resulting in organosulfate formation, and (4) observe temperature-driven partitioning of IVOCs.

We would like to encourage the community to apply the database on their own samples. Therefore we provide the database as db-files, msp-files, and csv-files which allows the application of the database with Compound Discoverer, MZmine 3 or every self-build solution. As a community effort, further input to the database is desirable to improve our understanding of sources and formation of secondary organic aerosol.

Data availability. Data from this work are freely available at zenodo.org. DOI: 10.5281/zenodo.6623244

Author contributions. MT wrote the manuscript, ALV and MS advised on manuscript writing. MT and ALV conceptualized the study, MT, FB, FG, and MS were responsible for the laboratory experiments, sample preparation, and measurements. Data evaluation was done by MT. ALV directed the project administration. All authors commented on the manuscript and contributed to the scientific discussion.

Competing interests. The authors declare that they have no conflict of interest.

465 *Financial support.* This work was supported by Emmy Noether-Programm (Deutsche Forschungsgemeinschaft), project number 410009325. This open-access publication was funded by the Goethe-University Frankfurt.

Acknowledgements. We thank Sophia Brilke, Dominik Stolzenburg, Paulus Bauer, and Paul Winkler (Faculty of Physics, University of Vienna), for ambient filter sampling and trace gas monitoring. We also thank the working group of Joachim Curtius (Institute for Atmospheric and Environmental Sciences, Goethe-University Frankfurt) for experimental support.

470 References

- Al-Naiema, I. M., Offenberg, J. H., Madler, C. J., Lewandowski, M., Kettler, J., Fang, T., and Stone, E. A.: Secondary Organic Aerosols from Aromatic Hydrocarbons and their Contribution to Fine Particulate Matter in Atlanta, Georgia, *Atmos. Environ.*, 223, <https://doi.org/10.1016/j.atmosenv.2019.117227>, 2020.
- Brüggemann, M., Xu, R., Tilgner, A., Kwong, K. C., Mutzel, A., Poon, H. Y., Otto, T., Schaefer, T., Poulain, L., Chan, M. N., and Herrmann, H.: Organosulfates in Ambient Aerosol: State of Knowledge and Future Research Directions on Formation, Abundance, Fate, and Importance, *Environ. Sci. Technol.*, 54, 3767–3782, <https://doi.org/10.1021/acs.est.9b06751>, 2020.
- Burkholder, J. B., Abbatt, J. P. D., Barnes, I., Roberts, J. M., Melamed, M. L., Ammann, M., Bertram, A. K., Cappa, C. D., Carlton, A. G., Carpenter, L. J., Crowley, J. N., Dubowski, Y., George, C., Heard, D. E., Herrmann, H., Keutsch, F. N., Kroll, J. H., McNeill, V. F., Ng, N. L., Nizkorodov, S. A., Orlando, J. J., Percival, C. J., Picquet-Varrault, B., Rudich, Y., Seakins, P. W., Surratt, J. D., Tanimoto, H., Thornton, J. A., Tong, Z., Tyndall, G. S., Wahner, A., Weschler, C. J., Wilson, K. R., and Ziemann, P. J.: The Essential Role for Laboratory Studies in Atmospheric Chemistry, *Environ. Sci. Technol.*, 51, 2519–2528, <https://doi.org/10.1021/acs.est.6b04947>, 2017.
- Chen, X., Xie, M., Hays, M. D., Edgerton, E., Schwede, D., and Walker, J. T.: Characterization of organic nitrogen in aerosols at a forest site in the southern Appalachian Mountains, *Atmos. Chem. Phys.*, 18, 6829–6846, <https://doi.org/10.5194/acp-18-6829-2018>, 2018.
- Chen, Y., Takeuchi, M., Nah, T., Xu, L., Canagaratna, M. R., Stark, H., Baumann, K., Canonaco, F., Prévôt, A. S. H., Huey, L. G., Weber, R. J., and Ng, N. L.: Chemical characterization of secondary organic aerosol at a rural site in the southeastern US: insights from simultaneous high-resolution time-of-flight aerosol mass spectrometer (HR-ToF-AMS) and FIGAERO chemical ionization mass spectrometer (CIMS) measurements, *Atmos. Chem. Phys.*, 20, 8421–8440, <https://doi.org/10.5194/acp-20-8421-2020>, 2020.
- Christoffersen, T., Hjorth, J., Horie, O., Jensen, N., Kotzias, D., Molander, L., Neeb, P., Ruppert, L., Winterhalter, R., Virkkula, A., Wirtz, K., and Larsen, B.: cis-pinic acid, a possible precursor for organic aerosol formation from ozonolysis of α -pinene, *Atmos. Environ.*, 32, 1657–1661, [https://doi.org/https://doi.org/10.1016/S1352-2310\(97\)00448-2](https://doi.org/https://doi.org/10.1016/S1352-2310(97)00448-2), 1998.
- Claeys, M. and Maenhaut, W.: Secondary Organic Aerosol Formation from Isoprene: Selected Research, Historic Account and State of the Art, *Atmosphere*, 12, <https://doi.org/10.3390/atmos12060728>, 2021.
- Claeys, M., Iinuma, Y., Szmigielski, R., Surratt, J. D., Blockhuys, F., Van Alsenoy, C., Böge, O., Sierau, B., Gómez-González, Y., Vermeylen, R., Van der Veken, P., Shahgholi, M., Chan, A. W. H., Herrmann, H., Seinfeld, J. H., and Maenhaut, W.: Terpenylic Acid and Related Compounds from the Oxidation of α -Pinene: Implications for New Particle Formation and Growth above Forests, *Environ. Sci. Technol.*, 43, 6976–6982, <https://doi.org/10.1021/es9007596>, 2009.
- Daellenbach, K. R., Kourtev, I., Vogel, A. L., Bruns, E. A., Jiang, J., Petäjä, T., Jaffrezo, J.-L., Aksoyoglu, S., Kalberer, M., Baltensperger, U., El Haddad, I., and Prévôt, A. S. H.: Impact of anthropogenic and biogenic sources on the seasonal variation in the molecular composition of urban organic aerosols: a field and laboratory study using ultra-high-resolution mass spectrometry, *Atmos. Chem. Phys.*, 19, 5973–5991, <https://doi.org/10.5194/acp-19-5973-2019>, 2019.
- Ditto, J. C., Barnes, E. B., Khare, P., Takeuchi, M., Joo, T., Bui, A. A. T., Lee-Taylor, J., Eris, G., Chen, Y., Aumont, B., Jimenez, J. L., Ng, N. L., Griffin, R. J., and Gentner, D. R.: An omnipresent diversity and variability in the chemical composition of atmospheric functionalized organic aerosol, *Commun. Chem.*, 1, 75, <https://doi.org/10.1038/s42004-018-0074-3>, 2018.
- Ehn, M., Kleist, E., Junninen, H., Petäjä, T., Lönn, G., Schobesberger, S., Dal Maso, M., Trimborn, A., Kulmala, M., Worsnop, D. R., Wahner, A., Wildt, J., and Mentel, T. F.: Gas phase formation of extremely oxidized pinene reaction products in chamber and ambient air, *Atmos. Chem. Phys.*, 12, 5113–5127, <https://doi.org/10.5194/acp-12-5113-2012>, 2012.

- Fiehn, O.: Metabolomics – the link between genotypes and phenotypes, *Plant Mol. Biol.*, 48, 155–171, <https://doi.org/10.1023/A:1013713905833>, 2002.
- 510 Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C., Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y., Schaap, M., Slowik, J. G., Spracklen, D. V., Vignati, E., Wild, M., Williams, M., and Gilardoni, S.: Particulate matter, air quality and climate: lessons learned and future needs, *Atmos. Chem. Phys.*, 15, 8217–8299, <https://doi.org/10.5194/acp-15-8217-2015>, 2015.
- Gao, Y., Hall, W. A., and Johnston, M. V.: Molecular Composition of Monoterpene Secondary Organic Aerosol at Low Mass Loading, *Environ. Sci. Technol.*, 44, 7897–7902, <https://doi.org/10.1021/es101861k>, 2010.
- 515 Glasius, M., Lahaniati, M., Calogirou, A., Di Bella, D., Jensen, N. R., Hjorth, J., Kotzias, D., and Larsen, B. R.: Carboxylic Acids in Secondary Aerosols from Oxidation of Cyclic Monoterpenes by Ozone, *Environ. Sci. Technol.*, 34, 1001–1010, <https://doi.org/10.1021/es990445r>, 2000.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., 520 Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, <https://doi.org/10.5194/acp-9-5155-2009>, 2009.
- Hammes, J., Lutz, A., Mentel, T., Faxon, C., and Hallquist, M.: Carboxylic acids from limonene oxidation by ozone and hydroxyl radicals: insights into mechanisms derived using a FIGAERO-CIMS, *Atmos. Chem. Phys.*, 19, 13 037–13 052, [https://doi.org/10.5194/acp-19-](https://doi.org/10.5194/acp-19-13037-2019) 525 13037-2019, 2019.
- HighChem LLC: mzCloud Advanced Mass Spectral Database, <https://www.mzcloud.org>, Online; accessed 10 August 2021, 2013–2021.
- Holzke, C., Dindorf, T., Kesselmeier, J., Kuhn, U., and Koppmann, R.: Terpene emissions from European beech (*shape Fagus sylvatica~L.*): Pattern and Emission Behaviour Over two Vegetation Periods, *J. Atmos. Chem.*, 55, 81–102, <https://doi.org/10.1007/s10874-006-9027-9>, 2006.
- 530 Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbazade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prévôt, A. S. H.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218–222, <https://doi.org/10.1038/nature13774>, 2014.
- Isaacman-VanWertz, G. and Aumont, B.: Impact of organic molecular structure on the estimation of atmospherically relevant physicochemical parameters, *Atmos. Chem. Phys.*, 21, 6541–6563, <https://doi.org/10.5194/acp-21-6541-2021>, 2021.
- 535 Jaoui, M., Leungsakul, S., and Kamens, R. M.: Gas and Particle Products Distribution from the Reaction of β -Caryophyllene with Ozone, *J. Atmos. Chem.*, 45, 261–287, <https://doi.org/10.1023/A:1024263430285>, 2003.
- Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, 540 V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., E., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimojo, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn,

- 545 A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525–1529, <https://doi.org/10.1126/science.1180353>, 2009.
- Johnson, C. H., Ivanisevic, J., and Siuzdak, G.: Metabolomics: beyond biomarkers and towards mechanisms, *Nat. Rev. Mol.*, 17, 451–459, <https://doi.org/10.1038/nrm.2016.25>, 2016.
- Kahnt, A., Iinuma, Y., Blockhuys, F., Mutzel, A., Vermeylen, R., Kleindienst, T. E., Jaoui, M., Offenberg, J. H., Lewandowski, M., Böge, O., Herrmann, H., Maenhaut, W., and Claeys, M.: 2-Hydroxyterpenylic Acid: An Oxygenated Marker Compound for α -Pinene Secondary Organic Aerosol in Ambient Fine Aerosol, *Environ. Sci. Technol.*, 48, 4901–4908, <https://doi.org/10.1021/es500377d>, 2014.
- 550 Kahnt, A., Vermeylen, R., Iinuma, Y., Safi Shalamzari, M., Maenhaut, W., and Claeys, M.: High-molecular-weight esters in α -pinene ozonolysis secondary organic aerosol: structural characterization and mechanistic proposal for their formation from highly oxygenated molecules, *Atmos. Chem. Phys.*, 18, 8453–8467, <https://doi.org/10.5194/acp-18-8453-2018>, 2018.
- Kang, E., Root, M. J., Toohey, D. W., and Brune, W. H.: Introducing the concept of Potential Aerosol Mass (PAM), *Atmos. Chem. Phys.*, 7, 5727–5744, <https://doi.org/10.5194/acp-7-5727-2007>, 2007.
- 555 Kari, E., Hao, L., Ylisirniö, A., Buchholz, A., Leskinen, A., Yli-Pirilä, P., Nuutinen, I., Kuusalo, K., Jokiniemi, J., Faiola, C. L., Schobesberger, S., and Virtanen, A.: Potential dual effect of anthropogenic emissions on the formation of biogenic secondary organic aerosol (BSOA), *Atmos. Chem. Phys.*, 19, 15 651–15 671, <https://doi.org/10.5194/acp-19-15651-2019>, 2019.
- Kendrick, E.: A Mass Scale Based on $\text{CH}_2 = 14.0000$ for High Resolution Mass Spectrometry of Organic Compounds., *Anal. Chem.*, 35, 2146–2154, <https://doi.org/10.1021/ac60206a048>, 1963.
- 560 Kenseth, C. M., Hafeman, N. J., Huang, Y., Dalleska, N. F., Stoltz, B. M., and Seinfeld, J. H.: Synthesis of Carboxylic Acid and Dimer Ester Surrogates to Constrain the Abundance and Distribution of Molecular Products in α -Pinene and β -Pinene Secondary Organic Aerosol, *Environ. Sci. Technol.*, 54, 12 829–12 839, <https://doi.org/10.1021/acs.est.0c01566>, 2020.
- Kiendler-Scharr, A., Mensah, A. A., Friese, E., Topping, D., Nemitz, E., Prevot, A. S. H., Äijälä, M., Allan, J., Canonaco, F., Canagaratna, M., Carbone, S., Crippa, M., Dall'Osto, M., Day, D. A., De Carlo, P., Di Marco, C. F., Elbern, H., Eriksson, A., Freney, E., Hao, L., Herrmann, H., Hildebrandt, L., Hillamo, R., Jimenez, J. L., Laaksonen, A., McFiggans, G., Mohr, C., O'Dowd, C., Otjes, R., Ovadnevaite, J., Pandis, S. N., Poulain, L., Schlag, P., Sellegri, K., Swietlicki, E., Tiitta, P., Vermeulen, A., Wahner, A., Worsnop, D., and Wu, H.-C.: Ubiquity of organic nitrates from nighttime chemistry in the European submicron aerosol, *Geophys. Res. Lett.*, 43, 7735–7744, <https://doi.org/https://doi.org/10.1002/2016GL069239>, 2016.
- 570 Kitanovski, Z., Grgić, I., Vermeylen, R., Claeys, M., and Maenhaut, W.: Liquid chromatography tandem mass spectrometry method for characterization of monoaromatic nitro-compounds in atmospheric particulate matter, *J. Chromatogr. A*, 1268, 35–43, <https://doi.org/https://doi.org/10.1016/j.chroma.2012.10.021>, 2012.
- Kourtchev, I., Doussin, J.-F., Giorio, C., Mahon, B., Wilson, E. M., Maurin, N., Pangui, E., Venables, D. S., Wenger, J. C., and Kalberer, M.: Molecular composition of fresh and aged secondary organic aerosol from a mixture of biogenic volatile compounds: a high-resolution mass spectrometry study, *Atmos. Chem. Phys.*, 15, 5683–5695, <https://doi.org/10.5194/acp-15-5683-2015>, 2015.
- 575 Krechmer, J. E., Coggon, M. M., Massoli, P., Nguyen, T. B., Crounse, J. D., Hu, W., Day, D. A., Tyndall, G. S., Henze, D. K., Rivera-Rios, J. C., Nowak, J. B., Kimmel, J. R., Mauldin, R. L., Stark, H., Jayne, J. T., Sipilä, M., Junninen, H., St. Clair, J. M., Zhang, X., Feiner, P. A., Zhang, L., Miller, D. O., Brune, W. H., Keutsch, F. N., Wennberg, P. O., Seinfeld, J. H., Worsnop, D. R., Jimenez, J. L., and Canagaratna, M. R.: Formation of Low Volatility Organic Compounds and Secondary Organic Aerosol from Isoprene Hydroxyhydroperoxide Low-NO
- 580 Oxidation, *Environ. Sci. Technol.*, 49, 10 330–10 339, <https://doi.org/10.1021/acs.est.5b02031>, 2015.

- Kristensen, K., Watne, A. K., Hammes, J., Lutz, A., Petäjä, T., Hallquist, M., Bilde, M., and Glasius, M.: High-Molecular Weight Dimer Esters Are Major Products in Aerosols from α -Pinene Ozonolysis and the Boreal Forest, *Environ. Sci. Technol. Lett.*, 3, 280–285, <https://doi.org/10.1021/acs.estlett.6b00152>, 2016.
- 585 Kroll, J. H., Donahue, N. M., Jimenez, J. L., Kessler, S. H., Canagaratna, M. R., Wilson, K. R., Altieri, K. E., Mazzoleni, L. R., Wozniak, A. S., Bluhm, H., Mysak, E. R., Smith, J. D., Kolb, C. E., and Worsnop, D. R.: Carbon oxidation state as a metric for describing the chemistry of atmospheric organic aerosol, *Nature Chem.*, 3, 133–139, <https://doi.org/10.1038/nchem.948>, 2011.
- Lambe, A. T., Ahern, A. T., Williams, L. R., Slowik, J. G., Wong, J. P. S., Abbatt, J. P. D., Brune, W. H., Ng, N. L., Wright, J. P., Croasdale, D. R., Worsnop, D. R., Davidovits, P., and Onasch, T. B.: Characterization of aerosol photooxidation flow reactors: heterogeneous oxidation, secondary organic aerosol formation and cloud condensation nuclei activity measurements, *Atmos. Meas. Tech.*, 4, 445–461, <https://doi.org/10.5194/amt-4-445-2011>, 2011.
- 590 Larsen, B. R., Di Bella, D., Glasius, M., Winterhalter, R., Jensen, N. R., and Hjorth, J.: Gas-Phase OH Oxidation of Monoterpenes: Gaseous and Particulate Products, *J. Atmos. Chem.*, 38, 231–276, <https://doi.org/10.1023/A:1006487530903>, 2001.
- Li, M., Karu, E., Brenninkmeijer, C., Fischer, H., Lelieveld, J., and Williams, J.: Tropospheric OH and stratospheric OH and Cl concentrations determined from CH₄, CH₃Cl, and SF₆ measurements, *npj Clim. Atmos. Sci.*, 1, 29, <https://doi.org/10.1038/s41612-018-0041-9>, 2018.
- 595 Li, Y., Pöschl, U., and Shiraiwa, M.: Molecular corridors and parameterizations of volatility in the chemical evolution of organic aerosols, *Atmos. Chem. Phys.*, 16, 3327–3344, <https://doi.org/10.5194/acp-16-3327-2016>, 2016.
- Liang, M., Xian, Y., Wang, B., Hou, X., Wang, L., Guo, X., Wu, Y., and Dong, H.: High throughput analysis of 21 perfluorinated compounds in drinking water, tap water, river water and plant effluent from southern China by supramolecular solvents-based microextraction coupled with HPLC-Orbitrap HRMS, *Environ. Pollut.*, 263, 114–139, <https://doi.org/10.1016/j.envpol.2020.114389>, 2020.
- 600 Liu, Y., Pereira, A. D. S., and Martin, J. W.: Discovery of C₅–C₁₇ Poly- and Perfluoroalkyl Substances in Water by In-Line SPE-HPLC-Orbitrap with In-Source Fragmentation Flagging, *Anal. Chem.*, 87, 4260–4268, <https://doi.org/10.1021/acs.analchem.5b00039>, 2015.
- Ma, J., Ungeheuer, F., Zheng, F., Du, W., Wang, Y., Cai, J., Zhou, Y., Yan, C., Liu, Y., Kulmala, M., Daellenbach, K. R., and Vogel, A. L.: Nontarget Screening Exhibits a Seasonal Cycle of PM_{2.5} Organic Aerosol Composition in Beijing, *Environ. Sci. Technol.*, <https://doi.org/10.1021/acs.est.1c06905>, 2022.
- 605 McDonald, B. C., de Gouw, J. A., Gilman, J. B., Jathar, S. H., Akherati, A., Cappa, C. D., Jimenez, J. L., Lee-Taylor, J., Hayes, P. L., McKeen, S. A., Cui, Y. Y., Kim, S.-W., Gentner, D. R., Isaacman-VanWertz, G., Goldstein, A. H., Harley, R. A., Frost, G. J., Roberts, J. M., Ryerson, T. B., and Trainer, M.: Volatile chemical products emerging as largest petrochemical source of urban organic emissions, *Science*, 359, 760–764, <https://doi.org/10.1126/science.aaq0524>, 2018.
- McFiggans, G., Mentel, T. F., Wildt, J., Pullinen, I., Kang, S., Kleist, E., Schmitt, S., Springer, M., Tillmann, R., Wu, C., Zhao, D., Hallquist, 610 M., Faxon, C., Le Breton, M., Hallquist, Å. M., Simpson, D., Bergström, R., Jenkin, M. E., Ehn, M., Thornton, J. A., Alfarra, M. R., Bannan, T. J., Percival, C. J., Priestley, M., Topping, D., and Kiendler-Scharr, A.: Secondary organic aerosol reduced by mixture of atmospheric vapours, *Nature*, 565, 587–593, <https://doi.org/10.1038/s41586-018-0871-y>, 2019.
- Müller, L., Reinnig, M.-C., Warnke, J., and Hoffmann, T.: Unambiguous identification of esters as oligomers in secondary organic aerosol formed from cyclohexene and cyclohexene/ α -pinene ozonolysis, *Atmos. Chem. Phys.*, 8, 1423–1433, <https://doi.org/10.5194/acp-8-1423-2008>, 2008.
- 615 Müller, L., Reinnig, M.-C., Naumann, K. H., Saathoff, H., Mentel, T. F., Donahue, N. M., and Hoffmann, T.: Formation of 3-methyl-1,2,3-butanetricarboxylic acid via gas phase oxidation of pinonic acid – a mass spectrometric study of SOA aging, *Atmos. Chem. Phys.*, 12, 1483–1496, <https://doi.org/10.5194/acp-12-1483-2012>, 2012.

- Nestorowicz, K., Jaoui, M., Rudzinski, K. J., Lewandowski, M., Kleindienst, T. E., Spólnik, G., Danikiewicz, W., and Szmigielski, R.:
620 Chemical composition of isoprene SOA under acidic and non-acidic conditions: effect of relative humidity, *Atmos. Chem. Phys.*, 18,
18 101–18 121, <https://doi.org/10.5194/acp-18-18101-2018>, 2018.
- Nozière, B., Kalberer, M., Claeys, M., Allan, J., D’Anna, B., Decesari, S., Finessi, E., Glasius, M., Grgić, I., Hamilton, J. F., Hoffmann, T.,
Iinuma, Y., Jaoui, M., Kahnt, A., Kampf, C. J., Kourtchev, I., Maenhaut, W., Marsden, N., Saarikoski, S., Schnelle-Kreis, J., Surratt, J. D.,
625 Szidat, S., Szmigielski, R., and Wisthaler, A.: The Molecular Identification of Organic Compounds in the Atmosphere: State of the Art
and Challenges, *Chem. Rev.*, 115, 3919–3983, <https://doi.org/10.1021/cr5003485>, 2015.
- Pankow, J. F. and Asher, W. E.: SIMPOL.1: a simple group contribution method for predicting vapor pressures and enthalpies of vaporization
of multifunctional organic compounds, *Atmos. Chem. Phys.*, 8, 2773–2796, <https://doi.org/10.5194/acp-8-2773-2008>, 2008.
- Peng, Z., Day, D. A., Stark, H., Li, R., Lee-Taylor, J., Palm, B. B., Brune, W. H., and Jimenez, J. L.: HO_x radical chemistry in ox-
idation flow reactors with low-pressure mercury lamps systematically examined by modeling, *Atmos. Meas. Tech.*, 8, 4863–4890,
630 <https://doi.org/10.5194/amt-8-4863-2015>, 2015.
- Peng, Z., Day, D. A., Ortega, A. M., Palm, B. B., Hu, W., Stark, H., Li, R., Tsigaridis, K., Brune, W. H., and Jimenez, J. L.: Non-OH
chemistry in oxidation flow reactors for the study of atmospheric chemistry systematically examined by modeling, *Atmos. Chem. Phys.*,
16, 4283–4305, <https://doi.org/10.5194/acp-16-4283-2016>, 2016.
- Pereira, K. L., Ward, M. W., Wilkinson, J. L., Sallach, J. B., Bryant, D. J., Dixon, W. J., Hamilton, J. F., and Lewis, A. C.: An
635 Automated Methodology for Non-targeted Compositional Analysis of Small Molecules in High Complexity Environmental Matri-
ces Using Coupled Ultra Performance Liquid Chromatography Orbitrap Mass Spectrometry, *Environ. Sci. Technol.*, 55, 7365–7375,
<https://doi.org/10.1021/acs.est.0c08208>, 2021.
- Pleil, J. D., Wallace, M. A. G., and McCord, J.: Beyond monoisotopic accurate mass spectrometry: ancillary techniques for identifying
unknown features in non-targeted discovery analysis, *J. Breath Res.*, 13, 012 001, <https://doi.org/10.1088/1752-7163/aae8c3>, 2018.
- 640 Qi, L., Vogel, A. L., Esmailirad, S., Cao, L., Zheng, J., Jaffrezo, J.-L., Fermo, P., Kasper-Giebl, A., Daellenbach, K. R., Chen,
M., Ge, X., Baltensperger, U., Prévôt, A. S. H., and Slowik, J. G.: A 1-year characterization of organic aerosol composition and
sources using an extractive electrospray ionization time-of-flight mass spectrometer (EESI-TOF), *Atmos. Chem. Phys.*, 20, 7875–7893,
<https://doi.org/10.5194/acp-20-7875-2020>, 2020.
- Riva, M., Budisulistiorini, S. H., Chen, Y., Zhang, Z., D’Ambro, E. L., Zhang, X., Gold, A., Turpin, B. J., Thornton, J. A., Canagaratna,
645 M. R., and Surratt, J. D.: Chemical Characterization of Secondary Organic Aerosol from Oxidation of Isoprene Hydroxyhydroperoxides,
Environ. Sci. Technol., 50, 9889–9899, <https://doi.org/10.1021/acs.est.6b02511>, 2016.
- Salvador, C. M. G., Tang, R., Priestley, M., Li, L., Tsiligiannis, E., Le Breton, M., Zhu, W., Zeng, L., Wang, H., Yu, Y., Hu, M., Guo, S., and
Hallquist, M.: Ambient nitro-aromatic compounds – biomass burning versus secondary formation in rural China, *Atmos. Chem. Phys.*,
21, 1389–1406, <https://doi.org/10.5194/acp-21-1389-2021>, 2021.
- 650 Schervish, M. and Donahue, N. M.: Peroxy radical chemistry and the volatility basis set, *Atmos. Chem. Phys.*, 20, 1183–1199,
<https://doi.org/10.5194/acp-20-1183-2020>, 2020.
- Schymanski, E. L., Jeon, J., Gulde, R., Fenner, K., Ruff, M., Singer, H. P., and Hollender, J.: Identifying Small Molecules via High Resolution
Mass Spectrometry: Communicating Confidence, *Environ. Sci. & Technol.*, 48, 2097–2098, <https://doi.org/10.1021/es5002105>, 2014.
- Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., Ng, N. L.,
655 Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop,

- D. R., Zaveri, R. A., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Rev. Geophys.*, 55, 509–559, <https://doi.org/https://doi.org/10.1002/2016RG000540>, 2017.
- 660 Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrakou, T., Müller, J.-F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmos. Chem. Phys.*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014.
- Surratt, J. D., Kroll, J. H., Kleindienst, T. E., Edney, E. O., Claeys, M., Sorooshian, A., Ng, N. L., Offenberg, J. H., Lewandowski, M., Jaoui, M., Flagan, R. C., and Seinfeld, J. H.: Evidence for Organosulfates in Secondary Organic Aerosol, *Environ. Sci. Technol.*, 41, 517–527, <https://doi.org/10.1021/es062081q>, 2007.
- 665 Surratt, J. D., Gómez-González, Y., Chan, A. W. H., Vermeylen, R., Shahgholi, M., Kleindienst, T. E., Edney, E. O., Offenberg, J. H., Lewandowski, M., Jaoui, M., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Organosulfate Formation in Biogenic Secondary Organic Aerosol, *J. Phys. Chem. A*, 112, 8345–8378, <https://doi.org/10.1021/jp802310p>, 2008.
- Szmigielski, R., Surratt, J. D., Gómez-González, Y., Van der Veken, P., Kourtchev, I., Vermeylen, R., Blockhuys, F., Jaoui, M., Kleindienst, T. E., Lewandowski, M., Offenberg, J. H., Edney, E. O., Seinfeld, J. H., Maenhaut, W., and Claeys, M.: 3-methyl-1,2,3-butanetricarboxylic acid: An atmospheric tracer for terpene secondary organic aerosol, *Geophys. Res. Lett.*, 34, <https://doi.org/https://doi.org/10.1029/2007GL031338>, 2007.
- 670 Thomsen, D., Elm, J., Rosati, B., Skønager, J. T., Bilde, M., and Glasius, M.: Large Discrepancy in the Formation of Secondary Organic Aerosols from Structurally Similar Monoterpenes, *ACS Earth Space Chem.*, 5, 632–644, <https://doi.org/10.1021/acsearthspacechem.0c00332>, 2021.
- Umweltbundesamt GmbH: Aktuelle Luftgütemesswerte des Bundes und der Länder, <http://luft.umweltbundesamt.at/pub/gmap/start.html>, Online; accessed on July 26, 2022, 2021.
- 675 van Eijck, A., Opatz, T., Taraborrelli, D., Sander, R., and Hoffmann, T.: New tracer compounds for secondary organic aerosol formation from β -caryophyllene oxidation, *Atmos. Environ.*, 80, 122–130, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2013.07.060>, 2013.
- Wang, Y., Ma, Y., Li, X., Kuang, B. Y., Huang, C., Tong, R., and Yu, J. Z.: Monoterpene and Sesquiterpene α -Hydroxy Organosulfates: Synthesis, MS/MS Characteristics, and Ambient Presence, *Environ. Sci. Technol.*, 53, 12 278–12 290, <https://doi.org/10.1021/acs.est.9b04703>, 680 2019.
- Wang, Y., Tong, R., and Yu, J. Z.: Chemical Synthesis of Multifunctional Air Pollutants: Terpene-Derived Nitrooxy Organosulfates, *Environ. Sci. Technol.*, 55, 8573–8582, <https://doi.org/10.1021/acs.est.1c00348>, 2021.
- Xu, L., Du, L., Tsona, N. T., and Ge, M.: Anthropogenic Effects on Biogenic Secondary Organic Aerosol Formation, *Adv. Atmos. Sci.*, 38, 1053–1084, <https://doi.org/10.1007/s00376-020-0284-3>, 2021.
- 685 Yasmeen, F., Vermeylen, R., Szmigielski, R., Iinuma, Y., Böge, O., Herrmann, H., Maenhaut, W., and Claeys, M.: Terpenylic acid and related compounds: precursors for dimers in secondary organic aerosol from the ozonolysis of α - and β -pinene, *Atmos. Chem. Phys.*, 10, 9383–9392, <https://doi.org/10.5194/acp-10-9383-2010>, 2010.
- Yasmeen, F., Szmigielski, R., Vermeylen, R., Gómez-González, Y., Surratt, J. D., Chan, A. W. H., Seinfeld, J. H., Maenhaut, W., and Claeys, M.: Mass spectrometric characterization of isomeric terpenoic acids from the oxidation of α -pinene, β -pinene, d-limonene, and Δ^3 -carene 690 in fine forest aerosol, *J. Mass Spectrom.*, 46, 425–442, <https://doi.org/https://doi.org/10.1002/jms.1911>, 2011.
- Yu, J., Cocker, D. R., Griffin, R. J., Flagan, R. C., and Seinfeld, J. H.: Gas-Phase Ozone Oxidation of Monoterpenes: Gaseous and Particulate Products, *J. Atmos. Chem.*, 34, 207–258, <https://doi.org/10.1023/A:1006254930583>, 1999.

Zhao, W., Kawamura, K., Yue, S., Wei, L., Ren, H., Yan, Y., Kang, M., Li, L., Ren, L., Lai, S., Li, J., Sun, Y., Wang, Z., and Fu, P.: Molecular distribution and compound-specific stable carbon isotopic composition of dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls in PM_{2.5} from Beijing, China, Atmos. Chem. Phys., 18, 2749–2767, <https://doi.org/10.5194/acp-18-2749-2018>, 2018.