

Response to Referee Comment #3

The referee's comments are presented in italic and our answers are written in plain text. Modifications of the manuscript, if any, are written in plain bold text.

The manuscript by O'Neill et al. present a simple linear equation between SMF and FMF which explains why SMF is generally greater than the FMF. The manuscript is thorough, clear and well written. I only have some minor/technical comments as listed below.

We thank the referee helpful feedback.

Comments:

1. line 70: "Kaku et al., 2014 (Ka)" mentioned only here and not included in the references.

We added the reference.

2. line 172 "SMF being generally $> \sim$; FMF" seems to have an issue with a symbol or character, at least on my system.

We removed " $> \sim$ ";. This was an accidental insertion.

3. line 200: Could the average be calculated from a single AOD observation or was there a requirement for a minimum number of observations?

A footnote was added to the sentence defining the "time window of ± 16 minutes":

"The number of AODs allowed to define a match in the time window could be as small as one (i.e. there was no minimum number)."

4. line 209: Why did you choose specifically these sites and not other sites with similar aerosol types (e.g. Alta Floresta)? Biomassa burning events are limited to specific seasons, so did you only consider the burning season in Mongu or did you use all the available data?

A relevant point. A sentence was added to the end of the small paragraph describing our choice of aerosol classes:

"Our choice of aerosol types was not intended to be comprehensive from the standpoint of investigating variations in say, different types of smoke aerosols or different types of dust aerosols: rather we sought to properly exercise the SMF versus FMF relationship by largely filling the admissible portion of the SMF versus FMF scattergram (see Figure 3)."

5. line 220: "M" seems to be mentioned here for the first time but it has not been defined anywhere.

"(large M)" was changed to "**(large solar airmass, M)**"

6. Figure 6: Please, clarify in the caption the meaning of the color in the points.

Somehow we failed to fix that obvious oversight. There is now a new 2nd sentence in the Figure 6 caption:

"The four colors represent the four different AERinv inflection points (r_0 values with units of μm)."

7. *Figure 9: "The orange filled large circle represents the only εf point... " I couldn't find any orange circle from the figures. Could you make it more apparent?*

The figure was incorrect : the present figure includes an **"orange-filled (red-rimmed) large circle"**.

8. *Figure A2: I would prefer if the full caption of the figure would be presented here so that the reader does not have to go to the supplement to find it.*

That was a mistake. The Figure A2 caption now refers to the caption of Figure A1 for details.

Relationship between the sub-micron fraction (SMF) and fine mode fraction (FMF) in the context of AERONET retrievals

Norman T. O'Neill¹, Keyvan Ranjbar², Liviu Ivănescu¹, Thomas F. Eck³, Jeffrey S. Reid⁴, David M. Giles³, Daniel Pérez-Ramírez⁵, Jai Prakash Chaubey⁶

- 5 ¹Centre d'applications et de recherches en télédétection (CARTEL), Université de Sherbrooke, Sherbrooke, J1K 2R1, Canada
²Flight Research Laboratory, National Research Council, Ottawa, ON, K1V 1J8, Canada
³NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA
⁴US Naval Research Laboratory, Monterey, CA 93943, USA
⁵Applied Physics Department, University of Granada, Granada 18071, Spain
10 ⁶Department of Physics and Atmospheric Sciences, Dalhousie University, Halifax, NS, B3H 4R2, Canada

Correspondence to: Norman T O'Neill (Norman.T.ONeill@usherbrooke.ca)

Abstract. The sub-micron (SM) aerosol optical depth (AOD) is an optical separation based on the fraction of particles below a specified cut off radius of the particle size distribution (PSD) at a given particle radius. It is fundamentally different from spectrally separated FM (fine mode) AOD. We present a simple (AOD-normalized) SM fraction versus FM fraction (SMF vs FMF) linear equation that explains the well-recognized empirical result of SMF generally being greater than the FMF. The AERONET inversion (AERinv) products (combined inputs of spectral AOD and sky radiance) and the Spectral Deconvolution Algorithm (SDA) products (input of AOD spectra) enable, respectively, an empirical SMF vs FMF comparison at similar (columnar) remote sensing scales across a variety of aerosol types.

SMF (AERinv derived) vs FMF (SDA derived) behavior is primarily dependent on the relative truncated portion (ε_c) of the coarse mode (CM) AOD associated with the cutoff portion of the CM PSD and, to a second order, the cutoff FM PSD and FM AOD (ε_f). The SMF vs FMF equation largely explains the SMF vs FMF behavior of the AERinv vs SDA products as a function of PSD cutoff radius (“inflection point”) across an ensemble of AERONET sites and aerosol types (urban industrial, biomass burning, dust, maritime and a mixed class of Arctic aerosols). The overarching dynamic was that the linear SMF vs FMF relation pivots clockwise about the approximate (SMF, FMF) singularity of (1, 1) in a “linearly inverse” fashion (slope and intercept of approximately $1 - \varepsilon_c$ and ε_c) with increasing cutoff radius. SMF vs FMF slopes and intercepts derived from AERinv and SDA retrievals confirmed the general domination of ε_c over ε_f in controlling that dynamic. A more general conclusion is the apparent confirmation that the optical impact of truncating modal (whole) PSD features can be detected by a SMF vs FMF analysis.

1 Introduction

30 Anderson et al, (2005) noted the “decades old observation that aerosol mass generally consists of two modes: (1) a mechanically produced coarse mode [CM] and (2) a fine mode [FM] produced by combustion and/or gas to particle

conversion”. Typical CM examples include wind-eroded desert dust and sea-salt while FM aerosols tend to be dominated by biomass-burning smoke and anthropogenic and biogenic (ABF) fine particles (the latter classification being proposed, for example, by Lynch et al., 2016). Given this speciated physical reality (and notwithstanding the known chemical and microphysical internally mixed changes that occur in aerosol properties as they are transported through the atmosphere) there is a certain level of justification in treating different species of aerosols as independent FM and CM particle size distributions (PSDs). The degree of rigour in this modal paradigm is, in fact, the subject of model analyses that compare prescribed bulk aerosol PSDs with sectional (binned) PSDs (see, for example, Mann et al., 2012 for the case of size-referenced PSDs and Kodros and Pierce, 2017 for the case of mass-referenced PSDs).

AERONET inversions (AERinv), derived from input spectral AODs and solar almucantar radiances (Dubovik and King, 2000), provide a comprehensive suite of microphysical and optical products (at ~ hourly temporal resolutions). The general tendency towards (particle-volume) PSD bimodality is notably evidenced in retrieved AERinv PSDs (see, for example, Figure 1 of Dubovik et al., 2002, Figure 11a of Eck et al., 2009 and Figure 3 of AboEl-Fetouh et al., 2020¹). A bi-modal optical representation of that PSD bimodality (without the requirement of having to specify the PSD shape of each mode) can largely determine the measurable (low spectral order) optical behavior of remote sensing data (O’Neill et al, 2001 and references cited therein). O’Neill et al. (2003) employed spectral AODs (sampling intervals ~ 3 minutes) as an input to their spectral deconvolution algorithm (SDA) to separate aerosols into (a) extensive (quantity dependent) FM and CM AOD components (τ_f and τ_c) of the total AOD (τ_a), (b) intensive (quantity independent) spectral derivatives of τ_f ($\alpha_f, \alpha'_f, \alpha''_f$, etc.) and τ_c ($\alpha_c, \alpha'_c, \alpha''_c$, etc.) as components of the total AOD spectral derivatives ($\alpha, \alpha', \alpha''$, etc.) as well as (c) semi-intensive FM and CM fractions (FMF and CMF represented respectively by $\eta = \tau_f/\tau_a$ and $1 - \eta = \tau_c/\tau_a$). If aerosols can be viewed as independent coarse and fine modal features then this purely spectral technique (~~which is generically assigned the FMF acronym~~) acts to separate those modal features in an optical sense.

The SMF (sub-micron fraction) is a microphysically determined alternative to the optically determined FMF: the division into fine and coarse components is effected by an explicit separation of the PSD at a cutoff radius (r_0) that typically ranges from ~ 0.4 to 1.1 μm across different types of aerosol volume sampling instruments^{2 3} (~~a similar radius range defines the AERinv case~~). ~~This approach, SMF separation~~ represents a moderate but significant difference relative to the optically based FMF separation. Anderson et al. (2005) describe the SMF as “...an operational definition ... to distinguish it from the theoretical concept of fine mode fraction, FMF.” Because the AERinv ~~approach algorithm~~ incorporates a cutoff radius division of the retrieved PSD into sub- and super-micron parameters (Dubovik and King, 2000) it is actually an SMF approach.

¹ The examples in these papers can include the apparent presence of more than two modes (notably what often appear to be two CM sub-modes). Such cases are discussed below.

² The expression “sub-micron” is generally “defocussed” (relative to a literally exact value of 1.0 μm) to encompass this approximate range of r_0 values.

³ It is worth noting that the in situ community almost universally refers to diameter rather than radius.

60 Comparisons with the AERONET SDA product provide a unique opportunity to empirically analyze the SMF vs FMF approaches at similar remote sensing (columnar) scales and across a global variety of aerosol types.

The literature on the direct comparison of the SMF vs FMF retrievals is sparse and oftentimes at the margins of significance. The significance problem relates to the fact that one may be forced to extract relatively subtle microphysical and optical changes in the face of SMF and FMF variations that are often limited in range (lack of extensive-parameter aerosol variation and / or types for example). The sparsity of SMF vs FMF investigations relates to limitations such as the rarity of
65 experiments designed for such a comparison or the sparsity of data sets whose modes can be readily separated (by source information and/or chemical identification for example).

In the most controlled experiments, volume-sampled SMF estimates are compared with FMF retrievals derived from applying the SDA to volume-sampled, 3-channel CRD (Cavity Ring Down) or 3-channel nephelometer “spectra” (Atkinson
70 et al., 2010, Kaku et al., 2014 (Ka) and Atkinson et al., 2018). In general we find, as expected, the SMF to be \sim FMF in Atkinson et al., 2010 and Ka: this is notably true at the near-IR (700 nm) wavelength in Ka (ACE-Asia data of their Figure 2)⁴. The cases where this is not true (the T1 “ext_sum” results of Atkinson et al., 2018) are likely attributable to the spectral sensitivity of the FMF: this is most severe in the case of just 3 channels (for example, Ka’s VOCALS 5% calibration correction of one 450 nm nephelometer channel transformed a case of SMF being substantially $<$ FMF to a corrected case of SMF \sim
75 FMF).

Lesser constrained experiments involve multi-altitude, volume-sampled SMF estimates (from nephelometer scattering coefficient & absorption coefficient devices) compared with layer and column SDA-derived FMF estimates from multi-altitude AOD and Δ AOD spectra acquired with an airborne sunphotometer (for example, Gassó & O’Neill, 2006 and Shinozuka et al., 2011; respectively represented by the acronyms GO and SHN). In these cases, the lesser degree of experimental control
80 (associated with multi-altitude flights) was somewhat offset by a more generous number of sunphotometer spectral bands (4 bands from the UV to the NIR in the GO case and 5 bands near the 5 bands employed in the AERONET SDA in the SHN case). The GO results were fairly coherent with SMF / FMF expectations (especially when the layered estimates were added to obtain columnar estimates) and quite marginal for SHN with a large spread of near-unity points about the SMF = FMF line (a situation with practically no significant SMF or FMF range and hence little relevant testing of their relationship). Analyses
85 of the FMF vs SMF relationship comparing satellite estimates of FMF⁵ with, for example, airborne estimates of SMF (Anderson et al., 2005) or coastal/island AERinv estimates of SMF (Kleidman et al., 2005) have proven difficult given the spectrally sensitive nature of the few bands employed in satellite-derived estimates of FMF and all the possible sources of

⁴ for which the fixed cutoff radius for the optically smaller particles (and thus more optically active in the sense of more strongly attenuating) results in a greater relative contribution to the FM AOD (by “optically smaller” we mean the ratio of particle size to wavelength has decreased because the wavelength has increased). The optically smaller particles are more optically active because they are ascending the right side of the anomalous diffraction peak (see, for example, O’Neill et al., 2005).

⁵ retrievals that generally employ prescribed and speciated bulk (modal) FM and CM PSDs (of constant shape and position as a function of radius) as a basis for fitting spectral AODs at a few wavelengths.

incoherencies between the satellite retrievals and the ground- or airborne estimates of SMF (differences in spatio-temporal sampling volumes for example).

90 A simple (approximate) relationship between the SMF and FMF will be presented ~~in section 2.2, equation (7a) below.~~
We seek to demonstrate that the ~~AERONET AERinv-derived value of~~ SMF and ~~the SDA-derived value of~~ FMF ~~versions~~ are
largely ~~governed~~ linked by that simple relationship and that fitting parameters extracted from their empirical comparison yields
insight into ~~the cutoff fraction of the SMF~~ their fundamental opto-physical dynamics. We argue that the similar columnar
95 scales as well as the diversity of AERONET aerosol types shared by the two retrievals facilitates the analysis of their 2nd order
intensive-parameter relationship. It is emphasized that the AERinv and SDA algorithms were developed independently and
share no explicit algorithmic links: what they do share is that the four spectral-AOD inputs to the AERinv represent 4 of the 5
spectral AODs employed as input to the SDA. What they do not share is the almucantar (angularly variable) radiance input
employed in the AERinv retrieval. It is notable that the AERONET AOD is accurately measured with an estimated uncertainty
of ~0.01 and 0.02 in the visible / NIR and the UV respectively (Eck et al., 1999) for an overhead sun (solar airmass ~~of~~ M =
100 1). This data quality enables both algorithms to, for example, retrieve notably consistent intensive parameter information such
as particle size. The consistency is also attributable to the AERinv ~~requiring a fit to~~ requirement that the retrieved AODs be
within 0.01 of the measured AOD spectra to within 0.01 AODs while the 2nd order spectral fit to the AODs employed in the
SDA has similar constraints (see, for small AODs (see example, Figure 4 of O'Neill et al., 2001).

2 Theoretical considerations


105 2.1 Size-cutoff integration versus modal integration

Figure 1 ~~illustrates the theoretical mechanical / optical framework of this paper (it is based on~~ generated from a simple
lognormal fit to a sample AERinv particle-volume PSD retrieval⁶). ~~illustrates the theoretical mechanical / optical framework~~
of this paper. That fit was applied to lend an air of empirical relevance ~~to~~ to this theoretical section: the discussion presented
here is otherwise of a general nature and is not mean-tintended to represent an algorithmic step of the AERinv retrievals (or of
110 the SDA for that matter). The FM and CM PSDs are represented by the red and blue lognormal curves respectively. The SMF-
type cutoff radius of r_0 is represented by the black dashed vertical line where the associated optical and microphysical bimodal
quantities are computed from both FM and CM PSDs to obtain sub-micron parameters to the left of r_0 and super-micron
parameters to the right of r_0 . The rest of the parameters in Figure 1 are defined immediately below as we develop the theoretical
framework.

⁶ The AERinv PSD particle-volume PSD ($dV/d \ln r$) is readily transformed to the particle-surface PSD of Figure 1 ($dS/d \ln r = 3/(4r) dV/d \ln r$). See the Figure 1 caption for details.

Letting prime variables refer to integrations which are carried out over size regimes of $(0, r_0)$ and (r_0, ∞) and unprimed variables refer to integrations carried out over entire modal features (over the entire FM or the entire CM PSDs) we can write the total AOD (τ_a) at some given reference wavelength as the sum of the fine and coarse total-modal AODs;

$$\tau_a = \tau_f + \tau_c \quad (1)$$



120 **Figure 1:** Illustrative, fine and coarse mode (red and blue) lognormal particle-surface size distributions ($dS/d \ln r$) and the Mie kernel
 (extinction efficiency Q_{ext}) for a wavelength of $0.5 \mu\text{m}$ and a refractive index of $1.44 - 0.0035i$ (by particle-surface we mean the projected
 particle surface area of πr^2). The cutoff radius⁷ (r_0) applies to a SMF type of computation. The total $dS/d \ln r$ (black curve) is (in order to
 have a level of grounding in reality) the sum of the FM and CM lognormal $dS/d \ln r$ curves that were individually manipulated to fit the
 mean of a series of AERONET and ground-based ($dV/d \ln r$) FM and CM sea-salt curves reported in Figure 8f of Reid et al. (2006). We
 125 base this illustration on $dS/d \ln r$ because it is more optically fundamental than the AERInv particle-volume PSD (the projected particle
 surface area is employed to normalize the extinction cross section to obtain the fundamental extinction efficiency). An analogous explanation
 (and a graph similar to that of Figure 2) would apply to the particle-volume PSD.

where (letting $\tau_x(r_1, r_2)$ refer to the optical integration from r_1 to r_2);

$$\tau_f = \tau_f(0, r_0) + \tau_f(r_0, \infty) \quad (2a)$$

130

$$\tau_c = \tau_c(0, r_0) + \tau_c(r_0, \infty) \quad (2b)$$

The analogous relationships in the domain of the cutoff size regimes are written;

$$\tau_a = \tau'_f + \tau'_c \quad (3)$$

$$\tau'_f = \tau_f(0, r_0) + \tau_c(0, r_0) \quad (4a)$$

$$\tau'_c = \tau_f(r_0, \infty) + \tau_c(r_0, \infty) \quad (4b)$$

⁷ "inflection point" as it is called in the AERONET documentation.

135 with an emphasis on the “conservation of τ_a ” ($\tau_f + \tau_c = \tau'_f + \tau'_{fc} = \tau_a$). The differences in the FM and CM (cutoff vs modal) integrations are respectively;

$$\begin{aligned}\tau'_f - \tau_f &= \Delta\tau_f = \tau_f(0, r_0) + \tau_c(0, r_0) - [\tau_f(0, r_0) + \tau_f(r_0, \infty)] \\ &= \tau_c(0, r_0) - \tau_f(r_0, \infty) \quad (5a)\end{aligned}$$

$$140 \quad \tau'_c - \tau_c = \Delta\tau_c = \tau_f(r_0, \infty) + \tau_c(r_0, \infty) - [\tau_c(0, r_0) + \tau_c(r_0, \infty)]$$

$$= \tau_f(r_0, \infty) - \tau_c(0, r_0) \quad (5b)$$

from which;

$$\Delta\tau_c = -\Delta\tau_f \quad (5c)$$

The last expression is nothing more than confirmation of the “conservation of τ_a ”.

2.2 SMF versus FMF

145 Using the definitions and relationships given above we can now define FMF and SMF respectively as;

$$\eta = \frac{\tau_f}{\tau_a} \quad \text{and} \quad \eta' = \frac{\tau'_f}{\tau_a} \quad (6)$$

Given $\tau_f + \tau_c = \tau'_f + \tau'_c$ we divide by τ_a to obtain;

$$\eta + \frac{\tau_c}{\tau_a} = \eta' + \frac{\tau'_c}{\tau_a}$$

and, with a little manipulation;

$$150 \quad \eta' = (1 - \varepsilon_c - \varepsilon_f)\eta + \varepsilon_c \quad (7a)$$

where ε_f and ε_c represent the pure truncation errors of the FM and CM PSDs (the 2nd terms of equations 5(a) and 5(b) normalized by τ_a);

$$\varepsilon_f = \frac{\tau_f(r_0, \infty)}{\tau_f} \quad \text{and} \quad \varepsilon_c = \frac{\tau_c(0, r_0)}{\tau_c} \quad (7b)$$

155 These two quantities represent intensive parameter (largely quantity independent) attributes⁸. Typically however, $\varepsilon_c \gg \varepsilon_f$ (the major part of the FM PSD is well displaced to the left of r_0 as per the illustration of Figure 1). This has much more of a cutoff impact on the blue-~~coloured~~ CM PSD of Figure 1 than on the red-~~coloured~~ FM PSD: the solid blue cutoff portion (ΔS_c) is a significant portion of the CM particle-surface density (S_c) and $\tau_c(0, r_0)$ is a significant optical depth portion of τ_c (i.e. both $\Delta S_c/S_c$ and ε_c are typically significant) while the analogous FM fractions above r_0 are relatively insignificant. This affirmation, which we will empirically demonstrate in the multi-station analysis below, is, in part, related to the fact that the FM PSD is \sim half the width of the CM PSD (Dubovik et al., 2002: in their paper “width” specifically refers to the σ value of fitted lognormal distributions).

⁸ by “largely quantity independent” we mean, from an empirical standpoint, that $\tau_f(r_0, \infty)$ and $\tau_c(0, r_0)$ are typically well correlated with τ_f and τ_c respectively.

Figure 2 is a plot of ε_c vs $\Delta S_c/S_c$ for a variety of retrievals from the sites listed in Table 1. All these cases involved, as per the Figure 1 illustration, the fitting of FM and CM lognormal curves to AERInv particle-volume PSDs (and a subsequent transformation to particle-surface parameters). These lognormal fits permit the explicit (Mie-based) calculations of the equation (7b) fractions. They represent an (AERONET-grounded) theoretical illustration of an expected strong correlation between cutoff optics withand cutoff mechanics (the correlation is less than monotonic because of variations in refractive index and the width of the lognormal curves for the different cases chosen in this illustration). We emphasize that automated lognormal fits to AERInv PSDs are not part of the empirical ~~analysis-process~~analyses presented in this paper: (nor do they have any role in the purely spectral SDA retrieval); rather the purpose of Figure 2 is to confirm the expected strong correlation between optical and microphysical CM cutoff fractions and thereby facilitate an understanding of ε_c 's role in the dynamics of equation (7a).

Table 1: Site coordinates, predominant aerosol type, sampling periods and retrieval numbers for the AERONET inversions and ~~the~~ SDA retrievals employed in this study

Sites	Latitude	Longitude	<u>Elevation</u> <u>lev.</u> (m)	Region	Aerosol class / type	Sampling period	<u>Number of retrievals</u> ^{aa} <u>N</u> ^a	<u>Sample citations</u>
GSFC	38° 59' 33" N	76° 50' 23" W	87	Maryland, USA	Urban-industrial	1994-2020	13070	<u>Holben et al (2001)</u>
Mongu	15° 15' 13" S	23° 9' 3" E	1047	Zambia	Biomass Burning	1997-2009	4751	<u>Eck et al (2001)</u>
Hamim	22° 58' 0" N	54° 18' 0" E	105	United Arab Emirates	Dust	2004-2007	2295	<u>Eck et al (2008)</u>
Solar Village	24° 54' 25" N	46° 23' 50" E	764	Saudi Arabia	Dust	1999-2012	14171	<u>Hamill et al (2016)</u>
Forth Crete	35° 19' 58" N	25° 16' 57" E	20	Greece	Maritime and Dust	2003-2017	4039	<u>Hamill et al (2016)</u>
Midway Island	28° 12' 35" N	177° 22' 42" W	20	Midway Island	Maritime	2001-2014	644	<u>Smirnov et al (2003)</u>
Lanai	20° 44' 6" N	156° 55' 18" W	20	Hawaii	Maritime	1997-2004	1175	<u>Reid et al (2006)</u>
PEARL ^b	80° 3' 13" N	86° 25' 1" W	615	Nunavut, Canada	Arctic	2007-2019	270	<u>Ranjbar et al (2019)</u>
Barrow	71° 18' 44" N	156° 39' 54" W	8	Alaska, USA	Arctic	1997-2020	351	<u>Stone et al (2014)</u>
Thule	76° 30' 58" N	68° 46' 8" W	225	Northern Greenland	Arctic	2007-2019	500	<u>AboEl-Fetouh et al (2020)</u>

^{aa} N = the number of retrievals. See the methodology section of the text for details on how the AERONET inversion and SDA retrievals were matched.

^b Polar Environment Arctic Research Lab. The acronym represents the total atmospheric research infrastructure at Eureka, Nunavut. The AERONET/AEROCAN site is more accurately referred to as the PEARL Ridge Lab.

The linear form of equation (7a) with its sub-unity slope of $(1 - \varepsilon_c - \varepsilon_f)(1 - \varepsilon_c - \varepsilon_f)$ and positive ε_c intercept is coherent with empirical results of η' being generally $> \eta$ (the well recognized empirical result, supported by the results presented below, of SMF being generally $> \sim$ FMF). Because ε_c generally dominates ε_f the linear η' vs η relation pivots clockwise about the point $(\eta, \eta') = (1, 1)$ in a “linearly inverse” fashion (slope and intercept of approximately $1 - \varepsilon_c$ and ε_c). Figure 3 shows a set of equation (7a) straight lines for representative ranges of different ε_c and ε_f values that were inspired by the ranges encountered in the empirical results that follow.

~~Cases of large c_f are relatively rare but would occur during intense FM events characterized by large amplitude FM PSDs that are shifted towards larger radii: illustrations include strong (large FM AOD) smoke events (see, for example, the sub-Arctic (Alaskan) smoke cases of AOD(440 nm) > 1 in Figure 9a of Eck et al., 2009 and strong FM pollution events enhanced by high relative humidity (see, for example, the PSDs corresponding to AOD(440 nm) > 2 in Figure 4 of Eck et al., 2020). However an increasing c_f presupposes that r_0 is fixed (as for volumetric surface sampling devices): the AERinv technique of setting the r_0 value to the minimum of the PSD (see below) results in a minimization of c_f .~~




Figure 2: Plot of ε_c vs $\Delta S_c / S_c$ for a variety of simulated optical depth retrievals. The lognormal FM and CM PSDs employed in these simulations were obtained by fitting their sum to retrieved AERinv PSDs over a variety of sites whose predominant aerosol type was urban, dust or marine (and employing the refractive indices provided by the retrieval product). The AERinv PSD retrievals were Version 2 for which r_0 was fixed at $0.6 \mu\text{m}$ and, accordingly, for which the ε_c vs $\Delta S_c / S_c$ regression would be only dependent on ε_c and 2nd order parameters such as refractive index (i.e. the plot is independent of r_0).




Figure 3: η' (SMF) vs η (FMF) lines of equation (7a) for three ε_c values and three ε_f values (0, 0.4, 0.8 and -0.1, 0 and 0.1 respectively). The $\varepsilon_f \varepsilon_c$ values were inspired by the range of empirical values seen in Figure 6 (and, in the case of ε_f by the range of values, whether real or artefactual, seen in Figure 7).

195 Cases of large ε_f are relatively rare but would occur during intense FM events characterized by large-amplitude FM PSDs that are shifted towards larger radii: illustrations include strong (large FM AOD) smoke events (see, for example, the sub-Arctic (Alaskan) smoke cases of AOD(440 nm) $> \sim 1$ in Figure 9a of Eck et al., 2009 and strong FM pollution events enhanced by high relative humidity (see, for example, the PSDs corresponding to AOD(440 nm) $> \sim 2$ in Figure 4 of Eck et al., 2020). However an increasing ε_f presupposes that r_0 is fixed (as for volumetric surface sampling devices): the AERInv
 200 technique of setting the r_0 value to the minimum of the PSD (see below) results in a minimization of ε_f .

3 Methodology

Hesaraki et al. (2017) give an overview of the AERInv and SDA with an emphasis on the fact that the former is a significantly more comprehensive algorithm whose low frequency sampling rate is appropriate for detailed climatological scale analyses (see, for example, Dubovik et al., 2002 and AboEl-Fetouh et al., 2020) while the high frequency sampling rate
 205 of the SDA is better suited to the detailed analysis of diurnal events (see, for example, Figure S5 of Saha et al., 2010). The SDA is readily applied to AOD spectra generated by starphotometers and moonphotometers (see, for example, Baibakov et al., 2015) while analogous AERInv products (requiring an almucantar scan) do not exist for nighttime conditions.


In this investigation we “matched” SDA to AERInv retrievals by employing averages of Version 3 Level 2.0 AODs (Giles et al., 2019) as inputs to the SDA if those AODs were within a time window of ± 16 minutes about the nominal AERInv

210 times⁹. Version 3 AERinv products (Sinyuk et al., 2020) were employed to derive estimates of SMF, τ_f' and τ_c' . The cutoff
radius for those products is actually defined as the minimum of the AERinv output particle-volume PSD ($dV/d\ln r$) with a
restriction that the radius bin centers of that minimum must be one of four AERinv bins (r_0 , which is referred to as an
“inflection point” in AERONET documentation is allocated a [AERinv](#) bin-center value of 0.439, 0.576, 0.756 or 0.992 μm).
The AERinv retrievals of τ_f' and τ_c' are interpolated to the SDA reference wavelength of 500 nm using a 2nd order (log-log)
215 space) spectral polynomial regressed to the τ_f' and τ_c' values at the four AERinv wavelengths of 440, 675, 870 and 1020 nm
(the same technique employed in the SDA).

A variety of AERONET sites (Figure 4), representing different types (classes) of aerosols were chosen to investigate
the SMF versus FMF relationship. These aerosol classes (Table 1) included sites known for FM urban-industrial aerosols
(GSFC in Greenbelt MD), FM biomass burning (Mongu, Zambia), CM dust (Crete, Hamim and Solar Village), CM maritime
220 aerosols (Midway Island and Lanai), a mixture of dust and marine aerosols (Forth Crete) and a mixture of high-Arctic
aerosols (PEARL and Thule) as well as low-Arctic aerosols (Barrow). Our choice of aerosol types was not intended to be
comprehensive from the standpoint of investigating variations in say, different types of smoke aerosols or different types of
dust aerosols: rather we sought to properly exercise the SMF versus FMF relationship by largely filling the admissible portion
of the SMF versus FMF scattergram (see Figure 3).

225 The Arctic category is a mixed class of aerosols: its inclusion in our table of aerosol types is more in terms of it
representing an important comparative test (relative to the applicability of southern latitude findings) in a region where the
aerosol signal is notably weak. Arctic illustrations of FMF vs SMF principles feature more in the analysis presented below
precisely because the signal is weak: the observation of a specific an independently-verified FMF vs SMF trend or characteristic
in the Arctic is an indicator of the robustness of that observation. This weak-signal robustness is something we often see for
230 Arctic retrievals (see, for example, the FM and CM results of AboEl-Fetouh et al., 2020): it may well be (at least in part)
attributable to the large solar zenith angles (large solar airmass, M) and attendant $1/M$ decrease in optical depth errors (a typical
finding over 20 years of AERONET Mauna Loa calibrations according to co-author Eck: see also Figure 2 of Karanikolas et
al., 2022 for an empirical validation of this $1/M$ error dependency).

⁹ The number of AODs allowed to define a match in the time window could be as small as one (i.e. there was no minimum number).



235 **Figure 4:** AERONET stations employed for generating the statistics employed in this paper. These stations were chosen to represent a regional variety of aerosols (pollution, biomass burning, dust mixtures, sea-salt mixtures and Arctic aerosols). See Table 1 for details.

4 Results

4.1 Frequency of Occurrence of inflection (r_0) points

Figure 5 shows the relative frequency of occurrence (FO) of the four different AERinv inflection (r_0) points for the different Table 1 classes. The relative importance of CM PSDs vs FM PSDs for the dust class sites of Solar Village and Hamim and the marine sites of Midway Island and Lanai push the AERinv minimum to smaller r_0 values (resulting in an FO dominance at the $0.439 \mu\text{m}$ inflection point). The marine sites are moderately less asymmetric (less pushed towards $0.439 \mu\text{m}$) than the dust sites because the CM PSDs for the dust sites tend to be of larger amplitude. There are, however, other factors at play that could have some impact on the FO distributions: for example, large FM sulfatic particles from Kiluea eruptions that might push back on the CM PSD dominance of marine particles at Lanai and/or non sphericity effects of dust that, when corrected (Dubovik et al., 2006) would produce a significantly larger FM PSD amplitude (a complication that does not impact the predominantly spherical marine particles).

240
245





Figure 5: Relative frequency occurrence (FO) distributions for the four different AERONET inflection points (τ_0 values with units of μm) for all Table 1 AERONET sites.

The two sites that are heavily influenced by strong FM PSDs (the Mongu biomass burning site and the GSFC urban-industrial site) tend (as suggested above in an ε_f context) to “push” the PSD minimum towards larger τ_0 values relative to the dust and marine sites (resulting in a more balanced FO curve in Figure 5). Eck et al. (2001) attributed this FM particle-size increase to coagulative effects for Mongu (see their Figure 6) while Eck et al. (2012) attributed the increase to the effects of hygroscopically induced FM particle growth at the GSFC site (see their Figure 17). Other smoke impacted and urban industrial sites show similar coagulative particle growth effects (see Figure 10b of Eck et al., 2019 for specific cases recorded in the southeast Asian tropical forest) and hygroscopic particle growth impacts (see, for example Figure 13 of Eck et al., 2005 for Beijing). We argue below that these large amplitude increases in FM particle size are, given the AERinv technique of variable inflection points, relatively minor in terms of producing significant ε_f values.

The Arctic sites show a FM PSD domination ~~that~~ which produces an FO distribution that is not unlike the Mongu distribution ~~for~~ in the singular case of Barrow while being much more skewed towards large τ_0 for PEARL and Thule. We





illustrate below (as part of a discussion on the variation of ε_c) that a systematic (seasonal), spring to summer inflection point increase can be attributed to the Arctic sites.



265 **Figure 5: Relative frequency occurrence (FO) distributions for the four different AERONET inflection points (r_g values) for all Table 1 AERONET sites.**

4.2 η' vs η (SMF vs FMF) scattergrams

270 Figure 6 shows η' vs η (SMF vs FMF) scattergrams representing four key aerosol types of Table 1 (scattergrams for the rest of the aerosol types and sites can be seen in Figure S1 of the supplementary material). The theoretical solid black lines of Figure 6 represent various values of ε_c in equation (7a): we chose to set ε_f to zero in tracing those lines because its value is, as indicated above, generally small (and to not obscure the graphs with 2nd order detail). We note (as per the [previous](#) discussion of the r_g -FO [curves/distributions](#)) that as the AERinv PSD minimum [radius \(inflection point\)](#) increases, the cutoff portion of the CM PSD increases (resulting in the transition from red to blue curves). The slopes tend to decrease (swing clockwise) with an attendant increase in the intercepts.






Figure 6: η' vs η (AERinv-derived SMF vs SDA-derived FMF) scattergrams for the sites of GSFC, Mongu, Lanai, and Hamim (respectively aerosol types of urban-industrial, biomass burning, marine and dust as per Table 1). The four colors represent the four different AERinv inflection points (r_0 values with units of μm). The remaining scattergrams for the other sites are shown in Figure S1. The solid black lines are those of equation (7a) with ϵ_f set to 1 (for the sake of the simplicity of the presentation coupled with the fact that ϵ_f has plays a relatively minor role), ϵ_f was set to 0.

The FO dominance of the 0.439 μm r_0 bin is most evident for the CM dominated dust class (Hamim) with, practically speaking, a single (small ϵ_c) red-coloured colored grouping of points being observable while the more balanced distributions of FO curves for GSFC and Mongu, provide four clear point groupings of points. In the extreme of CM cutoff, virtually all of the optically significant contributions of the CM PSD are out-off-eliminated and an asymptote of $\eta' \rightarrow 1$ with $\epsilon_c \rightarrow 1$ for all values of η is approached (the slope of equation (7a) approaches zero). That extreme CM cutoff condition is most evident for the blue, large- r_0 points of the GSFC scattergram) where a regression slope could be almost parallel to the η' axis. We did indeed find $r_0 = 0.992$ cases associated with near unity η' values for which most of the optically significant portion of the CM PSD was cut off while the FM and CM PSDs were not inordinately unbalanced in terms of amplitude (η was some sub-unity value of significance for which there were no apparent problems associated with the AOD spectra).

The small to large r_0 (red to blue) transformation translates, for example, into a classical seasonal pattern for the Arctic scattergrams of Figure S1: the springtime amplitude of the small-sized CM PSD (that AboEl-Fetouh et al., 2020, associated

with Asian dust) decreases progressively from spring to summer. This weakening of CM influence induces a spring to summer increase in the value of r_0 and, by extension, ε_c (see the seasonal r_0 histograms and derived table for Barrow in Figure S2).
295 This is an effect that is more noticeable in the Level 1.5 AERinv products of ibid than the Level 2.0 products of this paper¹⁰.

Within the theoretical context of equation (7a), the scattergrams of Figure 6 show (predominantly for the red-
~~coloured~~colored 0.439 μm inflection value of r_0), a minority of unphysical points below the $\eta' = \eta$ line (points for
which $\eta' < \eta$). An investigation into the most extreme cases of this inequality indicated that the wayward points were more
inclined to be associated with abnormally large values of η (rather than abnormally small values of η') and small AODs (<
300 0.05). Figure 10 of O'Neill et al. (2003) shows the noise sensitivity of η values to such small AODs if one assumes an RMS
AOD error of 0.01 in all bands (0.01 being <~ errors of AERONET field instruments). In general, this type of small-AOD
sensitivity was reported empirically by, for example, Eck et al. (1999) and O'Neill et al. (2000). The fundamental dynamic is
that a band-to-band AOD discontinuity <~ 0.01 can generate important perturbations in small-AOD curvature spectra and
produce significant outliers in the spectrally sensitive η values.

305 4.3 Analysis of the slopes and intercepts (derivation of ε_c and ε_f)

The η' vs η scattergrams support the hypothesis that there is a physical / optical interpretation that can be given to the
slope and intercept of equation (7a) (that equations (7a) and (7b) are theoretically relevant approximations). Figure 7a and 7b
show the regression slopes vs the intercepts derived for, respectively, all sites and all r_0 values of the Figure 6 and
supplementary material scattergrams ~~(with a coherent color scheme between the scattergrams and Figure 7a)~~¹¹ while Table 2
310 lists the associated regression statistics for the 0.439 μm inflection point. The large range of the intercept ~~variation~~ seen in
Figure 7 confirms that the cutoff portion of the CM PSD (ε_c of equation (7b)) is much more determinant in affecting the
interplay of η' vs η . The position of the colored circles relative to the dashed or dotted black lines (derived from equations
(7a) and ~~7(b)~~(7b)) visually support the Table 2 results of small ε_f (ε_f values being <~ 0.07 according to their positions between
the $\varepsilon_f = 0$ and $\varepsilon_f = 0.1$ grid lines ~~of Figure 7~~).

¹⁰ the Level 2.0 processing tends to eliminate springtime retrievals completely: the majority of eliminations are due to excessive residuals in the retrieved vs measured sky radiances that are, in turn, largely incited by the strong reflectance uncertainty of springtime snow (and its attendant impact on computed sky radiance) as well as the AERinv protocol of eliminating Level 2 retrievals if any snow is detected (by MODIS) within a 5 km radius of the site.

¹¹ with a coherent color scheme between the scattergrams and Figure 7b




Figure 7: Intercept vs slope plots for the η' vs η scattergrams as a function of (a)–(b) all the sites of Table 1 and (b) the cutoff r_0 value (AERinv inflection point radius) and (b) all the sites of Table 1. The black dashed or pointed lines represent the family of straight lines generated by equation (7a): Slope = $(1 - \epsilon_c - \epsilon_f)(1 - \epsilon_c - \epsilon_f)$ and Intercept = ϵ_c for values of $\epsilon_c = 0$ to 1 and three values of ϵ_f (0.0, 0.1 and 0.2).

The values corresponding to the five red-colored (0.439 μm inflection point) circles for the desert and marine sites (bottom right hand corner of Figure 7b) are however outside the physically coherent range of equation (7). The negative values of ϵ_f (see Table 2) can be inferred from the $\epsilon_f = 0, 0.1, 0.2$ grid of Figure 7b or, more visually, from the unphysical (superunity) slopes that arise from a regression through the red points of the desert and marine scattergrams of Figure 6. This incoherency is very likely the result of those regression lines representing a non-unique ensemble of ϵ_c and ϵ_f values. From an opto-physical standpoint, they vary as a function of the diversity of dust and marine CM PSDs that are predominantly

associated with the 0.439 μm inflection point (they roughly lie between the $\varepsilon_c = 0$ and $\varepsilon_c = 0.4$ boundaries of Figure 3: the $\eta = 1$ intercept ~~of the expression $(1 - \varepsilon_f)^{12}$~~ will (in consequence of a super-unity slope) yield negative values above the $\eta' = \eta$ intercept).

330 Non-systematic noise will also impact the derived ε_c and ε_f values. Scattergrams of τ'_f vs τ_f for the two sites that are largely dominated by strong FM variations (GSFC and Mongu) and which showed the most balanced (Figure 5) FO distributions are shown in Figure A1. One can observe, in the first instance, the point-dispersion reduction and the convergence towards the $\tau'_f = \tau_f$ line as both increase (as τ_a increases). The extensive (quantity-dependent) nature of those scattergrams complements the analogous $\eta' vs \eta$ semi-intensive scattergrams of Figure 6 by explicitly displaying the noise-like influences
335 of the amount of aerosol (τ'_f or τ_f) as well as r_0 . The $\tau'_f vs \tau_f$ dispersion of Figure A1 is nonetheless generally small: this underscores a hypothesis that the $\eta' vs \eta$ results presented above are robust 2nd order findings. The FO distributions of $\eta' vs \eta$ in Figure A2 effectively eliminate the extensive variations of Figure A1 with an attendant enhancement of ~~the~~ those 2nd order influences.

Table 2: η' vs η regression stats for $r_0 = 0.439 \mu\text{m}$ ^a

	m	b	R ²	$\sigma(\eta'_{res})$	$\sigma(m)$	$\sigma(b)$	N	ε_c	ε_f	$\sigma(\varepsilon_c)$	$\sigma(\varepsilon_f)$
GSFC	0.695	0.304	0.749	0.043	0.409	0.288	3924	0.304	9E-04	0.288	0.29
Mongu	0.42	0.541	0.641	0.038	0.314	0.256	2280	0.541	0.039	0.256	0.182
Hamim	1.015	0.076	0.969	0.025	0.172	0.069	2234	0.076	-0.091	0.069	0.158
Solar Village	0.99	0.065	0.912	0.043	0.314	0.12	13790	0.065	-0.055	0.12	0.291
Forth Crete	0.922	0.116	0.941	0.044	0.226	0.139	3125	0.116	-0.038	0.139	0.178
Midway	0.996	0.097	0.832	0.059	0.39	0.15	615	0.097	-0.093	0.15	0.36
Lanai	0.873	0.152	0.674	0.072	0.427	0.154	1032	0.152	-0.025	0.154	0.398
PEARL	N/A (N = 1)										
Barrow	0.34	0.604	0.447	0.053	0.462	0.341	44	0.604	0.057	0.341	0.311
Thule	0.259	0.66	0.196	0.038	0.482	0.405	8	0.66	0.081	0.405	0.262

^a See for example, Taylor (1997) for typical regression relationships. The ε_c and ε_f values as well as their standard deviation ($\sigma(\varepsilon_c)$ and $\sigma(\varepsilon_f)$) were derived from the regressed slope and intercept (“m” and “b”) using the slope and intercept expressions of equation (7a). The values of ($\sigma(\varepsilon_c)$ and $\sigma(\varepsilon_f)$) are ~~effective standard deviations~~ computed from effective standard deviations of m and b . By this we mean that the standard error (from sources such as Taylor, 1997) is multiplied by \sqrt{N} to yield $\sigma(m)$ and $\sigma(b)$: this transforms the unrealistically small m and b uncertainties into values that are more representative of the variation seen in the scattergrams of Figure 6. This change is coherent with the notion that those variations are more likely due to ε_c and ε_f being characterized by a systematic range of values for any given inflection point (~~r_0~~) value. Note that $\sigma(\eta'_{res})$ (η'_{res}) represents the standard deviation of the regression residuals (~~res~~).


340 The Figure 8 FO distributions of $\eta' - \eta$ vs τ_a more readily show the decreasing dispersion and the convergence towards the $\eta' = \eta$ ($\tau'_f = \tau_f$) line with increasing τ_a (increasing τ_f values for GSFC and Mongu). The broad left to right

¹² for which $\eta' = 1 - \varepsilon_f$

movement of the (red) FO peak values with increasing r_0 (notably in the case of GSFC) also clarifies an aspect that is not easily discernable in the highly correlated scattergrams of Figure A1: that an increase in r_0 is coarsely associated with η' values that approach η or hence, τ_f' values that approach τ_f (a trend that effectively drives the $\eta' = \eta$ ($\tau_f' = \tau_f$) convergence ~~incited~~induced by increasing τ_a).


345 Figure 9 shows the regression-derived ε_c and ε_f variation as a function of an artificial minimum ($\tau_{a,min}$) in the lower bound of the τ_a regression range (see the Figure 9 caption for more details). The result shows no strong ε_c dependency on $\tau_{a,min}$ and more variable but consistently small amplitudes of ε_f ¹³. Qualitatively, ~~this observation is~~these observations are not unexpected given the fairly persistent slope of the high FO (red-colored) ellipses of Figure A2).

350 The choice of a variable r_0 in the AERInv retrievals clearly means that the affirmation of increasing ε_f in the presence of strong FM events (i.e., what would be measured by a traditional fixed- r_0 device) cannot readily be observed with the AERInv retrievals.



~~Figure 8: $\eta' - \eta$ vs τ_a FO distributions for GSFC and Mongu.~~

¹³ except in the case of the (red) 0.439 μm r_0 (yellow-filled) value where the ε_f variations were more substantial. At the same time, the number of regressions points (N) rapidly decreases with increasing $\tau_{a,min}$ so that large ε_f (and ε_c) deviations are less statistically significant.



355 **Figure 9:** GSFC and Mongu variations of regression derived values of ϵ_c and ϵ_f as a function of a minimum value of total AOD at 500 nm ($\tau_{a,min}$) employed in the η' vs η regressions¹⁴ (to be clear, the regressions of Figure 6 and the regression derived ϵ_c and ϵ_f values of Figure 7 were obtained with no restrictions on the matched AERInv and SDA retrievals). Same colour scheme as Figure 6 and 7. The number of matched retrievals (N) are also shown (right hand axis). Result are not shown in cases where $N < 20$. **The orange filled large circle represents the only ϵ_f point of the $\tau_0 = 0.439 \mu\text{m}$ regressions that survived the $N < 20$ criterion and that did not yield a negative ϵ_f value.**

360

retrievals. Strong FM events basically push the FM PSD to larger radii (while the CM PSD remains relatively inactive¹⁵). The AERInv approach of selecting r_0 as the minimum of the PSD basically neuters the cutoff of significant optical portions of the FM PSD. The general tendency for r_0 to increase with increasing τ_f in Figure 8 is consistent with the concept of the FM PSD being pushed to the right as τ_f' and τ_f increase.

365

¹⁴ The decreasing point dispersion (the approach of τ_f' to τ_f) with increasing τ_f indicates that the regression derived precision of the ϵ_c and ϵ_f values should, for that reason alone, generally increase with larger values of τ_f . However, offsetting this type of influence on an increase in precision is the attendant decrease in τ_f , the decrease in the number of regression points and the squeeze in τ_f and τ_f' variability as the $\eta' = \eta$ singularity is approached.

¹⁵ and suffers from a per-particle extinction kernel (the green-colored Q_{ext} factor of Figure 1) that is weaker in CM radius range.




Figure 8: $\eta' - \eta$ vs τ_a FO distributions for GSF and Mongu.

AERinv PSDs can differ from the simple bi-modal paradigm incorporated in equations (1) through (7). The bi-modality of the Arctic CM PSDs of AboEl-Fetouh et al. (2020) were ascribed to a small-sized CM feature in the 1.3 μm AERinv bin (a systematic feature that they attributed to Asian dust) and a larger sized mode that might have been linked to local dust and/or sea-salt. Eck et al. (2012) reported a bi-modal FM PSD (their Figure 3) that they attributed to cloud processing of FM pollution (haze) aerosols. In such cases one can appeal to the optical equivalency of, for example, a bi-modal CM PSD to a single CM PSD whose curvature parameters become ~~curvature parameter~~ averages of those of the two CM components (see Appendix A for the two specific cases of a bimodal CM PSD with a unimodal FM PSD as well as a bimodal FM PSD with a unimodal CM PSD). This means the bi-modal expressions of O'Neill et al. (2001) still apply for the AERONET SDA product and thus that the FMF to SMF expressions (equations (1) to (7)) can still be used.




Figure 9: GSFC and Mongu variations of regression-derived values of ε_c and ε_f as a function of a minimum value of total AOD at 500 nm ($\tau_{a,min}$) employed in the η' vs η regressions¹⁶ (to be clear, Figure 9 represents an exercise in testing the sensitivity to $\tau_{a,min}$: the regressions of Figure 6 and the regression-derived ε_c and ε_f values of Figure 7 were obtained with no restrictions on the matched AERinv and SDA retrievals). Same colour scheme as Figure 6 and 7. The legend of the Mongu plot is the same as the GSFC legend. The number of matched retrievals (N) are also shown (right-hand axis). Result are not shown in cases where $N < 20$. The large, orange-filled (re-rimmed) circle in the GSFC case represents the only ε_f point of the $r_0 = 0.439 \mu\text{m}$ regressions that both survived the $N < 20$ rejection criterion and did not yield a negative ε_f value.

The FMF (SDA) approach is arguably the more fundamental approach for separating FM and CM optical contributions because it is intrinsically related to the modal nature of different types of aerosols (at least to the first order: modality becomes obscured, for example, with internal mixing of different aerosol types). The SMF (AERinv) approach is the more pragmatic approach as it is commonly and readily applied to microphysical surface and airborne measurements. It is however, as we have seen in results like those of Figure 6, very dependent on the selected cutoff radius.

390 5 Conclusions

We presented a simple SMF vs FMF (η' vs η) equation that enabled an understanding of the well recognized empirical result of SMF being larger than the FMF. This result has been reported for in-situ, satellite and ground-based remote sensing techniques: our focus was on an SMF vs FMF interpretation in the form of AERinv SMF vs SDA FMF retrievals. We pointed

¹⁶ The decreasing point dispersion (the approach of τ_f' to τ_f) with increasing τ_f indicates that the regression-derived precision of the ε_c and ε_f values should, for that reason alone, generally increase with larger values of τ_f . However, offsetting this type of influence on an increase in precision is the attendant decrease in τ_f , the decrease in the number of regression points and the squeeze in τ_f and τ_f' variability as the $\eta' = \eta$ singularity is approached.

out that these two AERONET products provide a unique opportunity to empirically compare the SMF and FMF approaches at
395 similar (columnar) remote sensing scales and across a shared global variety of aerosol types.

The SMF vs FMF equation largely captured the SMF vs FMF behavior of the AERinv vs SDA products as a function
of inflection point (τ_0) across an ensemble of AERONET sites and aerosol types (urban industrial, biomass burning, dust,
marine, maritime and Arctic). The SMF vs FMF behavior was primarily dependent on the intensive parameter of relative cutoff
portion of the CM PSD (ε_c) and, to a second order the relative cutoff portion of the FM PSD (ε_f). The overarching dynamic
400 was that the linear *SMF* vs *FMF* relation pivots clockwise about the point $(\eta, \eta') = (1, 1)$ in a “linearly inverse” fashion
(slope and intercept of approximately $1 - \varepsilon_c$ and ε_c) with increasing τ_0 . Derived SMF vs FMF slopes and intercepts confirmed
the general domination of ε_c over ε_f in controlling the “linear inverse” dynamic. The process of deriving and analyzing ε_c and
 ε_f values demonstrated an expected domination of FM optical depths for the urban pollution and biomass burning sites of
GSFC and Mongu and thus a convergence towards the $\tau'_f = \tau_f$ ($\eta' = \eta$) line as τ_a increased (the convergence of SDA FM
405 AODs towards AERinv FM AODs).

The more general conclusion resulting from this analysis is the apparent empirical confirmation that the influence of
PSD modal features can be detected by an indirect comparative analysis. While one would like to believe that this is true in
general, a more comprehensive event-level closure experiment employing, for example, multi-altitude microphysical and
optical measurements over a representative suite of AERONET instruments would do much to increase the level of confidence
410 in such a conclusion.

Appendix A

A.1 Frequency of occurrence analyses

The ~~retrievals~~retrieval results in this subsection are restricted to the two sites that are strongly impacted by historically large
variations in FM particles (GSFC and Mongu). These sites (details in Table 1) can experience τ_f values greater than unity.




Figure A1: τ_f' vs τ_f frequency of occurrence (FO) distributions for GSFC and Mongu. The FO colour scale is tied to variations on a logarithmic scale (with an attendant tendency to enhance the contributions of large FO values). $N(r_0)$ is the total FO at a given r_0 value. The decreasing dispersion cloud with increasing τ_f indicates (because of the τ_f' and τ_f log scales) a decreasing relative relative dispersion.




Figure A2: η' vs η FO distributions for GSFC and Mongu. See the caption of Figure S2A1 for FO and $N(r_0)$ details.

A.2 Dual-modal optical equivalency of a tri-modal size distribution

425 A.2.1 Fine mode and two coarse modes

One often sees a bi-modal coarse mode PSD and thus a tri-modal PSD. Treating the bi-modal CM PSD as an optically equivalent single mode CM PSD can be formalized as below (optical equivalent of equation (1a) of O'Neill et al., 2003):

$$\tau_a = \tau_f + \tau_c = \tau_f + \tau_{c1} + \tau_{c2} = \tau_f + \eta_{c1}\tau_c + \eta_{c2}\tau_c \quad (A1)$$

where $\eta_{c1} = \frac{\tau_{c1}}{\tau_c}$, and $\eta_{c2} = \frac{\tau_{c2}}{\tau_c}$. Accordingly;

$$430 \quad \tau_a = \tau_f + \eta_{c1}\tau_c + (1 - \eta_{c1})\tau_c \quad (A2)$$

Equations (3) of O'Neill et al. (2003) can be written as;

$$\begin{aligned} \alpha\tau_a &= \alpha_f\tau_f + \alpha_{c1}\tau_{c1} + \alpha_{c2}\tau_{c2} \\ \alpha\tau_a &= \alpha_f\tau_f + \tau_c(\alpha_{c1}\eta_{c1} + (1 - \eta_{c1})\alpha_{c2}) \\ &= \alpha_f\tau_f + \tau_c\langle\alpha_c\rangle \quad (A3) \end{aligned}$$

435 where $\langle\alpha_c\rangle = \alpha_{c1}\eta_{c1} + (1 - \eta_{c1})\alpha_{c2}$ represents the optical average of α_{c1} and α_{c2} . Equation (A3) is exactly the analogue of O'Neill et al., 2003). Differentiating and recalling that $d\tau/d\ln\lambda = -\alpha\tau$, one finds;

$$\begin{aligned} \alpha'\tau_a &= \alpha'_f\tau_f + \alpha'_{c1}\tau_{c1} + \alpha'_{c2}\tau_{c2} + \alpha^2\tau_a - \alpha_f^2\tau_f - \alpha_{c1}^2\tau_{c1} - \alpha_{c2}^2\tau_{c2} \\ &= \alpha'_f\tau_f + (\alpha'_{c1}\tau_{c1} + \alpha'_{c2}\tau_{c2}) + \{\alpha(\alpha\tau_a)\} - \alpha_f^2\tau_f - (\alpha_{c1}^2\frac{\tau_{c1}}{\tau_c} - \alpha_{c2}^2\frac{\tau_{c2}}{\tau_c})\tau_c \\ &= \alpha'_f\tau_f + \langle\alpha'_c\rangle\tau_c + \{\alpha\alpha_f\tau_f + \alpha\langle\alpha_c\rangle\tau_c\} - \alpha_f^2\tau_f - \langle\alpha_c^2\rangle\tau_c \\ 440 \quad &= \alpha'_f\tau_f + \langle\alpha'_c\rangle\tau_c + \alpha_f\tau_f(\alpha - \alpha_f) + \langle\alpha_c\rangle\tau_c(\alpha - \langle\alpha_c\rangle) + \langle\alpha_c\rangle^2\tau_c - \langle\alpha_c^2\rangle\tau_c \end{aligned}$$

but since $\alpha = \alpha_f\eta + (1 - \eta)\langle\alpha_c\rangle$ we can replace $(\alpha - \alpha_f)$ and $(\alpha - \langle\alpha_c\rangle)$ to obtain;

$$\begin{aligned} &= \alpha'_f\tau_f + \langle\alpha'_c\rangle\tau_c - \alpha_f\tau_f(1 - \eta)(\alpha_f - \langle\alpha_c\rangle) + \langle\alpha_c\rangle\tau_c\eta(\alpha_f - \langle\alpha_c\rangle) + \tau_c(\langle\alpha_c\rangle^2 - \langle\alpha_c^2\rangle) \\ &= \alpha'_f\tau_f + \langle\alpha'_c\rangle\tau_c - (1 - \eta)\tau_f(\alpha_f - \langle\alpha_c\rangle)^2 + \tau_c(\langle\alpha_c\rangle^2 - \langle\alpha_c^2\rangle), \text{ so that;} \\ &\alpha' = \eta\alpha'_f + (1 - \eta)\{\langle\alpha'_c\rangle + \langle\alpha_c\rangle^2 - \langle\alpha_c^2\rangle\} - (1 - \eta)\eta(\alpha_f - \langle\alpha_c\rangle)^2 \end{aligned}$$

445 These expressions are optically equivalent to equation (5) of O'Neill et al. (2003) with their equation (5) values being transformed according to;

$$\begin{aligned} \alpha_c &\leftrightarrow \langle\alpha_c\rangle \\ \alpha'_c &\leftrightarrow \langle\alpha'_c\rangle + \langle\alpha_c\rangle^2 - \langle\alpha_c^2\rangle \end{aligned}$$

where the average value of any parameter x_c is always;

$$450 \quad \langle x_c \rangle = x_{c1}\eta_{c1} + x_{c2}(1 - \eta_{c1})$$

A.2.2 Coarse mode and two fine modes

In this instance one can imagine two fine mode components: for example (i) large-sized FM smoke and (ii) smaller sized FM urban-industrial pollution. The algebra above for the single FM PSD and two CM PSDs can be employed here: everything is perfectly reversible (interchange index c with index f) if η is viewed as the ratio of coarse to total AOD. Then arriving at the final equations for α and α' one can return to the usual definition of η to obtain the same algebraic theorem. The classic equations for α and α' remain unchanged providing one employs the substitutions below;

$$\alpha_f \leftrightarrow \langle \alpha_f \rangle$$

$$\alpha'_f \leftrightarrow \langle \alpha'_f \rangle + \langle \alpha_f \rangle^2 - \langle \alpha_f^2 \rangle$$

where $\langle x \rangle = \eta_f x + (1 - \eta_f)x$ and $\eta_f = \tau_{f,1}/\tau_f$. Explicitly;

$$\alpha = \alpha_c(1 - \eta) + \eta\langle \alpha_f \rangle$$

$$\alpha' = (1 - \eta)\alpha'_c + \eta\{\langle \alpha'_f \rangle + \langle \alpha_f \rangle^2 - \langle \alpha_f^2 \rangle\} - (1 - \eta)\eta(\langle \alpha_f \rangle - \alpha_c)^2$$

Symbol and acronym glossary

AEROCAN	Federated Canadian subnetwork of AERONET run by Environment and Climate Change Canada (ECCC)
AERONET	Aerosol Robotic Network: World-wide NASA network of combined sunphotometer / sky-scanning radiometers manufactured by CIMEL Électronique. See http://aeronet.gsfc.nasa.gov/ for documentation and data downloads
AOD	Aerosol optical depth: The community uses "AOD" to represent anything from nominal aerosol optical depth which hasn't been cloud-screened to the conceptual (theoretical) interpretation of aerosol optical depth. In this paper we use it in the latter sense and apply adjectives as required.
AERinv	AERONET Inversion
SDA	Spectral Deconvolution Algorithm
CM	Coarse mode
ε_c	<u>Relative truncation error of the CM PSD (equation (7b))</u>
ε_f	<u>Relative truncation error of the FM PSD (equation (7b))</u>
η	<u>FMF (product of the SDA)</u>
η'	<u>SMF (product of the AERinv)</u>
FM	Fine mode
FO	Frequency of Occurrence
PSD	particle size distribution (precisely, notably the volume <u>AERinv particle-volume size distribution</u>), <u>$dV/d \ln r$ in the context of this paper</u>
x	<u>$x = a, f, \text{ or } c$ (total, fine mode or coarse mode)</u>
FMF, η	Fine Mode Fraction (an output parameter computed from SDA products)
SMF, η'	Sub-Micron Fraction (an output parameter computed from AERinv products)
r_0	Cut off radius (<u>the AERinv "inflection point"</u>) <u>separating a PSD into FM and CM PSDs. Units of μm</u>
$\tau_x \tau_a$	<u>Aerosol or FM or CM AOD</u> <u>Total aerosol optical</u> at 500 nm (SDA)
τ_c	<u>Coarse mode aerosol optical at 500 nm (SDA)</u>

τ_f	<u>Fine mode aerosol optical at 500 nm (SDA)</u>
τ'_a	<u>Total aerosol optical depth at 500 nm</u>
$\tau'_{\cancel{x}}\tau'_c$	Aerosol or FM or CM AOD <u>Coarse mode aerosol optical at 500 nm (using a cutoff radius of r_0 (derived AERinv product))</u>
$\epsilon_x\tau'_f$	Cut-off portion of the fine or coarse <u>Fine mode aerosol optical depth (e.f. equation 7) at 500 nm using a cutoff radius of r_0 (derived AERinv product)</u>

Author contributions

NTO led the conceptualization, methodology and formal analysis. All co-authors contributed to the revisions of the manuscript.
 465 TFE, JSR and DMG provided fundamental comments that resulted in substantial changes to the manuscript. KR, LI, DPR and JPC were instrumental in the strategical planning and operational processing of the data and the definition of the figures.

Data availability

All AERinv and SDA retrieval data are available from the [AERONET website](#).

Competing interests

470 The authors declare that they have no conflict of interest.

Acknowledgements

We acknowledge the generous permission of the AERONET PIs¹⁷ for the use of their AERinv and SDA retrieval data.

Financial support

This work was supported by CANDAC (the Canadian Network for the Detection of Atmospheric Change) through PAHA
 475 (Probing the Atmosphere of the High Arctic) funding (RGPCC-433842-2012), the NSERC Discovery Grant program (O'Neill, RGPIN-2017-05531), GSFC / NASA, notably GESTAR2 funding for O'Neill (SURA-GSTR-4100-NASA) and CSA's ESS-DA program (16SUASACIA) and CSA's FAST program (CASSAVA – PEARL project led by K. Strong of CANDAC).

References

AboEl-Fetouh, Y., N. T. O'Neill, K. Ranjbar, S. Hesaraki, I. Abboud, V. Fioletov, P. S. Sobolewski, Climatological-
 480 scale analysis of intensive and semi-intensive aerosol parameters derived from AERONET Arctic retrievals, Journal of Geophysical Research, 125(10), 2019JD031569, <https://doi.org/10.1029/2019JD031569>, 2020.

¹⁷ Brent Holben of AERONET/NASA (Goddard Space Flight Center) at the Hamim, GSFC, Lanai, Midway Island, Solar Village, Thule and Mongu sites, Ihab Abboud and Vitali Fioletov of AEROCAN/Environment and Climate Change Canada (ECCC) at the PEARL site, Richard Wagener of Brookhaven National Laboratory (BNL) at the Barrow site and Andrew Clive Banks of Hellenic Centre for Marine Research (HCMR) for the Forth Crete site.

Anderson, T.L., Wu, Y., Chu, D.A., Schmid, B., Redemann, J. and Dubovik, O., Testing the MODIS satellite retrieval of aerosol fine-mode fraction, *Journal of Geophysical Research: Atmospheres*, 110(D18), <https://doi.org/10.1029/2005JD005978>, 2005.

485 Atkinson, D.-B., P. Massoli, N.-T. O'Neill, P.-K. Quinn, D.-S. Covert, S. Brooks, and B. Lefer, Comparison of in situ and columnar spectral measurements during TexAQS-GoMACCS 2006: Testing parameterizations for estimating aerosol fine mode optical properties, *Atmos. Chem. Phys.*, 9, 17465-17494, <https://doi.org/10.5194/acp-10-51-2010>, 2010.

Atkinson, D.-B., M. Pekour, D. Chand, J.-G. Radney, K.-R. Kolesar, Q. Zhang, A. Setyan, N.-T. O'Neill, and C.-D. Cappa, Using spectral methods to obtain particle size information from optical data: applications to measurements from 490 CARES 2010, *Atmos. Chem. Phys.*, 18(8), 5499-5514, <https://doi.org/10.5194/acp-18-5499-2018>, 2018.

Baibakov, K., O'Neill, N.-T., Ivanescu, L., Duck, T.-J., Perro, C., Herber, A., Schulz, K.-H., and Schrems, O., Synchronous polar winter starphotometry and lidar measurements at a High Arctic station, *Atmos. Meas. Tech.*, 8, 3789-3809, <https://doi.org/10.5194/amt-8-3789-2015>, 2015.

Dubovik, O., and M.-D. King, A flexible inversion algorithm for retrieval of aerosol optical properties from Sun and 495 sky radiance measurements, *J. Geophys. Res.*, 105, 20,673– 20,696, <https://doi.org/10.1029/2000JD900282>, 2000.

Dubovik O., Holben B., Eck T.F., Smirnov A., Kaufman Y.J., King M.D., Tanré D., Slutsker I., Variability of absorption and optical properties of key aerosol types observed in worldwide locations. *Journal of the atmospheric sciences*, 59(3):590-608, [https://doi.org/10.1175/1520-0469\(2002\)059%3C0590:VOAAOP%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059%3C0590:VOAAOP%3E2.0.CO;2), 2002.

Dubovik O., Sinyuk A., Lapyonok T., Holben B.N., Mishchenko M., Yang P., Eck T.F., Volten H., Muñoz 500 O., Veihelmann B., Van der Zande W.J., Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of desert dust. *Journal of Geophysical Research: Atmospheres*, 111(D11208), <https://doi.org/10.1029/2005JD006619>, 2006.

~~Eck T.F., Holben B.N., Ward D.E., Dubovik O., Reid J.S., Smirnov A., Mukelabai M.M., Hsu N.C., O'Neill N.T., Slutsker I. Characterization of the optical properties of biomass burning aerosols in Zambia during the 1997 ZIBBEE field campaign. *Journal of Geophysical Research: Atmospheres*, 106(D4):3425-48, <https://doi.org/10.1029/2000JD900555>, 2001.~~

Eck, T.-F., Holben, B.-N., Reid, J.-S., Dubovik, O., Smirnov, A., O'Neill, N.-T., Slutsker, I., and Kinne, S.: Wavelength dependence of the optical depth of biomass burning, urban, and desert dust aerosols, *J. Geophys. Res.*, 104, 31333–31349, <https://doi.org/10.1029/1999JD900923>, 1999.

Eck T.F., Holben B.N., Ward D.E., Dubovik O., Reid J.S., Smirnov A., Mukelabai M.M., Hsu N.C., 510 O'Neill, N.T., Slutsker, I., Characterization of the Optical Properties of Biomass Burning Aerosols in Zambia during the 1997 ZIBBEE experiment, *J. Geophys. Res.*, Vol. 106, No. D4, 3425-3448, <https://doi.org/10.1029/2000JD900555>, 2001.

Eck T.F., Holben B.N., Dubovik O., Smirnov A., Goloub P., Chen H.B., Chatenet B., Gomes L., Zhang X.Y., Tsay S.C., Ji Q., Columnar aerosol optical properties at AERONET sites in central Eastern Asia and aerosol transport to the tropical Mid-Pacific, *J Geophys Res.*, 110, D06202, <https://doi.org/10.1029/2004JD005274>, 2005.

515 [Eck T.F., Holben B.N., Reid J.S., Sinyuk A., Dubovik O., Smirnov A., Giles D., O'Neill N.T., Tsay S.C., Ji Q., Al Mandoos A., Spatial and temporal variability of column-integrated aerosol optical properties in the southern Arabian Gulf and United Arab Emirates in summer, Journal of Geophysical Research: Atmospheres, 113\(D01204\), <https://doi.org/10.1029/2007JD008944>, 2008.](#)

Eck, T. F., B. N. Holben, E. J. Hyer, J. S. Reid, G.E. Shaw, A. Sinyuk, J. R. Vande Castle, F.S. Chapin, N.T. O'Neill, O. Dubovik, A. Smirnov, E. Vermote, J.S. Schafer, D. Giles, I. Slutsker, M. Sorokine, W. Newcomb, Optical Properties of Boreal Region Biomass Burning Aerosols In Alaska and Transport of Smoke to Arctic Regions, J. Geophys. Res., 114, D11201, <https://doi.org/10.1029/2008JD010870>, 2009.

525 Eck [TF,T.F.](#), Holben [BN,B.N.](#), Reid [JS,J.S.](#), Giles [DM,D.M.](#), Rivas [MA,M.A.](#), Singh [RP,R.P.](#), Tripathi [SN,S.N.](#), Bruegge [CJ,C.J.](#), Platnick [S,S.](#), Arnold [GT,G.T.](#), Krotkov [NA,N.A.](#), Fog-and cloud-induced aerosol modification observed by the Aerosol Robotic Network (AERONET), Journal of Geophysical Research: Atmospheres, 117(D07206), <https://doi.org/10.1029/2011JD016839>, 2012.

Eck [TF,T.F.](#), Holben [BN,B.N.](#), Giles [DM,D.M.](#), Slutsker [I,I.](#), Sinyuk [A,A.](#), Schafer [JS,J.S.](#), Smirnov [A,A.](#), Sorokin [M,M.](#), Reid [JS,J.S.](#), Sayer [AM,A.M.](#), Hsu [NC,N.C.](#), AERONET remotely sensed measurements and retrievals of biomass burning aerosol optical properties during the 2015 Indonesian burning season. Journal of Geophysical Research: Atmospheres, 530 124(8):4722-40, <https://doi.org/10.1029/2018JD030182>, 2019.

Eck [TF,T.F.](#), Holben [BN,B.N.](#), Kim [J,J.](#), Beyersdorf [AJ,A.J.](#), Choi [M,M.](#), Lee [S,S.](#), Koo [JH,J.H.](#), Giles [DM,D.M.](#), Schafer [JS,J.S.](#), Sinyuk [A,A.](#), Peterson [DA,D.A.](#), Influence of cloud, fog, and high relative humidity during pollution transport events in South Korea: Aerosol properties and PM2.5 variability. Atmospheric Environment, 232:117530, <https://doi.org/10.1016/j.atmosenv.2020.117530>, 2020.

535 Gassó, S. and O'Neill, N., Comparisons of remote sensing retrievals and in situ measurements of aerosol fine mode fraction during ACE-Asia, Geophysical research letters, 33(5), <https://doi.org/10.1029/2005GL024926>, 2006.

Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korokin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 database – automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements, Atmos. Meas. Tech., 12, 169–209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.

[Hamill, Patrick, Marco Giordano, Carlyne Ward, David Giles, Brent Holben, An AERONET-based aerosol classification using the Mahalanobis distance, Atmospheric Environment, Vol. 140, 213-233, ISSN 1352-2310, <https://doi.org/10.1016/j.atmosenv.2016.06.002>, 2016.](#)

545 Hesaraki, S., N. T. O'Neill, Lesins, G., Saha, A., R. V. Martin, V. E. Fioletov, K. Baibakov, I. Abboud, Polar summer comparisons of a chemical transport model with a 4-year analysis of fine and coarse mode aerosol optical depth retrievals over the Canadian Arctic, Atmosphere-Ocean, <https://doi.org/10.1080/07055900.2017.1356263>, 2017.

- 550 [Holben B.N., Tanre D., Smirnov A., Eck T.F., Slutsker I., Abuhassan N., Newcomb W.W., Schafer J.S., Chatenet B., Lavenu F.J., Kaufman Y.J., An emerging ground-based aerosol climatology: Aerosol optical depth from AERONET, *Journal of Geophysical Research: Atmospheres*, 106\(D11\):12067-97, <https://doi.org/10.1029/2001JD900014>, 2001.](#)
- [Kaku, K.C., J.S. Reid, N.T. O'Neill, P.K. Quinn, D.J. Coffman, and T.F. Eck, Verification and application of the extended spectral deconvolution algorithm \(SDA+\) methodology to estimate aerosol fine and coarse mode extinction coefficients in the marine boundary layer, *Atmos. Meas. Tech.*, 7, 3399-3412, <https://10.5194/amt-7-3399-2014>, 2014.](#)
- 555 Karanikolas A., Kouremeti N., Gröbner J., Egli L., Kazadzis S., Sensitivity of aerosol optical depth trends using long term measurements of different sun-photometers, *Atmos. Meas. Tech.*, 15, 5667–5680, <https://doi.org/10.5194/amt-15-5667-2022>, 2022.
- Kleidman, R.G., O'Neill, N.T., Remer, L.A., Kaufman, Y.J., Eck, T.F., Tanré, D., Dubovik, O. and Holben, B.N., Comparison of Moderate Resolution Imaging Spectroradiometer (MODIS) and Aerosol Robotic Network (AERONET) remote-sensing retrievals of aerosol fine mode fraction over ocean, *J. Geophys. Res. Atmos.*, 110, D22205, 560 <https://doi.org/10.1029/2005JD005760>, 2005.
- Kodros [JK,J.K.](#), Pierce [JR,J.R.](#), Important global and regional differences in aerosol cloud-albedo effect estimates between simulations with and without prognostic aerosol microphysics, *J. Geophys. Res. Atmos.*, 122, 4003– 4018, <https://doi.org/10.1002/2016JD025886>, 2017.
- 565 Lynch P., Reid [JS,J.S.](#), Westphal [DL,D.L.](#), Zhang J., Hogan [TF,T.F.](#), Hyer [EJ,E.J.](#), Curtis [CA,C.A.](#), Hegg [DA,D.A.](#), Shi [Y.](#), Campbell [JR,J.R.](#), Rubin [H,J.I.](#), An 11-year global gridded aerosol optical thickness reanalysis (v1.0) for atmospheric and climate sciences. *Geoscientific Model, Development*. 9(4):1489-522, <https://doi.org/10.5194/gmd-9-1489-2016>, 2016.
- 570 Mann [GW,G.W.](#), Carslaw [KS,K.S.](#), Ridley [DA,D.A.](#), Spracklen [DV,D.V.](#), Pringle [KJ,K.J.](#), Merikanto J., Korhonen H., Schwarz [JP,J.P.](#), Lee [LA,L.A.](#), Manktelow [PT,P.T.](#), Woodhouse [MT,M.T.](#), Intercomparison of modal and sectional aerosol microphysics representations within the same 3-D global chemical transport model. *Atmos. Chem. Phys., ACP*, 12, 4449-76, <https://doi.org/10.5194/acp-12-4449-2012>, 2012.
- O'Neill, N.-T. , Eck, T.-F., Holben, B.-N., Smirnov, A., Dubovik, O., A. Royer, Bi-modal size distribution influences on the variation of Angstrom derivatives in spectral and optical depth space, *J. Geophys. Res.*, 106(-D9), 9787–9806, <https://doi.org/10.1029/2000JD900245>, 2001.
- 575 O'Neill, N. T., T.-F., Eck, A. Smirnov, B.-N., Holben, S. Thulesiraman, Spectral discrimination of coarse and fine mode optical depth, *J. Geophys. Res.*, 108, 4559, D17, <https://doi.org/10.1029/2002JD002975>, 2003.
- [Ranjbar K., O'Neill N.T., Lutsch E., McCullough E.M., AboEl-Fetouh Y., Xian P., Strong K., Fioletov V.E., Lesins G., Abboud I., Extreme smoke event over the high Arctic, *Atmospheric Environment*, 218, 117002, <https://doi.org/10.1016/j.atmosenv.2019.117002>, 2019.](#)
- 580 Reid, J.-S., B. Brooks, K.-K. Crahan, D.-A. Hegg, T.-F. Eck, N.-T. O'Neill, G. de Leeuw, E.-A. Reid and K.-A. Anderson, Reconciliation of coarse mode sea-salt aerosol particle size measurements and parameterizations at a subtropical ocean receptor site, *J. Geophys. Res.*, 111, D02202, <https://doi.org/10.5194/acp-12-4449-2012>, 2006.

Saha, A., N.-T. O'Neill, E. Eloranta, R. S. Stone, T.-F. Eck, S. Zidane, D. Daou, A. Lupu, G. Lesins, M. Shiobara, L.-J. B. McArthur, Pan-Arctic sunphotometry during the ARCTAS-A campaign of April 2008, *Geophys. Res. Lett.*, 37, L05803, <https://doi.org/10.1029/2009GL041375>, 2010.

585 Shinozuka, Y., Redemann, J., Livingston, J.M., Russell, P.B., Clarke, A.D., Howell, S.G., Freitag, S., O'Neill, N.T., Reid, E.A., Johnson, R. and Ramachandran, S., ~~2011~~. Airborne observation of aerosol optical depth during ARCTAS: vertical profiles, inter-comparison and fine-mode fraction, *Atmos. Chem. Phys.*, 11, 3673–3688, <https://doi.org/10.5194/acp-11-3673-2011>, 2011.

Sinyuk A., Holben ~~BN~~, B.N., Eck ~~TF~~, T.F., Giles ~~DM~~, D.M., Slutsker I., Korkin S., Schafer ~~JS~~, J.S., Smirnov A., Sorokin M., Lyapustin A., The AERONET Version 3 aerosol retrieval algorithm, associated uncertainties and comparisons to Version 2, *Atmos. Meas. Tech.*, 13, 3375–3411, <https://doi.org/10.5194/amt-13-3375-2020>, 2020.

Smirnov A., Holben B.N., Eck T.F., Dubovik O., Slutsker I., Effect of wind speed on columnar aerosol optical properties at Midway Island. *Journal of Geophysical Research: Atmospheres*, 108(D24), 4802, <https://doi.org/10.1029/2003JD003879>, 2003.

595 Stone R.S., Sharma S., Herber A., Eleftheriadis K., Nelson D.W., A characterization of Arctic aerosols on the basis of aerosol optical depth and black carbon measurements, *Elementa: Science of the Anthropocene*. <https://doi.org/10.12952/journal.elementa.000027>, 2014.

Taylor, John R., *An Introduction to Error Analysis: The Study of Uncertainties in Physical Measurements*, 2nd edition, University Science Books, <http://www.worldcat.org/title/301634764>, 1997.