

The CALIPSO version 4.5 stratospheric aerosol subtyping algorithm

Jason L. Tackett¹, Jayanta Kar², Mark A. Vaughan¹, Brian J. Getzewich¹, Man-Hae Kim³, Jean-Paul Vernier⁴, Ali H. Omar¹, Brian E. Magill², Michael C. Pitts¹, David M. Winker¹

¹NASA Langley Research Center, Hampton, VA, USA

5 ²Science Systems and Applications, Inc., Hampton, VA, USA

³Seoul National University, Seoul, South Korea

⁴National Institute of Aerospace Associates, Hampton, VA, USA

Correspondence to: Jason L. Tackett (jason.l.tackett@nasa.gov)

Abstract.

10 The accurate classification of aerosol types injected into the stratosphere is important to properly characterize their chemical and radiative impacts within the Earth climate system. The updated stratospheric aerosol subtyping algorithm used in the Version 4.5 (V4.5) release of the Cloud Aerosol Lidar with Orthogonal Polarization (CALIOP) level 2 data products now delivers more comprehensive and accurate classifications than its predecessor. The original algorithm identified four aerosol subtypes for layers detected above the tropopause: volcanic ash, smoke, sulfate/other, and polar stratospheric aerosol (PSA).

15 In the revised algorithm, sulfates are separately identified as a distinct, homogeneous subtype and the diffuse, weakly scattering layers previously assigned to the sulfate/other class are recategorized as a fifth “unclassified” subtype. By making two structural changes to the algorithm and revising two thresholds, the V4.5 algorithm improves the ability to discriminate between volcanic ash and smoke from pyrocumulonimbus injections, improves the fidelity of the sulfate subtype, and more accurately reflects the uncertainties inherent in the classification process. The 532 nm lidar ratio for volcanic ash was also

20 revised to a value more consistent with the current state of knowledge. This paper briefly reviews the previous version of the algorithm (V4.1/V4.2), then fully details the rationale and impact of the V4.5 changes on subtype classification frequency for specific events where the dominant aerosol type is known based on literature. Classification accuracy is best for volcanic ash due to its characteristically high depolarization ratio. Smoke layers in the stratosphere are also classified with reasonable accuracy, though during the daytime a substantial fraction are misclassified as ash. It is also possible for mixtures of ash and

25 sulfate to be misclassified as smoke. The V4.5 sulfate subtype accuracy is less than that for ash or smoke, with sulfates being misclassified as smoke about one-third of the time. However, because exceptionally tenuous layers are now assigned to the unclassified subtype and the revised algorithm levies more stringent criteria for identifying an aerosol as sulfate, it is more likely that layers labeled as this subtype are in fact sulfate compared to those given the sulfate/other classification in the previous data release.

30 1 Introduction

Injections of aerosol into the stratosphere have important impacts on the chemistry in the upper atmosphere and affect the Earth's radiative energy balance (Kremser et al., 2016). Explosive volcanic eruptions can inject large amounts of ash and sulfur dioxide (SO₂) gas into the lower and middle stratosphere. Ash, which is most often composed of silicates, can remain in the atmosphere for weeks or, in extreme cases months, as in the case of the 2014 Kelud eruption (Vernier et al., 2016).
35 SO₂ reacts with the hydroxyl radical OH in the stratosphere through photochemistry, forming sulfuric acid (H₂SO₄) nuclei which grow by condensation and coagulation into larger sulfate aerosols (Kremser et al., 2016). The radiative and chemical impacts of sulfate in the stratosphere can be significant (e.g., Stone et al., 2017). A third aerosol type in the upper troposphere and lower stratosphere (UTLS) is becoming widely recognized as important: smoke from pyrocumulonimbus (pyroCb) events within intense wildfires (Fromm et al., 2010). PyroCb events inject smoke to altitudes of 20 km or higher
40 due to the buoyancy from the intense heat of the fire and meteorological conditions that favor the development of deep convection, specifically moisture at mid-levels which accelerates the upward motion (Peterson et al. 2017; Fromm et al., 2019). Evidence exists that absorption of solar radiation can cause smoke to self-loft even higher (de Laat et al., 2012; Khaykin et al., 2020; Yu et al., 2019). The amount of smoke injected by pyroCb activity into the UTLS is comparable to volcanic levels (Peterson et al., 2018). Recent evidence also suggests that self-lofting caused smoke from Siberian wildfires
45 to enter the UTLS in the summer/autumn of 2019 without the need for pyroconvection (Ohneiser et al., 2021), though this possibility is still under investigation among the community (e.g., Boone et al., 2022). Finally, ammonium nitrate particles can reach the UTLS within the Asian Tropopause Aerosol Layer by way of convection in the Asian Monsoon region (Vernier et al., 2018; Höpfner et al. 2019).

The Cloud Aerosol Lidar with Orthogonal Polarization (CALIOP) onboard the Cloud-Aerosol Lidar and Infrared
50 Pathfinder Satellite Observation (CALIPSO) platform was launched in April 2006 and is well suited to observe aerosol in the lower stratosphere (Winker et al., 2009; Vernier et al., 2009). Being a vertical profiling lidar operating at two wavelengths (532 and 1064 nm), CALIOP can measure plume altitudes with high precision; data are reported at vertical resolutions of 60–300 m for stratospheric altitudes (Hunt et al., 2009). Depolarization ratio measurements at 532 nm also provide critical information on particle shape, of which the CALIOP cloud-aerosol discrimination and aerosol subtyping algorithms take full
55 advantage (Kim et al., 2018; Liu et al., 2019). There are two important reasons for CALIOP level 2 algorithms to accurately discriminate between stratospheric aerosol types. First, identifying different aerosol types allows researchers to characterize the abundance of these types and to quantify their disparate radiative, chemical, and dynamical impacts – the scientific motivation. Second, the CALIOP retrieval scheme relies on its aerosol subtyping algorithm to select an appropriate lidar ratio, which is required to accurately retrieve extinction and to correct for overlying attenuation when retrieving optical
60 properties of underlying layers – the algorithmic motivation.

Given these compelling motivations, the CALIPSO project included new stratospheric aerosol subtypes for aerosol layers detected above the tropopause in the 2016 release of the version 4.1 level 2 data products (V4.1). These subtypes

included volcanic ash, sulfate/other, smoke, and polar stratospheric aerosol (PSA). Details of the initial stratospheric aerosol subtyping algorithm are given in Kim et al. (2018). In short, the algorithm discriminated between volcanic ash, smoke, and “sulfate/other” based on empirically derived thresholds of estimated particulate depolarization ratio at 532 nm and total attenuated backscatter color ratio (the ratio of attenuated backscatters at 1064 nm and 532 nm). Volcanic ash particles, aspherical in nature, exhibit a higher depolarization signature than sulfate aerosols which are spherical (Pueschel 1996), yielding low depolarization ratios (Vernier et al., 2013). Smoke observed in the stratosphere from pyroCb events can also be depolarizing (Haarig et al., 2018; Ohneiser et al., 2020), but to a lesser degree than ash. Layers with low 532 nm integrated attenuated backscatter (i.e., less than 0.001 sr^{-1}) were classified as sulfate/other. These tenuous layers represent the “other” fraction of the combined class. They were combined with sulfates under the assumption that the long residence time of sulfate aerosol would eventually yield low attenuated backscatter returns. Meanwhile, layers classified as aerosol at exceptionally low temperatures over the polar regions during polar stratospheric cloud (PSC) season were classified as PSA.

Since the initial release of the V4.1 level 2 data products, it became clear that some refinements were necessary. One significant example is the poor accuracy of discriminating between volcanic ash and smoke. During the massive Pacific Northwest (PNW) event in August 2017 (Peterson et al., 2018), smoke injected into the lower stratosphere was highly depolarizing (Haarig et al., 2018), exceeding the depolarization threshold used by the V4.1 algorithm to separate smoke and ash. As a consequence, over 58 % of the aerosol layers detected from this event were misclassified as volcanic ash. An example CALIOP observation of one of the earliest smoke plumes detected from the event (Torres et al., 2020) is shown in Fig. 1, illustrating the dominance of ash misclassification.




Figure 1. PyroCb smoke plume from the PNW event on 18 August 2017 at ~7:45 UTC over Quebec Province, Canada: (a) 532 nm total attenuated backscatter, (b) 532 nm volume depolarization, and (c) V4.1 aerosol subtype classification from the level 2 vertical feature mask. Inset map shows the CALIOP ground track.

Another area needing refinement was the sulfate/other class. This subtype is ambiguous for practical applications because it is shared both by layers that could legitimately be sulfate and by weakly scattering layers that could be any aerosol type. In addition, the 532 nm integrated attenuated backscatter (γ'_{532}) threshold used to identify these weakly scattering layers was too high for the features commonly detected in the stratosphere by CALIOP. Some 75 % of all stratospheric aerosol layers detected in V4.1 were classified as sulfate/other due to this threshold. For example, Fig. 2 shows several plumes from the June 2011 Puyehue-Cordón Caulle eruption where the layer depolarization ratios are elevated, indicating the presence of ash. However, some portions are classified as sulfate/other because their integrated attenuated backscatter falls below the threshold. A more reasonable classification for these layers would be ash.

Finally, studies have shown that the most common lidar ratio for volcanic ash may be larger than the default lidar ratio used for the ash subtype in V4.1 (e.g., Prata et al., 2017). Based on these observations, the stratospheric aerosol subtyping algorithm and lidar ratio assignments have been updated for the V4.5 level 2 data release.




Figure 2. Volcanic ash plume from the June 2011 Puyehue-Cordón Caulle eruption observed on 16 June 2011 at ~5:45 UTC, southwest of Argentina: (a) 532 nm total attenuated backscatter, (b) 532 nm volume depolarization ratio, and (c) V4.1 aerosol subtype classification from the level 2 vertical feature mask. Inset map shows the CALIOP ground track.

The organization of this paper is as follows. First, we describe the datasets used for our analysis in Sect. 2. Next, we review the V4.1/V4.2 stratospheric aerosol subtyping algorithm in Sect. 3 and then in Sect. 4 we describe the changes implemented to the algorithm for V4.5. We provide a statistical summary of layer classifications between the versions in Sect. 4.6 to demonstrate the improvements of the revised algorithm. Section 5 is a performance assessment for select events dominated by volcanic ash, sulfate, and smoke to explore the classification fidelity. Concluding remarks are given in Sect. 6.

2 Data used

110 The data used for our analysis is from CALIOP onboard CALIPSO which has been operating since June 2006. CALIOP is an elastic backscatter lidar measuring vertical profiles of attenuated backscatter at 532 nm and 1064 nm, with depolarization capability at 532 nm (Hunt et al., 2009; Winker et al., 2009). Level 2 retrievals detect features using a threshold-based algorithm after averaging the lidar backscatter signal to multiple horizontal resolutions (Vaughan et al., 2009). The horizontal averaging resolutions for layer detection are 1/3 km, 1 km, 5 km, 20 km, and 80 km. Following detection, the
115 cloud-aerosol discrimination (CAD) algorithm classifies each layer as either cloud or aerosol based on the layer geolocation and measured optical properties (Liu et al., 2019). Layers classified as aerosol are further classified as either tropospheric aerosol or stratospheric aerosol, depending on their altitudes with respect to the tropopause, which is obtained from the Modern-Era Retrospective analysis for Research Applications, Version 2 (MERRA-2) meteorological reanalysis product (Gelaro et al., 2017). Specifically, aerosol layers having a 532 nm attenuated backscatter centroid below the tropopause are
120 identified as one of seven different tropospheric aerosol subtypes: clean marine, dust, polluted continental/smoke, clean continental, polluted dust, elevated smoke, or dusty marine (Kim et al., 2018). Aerosol layers having a centroid above the tropopause are assigned to one of the stratospheric aerosol subtypes that are the subject of this paper.

CALIOP level 2 retrievals are reported in a variety of data products. The stratospheric aerosol subtyping algorithm is derived from layer properties reported in the level 2 aerosol layer product, and it is this same product that we primarily use
125 to evaluate the performance of the algorithm. The level 2 vertical feature mask, level 2 aerosol profile, and level 1B products are also used for demonstrating individual case studies.

To characterize the performance of the previous V4.1 stratospheric aerosol subtyping algorithm, we use the V4.2 level 2 data products. The V4.1 level 2 data products were replaced with V4.2 in 2018 to add a new science data set (SDS) reporting the minimum laser energy in each 80 km horizontal segment (the fundamental level 2 processing interval). This
130 new SDS was added to assist in quality screening data affected by low laser energy shots which began occurring primarily over the South Atlantic Anomaly (SAA) region in mid-2016 (CALIPSO Data Advisory Page, 2018). To reliably exclude affected level 2 retrievals in our analysis of V4.1 and V4.5 data, we use this SDS to impose the requirement that all laser pulses have at least 60 mJ within any 80 km horizontal segment. Note that beyond adding the new SDS, there is no difference between V4.1 and V4.2 in these level 2 products. In particular, the stratospheric aerosol subtyping algorithm is
135 identical in both data releases. Because V4.1 level 2 data is no longer publicly distributed, for the remainder of the paper we will refer only to V4.2.

To assess the performance of the revised algorithm, we use a pre-release test version of the V4.5 level 2 products. This test version accurately reflects the behavior of stratospheric aerosol subtyping in the official V4.5 level 2 release, planned for early 2023. Updates to the V4.5 level 2 algorithms – beyond those described in this paper – are primarily related
140 to tropospheric feature classification, boundary layer cloud clearing, and optical depth retrievals that are fully independent of aerosol subtyping (Tackett et al., 2021, 2022; Ryan et al., 2022). These updates have no impact on our analysis of

stratospheric aerosol detections. The V4.51 level 1B data used as input for the V4.5 level 2 products also contains several updates relative to the previous release, described in the CALIPSO Lidar Level 1 V4.51 Data Quality Statement (2022): The 532 nm daytime and 1064 nm calibration algorithms now mitigate the influence of low energy laser shots on the derived calibration coefficients. This corrects biases and reduces calibration uncertainty in these channels at SAA latitudes ($\sim 15^\circ$ S to 30° S) since the onset of low energy shots in mid-2016. Small corrections have also been made to the 1064 nm baseline shape (the shape of the profile measured by the detectors when the laser is not firing) having negligible impacts on the current analysis. Lastly, adjustments have been made to the polarization gain ratios to properly account for day and night differences rather than use the same value regardless of lighting conditions. The primary impact of these adjustments is to increase nighttime depolarization ratios by $\sim 4\%$ and decrease daytime depolarization ratios by $\sim 1\%$.

As a final note, we emphasize that the “layers” discussed throughout this paper are those detected by CALIOP at 5 km, 20 km, and 80 km resolutions. Each layer is only counted once regardless of horizontal extent because our intention is to characterize the classification frequencies for the unique layers that are input to the subtyping algorithm. As such, the true geospatial extent of aerosol from each event is not explicitly represented.

155 **3 Summary of the V4.2 stratospheric aerosol subtyping algorithm**

The V4.2 stratospheric aerosol subtyping algorithm, documented by Kim et al. (2018), evaluates several CALIOP measurables to identify four subtypes: volcanic ash, sulfate/other, smoke, and polar stratospheric aerosol (PSA). The method for discriminating between volcanic ash, sulfate, and smoke is based on an empirical analysis of joint distributions of estimated particulate depolarization ratio and feature integrated attenuated backscatter color ratio derived from the level 2 aerosol layer product for manually classified aerosol layers from specific events where the dominant aerosol type is well documented in the literature. Whereas manual classification is far more error-prone in the troposphere where multiple aerosol types often coexist, stratospheric aerosol events such as volcanic eruptions and large pyroCb injections tend to be episodic and their compositions are characterized by various independent methods. The events selected for the joint distributions are the only significant contributors to stratospheric aerosol loading at the time they are sampled, typically during the first 30 days after event initiation. Plumes are tracked manually in CALIOP imagery over successive days and their latitude/longitude/altitude boundaries are recorded for each CALIOP granule. The “plume boundaries” we select are rectangles of altitude \times along-track distance that encompass the plume, plus a ~ 1 km buffer of “clear-air” where no other features are detected. In order to avoid cloud contamination, plumes near or in contact with high altitude cirrus or overshooting cloud tops are excluded. All layers detected in the level 2 aerosol layer product within the rectangular plume boundaries contribute to the joint distributions. The full list of CALIOP granules and plume boundaries for all events analyzed for V4.2 and V4.5 development is reported in the Supplement.

The first dimension of the joint distribution relies on the depolarization sensitivity of CALIOP. An elevated depolarization ratio is an excellent discriminator for identifying non-spherical particles such as volcanic ash, dust, and cirrus

relative to spherical particles such as sulfate aerosol. Volume depolarization ratio (δ_v) is calculated from the ratio of
 175 CALIOP measurements of attenuated backscatter, $\beta' = (\beta_m + \beta_p) T_m^2 T_{O_3}^2 T_p^2$, measured perpendicular and parallel to the plane
 of the emitted pulse, which was originally linearly polarized:

$$\delta_v = \frac{\beta'_{\perp}}{\beta'_{\parallel}} = \frac{(\beta_{m,\perp} + \beta_{p,\perp}) T_{m,\perp}^2 T_{O_3,\perp}^2 T_{p,\perp}^2}{(\beta_{m,\parallel} + \beta_{p,\parallel}) T_{m,\parallel}^2 T_{O_3,\parallel}^2 T_{p,\parallel}^2} \quad (1)$$

180 Attenuated backscatter is comprised of molecular and particulate backscatter (β_m and β_p) and two-way transmittances due
 to molecular, ozone, and particulate extinction (T_m^2 , $T_{O_3}^2$, and T_p^2).

Because δ_v contains contributions from both molecular and particulate backscattering, it can under-represent the
 particulate depolarization of weakly scattering features. In order to correct for the molecular contribution, CALIOP aerosol
 subtyping uses an estimated particulate depolarization ratio (δ_p^{est}) according to Eq. (2) (Omar et al., 2009). Here, the mean
 185 attenuated scattering ratio (Kar et al., 2018) is defined as $R' = \langle \beta'(z) / \beta'_m(z) \rangle$, where the angle brackets indicate averaging
 over the vertical extent of a layer. The molecular attenuated backscatter, $\beta'_m = \beta_m T_m^2 T_{O_3}^2$, is computed from the MERRA-2
 model and the molecular depolarization ratio, δ_m , is 0.00366 at 532 nm (Hostetler et al., 2006).


$$\delta_p^{est} = \frac{\delta_v [(R' - 1)(1 + \delta_m) + 1] - \delta_m}{(R' - 1)(1 + \delta_m) + \delta_m - \delta_v} \quad (2)$$

190

This is an estimate of the true particulate ratio δ_p because Eq. (2) uses the R' term that still contains the two-way
 particulate transmittance T_p^2 within the layer: $R' = R T_p^2$, where $R = \langle (\beta_m + \beta_p) / \beta_m \rangle$. The true δ_p can be calculated from
 Eq. (2) if R is used rather than R' (e.g., Cairo et al., 1999). However, the value of T_p^2 cannot be retrieved at the point in
 CALIOP data processing where layer classification occurs because retrieving T_p^2 requires a lidar ratio which is assigned only
 195 after layer classification is complete. Therefore, as shown in Fig. 3a, δ_p^{est} will overestimate δ_p primarily for strongly
 attenuating layers having high depolarization. The overestimate is particularly acute for layers with low scattering ratios.

The primary benefit of using δ_p^{est} instead of δ_v for aerosol classification is demonstrated by Fig. 3b. At low
 scattering ratios, the molecular contribution dominates δ_v causing large underestimates of the true δ_p , particularly for
 strongly depolarizing layers. A sizeable fraction of depolarizing aerosol layers is susceptible to underestimation in the

200 CALIOP data record. For example, 50 % of stratospheric volcanic ash layers considered in this paper have $R' < 2.6$ at night which would yield underestimates of > 50 % in δ_p , leading to difficulty in differentiating between aerosol with lower and higher depolarizations. The compromise of using δ_p^{est} instead of δ_v is that additional systematic and random errors are propagated into δ_p^{est} (primarily random errors in R') which can cause errors in aerosol subtyping for weakly scattering layers. Despite this, the CALIOP level 2 algorithms use δ_p^{est} for aerosol subtyping to overcome the known underestimate
 205 that would arise from the molecular contribution to Eq. (1).



210 **Figure 3.** (a) Estimated particulate depolarization ratio as a function of optical depth for four values of the true particulate depolarization ratio in layers having unattenuated scattering ratios of 5 (solid lines) and 10 (dashed lines). (b) Volume depolarization ratio as a function of scattering ratio for four values of true particulate depolarization ratio.

The second dimension of the joint distribution is feature integrated attenuated backscatter color ratio, χ' , which can give qualitative information about particle size. This quantity is ratio of the feature integrated attenuated backscatters (γ'_λ), computed separately at both lidar wavelengths between layer top and base, and is derived as follows:

215

$$B_{\lambda,k} = \frac{B'_\lambda(z_k)}{T_{m,\lambda}^2(z_k)T_{O_3,\lambda}^2(z_k)} \quad (3)$$

$$g_\lambda = \frac{1}{2} \sum_{k=top+1}^{base} (z_{k-1} - z_k) (B_{\lambda,k-1} + B_{\lambda,k}) \quad (4)$$

$$\gamma'_\lambda = g_\lambda - \left[\frac{1}{2} (z_{top} - z_{base}) (B_{\lambda,top} + B_{\lambda,base}) \right] \quad (5)$$

$$\chi' = \gamma'_{1064} / \gamma'_{532} \quad (6)$$

220

This formulation corrects the attenuated backscatter coefficients for molecular and ozone attenuation and applies an approximate correction for molecular scattering (Vaughan et al., 2005), which is important for low optical depth layers typically found in the stratosphere. Molecular and ozone two-way transmittances T_m^2 and $T_{O_3}^2$ are obtained from the MERRA-2 model.


225

The joint distributions used for the V4.2 release are shown in Fig. 4 and the contributing events are documented in Table 1. Note that only a subset of the events in Table 1 contributed to the V4.2 analysis; more events were added for V4.5 (discussed in Sect. 4). The contributions for volcanic ash and sulfate in V4.2 were dominated by the June 2011 Puyehue-Cordón Caulle eruption and the August 2008 Kasatochi eruption, respectively. Very few stratospheric smoke events had been observed when these distributions were first constructed in 2015 during development for the V4.2 release. The primary contributor was the February 2009 “Black Saturday” bushfires in Australia where smoke reached altitudes of 19 km in the southern mid-latitudes (Siddaway and Petelina, 2011). In order to more fully sample δ_p^{est} and χ' for smoke, several high-

230

altitude, yet not stratospheric, smoke events were included in the joint distributions. From this, two populations emerged: smoke with low depolarization in the troposphere and higher depolarization in the stratosphere (Fig. 4). As we shall discuss later, enhanced depolarization is a common feature of smoke reaching the stratosphere associated with pyroCb activity, though we had not fully appreciated this fact during the V4.2 development.

235



240

Figure 4. Joint distributions used in V4.2 development: estimated particulate depolarization ratio (δ_p^{est}) and feature integrated attenuated backscatter color ratio (χ') for manually classified layers during events dominated by the aerosol type indicated in the title. Only layers with $\gamma'_{532} > 0.001 \text{ sr}^{-1}$ contribute. Histograms of layer numbers (N) are min-max normalized. Black lines indicate discrimination thresholds in V4.2 and red text indicates the algorithm classification.

Using the joint distributions in Fig. 4, thresholds were established for ash, sulfate, and smoke that minimized their overlap in measurement space. The flowchart in Fig. 5 shows how these thresholds are used in the V4.2 release. Stratospheric aerosol layers having $\delta_p^{est} > 0.15$ are classified as ash. Separate branches exist to capture smoke with low depolarization ($\delta_p^{est} < 0.075$) and with high depolarization ($0.075 < \delta_p^{est} < 0.15$), depending on the layer color ratio. Sulfate is identified for layers having low δ_p^{est} and low χ' .

Note that the name given to the V4.2 subtype containing sulfate is “sulfate/other”. The “other” contributions to this subtype are layers having low integrated attenuated backscatter. The ability to accurately discriminate between aerosol types using depolarization and color ratio requires measurements having sufficient backscatter magnitudes. This is especially relevant in the stratosphere where exceptionally low optical depths are common. Therefore, a test was placed prior to the determination of ash, sulfate, or smoke in the V4.2 flowchart (Fig. 5) to weed out weakly scattering features using the feature integrated attenuated backscatter at 532 nm (γ'_{532}) defined by Eq. (5).

In the V4.2 data release, all stratospheric aerosol layers having $\gamma'_{532} < 0.001 \text{ sr}^{-1}$ (hereafter named “low- γ' layers”) that were not previously classified as PSA are assigned to the sulfate/other subtype. The rationale for combining sulfate and low- γ' layers was because we assumed volcanic sulfate would eventually become weakly scattering over time since this aerosol type tends to persist in the stratosphere for weeks to months after injection, all-the-while becoming increasingly diffuse and hence decreasing in optical depth.

V4.1/V4.2 Stratospheric Aerosol Subtyping Flowchart




Figure 5. Flowchart for stratospheric aerosol subtyping in V4.2.

The fourth subtype given by the stratospheric aerosol subtyping algorithm is for polar stratospheric aerosol. The PSA type is meant to assign a reasonable classification for aerosol layers identified in regions where PSC formation is likely. The method to identify PSA is identical for both the V4.2 and V4.5 releases, so it is only briefly summarized here. PSA layers are identified first by evaluating the layer midpoint temperature for stratospheric aerosol layers during polar-winter PSC seasons. Based on the climatology of Poole and Pitts (1994), PSC season is assumed to be December to February for the Arctic and May to October for the Antarctic. A latitude threshold of 50° confines PSA classifications to the appropriate pole and encompasses the expected range of PSC/PSA formation. Temperature information is provided by the MERRA-2 model. The -70°C layer midpoint temperature maximum is based on the midpoint temperatures of stratospheric aerosol layers detected by CALIOP in these regions/seasons (see Fig. 5 of Kim et al. (2018)). It also agrees with the observed temperatures for PSC formation of Rosen et al. (1997).

The classification of PSA is the first decision in the stratospheric aerosol typing flowchart in Fig. 5. This step confines the PSA classification to only those geographical regions and seasons where they are expected to exist. These layers are often detected adjacent to features classified as cloud and have depolarization levels that overlap with the expected ranges for sulfate, smoke, and ash (Fig. 6). It is quite possible that some fraction of these are CAD misclassifications along the fringes of clouds rather than legitimate aerosol (Liu et al., 2019). The PSA classification prevents these layers from being misclassified as volcanic ash or sulfate when none exists. The flip side of this is that when true volcanic aerosol enters these regions during PSC season, it may not be classified correctly. An example is considered in Sect. 5.1. The confidence in the PSA classification is considered low at this time and its accuracy is not evaluated further in this paper. Users investigating PSC observations by CALIOP are instead referred to the level 2 polar stratospheric cloud mask product which is specialized for this purpose (NASA/LARC/SD/ASDC, 2016a; Pitts et al., 2018).




Figure 6. Same joint distributions as Fig. 4, but for layers classified as PSA in the southern hemisphere in 2013, based on V4.5 data.

285 We now turn our attention to the changes made to the stratospheric aerosol subtyping algorithm for the V4.5 release and how they improve the shortcomings in the previous V4.2 release.

4 The version 4.5 stratospheric aerosol subtyping algorithm

In Sect. 1 we highlighted the need to improve several aspects of the stratospheric aerosol subtyping algorithm. The goals of the revisions implemented in V4.5 are therefore to:

290

- Improve discrimination between volcanic ash and smoke in the stratosphere.
- Remove the ambiguity in the sulfate/other subtype.
- Reduce the number of layers classified as low- γ' .
- Update the volcanic ash lidar ratio.

295

In order to accomplish this, two structural changes were made to the algorithm and two thresholds were adjusted. The new flowchart for V4.5 is shown in Fig. 7. As in the previous release, discrimination between volcanic ash, sulfate, and smoke is determined by an empirical analysis of joint distributions of depolarization and color ratios from events where the aerosol type is known. The joint distributions used for the V4.5 analysis are shown in Fig. 8. More events have been added since the V4.2 analysis so that we can better characterize the depolarization and color ratio for these types (denoted by daggers in Table 1). The new events include two volcanic eruptions: sulfate layers from the July 2009 Sarychev Peak eruption and ash layers from the April 2015 Mount Calbuco eruption. The plume of volcanic aerosol from Sarychev Peak primarily consisted of sulfate based on AIRS retrievals and in-situ aircraft measurements (Prata et al., 2014; Andersson et al., 2013). The Mount Calbuco eruption, discussed in detail in Sect. 4.1, injected large quantities of ash and a lesser amount of SO₂ into the atmosphere (Marzano et al., 2018) resulting in sulfate formation about a month after the eruption (Bègue et al., 2017). To sample the ash component in the joint distribution, we only selected layers within the first two weeks of the eruption; the CALIOP depolarization was elevated for these layers, consistent with ash. Four new wildfire events were also added which had a critical influence on the revisions that were made: in particular, smoke layers from the August 2017 Pacific Northwest (PNW) event and the December 2019/January 2020 Australian New Year (ANY) event. These events injected large amounts of depolarizing smoke into the stratosphere (Peterson et al., 2018; Hu et al., 2019; Ohneiser et al., 2020). Stratospheric smoke layers were also added from the December 2006 Australian bushfires (Dirksen et al., 2009) and from the July 2014 North American wildfires. The much larger number of stratospheric smoke layers now contributing to the joint distribution analysis allows us to exclude the tropospheric smoke layers that previously influenced the V4.2 threshold selections.

315

V4.5 Stratospheric Aerosol Subtyping Flowchart





Figure 7. Flowchart for stratospheric aerosol subtyping in the V4.5.



320 **Figure 8.** Joint distributions used in V4.5 development; same description as Fig. 4, but only stratospheric aerosol layers with $\gamma'_{532} > 0.001$ sr^{-1} contribute. Black lines indicate discrimination thresholds in V4.5 and red text indicates the algorithm classification.

325 **Table 1.** Manually classified events used to establish thresholds to discriminate between volcanic ash, volcanic sulfate, and smoke in V4.5. The number of unique stratospheric aerosol layers detected in V4.5 is given along with the fraction of these layers in night granules, and the dominant aerosol type. Only layers with $\gamma'_{532} > 0.001$ sr^{-1} contribute. The individual CALIOP granule names and latitude/longitude/altitude information for these layers are given in the Supplement.

N layers	Fraction	Event	Dominant aerosol type
----------	----------	-------	-----------------------

in V4.5	at night (%)		
2528	100	Puyehue-Cordón Caulle eruption, June 2011	Volcanic ash (Bignami et al., 2014)
687	50	†Calbuco eruption, April 2015	Volcanic ash (Marzano et al., 2018)
36	100	Chaitén eruption, May 2008	Volcanic ash (Prata et al., 2010)
3095	29	†Sarychev eruption, June 2009	Sulfate (Prata et al., 2017)
2148	100	Kasatochi eruption, August 2008	Sulfate (Krotkov et al., 2010)
186	100	Nabro eruption, June 2011	Sulfate (Theys et al., 2013)
63	100	Okmok eruption, July 2008	Sulfate (Prata et al., 2010)
6918	75	†Australian New Year (ANY) event, Dec. 2019/Jan. 2020	Smoke (Kablick et al., 2020)
5016	63	†Pacific Northwest (PNW) event, August 2017	Smoke (Peterson et al., 2018)
2177	11	†North American wildfires, July 2014	Smoke
1720	63	†Australian bushfires, December 2006	Smoke (Dirksen et al., 2009)
760	100	Black Saturday Australian bushfires, February 2009	Smoke (Siddaway and Petelina, 2011)
187	100	Siberian wildfires, May 2012	Smoke
15	100	Siberian wildfires, June 2007	Smoke
0	0	*Canadian wildfires, July–August 2007	Smoke

†New events added since the V4.2 development of the stratospheric aerosol typing algorithm described in Kim et al., 2018.

* Event is exclusively comprised of tropospheric layers used in V4.2 development, but not V4.5.

4.1 Color ratio test for smoke removed

330 The new joint distributions in Fig. 8 show that excluding tropospheric smoke layers from the sample population had an important impact: the population of smoke with low depolarization and higher color ratio is no longer prominent (notice that the secondary mode labeled “tropospheric smoke” in Fig. 4 is not evident in Fig. 8). Recent literature has confirmed the enhanced depolarization from smoke lofted by pyroCb events. The analysis of Christian et al. (2020) demonstrated increased depolarization and decreased color ratio for fresh pyroCb plumes at high altitudes compared to lower altitudes. This is corroborated by Fig. 9, which shows higher values of δ_p^{est} for stratospheric smoke layers used for the V4.5 joint distribution compared to the tropospheric smoke layers contributing to the V4.2 joint distribution.

335




Figure 9. Frequency distributions of estimated particulate depolarization ratio (δ_p^{est}) for stratospheric smoke layers in Table 1 (red) and tropospheric smoke layers contributing to the V4.2 joint distributions (496 layers from the May 2012 Siberian wildfires and 36 layers from July–August 2007 Canadian wildfires). Only layers with $\gamma'_{532} > 0.001 \text{ sr}^{-1}$ contribute.

340

These observations, primarily representing pyroCb events, suggest that smoke injected to extremely high altitudes contains particles that are aspherical and smaller (based on enhanced depolarization and lower color ratios, respectively) compared to smoke injected to lower altitudes. The cause of aspherical particles in smoke plumes from these pyroCb events is an active area of research (e.g., Gialitaki et al., 2020; Haarig et al., 2018; Kablick et al., 2018; Sicard et al., 2019).

345

However, the message is clear. Smoke reaching the stratosphere in these events typically depolarizes the CALIOP backscatter signal more than smoke that is confined to the troposphere. Based on this information, we removed the color ratio test, which previously was employed to capture the low depolarization, higher color ratio smoke that is characteristic of tropospheric events. Because smoke layers having low depolarization/high color ratio are not routinely observed in the stratosphere, then there is no need to search for them. This marks the first structural change. The V4.5 stratospheric aerosol subtyping algorithm now strictly relies on depolarization ratio to discriminate between ash, sulfate, and smoke.

350

A caveat is the possibility of misclassification by relying solely on depolarization ratio to discriminate between sulfate and smoke. Recent research by Ohneiser et al. (2021) hypothesizes that smoke from Siberian wildfires in 2019 self-lofted from the troposphere into the UTLS. Because this smoke was aged and not of pyroCb origin, its depolarization was low (< 0.05) causing the CALIOP V4.2 stratospheric aerosol subtyping algorithm to misclassify these smoke layers as sulfate (Ansmann et al., 2021). The subtyping algorithm will continue to struggle in these cases in V4.5 due to the similarly low depolarization ratios of tropospheric smoke and sulfate. It is currently unknown how often smoke plumes reach the UTLS exclusively by self-lifting.

355

Removing the color ratio test has the benefit of allowing more sulfate layers to be classified correctly as sulfate rather than smoke. This improves the accuracy of the retrieved extinction for these sulfate layers because the correct lidar ratio of 50 sr will be selected rather than 70 sr (Sect. 4.5), thereby avoiding a 40 % overestimate in aerosol optical depth

360

(AOD). Conversely, there will be smoke layers misclassified as sulfate due to the overlap in the δ_p^{est} distributions for these two types, yielding a 40 % underestimate in AOD due to selecting the lower lidar ratio. The relative frequencies of correct and incorrect classification for sulfate and smoke will be evaluated in Sect. 4.6.

4.2 Depolarization ratio threshold between smoke and ash increased

365 The estimated particulate depolarization ratio (δ_p^{est}) threshold to discriminate between ash and smoke was 0.15 in the V4.2
 release. This threshold worked well for the 2009 Black Saturday Australian bushfires which had particulate depolarization
 ratios around 0.10 to 0.15. However, depolarization ratios were higher for stratospheric smoke layers from the PNW event in
 August 2017 (Fig. 10a). European lidar systems observed 532 nm particulate depolarization ratios ranging from 0.15 to 0.2
 in the two weeks following the event (Ansmann et al, 2018; Haarig et al., 2018; Khaykin et al., 2018; Hu et al., 2019). Figure
 370 10b shows the distributions of CALIOP estimated particulate depolarization for all manually classified stratospheric smoke
 and volcanic ash events listed in Table 1. The markedly larger depolarization of layers from the PNW event compared to
 previous stratospheric smoke events caused a large frequency of misclassification in V4.2: whereas ~25 % of all smoke
 events excluding the PNW event are misclassified as volcanic ash with the 0.15 threshold, a whopping 58 % of the smoke
 layers detected during the PNW event were misclassified as ash.

375




Figure 10. Frequency distributions of estimated particulate depolarization ratio (δ_p^{est}) for layers identified in Table 1, comparing (a) three major stratospheric smoke events and (b) all stratospheric smoke events to all volcanic ash events. Only layers with $\gamma'_{532} > 0.001 \text{ sr}^{-1}$ contribute.

380 In order to better discriminate between stratospheric smoke and volcanic ash in V4.5, the δ_p^{est} threshold between
 these types was increased from 0.15 to 0.25. This is roughly the minimum overlap between the distributions in Fig. 10b.
 Doing so reduces the amount of smoke layers misclassified as ash in the PNW event to 9 % (Sect. 4.6). The example from
 this event in Fig. 11a shows that the dominant classification is now smoke rather than ash (cf. Fig. 1). A more thorough
 assessment of the classification performance is given in Sect. 5.

385




Figure 11. Aerosol subtyping after V4.5 revisions for (a) pyroCb smoke plume from the PNW event (cf. Fig. 1) and (b) volcanic ash plume from the June 2011 Puyehue-Cordón Caulle eruption (cf. Fig. 2).

4.3 Sulfate/other separated into two types: sulfate and unclassified

390 The second structural change to the stratospheric aerosol subtyping algorithm is the separation of the “sulfate/other”
 classification into two separate types: sulfate and “unclassified”. Previously, both low- γ' layers and layers meeting the sulfate
 criteria were given the sulfate/other classification. This hampered the ability to easily identify volcanic sulfate layers because
 low- γ' layers can be any aerosol type, provided γ'_{532} is sufficiently low. Returning to the example in the introduction, Fig.
 11b shows that the majority of the ash plume is now correctly classified whereas before it was classified as sulfate/other (cf.
 395 Fig. 2). This is a result of separating the low- γ' branch and also lowering the γ'_{532} threshold (discussed next). The
 nomenclature for the low- γ' branch has also changed to unclassified to emphasize that no attempt has been made to classify
 the subtype based on its depolarization. This subtype serves as a catch-all for aerosol having insufficient backscatter to yield
 confident classifications of volcanic ash, sulfate, or smoke. We will characterize unclassified layers in Sect. 5.4.

4.4 Low- γ' threshold decreased

400 The second threshold adjustment was for low- γ' layers, now identified as “unclassified”. At low signal levels, δ_p^{est}
 becomes prone to increases in systematic and random errors along with the increased risk of overestimates as discussed in
 Sect. 3. Previously the threshold was $\gamma'_{532} < 0.001 \text{ sr}^{-1}$ which caused 75 % of all stratospheric aerosol layers detected from
 June 2006 – December 2018 to be classified as sulfate/other. In the V4.5 release, this threshold has been lowered to assign
 the unclassified subtype to layers having γ'_{532} in the lowest quartile of the June 2006 – December 2018 stratospheric aerosol
 405 distribution, corresponding to $\gamma'_{532} < 0.0003 \text{ sr}^{-1}$ at day and $\gamma'_{532} < 0.00025 \text{ sr}^{-1}$ at night. The lowest-quartile γ'_{532} metric was
 selected based on the rapid increase in relative uncertainty in δ_p^{est} as a function of decreasing γ'_{532} (Fig. 12). Based on the

low- γ' thresholds in V4.5, the average relative uncertainty in δ_p^{est} is less than 250 % for layers classified as ash, sulfate, and smoke at all detection resolutions – except for layers detected at 80 km resolution in daytime which typically have uncertainties of ~ 400 %. Classifications for daytime, 80 km resolution stratospheric aerosol layers that are not assigned the unclassified subtype should be interpreted with caution. The average relative uncertainty for stratospheric layers detected at 5 km resolution is < 50 %.




Figure 12. Average relative uncertainty in estimated particulate depolarization ratios as a function of layer integrated attenuated backscatter (γ'_{532}) for all stratospheric aerosol layers detected between June 2006 – December 2018, segregated by horizontal averaging required for detection. Solid lines are nighttime detections and dashed lines are daytime detections. Layers classified as PSA, $|\text{CAD Score}| < 20$ (indicating no confidence in cloud-aerosol discrimination), or top altitudes above 20 km are excluded.

For reference, the average relative uncertainty in δ_p^{est} was less than 200 % for stratospheric aerosol layers having γ'_{532} above the previous low- γ' threshold in V4.2. With the new thresholds, layers detected at 5 km and 20 km horizontal resolution, regarded as robustly scattering features, will almost certainly be subtyped as something other than unclassified, as will approximately half of the 80 km resolution layers. This reduces the frequency of unclassified aerosol layers to 25 % with a corresponding increase of 50 percentage points in the relative uncertainty of δ_p^{est} which is not expected to noticeably degrade the fidelity of the subtyping algorithm. Rather, we consider the increase in opportunities to classify stratospheric aerosol layers outweighs the increase in uncertainty.

If we assume a lidar ratio of 50 sr for sulfates (i.e., as in Kim et al. (2018)), we can translate the γ'_{532} thresholds into approximate optical depth thresholds. Using the V4.2 threshold of 0.001 sr^{-1} caused all layers with optical depths less than ~ 0.053 to be classified as sulfate/other. By contrast, the revised γ'_{532} threshold of 0.0003 sr^{-1} translates into an optical depth threshold of ~ 0.015 , so that only those layers with optical depths less than ~ 0.015 are identified as unclassified.

430 4.5 Lidar ratio for ash increased

As emphasized in the introduction, the algorithmic motivation for improving aerosol subtyping is to ensure that a representative lidar ratio is selected for each subtype, thereby yielding accurate extinction retrievals. Because CALIOP retrievals operate from top-down, any errors in the retrieved extinction due to incorrect lidar ratio selection will propagate into underlying layers. The previous subsections described the improvements we made to increase the likelihood that the correct stratospheric aerosol subtype (i.e., correct lidar ratio) will be selected. However, it is also important that the lidar ratio for these subtypes are representative of what is observed in nature. A full, detailed accounting of error propagation in CALIOP extinction retrievals is given in Young et al., 2013, including the impacts of incorrect lidar ratio selection. One key take-away from that paper is that the relative error in retrieved AOD equals the relative error in the lidar ratio for layers having low optical depths, typical of aerosol. A second key take-away is that the error in retrieved AOD due to incorrect lidar ratio selection behaves as a systematic bias in the retrieval of optical depth for underlying layers. The magnitude of the error in AOD for the underlying layer depends on its R' , AOD, and the magnitude of the error in lidar ratio of the overlying layer. In general, though, the sign of the AOD bias in the underlying layer is the same as the sign of the error in lidar ratio selection of the overlying layer (i.e., selecting too low of a lidar ratio causes an underestimate in the AOD retrieved for layers at lower altitudes). Feature detection accuracy for lower layers can also be degraded if overlying attenuation is not correctly accounted for. Clearly, accurate lidar ratio selection is critical for elastic backscatter lidar retrievals.

The stratospheric aerosol lidar ratios assignments for V4.5 are shown in Table 2. The same values were used in V4.2 for sulfate (previously sulfate/other), smoke, and PSA, as justified by Kim et al., 2018. The lidar ratio for the unclassified subtype is based on extinction retrieval considerations. As discussed in Sect. 4.3, unclassified layers can be any subtype provided their γ'_{532} is sufficiently low. In order to reduce the impact of extinction retrieval errors that propagate into underlying layers, it is better to use a lidar ratio that is too low rather than too high when accurate knowledge of the subtype is unavailable (Young et al., 2013). Therefore, the unclassified subtype shares the same lidar ratios as sulfate because these are the lowest lidar ratios expected for non-PSA stratospheric aerosol layers.

The 532 nm lidar ratio for ash was increased in V4.5. In the previous release, the default lidar ratio for volcanic ash was set to 44 sr at both 532 nm and 1064 nm, matching the lidar ratios of the tropospheric dust subtype (Kim et al., 2018). This choice was motivated by similarities between the size distributions of volcanic ash and dust (Winker et al., 2012), and to avoid discontinuities in extinction retrievals between ash above and below the tropopause (because ash and dust are depolarizing, volcanic ash will be misclassified as dust in the troposphere). However, several analyses suggest that the 532 nm lidar ratio for ash is higher. Ash plumes from the April 2010 Eyjafjallajökull eruption, observed by EARLINET lidar measurements, revealed 532 nm lidar ratios over Germany from 45–60 sr (Ansmann et al., 2010; Groß et al., 2012). Higher 532 nm lidar ratios for the plume were observed over Italy, ranging from 50–92 sr (Mona et al., 2012), and over Greece, ranging from 44 to 88 sr (Kokkalis et al., 2013). These latter studies suggest that relative humidity and/or ageing may have played a role in the variability. Prata et al. (2017) used CALIOP constrained retrievals to characterize ash 532 nm lidar ratios

from the 2011 Puyehue-Cordón Caulle eruption. The constrained retrieval method computes transmittance if there is clear air above and below a feature, thus allowing a measurement of the layer lidar ratio at 532 nm. Their analysis reveals a median ash 532 nm lidar ratio of 67 sr for this event.

Table 2. Lidar ratios for stratospheric aerosol subtypes in V4.5.

Aerosol Subtype	S ₅₃₂ (sr)	S ₁₀₆₄ (sr)
Volcanic ash	61 ± 17	44 ± 13
Sulfate	50 ± 18	30 ± 14
Smoke	70 ± 16	30 ± 18
Unclassified	50 ± 18	30 ± 14
Polar stratospheric aerosol	50 ± 20	25 ± 10

Based on the growing consensus in the literature that the 532 nm lidar ratio for volcanic ash is typically larger than the value used in V4.2, we have revised the 532 nm lidar ratio in the V4.5 release. Following the method of Prata et al. (2017), Fig. 13 shows the 532 nm lidar ratios from CALIOP constrained retrievals for ash layers from several eruptions including the Puyehue-Cordón Caulle (2011), Kelud (2014), Sarychev (2009), and Calbuco (2015). In order to remove no-confidence retrievals and any possible cloud contamination, only layers with retrieved lidar ratio uncertainty < 100 % and |CAD score| > 20 contribute to the histogram. Using the mean and standard deviation of lidar ratios derived from this analysis, the default 532 nm lidar ratio is increased to 61 ± 17 sr, consistent with values reported in the literature. This will increase the retrieved AOD for ash layers by ~39 % and prevent underestimates of optical depth for underlying layers. Because knowledge of 1064 nm lidar ratios is not as broad in the literature, the default 1064 nm lidar ratio for ash will not be changed for the V4.5 release.




Figure 13. Distribution of 532 nm lidar ratios retrieved from constrained retrievals of ash layers from several volcanic events (see text). Lines indicate default values for ash in V4.2 and V4.5.


Recent ground-based lidar and lidar-photometer retrievals of smoke arising from pyroCb events have measured higher 1064 nm lidar ratios than the default value used by CALIOP (30 sr), with values ranging from 80 – 120 sr (Haarig et al., 2018; Ohneiser et al., 2022). The 1064 nm lidar ratio used for all smoke layers in V4.2, and carried forward into V4.5, is based on AERONET retrievals of tropospheric smoke (Sayer et al., 2014). Microphysical differences likely exist between smoke injected into the stratosphere from pyroCb events and smoke residing in the troposphere from less explosive events, so lidar ratio differences are plausible. We plan to reevaluate 1064 nm lidar ratios for stratospheric smoke and ash in a future data release.

4.6 Change in classifications between V4.2 and V4.5

The changes in classifications due to V4.5 revisions are summarized in Fig. 14 based on all the manually classified events in Table 1. Here we are comparing unique layers detected within the plume boundaries reported in the Supplement. Because each rectangular boundary contains a buffer of clear-air around the actual plume (Sect. 3), it is not necessary to perfectly match layer top and base altitudes between versions. We are only concerned with the classifications within those boundaries, and the buffer allows for any differences in layer detection that may occur. The hatched bars in Fig. 14 indicate low- γ' features (“other” in V4.2 and unclassified in V4.5). One obvious change for ash-dominant events is the radical reduction of the sulfate/other subtype and accompanying increase in ash classifications. This is a direct consequence of reducing the threshold for low- γ' layers; the new ash layers have γ'_{532} somewhere between the old and new qualifying thresholds. Evidently, δ_p^{est} is still an excellent discriminator for ash even as γ'_{532} decreases, bolstering our confidence in our decision to reduce the γ'_{532} threshold. There is a minor increase in smoke misclassifications where δ_p^{est} for some ash layers is just low enough to resemble that of depolarizing smoke.

For the sulfate-dominant events in V4.2, a substantial fraction of sulfate/other classifications were low- γ' layers (hatched bars in Fig. 14). A small number of these low- γ' layers became unclassified in V4.5 after separating the sulfate/other class. Meanwhile a larger number became classified as sulfate because reducing the γ'_{532} threshold and removing the color ratio test allowed more opportunities for the sulfate classification. The net effect is that sulfate is the dominant subtype given for these events. There is a small increase in smoke misclassification for at least three reasons. (1) These layers could be mixtures of sulfate and ash, yielding moderate values of δ_p^{est} (discussed further in Sect. 5.2). (2) The increased variability in δ_p^{est} due to the greater influence of random and systematic errors for layers having γ'_{532} in between the old and new low- γ' thresholds allows more opportunities for sulfate layers to exceed the $\delta_p^{est} > 0.075$ threshold. (3) A small number of ash classifications changed to smoke due to the increase in δ_p^{est} threshold separating these types. Even though smoke misclassification increases for these sulfate-dominant events, eliminating the color ratio test improved the accuracy. For all the sulfate-dominant events shown in Fig. 14, the V4.5 classification frequency for sulfate and smoke is 70 % and 28 %, respectively. Retaining the color ratio test would have yielded classification frequencies of 58 % and 41 %, respectively. Given the limited CALIOP observables, discriminating sulfate from smoke will always be a challenge, and hence this seemingly modest improvement represents a welcome and useful increase in classification accuracy. There remains a substantial overlap in the δ_p^{est} distributions for these two types, which inherently reduces the ability to discriminate sulfate with a high degree of accuracy.

Classifications for pyroCb-smoke dominated events show a marked improvement. The frequency of layers classified as smoke increased while misclassifications as other types decreased. As previously mentioned, layers from the PNW event were primarily misclassified as ash in V4.2 (~58 %). Now smoke is the dominant classification (~85 %), with an ash misclassification rate around 9 %. A similar reduction in ash misclassification occurs in the other stratospheric smoke events highlighted in Fig. 14. Lowering the low- γ' threshold also improves smoke classification by moving many layers previously classified as sulfate/other to the smoke subtype. Just as with volcanic ash, the elevated depolarization ratios permit recognition of these layers as depolarizing smoke despite having lower γ'_{532} . There is a small increase in sulfate misclassifications for smoke layers having $\delta_p^{est} < 0.075$ due to the removal of the color ratio test.



530 **Figure 14.** Stratospheric aerosol subtype classification frequency for events dominated by ash, sulfate, and smoke for V4.2 (left) and V4.5 (right) based on manually identified layers in Table 1. Low- γ' layers, based on the γ' threshold for the relevant version, are indicated by hatched bars.

5 Performance assessment

535 We now take a closer look at the geographical and time-evolution of classifications for specific stratospheric aerosol events to assess the performance of the algorithm. Whereas the previous section summarized classifications for manually identified volcanic ash, sulfate, and smoke layers, we now evaluate the classifications for all aerosol layers detected in the stratosphere following major aerosol injections. As before, we have selected events where the dominant aerosol subtype is known based on literature sources. Broadening our evaluation to include all aerosol layers detected in the stratosphere rather than manually identified layers gives a sense of the fidelity of the algorithm in the wide range of scenes that CALIPSO encounters. Subtype classification frequencies and depolarization ratio statistics are for night and day layer detections collectively unless otherwise noted.

540

5.1 Ash dominated events

The stratospheric aerosol typing algorithm performs exceptionally well at identifying volcanic ash. One event dominates the CALIPSO record for this aerosol type: the Puyehue-Cordón Caulle eruption on 4 June 2011. Located in southern Chile (40.6° S, 72.1° W), the volcano injected an estimated ~0.4 Tg of ash into the atmosphere of the southern hemisphere (Bignami et al., 2014) to altitudes of 12–14 km (Ulke et al., 2016). The plume circumnavigated the globe, affecting air traffic in multiple countries (Wunderman 2012). A strong signature of ash was evident based on elevated CALIOP depolarization ratios and ash retrievals by MODIS and IASI (Klüser et al., 2013; Vernier et al., 2013; Bignami et al., 2014; Maes et al., 2016; Prata et al., 2017, 2020; Christian et al., 2020).

CALIOP detected ash primarily after 15 June 2011. (The CALIPSO payload was turned off due to adverse space weather during 6–15 June). Stratospheric aerosol layer detections in Fig. 15a show that the layers detected during 15–28 June 2011 span all longitudes, primarily south of 30° S. These layers were mainly detected from 8 to 14 km (Fig. S1a). The median δ_p^{est} is 0.34 for all stratospheric aerosol layers in the southern hemisphere (Fig. 16), excluding those given the PSA subtype. Therefore, ash is the dominant subtype assigned at a frequency of 84.3 %, with smoke and sulfate classifications at rates of 3.3 % and 0.2 %, respectively. Smoke misclassifications are expected as well, given the overlap in distributions of δ_p^{est} between the volcanic ash and depolarizing smoke regimes. In addition, the daily median δ_p^{est} of detected layers steadily decreased from 0.35 to 0.25 during days +15 to +45 after 4 June (–0.02 /week), possibly contributing to the number of smoke misclassifications. Nonetheless, ash remained the dominant aerosol subtype classification for over 45 days past the eruption (Fig. 15b).





Figure 15. For ash-dominated events in V4.5; (left) locations of subtype classifications following first CALIOP detection and (right) time-history summation of subtype classifications. Volcano locations denoted by red triangles. Hatched areas indicate missing CALIOP data. The payload was down for testing and spacecraft maneuvers during part of 27 April 2015, causing the reduced number of layer detections on that date in panel (d).

565

The time period of the Puyehue-Cordón Caulle eruption coincided with the beginning of PSC season over Antarctica and many PSA classifications are evident in Fig. 15a. During the first two weeks after the main eruption, the volcanic ash and PSA classifications were mostly separated in altitude. Figure 17 compares the layer top altitudes of ash and PSA classifications south and north of 50° S to that of PSA classifications which can only be south of 50° S. Ash layers are confined below 14 km whereas 64 % of the PSA classifications are above this altitude. These higher altitude layers are likely legitimate PSA classifications due to their low temperatures (-70 °C) and low depolarization ratios (median ~ 0.02), consistent with liquid supercooled ternary solution droplets (Pitts et al., 2011). The accuracy is questionable for the remaining 36 % of PSA classifications below 14 km, accounting for 5 % of all layers detected at these altitudes. These layers have a median depolarization of 0.32, consistent with ash, though some mixtures of PSC particles also have elevated depolarization ratios (Pitts et al., 2011). Given the coincident altitude with the ash plume, some of these layers are likely to be ash misclassified as PSA.

570

575



580 **Figure 16.** Estimated particulate depolarization ratio for all unique stratospheric aerosol layers detected in the first two weeks following first CALIOP observation of the volcanic plume. PSA and layers with $|\text{CAD score}| < 20$ are excluded.




Figure 17. Top altitudes for layers classified as ash and PSA during 15–28 June 2011.

585

The second major volcanic ash event we evaluate is the Mount Calbuco eruption on 22–23 April 2015. Also a Chilean volcano, Mount Calbuco (41.3° S , 72.6° W) injected an estimated 3 Tg of volcanic ash (Marzano et al., 2018) and 0.2–0.4 Tg of SO_2 with initial plume heights reaching 18–21 km (Pardini et al., 2018; Zhu et al., 2018). Lidar and AERONET observations from São Paulo acquired just days after the eruption indicated the presence of both sulfate and ash
 590 (Lopes et al., 2019), though the bulk of sulfate formation did not complete until the second half of May (Bègue et al., 2017). During the first two weeks, however, the presence of volcanic ash in the plume was confirmed by elevated lidar depolarization ratios (Klekociuk et al., 2020) and by negative $10.06 - 12.05 \mu\text{m}$ brightness temperature differences (Prata, 1989) measured by the Imaging Infrared Radiometer, also on board the CALIPSO platform (Fig. S2).

CALIOP detections of stratospheric aerosol layers during this time are primarily along $30^\circ \pm 10^\circ \text{ S}$ from Chile to
 595 the western coast of Australia (Fig. 15c). Most of these layers were detected between 12 and 22 km (Fig. S1b). The median

δ_p^{est} for stratospheric aerosol layers from Calbuco was slightly lower than Puyehue-Cordón Caulle, at 0.31 for the first two weeks (Fig. 16). These layers experienced a more rapid decline in depolarization of -0.10 /week. This decline was also observed by the Cloud-Aerosol Transport System (CATS) lidar at 1064 nm (Christian et al., 2020). Ash classifications are the dominant CALIOP subtype identified during this event, at 69.4 % (Fig. 15d). Sulfate classifications accounted for 2.7 % of layers detected during these two weeks, which is reasonable based on AERONET observations consistent with sulfate over Chile (Lopes et al., 2019). Due in part to the broader distribution of δ_p^{est} compared to Puyehue-Cordón Caulle, smoke misclassifications occur at a higher frequency of 21.1 % beginning ~ 5 days following the eruption. Notably, while the median δ_p^{est} decreased in days +1 to +5 following the eruption compared to days +6 to +10, the breadth of the δ_p^{est} distribution remained roughly the same (median absolute deviation (MAD) of 0.057 and 0.053, respectively). It is possible that these smoke misclassifications are mixtures of ash and sulfate in the same air mass measured by CALIOP, resulting in intermediate δ_p^{est} values between the two aerosol types. This cannot be definitively established with CALIOP measurements alone, however.

5.2 Sulfate dominated events

Two major stratospheric sulfate events are selected for assessment. The first is the Nabro stratovolcano in Eritrea (13.37° N, 41.7° E) which erupted on 12–13 June 2011, injecting an estimated 1.5 Tg of SO₂ into the upper troposphere-lower stratosphere (Clarisse et al., 2014). A second injection into the stratosphere on 16 June was inferred based on geostationary and limb-profiling satellite data (Fromm et al. 2014). The SO₂ plume initially traveled east and then followed the Asian summer monsoon anticyclonic circulation over northern Africa, the Middle East, and Asia for the first two weeks (Fairlie et al., 2014). Sulfate aerosol then transported to the rest of the northern hemisphere over July and August.

We focus on the region shown in Fig. 18a. CALIOP detected the majority of the stratospheric plume between 13 to 19 km during the two weeks following the initial eruption (Fig. S3a). The median δ_p^{est} for these layers is the smallest of all volcanic events evaluated, at 0.021 (Fig. 16), consistent with various ground based lidars that also measured small depolarization from the Nabro plume, (Zhuang and Yi, 2016; Noh et al., 2017). This is indicative of sulfate aerosol. Due to the low δ_p^{est} , CALIOP classifies 82.6 % of these layers as sulfate. Smoke and ash classifications account for 5.4 % and 1.7 %, respectively, consistent with observations suggesting only a small ash component to the eruption (Clarisse et al. 2014).




Figure 18. Same as Fig. 15, but for sulfate-dominated events.


625

The second sulfate-dominated event we evaluate had a slightly larger ash component. Kasatochi, an island volcano along the Aleutian arc (52.17° N, 175.51° W), erupted on 7–9 August 2008, injecting SO_2 and ash up to ~ 15 km in altitude (Waythomas et al. 2010). In subsequent days, signatures of volcanic ash and SO_2 were observed spreading eastward over the Pacific by MODIS, AVHRR, AIRS, and OMI (Corradini et al., 2010; Krotkov et al. 2010). An analysis of AIRS measurements by Prata et al. (2010) suggests that the total ash mass injected was approximately 25 % smaller than the mass of SO_2 . Based on passive imager retrievals and modeling analyses, the greatest fraction of this ash is believed to have settled out of the plume during the first week following the eruption (Martinsson et al., 2009; Guffanti et al., 2010; Langmann et al., 2010). The most long-lasting component of the emission was SO_2 and subsequent sulfate aerosol that persisted for over two months. At the time, it was the largest injection of SO_2 into the atmosphere in over 17 years, with SO_2 mass estimates of 1.2–1.7 Tg (Kristiansen et al. 2010; Prata et al. 2010).

The majority of stratospheric aerosol layers detected by CALIOP in the two weeks following its first detection of the plume on 10 August were above 30° N (Fig. 18c), at altitudes of 9–14 km (Fig. S3b). The median δ_p^{est} of layers in this altitude range is around 0.052, leading to a sulfate classification rate of 67.6 %. The smoke and ash classification rates are 24.4 % and 5.8 %, respectively. A secondary peak with a smaller number of layers around 16 to 18 km was also detected (Fig. S3b). These layers have a somewhat higher median δ_p^{est} of ~ 0.062 , yielding sulfate, smoke, and ash classification rates of 52.3 %, 37.8 %, and 3.3 % respectively. Figure 19 shows two examples of these higher altitude plumes over the eastern

640

Pacific on 14 and 15 August, having median δ_p^{est} values of 0.161 and 0.109. Due to these elevated depolarization ratios, the dominant classification was smoke, but it is likely that these are mixtures of ash and sulfate, with the ash component larger in the 14 August observation. All told, the time series of stratospheric aerosol classification shows an appearance of smoke classifications coincident in time with the appearance of sulfate classifications (Fig. 18d). The altitudes of layers given these classifications are also the same (Fig. S3b), suggesting these plumes are from the same event. Taken together with broader δ_p^{est} distribution compared to the 2011 Nabro eruption (Fig. 16) these smoke classifications could indicate mixtures of ash and sulfate.



650

Figure 19. Possible ash-sulfate mixtures from the 2008 Kasatochi eruption; 532 nm volume depolarization ratio and V4.5 aerosol type classifications from the level 2 aerosol profile product on (a, b) 14 August 2008 at ~11:30 UTC, and (c, d) 15 August 2008 at ~10:30 UTC. Estimated particulate depolarization ratios for select layers are indicated by callouts in upper panels. Inset maps show CALIOP ground track.

655 5.3 Smoke dominated events

In recent years, two major wildfire events demonstrated the massive influence pyroCb activity can have on stratospheric aerosol loading. On 12 August 2017, a series of pyroCbs occurred in northern Washington state, United States and British Columbia, Canada. Dubbed the “Pacific Northwest (PNW) event”, Peterson et al. (2018) estimated that 0.1–0.3 Tg of aerosol mass was injected into the stratosphere on this day. CALIPSO initially measured the plume at 12–14 km on 14 August over northeastern Canada. Several lidar systems over Europe detected the plume over Europe by as early as 10 days later at altitudes spanning 15 to 20 km (Ansmann et al., 2018; Haarig et al., 2018; Khaykin et al., 2018; Hu et al., 2019).

660

Analysis of CALIPSO observations by Khaykin et al. (2018) show the smoke plume had circumnavigated the globe by 30 August, affecting the stratosphere in the entire northern hemisphere above 30° N.

During the first two weeks of the PNW event, the CALIOP median δ_p^{est} was 0.157 for all stratospheric aerosol layers detected in the northern hemisphere. Consequently, for night & day, 76.6 % were classified as smoke, with 9.1 % and 10.4 % misclassified as sulfate and ash, respectively. During this timeframe, these stratospheric smoke layers were primarily detected over northeast Canada, across the north Atlantic, and well into northern Asia (Fig. 20a) at altitudes of 9 to 19 km (Fig. S4a). The time series of smoke layers classification shows a maximum 10–15 days after the initial event (Fig. 20b). Ash misclassifications primarily occur in daytime orbits where additional solar noise broadens the variability of δ_p^{est} , more than doubling the MAD (Fig. 21a). As a result, 22.1 % ash misclassifications occur at day compared to 1.8 % at night. Smoke classifications fare much better at night, at a frequency of 90.1 %.




Figure 20. Same as Fig. 15, but for smoke-dominated events. PyroCb locations denoted by red ovals. Approximate area affected by SAA denoted by yellow polygon.

A second major wildfire event occurred just 2.5 years later in southeastern Australia. From 29 December 2019–4 January 2020, a series of massive pyroCbs injected smoke as high as 16 km (Kablick et al., 2020). Dubbed the 2019/2020 Australian New Year (ANY) event, preliminary estimates of the injected aerosol mass are even larger than the PNW event, ranging from 0.2–0.9 Tg (Peterson et al., 2019; Khaykin et al., 2020). The smoke plumes primarily traveled eastward during the first month, ultimately ascending to heights of over 30 km in February as the smoke absorbed solar radiation, heating the surrounding air and affecting atmospheric dynamics locally (Allen et al., 2020; Kablick et al., 2020). Depolarization of the pyroCb plume from the ANY event was also elevated relative to tropospheric smoke. A Raman lidar in Punta Arenas, Chile

measured 532 nm depolarization ratios of 0.14–0.22 during January 2020 (Ohneiser et al., 2020) with indications of an
 685 increase in depolarization with time (Christian et al., 2020).




Figure 21. Night and daytime estimated particulate depolarization ratio distributions for all stratospheric aerosol layers detected during the first two weeks after (a) the PNW event (northern hemisphere) and (b) the ANY event (southern hemisphere). Low- γ' , PSA, and layers with $|\text{CAD score}| < 20$ are excluded.
 690

During the first two weeks of January 2020, CALIOP detected stratospheric smoke layers primarily over the southern Pacific Ocean yet spanning all longitudes (Fig. 20c) at altitudes of 11 to 22 km (Fig. S4b). Median δ_p^{est} was 0.125 for stratospheric aerosol layers in the southern hemisphere, somewhat smaller than the Raman lidar measurements in Chile.
 695 However, the distributions of δ_p^{est} are quite broad, in particular during daytime (Fig. 21b). Owing to the elevated values of δ_p^{est} , the night & day stratospheric smoke classification frequency was 73.1 %, with misclassification frequencies of 15.5 % and 9.2 % for sulfate and ash, respectively. As with the PNW event, the daytime δ_p^{est} distribution was broader compared to night with a strong skew toward larger values. Consequently, ash misclassifications for the ANY event are more frequent in the daytime, at a rate of 26.9 % compared to 0.6 % at night. The most influential factor driving the broader daytime δ_p^{est}
 700 distribution is sunlight reflecting from high albedo targets at lower altitudes such as stratocumulus in the planetary boundary layer and snow-covered surfaces (e.g., Antarctica in the ANY event). This reflected sunlight enhances noise throughout the profile overhead, thereby increasing the variability of depolarization ratio measurements. The nighttime distributions of δ_p^{est} in Fig. 21 are expected to more closely resemble natural variability, the cause of which is an active area of research (Haarig et al., 2018).

Returning to the geographic distribution of smoke from the ANY event, the map in Fig. 20c shows during the first two weeks, most smoke layers are detected over the southern Pacific Ocean as far north as the equator, though most are detected south of 30° S at all longitudes. Note that the majority of layers detected in the SAA are excluded from our analysis by the minimum laser energy requirement (Sect. 2) we impose to avoid the detrimental influence of low laser energy shots
 705

that are prevalent in this region since mid-2016 (CALIPSO Data Advisory Page, 2018). The maximum number of smoke
 710 classifications occurs during the first 4 weeks, with an increase in unclassified low- γ' layers (Fig. 20d). These low- γ' layers
 become the dominant classification in early February 2020. However, the depolarization ratios remained notable. Figure 22
 shows the coherent “bubble” of smoke southwest of the southern tip of Chile on 31 January that was examined by Allen et
 al. (2020) and Khaykin et al. (2020). They found this bubble rose to 35 km in subsequent months due to dynamics associated
 with absorption of solar radiation. The CALIOP CAD algorithm struggled with this scene, classifying much of the feature as
 715 ice cloud due to the elevated depolarization and spread in moderate color ratio values (median \pm MAD of 0.44 ± 0.23). The
 CAD probability density functions have a fair amount of overlap in the color ratio dimension for features with elevated
 depolarization ratios at high altitudes (Liu et al., 2018), thereby contributing to these CAD errors. However, most of the
 layers classified as aerosol in V4.5 are correctly given the smoke subtype (65 %), whereas ash was the dominant subtype
 given in V4.2, with 8 % smoke.

720





Figure 22. PyroCb smoke plume from ANY event, 31 January 2020 at ~6:30 UTC: (a) 532 nm total attenuated backscatter, (b) 532 nm
 volume depolarization ratio, and (c) V4.5 aerosol subtype and cloud classification from the level 2 aerosol profile product. Inset map
 shows CALIOP ground track.

725 5.4 Unclassified layers and false-positive feature detections

For features identified as stratospheric aerosols by the CALIOP CAD algorithm, the frequency of unclassified layers is
 bound by the low- γ' threshold at the high end and by the feature detection sensitivity at the low end. Regardless of the actual
 aerosol type, all stratospheric aerosol layers can be assigned this classification because eventually, due to sedimentation and
 diffusion, their concentrations in the atmosphere will decline until they are no longer detectable by the CALIOP feature
 730 finder. By design, the lowest-quartile γ'_{532} metric causes 25 % of stratospheric aerosol layers in the CALIOP data record to
 be unclassified on average (Sect. 4.4). Though the number of unclassified stratospheric aerosols peaks legitimately during
 major events, there is a “background” number reported during quiescent periods associated with false layer detections (Fig.

23). As a first-order estimate, during the year 2013 when there were no major stratospheric aerosol injections, suspected false-positive feature detections (any layer detected above 20 km, excluding PSA) occurred in 0.4 % of profiles at night and 1 % at day, globally. These layers are primarily located inside the SAA. Outside the SAA, a relatively small number of layers are randomly distributed over the globe (Fig. S5). Because they do not occur within specific latitude bands, as would legitimate layers associated with specific events, they are likely due to false layer detections caused by radiation-induced current spikes in the 532 nm channel detectors (Hunt et al., 2009; e.g., Fig. S6) or enhanced background noise from sunlight reflecting off underlying clouds in the daytime (e.g., Fig. S7). The aerosol subtypes of these false layer detections at night are 54 % ash, 30 % sulfate, 5 % smoke, and 11 % unclassified. During the day, these frequencies change to 84 % ash, <1 % sulfate, 3 % smoke, and 13 % unclassified. The propensity for ash classification is due to excess solar background noise broadening the distribution of depolarization values to create artificially high values and from cases of radiation-induced current spikes which only affect the 532 nm perpendicular channel, but not the parallel channel. False layer detections above 20 km have characteristically low CAD scores: 97 % have $|CAD| < 20$ indicating no confidence in cloud-aerosol discrimination accuracy, which is expected since these layers are caused by noise excursions, and being neither aerosol nor cloud, receive a low CAD score. They can be readily rejected using the CAD score as we have done throughout this paper.



750 **Figure 23.** Number of unclassified layers in the northern (blue) and southern (red) hemisphere, excluding the SAA and layers with $|CAD$ score < 20 . Computed using V4.2 integrated attenuated backscatter with V4.5 low- γ' thresholds, so the figure serves as a close approximation to V4.5.

6. Conclusion

The stratospheric aerosol subtyping algorithm has been updated for the V4.5 release of the CALIOP level 2 data products. Following the previous V4.2 release, it became clear that several aspects of the newly introduced stratospheric aerosol subtyping algorithm needed further refinement. This paper describes the changes the CALIPSO project has implemented to improve aerosol subtyping in the stratosphere and characterized the performance of the refined algorithm based on well-755 documented events. The changes include: removing the use of integrated attenuated backscatter color ratio, so that the

algorithm now discriminates between volcanic ash, sulfate, and smoke solely based on depolarization; increasing the depolarization threshold to discriminate between volcanic ash and smoke; separating the V4.2 sulfate/other subtype into sulfate and unclassified subtypes; lowering the low- γ' threshold for identifying weakly scattering, unclassified, layers; and increasing the 532 nm lidar ratio for volcanic ash to a value consistent with the current state of knowledge. As a consequence, these changes improve the discrimination capability between volcanic ash and smoke by better accounting for the depolarizing nature of smoke often observed for layers associated with pyroCb activity. Sulfate classifications now solely identify layers having low depolarization ratios, a characteristic of sulfate aerosol. Our analysis also postulates that volcanic layers classified as smoke can indicate mixtures of sulfate and ash. Finally, weakly backscattering features have been relegated to a new unclassified subtype for which the signal-to-noise ratio is considered insufficient to reliably discern the true type.

The performance of the revised algorithm is very good for volcanic ash layers, with 84 % correctly classified during the ash-dominated Puyehue-Cordón Caulle eruption of 2011. This is no surprise, given the strongly depolarizing nature of volcanic ash. Sulfate classifications are also dominant for events having a strong sulfate component. However, the interpretation of the CALIOP stratospheric aerosol classification requires some extra care for sulfate-dominated scenes with some ash component. Sulfate/ash mixtures are misclassified as smoke for nearly one-third of these layers and there exists the possibility for legitimate smoke layers to be misclassified as sulfate due to the overlap in the depolarization ratio distributions for these two types (a combination of natural variability and measurement noise). Smoke classification performance for events dominated by pyroCb activity was also very good, with most layers classified as smoke. There remains a moderate number of smoke layers that are misclassified as sulfate and ash. In particular, misclassification frequencies of smoke as volcanic ash are substantially higher at day than at night (~27 % vs. 1 %, respectively), due to reflected sunlight from lower altitude high-albedo features that adds substantial noise to the column, broadening the distribution of depolarization ratios. Additionally, any smoke transported from the troposphere into the UTLS by self-lofting rather than pyroCb activity will likely be misclassified as sulfate due to their similarly low values of depolarization. Researchers should be aware of these potential artifacts when performing automated analyses with CALIOP V4.5 level 2 data.

It is important to recognize that although the aerosol subtyping algorithm performs very well for ash, sulfate, and depolarizing smoke in the stratosphere, aerosol subtyping is less satisfactory for these same aerosol types below the tropopause largely because no attempt is made to identify them in the troposphere. There, volcanic ash will inexorably be misclassified as dust, depolarizing smoke mostly misclassified as polluted dust, and volcanic sulfate misclassified as elevated smoke (Kim et. al, 2018). These misclassifications occur because it is difficult to discriminate among these aerosol types in a robust automated manner given the limited number of CALIOP observables. The critical information for the CALIOP stratospheric aerosol subtyping algorithm is the high altitude of the tropopause, which most often rules out the possibility of all but a few subtypes. More sophisticated instrumentation will improve discrimination capability in the troposphere, such as high spectral resolution lidar with depolarization sensitivity at 355 nm, 532 nm, and 1064 nm (e.g., as in

Burton et al., 2015) or combined lidar plus passive instrument retrievals. Additionally, combining SO₂ and CO measurements from other sensors could help differentiate between ash/sulfate mixtures and smoke. Given the aviation hazards posed by volcanic ash and the climate implications of sulfate and stratospheric smoke injections, space-based lidar retrievals stand to provide valuable vertically resolved information to disaster response agencies and climate modelers. Our hope is that this work provides a meaningful steppingstone toward more sophisticated solutions in future missions.

Data availability

CALIOP data are available through the NASA Langley Research Center Atmospheric Science Data Center (ASDC), <https://asdc.larc.nasa.gov/> (last access: 7 January 2022): V4.1 level 1B (NASA/LARC/SD/ASDC 2016b); V4.2 level 2 aerosol layer, aerosol profile, and vertical feature mask products (NASA/LARC/SD/ASDC 2018a, 2018b, 2018c); V4.5 level 1B (NASA/LARC/SD/ASDC 2022a). The V4.5 level 2 aerosol layer, aerosol profile, and vertical feature mask products will be released in early 2023 and made available through the ASDC.

Author contributions

The CALIPSO stratospheric aerosol subtyping algorithm described in this paper was conceived by JT, MV, JK, AO, and DW, with contributions from the CALIPSO mission team, JV, MP, and Lamont Poole. Formal analysis and visualizations were prepared by JT, except for Sect. 4.5 which was prepared by JK. Writing was led by JT with extensive contributions by MV, JK, MK, and JV. All V4.5 pre-release data was generated by BG. CALIPSO V4.5 level 2 software modifications for the stratospheric aerosol subtyping algorithm were performed by BM with support from the CALIPSO Data Management Team.

Acknowledgements

Volcano geographic locations are acquired from the Smithsonian Institution National Museum of Natural History Global Volcanism Program <https://volcano.si.edu/>. We sincerely thank the anonymous referees for their thorough assessment which helped improved the clarity of the manuscript.

References

Allen, D. R., Fromm, M. D., Kablick III, G. P., and Nedoluha, G. E.: Smoke with Induced Rotation and Lofting (SWIRL) in the Stratosphere, *J. Atmos. Sci.*, 77, 4297-4316, <https://doi.org/10.1175/JAS-D-20-0131.1>, 2020.

Andersson, S. M., Martinsson, B. G., Friberg, J., Brenninkmeijer, C. A. M., Rauthe-Schöch, A., Hermann, M., van Velthoven, P. F. J., and Zahn, A.: Composition and evolution of volcanic aerosol from eruptions of Kasatochi, Sarychev and
820 Eyjafjallajökull in 2008–2010 based on CARIBIC observations, *Atmos. Chem. Phys.*, 13, 1781–1796, <https://doi.org/10.5194/acp-13-1781-2013>, 2013.

Ansmann, A., Tesche, M., Groß, S., Freudenthaler, V., Seifert, P., Hiebsch, A., Schmidt, J., Wandinger, U., Mattis, I., Müller, D., and Wiegner, M.: The 16 April 2010 major volcanic ash plume over Central Europe: EARLINET lidar and
825 AERONET photometer observations at Leipzig and Munich, Germany. *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010GL043809>, 2010.

Ansmann, A., Baars, H., Chudnovsky, A., Mattis, I., Veselovskii, I., Haarig, M., Seifert, P., Engelmann, R., and Wandinger, U.: Extreme levels of Canadian wildfire smoke in the stratosphere over central Europe on 21–22 August 2017, *Atmos.*
830 *Chem. Phys.*, 18, 11831–11845, <https://doi.org/10.5194/acp-18-11831-2018>, 2018.

Ansmann, A., Ohneiser, K., Chudnovsky, A., Baars, H., and Engelmann, R.: CALIPSO aerosol-typing scheme misclassified stratospheric fire smoke: case study from the 2019 Siberian wildfire season, *Front. Environ. Sci.*, 21, 769852, <https://doi.org/10.3389/fenvs.2021.769852>, 2021.

835

Bègue, N., Vignelles, D., Berthet, G., Portafaix, T., Payen, G., Jégou, F., Benchérif, H., Jumelet, J., Vernier, J.-P., Lurton, T., Renard, J.-B., Clarisse, L., Duverger, V., Posny, F., Metzger, J.-M., and Godin-Beekmann, S.: Long-range transport of stratospheric aerosols in the Southern Hemisphere following the 2015 Calbuco eruption, *Atmos. Chem. Phys.*, 17, 15019–15036, <https://doi.org/10.5194/acp-17-15019-2017>, 2017.

840

Bignami, C., Corradini, S., Merucci, L., de Michele, M., Raucoules, D., de Astis, G., Stramondo, S., and Piedra, J.: Multisensor satellite monitoring of the 2011 Puyehue-Cordón Caulle eruption, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 7, 2786–2796, <https://doi.org/10.1109/JSTARS.2014.2320638>, 2014.

845 Boone, C., Bernath, P. F., Labelle, K., Crouse, J.: Stratospheric Aerosol Composition Observed by the Atmospheric Chemistry Experiment Following the 2019 Raikoke Eruption, *J. Geophys. Res.-Atmos.*, 127, e2022JD036600, <https://doi.org/10.1029/2022JD036600>, 2022.

Burton, S. P., Hair, J. W., Kahnert, M., Ferrare, R. A., Hostetler, C. A., Cook, A. L., Harper, D. B., Berkoff, T. A., Seaman,
850 S. T., Collins, J. E., Fenn, M. A., and Rogers, R. R.: Observations of the spectral dependence of linear particle depolarization

ratio of aerosols using NASA Langley airborne High Spectral Resolution Lidar, *Atmos. Chem. Phys.*, 15, 13453–13473, <https://doi.org/10.5194/acp-15-13453-2015>, 2015.

855 Cairo, F., Di Donfrancesco, G., Adriani, A., Pulvirenti, L., and Fierli, F.: Comparison of various linear depolarization parameters measured by lidar, *Appl. Optics*, 38, 4425–4432, <https://doi.org/10.1364/AO.38.004425>, 1999.

CALIPSO Data Advisory Page,

https://www-calipso.larc.nasa.gov/resources/calipso_users_guide/advisory.php, last access: 3 October 2022, 2018.

860 CALIPSO Lidar Level 1 V4.51 Data Quality Statement,

https://www-calipso.larc.nasa.gov/resources/calipso_users_guide/qs/cal_lid_l1_v4-51_qs.php, last access: 3 October 2022, 2022.

865 Christian, K., Yorks, J., and Das, S.: Differences in the Evolution of Pyrocumulonimbus and Volcanic Stratospheric Plumes as Observed by CATS and CALIOP Space-Based Lidars, *Atmosphere*, 11, 1035, <https://doi.org/10.3390/atmos11101035>, 2020.

Clarisse, L., Coheur, P.-F., Theys, N., Hurtmans, D., and Clerbaux, C.: The 2011 Nabro eruption, a SO₂ plume height analysis using IASI measurements, *Atmos. Chem. Phys.*, 14, 3095–3111, <https://doi.org/10.5194/acp-14-3095-2014>, 2014.

870

Corradini, S., Merucci, L., Prata, A. J., and Piscini, A.: Volcanic ash and SO₂ in the 2008 Kasatochi eruption: Retrievals comparison from different IR satellite sensors, *J. Geophys. Res.*, 115, D00L21, <https://doi:10.1029/2009JD013634>, 2010.

875 de Laat, A. T. J., Stein Zweers, D. C., Boers, R., and Tuinder, O. N. E.: A solar escalator: Observational evidence of the self-lifting of smoke and aerosols by absorption of solar radiation in the February 2009 Australian Black Saturday plume, *J. Geophys. Res.*, 117, D04204, <https://doi.org/10.1029/2011JD017016>, 2012.

880 Dirksen, R. J., Boersma, K. F., de Laat, A. T. J., Stammes, P., van der Werf, G. R., Val Martin, M., , and Kelder, H. M.: An aerosol boomerang: rapid around-the-world transport of smoke from the December 2006 Australian forest fires observed from space, *J. Geophys. Res.*, 114, D21201, <https://doi:10.1029/2009JD012360>, 2009.

Fairlie, T. D., Vernier, J.-P., Natarajan, M., and Bedka, K. M.: Dispersion of the Nabro volcanic plume and its relation to the Asian summer monsoon, *Atmos. Chem. Phys.*, 14, 7045–7057, <https://doi:10.5194/acp-14-7045-2014>, 2014.

- 885 Fromm, M., Kablick III, G., Nedoluha, G., Carboni, E., Grainger, R., Campbell, J., and Lewis, J.: Correcting the record of volcanic stratospheric aerosol impact: Nabro and Sarychev Peak, *J. Geophys. Res.-Atmos.*, 119, 10343–10364, <https://doi.org/10.1002/2014JD021507>, 2014.
- Fromm, M., Lindsey, D. T., Servranckx, R., Yue, G., Trickl, T., Sica, R., Doucet, P., and Godin-Beekmann, S. E.: The
890 untold story of pyrocumulonimbus, *Bull. Am. Meteorol. Soc.*, 91, 1193–1209, <https://doi.org/10.1175/2010bams3004.1>, 2010.
- Fromm, M., Peterson D., and Di Girolamo, L.: The primary convective pathway for observed wildfire emissions in the upper troposphere and lower stratosphere: a targeted reinterpretation, *J. Geophys. Res. Atmos.*, 124, 13254–13272, <https://doi.org/10.1029/2019JD031006>, 2019.
- 895 Gelaro, R., McCarty, W., Suarez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., Da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Markova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. C., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications,
900 Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Gialitaki, A., Tsekeri, A., Amiridis, V., Ceolato, R., Paulien, L., Kampouri, A., Gkikas, A., Solomos, S., Marinou, E., Haarig, M., Baars, H., Ansmann, A., Lapyonok, T., Lopatin, A., Dubovik, O., Groß, S., Wirth, M., Tschla, M., Tsikoudi, I., and Balis, D.: Is the near-spherical shape the “new black” for smoke?, *Atmos. Chem. Phys.*, 20, 14005–14021,
905 <https://doi.org/10.5194/acp-20-14005-2020>, 2020.
- Groß, S., Freudenthaler, V., Wiegner, M., Gasteiger, J., Geiß, A., and Schnell, F.: Dual-wavelength linear depolarization ratio of volcanic aerosols: Lidar measurements of the Eyjafjallajökull plume over Maisach, Germany, *Atmos. Environ.*, 48, 85–96, <https://doi.org/10.1016/j.atmosenv.2011.06.017>, 2012.
910
- Guffanti, M., Schneider, D. J., Wallace, K. L., Hall, T., Bensimon, D. R., and Salinas, L. J.: Aviation response to a widely dispersed volcanic ash and gas cloud from the August 2008 eruption of Kasatochi, Alaska, USA, *J. Geophys. Res.*, 115, D00L19, <https://doi.org/10.1029/2010JD013868>, 2010.
- 915 Haarig, M., Ansmann, A., Baars, H., Jimenez, C., Veselovskii, I., Engelmann, R., and Althausen, D.: Depolarization and lidar ratios at 355, 532, and 1064 nm and microphysical properties of aged tropospheric and stratospheric Canadian wildfire smoke, *Atmos. Chem. Phys.*, 18, 11847–11861, <https://doi.org/10.5194/acp-18-11847-2018>, 2018.

- Höpfner, M., Ungermann, J., Borrmann, S., Wagner, R., Spang, R., Riese, M., Stiller, G., Appel, O., Batenburg, A. M.,
920 Bucci, S., Cairo, F., Dragoneas, A., Friedl-Vallon, F., Hünig, A., Johansson, S., Krasauskas, L., Legras, B., Leisner, T.,
Mahnke, C., Möhler, O., Molleker, S., Müller, R., Neubert, T., Orphal, J., Preusse, P., Rex, M., Saathoff, H., Stroh, F.,
Weigel, R., and Wohltmann, I.: Ammonium nitrate particles formed in upper troposphere from ground ammonia sources
during Asian monsoons, *Nat. Geosci.*, 12, 608–612, <https://doi.org/10.1038/s41561-019-0385-8>, 2019.
- 925 Hostetler, C. A., Liu, Z., Reagan, J., Vaughan, M., Winker, D., Osborn, M., Hunt, W. H., Powell, K. A., and Trepte, C.:
CALIOP Algorithm Theoretical Basis Document, Calibration and Level 1 Data Products, PC-SCI-201, NASA Langley
Research Center, Hampton, VA 23681, 66 pp., http://www-calipso.larc.nasa.gov/resources/project_documentation.php (last
access: 3 September 2021), 2006.
- 930 Hu, Q., Goloub, P., Veselovskii, I., Bravo-Aranda, J.-A., Popovici, I. E., Podvin, T., Haeffelin, M., Lopatin, A., Dubovik, O.,
Pietras, C., Huang, X., Torres, B., and Chen, C.: Long-range-transported Canadian smoke plumes in the lower stratosphere
over northern France, *Atmos. Chem. Phys.*, 19, 1173–1193, <https://doi.org/10.5194/acp-19-1173-2019>, 2019.
- Hunt, W., Winker, D., Vaughan, M., Powell, K., Lucker, P., and Weimer, C.: CALIPSO lidar description and performance
935 assessment, *J. Atmos. Ocean. Tech.*, 26, 1214–1228, <https://doi.org/10.1175/2009JTECHA1223.1>, 2009.
- Kablick, G. P., Fromm, M. D., Miller, S. D., Partain, P., Peterson, D., Lee, S., Zhang, Y., Lambert, A., Li, Z.: The Great
Slave Lake pyroCb of 5 August 2014: Observations, simulations, comparisons with regular convection, and impact on UTLS
water vapor, *J. Geophys. Res. Atmos.*, 123, 12,332–12,352, <https://doi.org/10.1029/2018JD028965>, 2018.
940
- Kablick, G. P., Allen, D. R., Fromm, M. D., and Nedoluha, G. E.: Australian pyroCb smoke generates synoptic-scale
stratospheric anticyclones, *Geophys. Res. Lett.*, 47, e2020GL088101, <https://doi.org/10.1029/2020GL088101>, 2020.
- Kar, J., Vaughan, M. A., Lee, K.-P., Tackett, J. L., Avery, M. A., Garnier, A., Getzewich, B. J., Hunt, W. H., Josset, D., Liu,
945 Z., Lucker, P. L., Magill, B., Omar, A. H., Pelon, J., Rogers, R. R., Toth, T. D., Trepte, C. R., Vernier, J.-P., Winker, D. M.,
and Young, S. A.: CALIPSO lidar calibration at 532 nm: version 4 nighttime algorithm, *Atmos. Meas. Tech.*, 11, 1459–
1479, <https://doi.org/10.5194/amt-11-1459-2018>, 2018.
- Khaykin, S. M., Godin-Beekmann, S., Hauchecorne, A., Pelon, J., Ravetta, F., and Keckut, P.: Stratospheric smoke with
950 unprecedentedly high backscatter observed by lidars above southern France, *Geophys. Res. Lett.*, 45, 1639–1646,
<https://doi.org/10.1002/2017GL076763>, 2018.

- 955 Khaykin, S., Legras, B., Bucci, S., Sellitto, P., Isaksen, L., Tence, F., Bekki, S., Bourassa, A. E., Rieger, L. A., Zawada, D., Jumelet, J., and Godin-Beekmann, S.: The 2019/20 Australian wildfires generated a persistent smoke-charged vortex rising up to 35 km altitude, *Commun. Earth Environ.*, 1, 22, <https://doi.org/10.1038/s43247-020-00022-5>, 2020.
- 960 Kim, M.-H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Trepte, C. R., Hu, Y., Liu, Z., Poole, L. R., Pitts, M. C., Kar, J., and Magill, B. E.: The CALIPSO version 4 automated aerosol classification and lidar ratio selection algorithm, *Atmos. Meas. Tech.*, 11, 6107–6135, <https://doi.org/10.5194/amt-11-6107-2018>, 2018.
- Klekociuk, A. R., Ottaway, D. J., MacKinnon, A. D., Reid, I. M., Twigger, L. V. and Alexander, S. P.: Australian Lidar Measurements of Aerosol Layers Associated with the 2015 Calbuco Eruption, *Atmosphere*, 11, 124, <https://doi.org/10.3390/atmos11020124>, 2020.
- 965 Klüser, L., Erbertseder, T., and Meyer-Arne, J.: Observation of volcanic ash from Puyehue–Cordón Caulle with IASI, *Atmos. Meas. Tech.*, 6, 35–46, <https://doi.org/10.5194/amt-6-35-2013>, 2013.
- 970 Kokkalis, P., Papayannis, A., Amiridis, V., Mamouri, R. E., Veselovskii, I., Kolgotin, A., Tsaknakis, G., Kristiansen, N. I., Stohl, A., and Mona, L.: Optical, microphysical, mass and geometrical properties of aged volcanic particles observed over Athens, Greece, during the Eyjafjallajökull eruption in April 2010 through synergy of Raman lidar and sunphotometer measurements, *Atmos. Chem. Phys.*, 13, 9303–9320, <https://doi.org/10.5194/acp-13-9303-2013>, 2013.
- 975 Kremser, S., Thomason, L. W., Hobe, M., Hermann, M., Deshler, T., Timmreck, C., Toohey, M., Stenke, A., Schwarz, J. P., Weigel, R., Fueglistaler, S., Prata, F. J., Vernier, J.-P., Schlager, H., Barnes, E. J., Antuna-Marrero, J.-C., Fairlie, D., Palm, M., Mahieu, E., Notholt, J., Rex, M., Bingen, C., Vanhellefont, F., Bourassa, A., Plane, J. M. C., Klocke, D., Carn, S. A., Clarisse, L., Trickl, T., Neely, R. D., James, A., Rieger, L., Wilson, C. J., and Meland, B.: Stratospheric aerosol – Observations, processes, and impact on climate, *Rev. Geophys.*, 54, 278–335, <https://doi.org/10.1002/2015RG000511>, 2016.
- 980 Kristiansen, N. I., Stohl, A., Prata, A. J., Richter, A., Eckhardt, S., Seibert, P., Hoffmann, A., Ritter, C., Bitar, L., Duck, T. J., and Stebel, K.: Remote sensing and inverse transport modelling of the Kasatochi eruption sulphur dioxide cloud, *J. Geophys. Res.*, 115, D00L16, <https://doi.org/10.1029/2009JD013286>, 2010.
- Krotkov, N. A., Schoeberl, M. R., Morris, G. A., Carn, S., and Yang, K.: Dispersion and lifetime of the SO₂ cloud from the August 2008 Kasatochi eruption, *J. Geophys. Res.*, 115, D00L20, <https://doi.org/10.1029/2010JD013984>, 2010.
- 985

- Langmann, B., Zaksek, K., and Hort, M.: Atmospheric distribution and removal of volcanic ash after the eruption of Kasatochi volcano: A regional model study, *J. Geophys. Res.*, 115, D00L06, <https://doi.org/10.1029/2009JD013298>, 2010.
- 990 Liu, Z., Kar, J., Zeng, S., Tackett, J., Vaughan, M., Avery, M., Pelon, J., Getzewich, B., Lee, K.-P., Magill, B., Omar, A., Lucker, P., Trepte, C., and Winker, D.: Discriminating between clouds and aerosols in the CALIOP version 4.1 data products, *Atmos. Meas. Tech.*, 12, 703–734, <https://doi.org/10.5194/amt-12-703-2019>, 2019.
- Lopes, F. J. S., Silva, J. J., Antuña Marrero, J. C., Taha, G. and Landulfo, E.: Synergetic Aerosol Layer Observation After the 2015 Calbuco Volcanic Eruption Event, *Remote Sensing*, 11, 195, <https://doi.org/10.3390/rs11020195>, 2019.
- 995 Maes, K., Vandenbussche, S., Klüser, L., Kumps, N., and De Mazière, M.: Vertical Profiling of Volcanic Ash from the 2011 Puyehue Cordón Caulle Eruption Using IASI, *Remote Sensing*, 8, 103, <https://doi.org/10.3390/rs8020103>, 2016.
- Martinsson, B. G., Brenninkmeijer, C. A. M., Carn, S. A., Hermann, M., Heue, K.-P., Velthoven, P. F. J. V., and Zahn, A.: 1000 Influence of the 2008 Kasatochi volcanic eruption on sulfurous and carbonaceous aerosol constituents in the lower stratosphere, *Geophys. Res. Lett.*, 36, L12813, <https://doi.org/10.1029/2009GL038735>, 2009.
- Marzano, F., Corradini, S., Mereu, L., Kylling, A., Montopoli, M., Cimini, D., Merucci, L., and Stelitano, D.: Multisatellite Multisensor Observations of a Sub-Plinian Volcanic Eruption: The 2015 Calbuco Explosive Event in Chile, *IEEE 1005 Transactions on Geoscience and Remote Sensing*, 56, 2597–2612, <https://doi.org/10.1109/TGRS.2017.2769003>, 2018.
- Mona, L., Amodeo, A., D'Amico, G., Giunta, A., Madonna, F., and Pappalardo, G.: Multi-wavelength Raman lidar observations of the Eyjafjallajökull volcanic cloud over Potenza, southern Italy, *Atmos. Chem. Phys.*, 12, 2229–2244, <https://doi.org/10.5194/acp-12-2229-2012>, 2012.
- 1010 NASA/LARC/SD/ASDC: CALIPSO Lidar Level 2 Polar Stratospheric Clouds (PSC) Mask, Provisional V1-11. NASA Langley Atmospheric Science Data Center DAAC [Data set], https://doi.org/10.5067/CALIOP/CALIPSO/CAL_LID_L2_PSCMask-Prov-V1-11, 2016a.
- 1015 NASA/LARC/SD/ASDC: CALIPSO Lidar Level 1B profile data, V4-10, NASA Langley Atmospheric Science Data Center DAAC [Data set], https://doi.org/10.5067/CALIOP/CALIPSO/LID_L1-STANDARD-V4-10, 2016b.

- 1020 NASA/LARC/SD/ASDC: CALIPSO Lidar Level 2 5 km Aerosol Layer Data, V4-20, NASA Langley Atmospheric Science Data Center DAAC [Data set], https://doi.org/10.5067/CALIOP/CALIPSO/LID_L2_05KMALAY-STANDARD-V4-20, 2018a.
- NASA/LARC/SD/ASDC: CALIPSO Lidar Level 2 Aerosol Profile, V4-20, NASA Langley Atmospheric Science Data Center DAAC [Data set], https://doi.org/10.5067/CALIOP/CALIPSO/LID_L2_05KMAPRO-STANDARD-V4-20, 2018b.
- 1025 NASA/LARC/SD/ASDC: CALIPSO Lidar Level 2 Vertical Feature Mask (VFM), V4-20, NASA Langley Atmospheric Science Data Center DAAC [Data set], https://doi.org/10.5067/CALIOP/CALIPSO/LID_L2_VFM-STANDARD-V4-20, 2018c.
- NASA/LARC/SD/ASDC: CALIPSO Lidar Level 1B profile data, V4-51, NASA Langley Atmospheric Science Data Center DAAC [Data set], https://doi.org/10.5067/CALIOP/CALIPSO/CAL_LID_L1-Standard-V4-51, 2022a.
- 1030 Noh, Y.M., Dong, H.S., Müller, D.: Variation of the vertical distribution of Nabro volcano aerosol layers in the stratosphere observed by LIDAR, *Atmos. Environ.*, 154, 1-8, <https://doi.org/10.1016/j.atmosenv.2017.01.033>, 2017.
- 1035 Ohneiser, K., Ansmann, A., Baars, H., Seifert, P., Barja, B., Jimenez, C., Radenz, M., Teisseire, A., Floutsi, A., Haarig, M., Foth, A., Chudnovsky, A., Engelmann, R., Zamorano, F., Bühl, J., and Wandinger, U.: Smoke of extreme Australian bushfires observed in the stratosphere over Punta Arenas, Chile, in January 2020: optical thickness, lidar ratios, and depolarization ratios at 355 and 532 nm, *Atmos. Chem. Phys.*, 20, 8003–8015, <https://doi.org/10.5194/acp-20-8003-2020>, 2020.
- 1040 Ohneiser, K., Ansmann, A., Chudnovsky, A., Engelmann, R., Ritter, C., Veselovskii, I., Baars, H., Gebauer, H., Griesche, H., Radenz, M., Hofer, J., Althausen, D., Dahlke, S., and Maturilli, M.: The unexpected smoke layer in the High Arctic winter stratosphere during MOSAiC 2019–2020, *Atmos. Chem. Phys.*, 21, 15783–15808, <https://doi.org/10.5194/acp-21-15783-2021>, 2021.
- 1045 Ohneiser, K., Ansmann, A., Kaifler, B., Chudnovsky, A., Barja, B., Knopf, D. A., Kaifler, N., Baars, H., Seifert, P., Villanueva, D., Jimenez, C., Radenz, M., Engelmann, R., Veselovskii, I., and Zamorano, F.: Australian wildfire smoke in the stratosphere: the decay phase in 2020/2021 and impact on ozone depletion, *Atmos. Chem. Phys.*, 22, 7417–7442, <https://doi.org/10.5194/acp-22-7417-2022>, 2022.
- 1050

- Pardini, F., Burton, M., Arzilli, F., La Spina, G., and Polacci, M.: SO₂ emissions, plume heights and magmatic processes inferred from satellite data: The 2015 Calbuco eruptions, *J. Volcanol. Geoth. Res.*, 361, 12–24, <https://doi.org/10.1016/j.jvolgeores.2018.08.001>, 2018.
- 1055 Peterson, D. A., Campbell, J. R., Hyer, E. J., Fromm, M. D., Kablick, G. P., Cossuth, J. H., and DeLand, M. T.: Wildfire-driven thunderstorms cause a volcano-like stratospheric injection of smoke, *Climate and Atmospheric Science*, 1, 30, <https://doi.org/10.1038/s41612-018-0039-3>, 2018.
- Peterson, D. A., Hyer, E., Campbell, J., Fromm, M., Bennese, C., Berman, M., and Van, T.: Quantifying the impact of
1060 intense pyroconvection on stratospheric aerosol loading. American Geophysical Union 2019 Fall Meeting, San Francisco, CA, 9-13 December 2019, Abstract GC11F-1150, <https://agu.confex.com/agu/fm19/meetingapp.cgi/Paper/510480>, 2019.
- Peterson, D. A., Hyer, E. J., Campbell, J. R., Solbrig, J. E., and Fromm, M. D.: A conceptual model for development of
1065 intense pyrocumulonimbus in western North America, *Mon. Weather Rev.*, 145, 2235–2255, <https://doi.org/10.1175/MWR-D-16-0232.1>, 2017.
- Pitts, M. C., Poole, L. R., Dörnbrack, A., and Thomason, L. W.: The 2009–2010 Arctic polar stratospheric cloud season: a
1070 CALIPSO perspective, *Atmos. Chem. Phys.*, 11, 2161–2177, <https://doi.org/10.5194/acp-11-2161-2011>, 2011.
- Pitts, M. C., Poole, L. R., and Gonzalez, R.: Polar stratospheric cloud climatology based on CALIPSO spaceborne lidar
measurements from 2006 to 2017, *Atmos. Chem. Phys.*, 18, 10881–10913, <https://doi.org/10.5194/acp-18-10881-2018>,
2018.
- 1075 Poole, L. R. and Pitts, M. C.: Polar stratospheric cloud climatology based on Stratospheric Aerosol Measurement II
observations from 1978 to 1989, *J. Geophys. Res.*, 99, 13083–13089, <https://doi.org/10.1029/94JD00411>, 1994.
- Prata, A. J., Gangale, G., Clarisse, L., and Karagulian, F.: Ash and sulfur dioxide in the 2008 eruptions of Okmok and
Kasatochi: Insights from high spectral resolution satellite measurements, *J. Geophys. Res.*, 115, D00L18,
1080 <https://doi.org/10.1029/2009JD013556>, 2010.
- Prata, A. J.: Infrared radiative transfer calculations for volcanic ash clouds, *Geophys. Res. Lett.*, 16, 1293–1296, 1989

- 1085 Prata, A. T., Young, S. A., Siems, S. T., and Manton, M. J.: Lidar ratios of stratospheric volcanic ash and sulfate aerosols retrieved from CALIOP measurements, *Atmos. Chem. Phys.*, 17, 8599–8618, <https://doi.org/10.5194/acp-17-8599-2017>, 2017.
- 1090 Pueschel, R. F.: Stratospheric aerosols: Formation, properties, effects, *Journal of Aerosol Science*, 27, 383-402, [https://doi.org/10.1016/0021-8502\(95\)00557-9](https://doi.org/10.1016/0021-8502(95)00557-9), 1996.
- Rosen, J. M., Kjöme, N. T., Larsen, N., Knudsen, B. M., Kyrö, E., Kivi, R., Karhu, J., Neuber, R., and Beninga, I.: Polar stratospheric cloud threshold temperatures in the 1995–1996 arctic vortex, *J. Geophys. Res.*, 102, 28195–28202, <https://doi.org/10.1029/97JD02701>, 1997.
- 1095 Ryan, R., Vaughan, M., Rodier, S. D., Getzewich, B. J., and Winker, D. M.: Column Optical Depths (COD) Derived from CALIOP Ocean Surface Returns, 30th International Laser Radar Conference, Virtual, 26 June-1 July 2022, Paper S01_P09_Ryan, <https://meeting-info.org/wp-content/uploads/elementor/forms/6299271aeacb2.pptx?6bfec1&6bfec1>, last access 4 October 2022, 2022.
- 1100 Sayer, A. M., Hsu, N. C., Eck, T. F., Smirnov, A., and Holben, B. N.: AERONET-based models of smoke-dominated aerosol near source regions and transported over oceans, and implications for satellite retrievals of aerosol optical depth, *Atmos. Chem. Phys.*, 14, 11493–11523, <https://doi.org/10.5194/acp-14-11493-2014>, 2014.
- 1105 Sicard, M., Granados-Muñoz, M. J., Alados-Arboledas, L., Barragán, R., Bedoya-Velásquez, A. E., Benavent-Oltra, J. A., Bortoli, D., Comerón, A., Córdoba-Jabonero, C., Costa, M. J., del Águila, A., Fernández, A. J., Guerrero-Rascado, J. L., Jorba, O., Molero, F., Muñoz-Porcar, C., Ortiz-Amezcu, P., Papagiannopoulos, N., Potes, M., Pujadas, M., Rocadenbosch, F., Rodríguez-Gomez, A., Román, R., Salgado, R., Salgueiro, V., Sola, Y., and Yela, M.: Ground/space, passive/active remote sensing observations coupled with particle dispersion modelling to understand the inter-continental transport of wildfire smoke plumes, *Remote Sens Environ.*, 232, 111294, <https://doi.org/10.1016/j.rse.2019.111294>, 2019.
- 1110 Siddaway, J. M. and Petelina, S. V.: Transport and evolution of the 2009 Australian Black Saturday bushfire smoke in the lower stratosphere observed by OSIRIS on Odin, *J. Geophys. Res.*, 116, D06203, <https://doi.org/10.1029/2010JD015162>, 2011.
- 1115 Stone, K. A., Solomon, S., Kinnison, D. E., Pitts, M. C., Poole, L. R., Mills, M. J., Schmidt, A., Neely III, R. R., Ivy, D., Schwartz, M. J., Vernier, J. P., Johnson, B. J., Tully, M. B., Klekocius, A. R., König-Langlo, G., and Hagiya, S.: Observing

- the impact of Calbuco volcanic aerosols on south polar ozone depletion in 2015, *J. Geophys. Res.-Atmos.*, 122, 11862–11879, <https://doi.org/10.1002/2017JD026987>, 2017.
- 1120 Tackett, J., Vaughan, M., Lambeth, J., Garnier, A.: Critical Improvements to CALIOP Boundary Layer Cloud-Clearing in Version 4.5, CloudSat/CALIPSO Annual Science Program Review, Fort Collins, CO, 12-14 September 2022, Paper 10 Day 1, <https://sites.google.com/view/ccstm-2022/home>, last access 4 October 2022, 2022.
- 1125 Tackett, J. L., Vaughan, M. A., Lee, K-P. A., Kar, J., and Trepte, C. R.: Improvements in CALIOP Smoke Optical Depth over Clouds, American Meteorological Society 101st Annual Meeting, Virtual, 10-15 January 2021, Paper 381852, <https://ams.confex.com/ams/101ANNUAL/meetingapp.cgi/Paper/381852>, last access: 4 October 2022, 2021.
- 1130 Theys, N., Campion, R., Clarisse, L., Brenot, H., van Gent, J., Dils, B., Corradini, S., Merucci, L., Coheur, P.-F., Van Roozendael, M., Hurtmans, D., Clerbaux, C., Tait, S., and Ferrucci, F.: Volcanic SO₂ fluxes derived from satellite data: a survey using OMI, GOME-2, IASI and MODIS, *Atmos. Chem. Phys.*, 13, 5945–5968, <https://doi.org/10.5194/acp-13-5945-2013>, 2013.
- 1135 Torres, O., Bhartia, P. K., Taha, G., Jethva, H., Das, S., Colarco, P., Krotkov, N., Omar, A., and Ahn, C.: Stratospheric Injection of Massive Smoke Plume from Canadian Boreal Fires in 2017 as seen by DSCOVR-EPIC, CALIOP and OMPS-LP Observations, *J. Geophys. Res.-Atmos.*, 125, e2020JD032579, <https://doi.org/10.1029/2020JD032579>, 2020.
- 1140 Ulke, A. G., Torres Brizuela, M. M., Raga, G. B., and Baumgardner, D.: Aerosol properties and meteorological conditions in the city of Buenos Aires, Argentina, during the resuspension of volcanic ash from the Puyehue-Cordón Caulle eruption, *Nat. Hazards Earth Syst. Sci.*, 16, 2159–2175, <https://doi.org/10.5194/nhess-16-2159-2016>, 2016.
- Vaughan, M. A., Powell, K. A., Winker, D. M., Hostetler, C. A., Kuehn, R. E., Hunt, W. H., Getzewich, B. J., Young, S. A., Liu, Z., and McGill, M. J.: Fully automated detection of cloud and aerosol layers in the CALIPSO lidar measurements, *J. Atmos. Ocean. Tech.*, 26, 2034–2050, <https://doi.org/10.1175/2009JTECHA1228.1>, 2009.
- 1145 Vaughan, M. A., Winker, D. M., and Powell, K. A.: CALIOP Algorithm Theoretical Basis Document Part 2: Feature Detection and Layer Properties Algorithms, available at: https://www-calipso.larc.nasa.gov/resources/pdfs/PC-SCI-202_Part2_rev1x01.pdf, 2005.

- Vernier, J. P., Pommereau, J. P., Garnier, A., Pelon, J., Larsen, N., Nielsen, J., Christensen, T., Cairo, F., Thomason, L. W.,
1150 Leblanc, T., and McDermid, I. S.: Tropical stratospheric aerosol layer from CALIPSO lidar observations, *J. Geophys. Res.-
Atmos.*, 114, D00H10, <https://doi.org/10.1029/2009jd011946>, 2009.
- Vernier, J.-P., Fairlie, T. D., Murray, J. J., Tupper, A., Trepte, C., Winker, D., Pelon, J., Garnier, A., Jumelet, J., Pavolonis,
1155 M., Omar, A. H., and Powell, K. A.: An advanced system to monitor the 3D structure of diffuse volcanic ash clouds, *J. Appl.
Meteorol. Clim.*, 52, 2125–2138, <https://doi.org/10.1175/JAMC-D-12-0279.1>, 2013.
- Vernier, J.-P., Fairlie, T. D., Deshler, T., Natarajan, M., Knepp, T., Foster, K., Wienhold, F. G., Bedka, K. M., Thomason,
L., and Trepte, C.: In situ and space-based observations of the Kelud volcanic plume: The persistence of ash in the lower
stratosphere, *J. Geophys. Res. Atmos.*, 121, 11104–11118, <https://doi.org/10.1002/2016JD025344>, 2016.
1160
- Vernier, J., Fairlie, T.D., Deshler, T., Venkat Ratnam, M., Gadhavi, H., Kumar, B. S., Natarajan, M., Pandit, A. K., Akhil
Raj, S. T., Hemanth Kumar, A., Jayaraman, A., Singh, A. K., Rastogi, N., Sinha, P. R., Kumar, S., Tiwari, S., Wegner, T.,
Baker, N., Vignelles, D., Stenchikov, G., Shevchenko, I., Smith, J., Bedka, K., Kesarkar, A., Singh, V., Bhate, J., Ravikiran,
V., Durga Rao, M., Ravindrababu, S., Patel, A., Vernier, H., Wienhold, F. G., Liu, H., Knepp, T. N., Thomason, L.,
1165 Crawford, J., Ziemba, L., Moore, J., Crumeyrolle, S., Williamson, M., Berthet, G., Jegou, F., and Renard, J.: BATAL: The
Balloon Measurement Campaigns of the Asian Tropopause Aerosol Layer, *B. Am. Meteorol. Soc.*, 99, 955–973,
<https://doi.org/10.1175/BAMS-D-17-0014.1>, 2018.
- Waythomas, C. F., Scott, W. E., Prejean, S. G., Schneider, D. J., Izbekov, P., and Nye, C. J.: The 7–8 August 2008 eruption
1170 of Kasatochi Volcano, central Aleutian Islands, Alaska, *J. Geophys. Res.*, 115, <https://doi:10.1029/2010JB007437>, 2010.
- Wunderman, R. (Ed.): Report on Puyehue-Cordon Caulle (Chile), Global Volcanism Program, Bulletin of the Global
Volcanism Network, 37:3, Smithsonian Institution. <https://doi.org/10.5479/si.GVP.BGVN201203-357150>, 2012 (last access:
3 October 2022).
1175
- Winker, D. M., Liu, Z., Omar, A., Tackett, J., and Fairlie, D.: CALIOP observations of the transport of ash from the
Eyjafjallajökull volcano in April 2010, *J. Geophys. Res.*, 117, D00U15, <https://doi.org/10.1029/2011JD016499>, 2012.
- Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the
1180 CALIPSO mission and CALIOP data processing algorithms, *J. Atmos. Ocean. Tech.*, 26, 2310–2323,
<https://doi:10.1175/2009JTECHA1281.1>, 2009.

- Young, S. A., Vaughan, M. A., Kuehn, R. E., and Winker, D. M.: The retrieval of profiles of particulate extinction from Cloud–Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) data: Uncertainty and error sensitivity analyses, *J. Atmos. Ocean. Tech.*, 30, 395–428, <https://doi.org/10.1175/JTECH-D-12-00046.1>, 2013.
- 1185
- Yu, P., Toon, O. B., Bardeen, C. G., Zhu, Y., Rosenlof, K. H., Portmann, R. W., Thornberry, T. D., Gao, R. S., Davis, S. M., Wolf, E. T., de Gouw, J., Peterson, D. A., Fromm, M. D., and Robock, A.: Black carbon lofts wildfire smoke high into the stratosphere to form a persistent plume, *Science*, 365, 587–590, <https://doi.org/10.1126/science.aax1748>, 2019.
- 1190
- Zhu, Y., Toon, O. B., Kinnison, D., Harvey, V. L., Mills, M. J., Bardeen, C. G., Pitts, M., Begue, N., Renard, J.-B., Berthet, G., and Jegou, F.: Stratospheric Aerosols, Polar Stratospheric Clouds, and Polar Ozone Depletion After the Mount Calbuco Eruption in 2015, *J. Geophys. Res.-Atmos.*, 123, 12308–12331, <https://doi.org/10.1029/2018JD028974>, 2018.
- 1195
- Zhuang, J. and Yi, F.: Nabro aerosol evolution observed jointly by lidars at a mid-latitude site and CALIPSO, *Atmos. Environ.*, 140, 106–116, <https://doi.org/10.1016/j.atmosenv.2016.05.048>, 2016.