

Comparison of atmospheric CO, CO₂ and CH₄ measurements at Schneefernerhaus and the mountain ridge at Zugspitze

Antje Hoheisel¹, Cedric Couret², Bryan Hellack², and Martina Schmidt¹

¹Institute of Environmental Physics, Heidelberg University, Heidelberg, Germany

²German Environment Agency UBA, Germany

Correspondence: Antje Hoheisel (antje.hoheisel@iup.uni-heidelberg.de)

Abstract. The CO, CO₂, and CH₄ mole fractions have been measured since 2002 at the Environmental Research Station Schneefernerhaus, which is located approximately 300 m below the summit of Mount Zugspitze in the German Alps. Although the station is located remotely at an altitude of 2650 m a.s.l., local pollution events by snow blowers and snow groomers can be detected in the high temporal resolution time series of seconds or minutes. Therefore, a time-consuming flagging process, carried out manually by the station manager, is necessary.

To examine local influences and the effectiveness of data flagging, a 290 m long intake line to the higher Zugspitz ridge was used to measure CO, CO₂ and CH₄ mole fractions at a potentially less polluted location between October 2018 and October 2020. The comparison of these two time series shows that the mountain ridge measurement is almost unaffected by local pollution. It also demonstrates that the influence of local pollution events on the Schneefernerhaus measurements is successfully removed by the station manager. Only a small deviation up to 0.24 ppm can be observed during the day between the CO₂ time series of Schneefernerhaus and the mountain ridge in winter, probably due to anthropogenic sources.

1 Introduction

Since the beginning of the industrial era, the mole fraction of greenhouse gases, such as CO₂ and CH₄, has increased strongly (Etheridge et al., 1996, 1998; Dlugokencky and Tans, 2022; Dlugokencky, 2022). The atmospheric mole fractions of CO₂ and CH₄ in preindustrial times are determined by measuring, for example, air enclosed in ice cores (Etheridge et al., 1996, 1998), and since the second half of the 20th century (e.g., Keeling et al., 1976), direct atmospheric measurements of CO₂ and CH₄ have been possible. Historically, most measurement stations are located on coasts, such as Mace Head, Ireland (Bousquet et al., 1996), island mountains, such as Mauna Loa, Hawaii (Keeling et al., 1976) or Izana, Spain (Navascues and Rus, 1991; Gomez-Pelaez et al., 2019) or continental mountains, such as Mount Cimone, Italy (Ciattaglia, 1983), Jungfrauoch, Switzerland (Sturm et al., 2005; Schibig et al., 2016) or Schauinsland, Germany (Schmidt et al., 2003). Thus, long-term records from remote stations play an important role in improving understanding of the global carbon cycle and the impact of greenhouse gases on global warming.

Trace gas measurements of CO₂ have been performed at Mount Zugspitze since 1981 at different locations (Reiter, 1986; Yuan et al., 2019). First, at a pedestrian tunnel (ZPT), approximately 250 m below the summit, then at the terrace of the summit

25 (ZUG) and since 2002 at the Environmental Research Station Schneefernerhaus (ZSF). The scientific program at Schnee-
fernerhaus is operated by several German research institutes with the aim of monitoring the physical and chemical properties
of the atmosphere and analysing various processes that influence the weather and climate (UFS, 2020). At Schneefernerhaus
the measurements of CO, CO₂ and CH₄ mole fractions are performed by the German Environmental Agency (UBA - Umwelt-
bundesamt). Zugspitze greenhouse gas measurement program is part of the Global Atmosphere Watch (GAW) program and
30 joined the atmospheric network ICOS (Integrated Carbon Observation System) in 2021.

Due to the high elevation of mountain stations, the measured ambient air is less affected by regional and local influences
compared to urban stations. In recent years, several studies have analysed the CO₂ and CH₄ measurements performed at
Zugspitze (e.g., Yuan et al., 2018, 2019; Giemsa et al., 2019), also in combination with $\delta^{13}\text{C-CO}_2$ ratios (e.g., Ghasemifard
et al., 2019a, b). However, recent studies of greenhouse gas measurements with a temporal resolution of seconds to minutes
35 have shown, that local influences of anthropogenic activities can be seen in several mountain station time series, such as Pic du
Midi, where a small sewage treatment facility near the air intake of the analyser causes local CH₄ peaks (El Yazidi et al., 2018),
or Jungfraujoch, where the CO₂ measurement shows an influence of local anthropogenic activities, potentially by visitors and
tourism (Affolter et al., 2021).

Since 2012, new measurement techniques with a temporal resolution of nearly 1 Hz have been used at Schneefernerhaus to
40 measure atmospheric CO, CO₂ and CH₄ mole fractions. This enabled the detection of local pollution events in the measured
records, which could not be seen to this extent in previous measurements with lower time resolution. In particular, in winter,
snow groomers and gasoline snow blowers lead to strong CO peaks in the Schneefernerhaus time series, which must be flagged
manually to prevent an influence on the records. The German Meteorological Service (DWD - Deutscher Wetterdienst) has
also found that its measurements of radon (²²²Rn) activity at Schneefernerhaus are partially contaminated by local geogenic
45 sources of radon. Therefore, since 2014 they have used an ambient air inlet at the mountain ridge approximately 150m uphill
Schneefernerhaus for their measurements of ²²²Rn (Frank et al., 2017). As the local sources influencing the measurements of
CO, CO₂ and CH₄ at Schneefernerhaus are presumably near the station, a change in the location of the air intake line could also
reduce the influence on the CO, CO₂ and CH₄ time series. A new inlet line, installed from Schneefernerhaus to the mountain
ridge in 2018, allowed for the measurement of CO, CO₂ and CH₄ mole fractions in ambient air collected at the mountain ridge
50 since October 2018.

In our study, a two-year long comparison measurement between ambient air of Schneefernerhaus and of the mountain ridge
is analysed. The main focus of this study is to characterise the local pollution events and to compare the influence of these
events on the Schneefernerhaus and mountain ridge measurements.

2 Methods

55 2.1 Site description

Mount Zugspitze is Germany's highest mountain and its summit is 2962 m above sea level. It is located in southern Germany
in the northern Alps at the border with Austria (Fig. 1). The surrounding area consists largely of bare land, forests and pas-

tures. Urban areas occur less frequently. Around 11 km northeast of Mount Zugspitze lies the city of Garmisch-Partenkirchen (~27 000 inhabitants). Innsbruck (~131 000 inhabitants, 35 km southeast) and Munich (~1 509 000 inhabitants, 90 km north-east) are the largest cities within 100 km.

Atmospheric greenhouse gas measurements are carried out by the German Environment Agency (UBA) at the Environmental Research Station Schneefernerhaus (47° 25' 00" N, 10° 58' 46" E), which is approximately 300 m below the summit of Mount Zugspitze (Fig. 1, right panel). The air inlet for the measurements at Schneefernerhaus (ZSF, 2669 m a.s.l.) is installed at the research terrace on the fifth floor (Fig. 1, right panel). The analysis of recent high temporal resolution trace gas measurements has shown that the measured records at Schneefernerhaus are occasionally influenced by local pollution.

Possible sources can be found at Schneefernerhaus itself, at the rack railway tunnel from Schneefernerhaus to the valley, at the summit of Zugspitze and at the glacier plateau Zugspitzplatt, which is a ski resort in winter. Zugspitzplatt is located approximately 100 m below Schneefernerhaus. Each year more than 500 000 tourists (Bayerische Zugspitzbahn Bergbahn AG) visited the summit of Zugspitze and Zugspitzplatt, which can be reached from the valley by cable cars and rack railways. Furthermore, working activities at Schneefernerhaus, such as the usage of gasoline snow blowers, can influence the measurement. To keep local impacts low, electric snow blowers are generally used. However, after heavy snowfall, gasoline-powered equipment is needed. On weekdays, 12 people normally work at Schneefernerhaus during regular working hours.

In 2018, the German Meteorological Service (DWD) replaced an old and broken inlet line from Schneefernerhaus to the mountain ridge (ZGR, 2825 m a.s.l.) with a 290 m long new inlet line made of stainless steel (Fig. 1). Since October 2018, it has been possible to simultaneously measure the mole fraction of CO, CO₂ and CH₄ in the ambient air of Schneefernerhaus and of the mountain ridge above Schneefernerhaus.

2.2 Experimental setup

Since October 2018, the mole fractions of CO, CO₂ and CH₄ in ambient air at Zugspitze have been measured with three analysers, which are installed in the Research Station Schneefernerhaus and are connected to one of two inlet lines. One leads to the research terrace of Schneefernerhaus (ZSF, 2669 m a.s.l.) and the other to the mountain ridge (ZGR, 2825 m a.s.l.). In addition to the ambient air of Schneefernerhaus or the mountain ridge, the analysers simultaneously measure the same calibration and target gases for quality control (see Sect. 2.3 for further details). The measuring routine is controlled by the same multiposition rotary valve (model: EMT2SF6MWE, Valco Vici, Switzerland), over which all three instruments receive the measuring samples. The experimental setup is shown in Fig. A1.

The CO₂ and CH₄ mole fractions in ambient air of Schneefernerhaus are measured with a cavity ring-down spectroscopy (CRDS) G2301 analyser (Picarro, Inc., Santa Clara, CA) and the CO mole fractions with an off-axis integrated cavity output spectroscopy (OA-ICOS) EP30 analyser (LGR, Los Gatos Research). Ambient air is pumped with a flow rate of 500 L min⁻¹ from the research terrace on the fifth floor through the glass inlet at a height of 2.5 m above the terrace. To avoid freezing, the top of the glass inlet is heated. Part of the flow is then passed through a ~350 ml cold trap (Gaßner Glastechnik GmbH, Germany) to dry the ambient air and, thus, reduce the influence of water vapour on the measurement. The fluid bath of the cold trap is filled with silicon oil (model: M90.055/03, Huber Kältemaschinenbau AG, Germany) and is cooled to -80°C with

a cryogenic cooler (model: TC100E, Huber Kältemaschinenbau AG, Germany). The dried air is then measured by the CRDS G2301 and the OA-ICOS EP30 analysers, with the flow rate controlled to 0.2 and 0.4 L min⁻¹, respectively. Therefore, the residence time of air in the setup from the inlet at the research terrace to the analysers is only 35 s.

95 The CO, CO₂ and CH₄ mole fractions in ambient air of the mountain ridge are measured with a CRDS G2401 analyser (Picarro, Inc., Santa Clara, CA). The air is pumped through a 290 m stainless steel tube (1.2 cm o.d.) from the mountain ridge to Schneefernerhaus with a flow rate of 16 L min⁻¹. The air inlet at the mountain ridge is protected from rain, but not heated. A part of this air flow is then used by the CRDS G2401 analyser. Since 5 February 2019 this ambient air has been dried using the same drying system as the one used for the Schneefernerhaus measurements. The residence time of ambient air
100 from the mountain ridge to the analyser is 6 min 40 s. Therefore, to account for the different residence times between the Schneefernerhaus and mountain ridge measurements, the CRDS G2401 data averaged over one minute are shifted by -6 minutes.

On 16 July 2019 the setup was modified (see Fig. A1) by relocating two pumps out of the direct flow path of the three analysers. During this study, the UBA air quality measuring station on Zugspitze joined the atmospheric network ICOS (Inte-
105 grated Carbon Observation System). In the course of the ICOS labelling process, the CRDS G2301 and the OA-ICOS EP30 analysers were sent to the ICOS Atmosphere Thematic Centre (ATC) Metrology Laboratory for validation. Therefore, two large data gaps occurred in the measured time series. From 9 May 2019 to 8 August 2019, no CO, CO₂ and CH₄ mole fractions were measured at the mountain ridge and between 1 January 2020 and 8 July 2020, no CO mole fractions were measured at Schneefernerhaus.

110 2.3 Calibration, correction and quality control

The mole fractions of CO, CO₂ and CH₄ in ambient air of Schneefernerhaus and the mountain ridge were measured with a temporal resolution of 5 to 10 seconds with two CRDS analysers (G2301, G2401) and one OA-ICOS EP30 analyser. These measurements are averaged over one minute. The one-minute averaged values of the ambient air time series are calibrated
115 (CO: WMO X2014A, CO₂: WMO X2019, CH₄: WMO X2004A). Therefore, every three days, all analysers measure the same low and high working standard cylinders for 15 minutes. The four reference gases provided by the ESRL's Global Monitoring Laboratory (GML) of the National Oceanic and Atmospheric Administration (NOAA) are analysed every two months for 30 min. These WMO reference gases span the CO mole fraction from 124 to 269 ppb, the CO₂ mole fraction from 379 to 430 ppm and the CH₄ mole fraction from 1835 to 2120 ppb. Unless otherwise specified, the calibrated average values over one
120 minute were used in the analysis of the time series.

In addition to calibration cylinders, a target cylinder is simultaneously measured every three days by all analysers for quality control. The calibrated target measurements of CO, CO₂ and CH₄ were stable over time and there is no significant mean difference between the G2301/EP30 and G2401 analyses for all three species. Table 1 shows the average and standard deviation of the CO, CO₂ and CH₄ mole fractions for the two-year comparison for each analyser and their difference. Furthermore, the
125 CO, CO₂ and CH₄ mole fractions in ambient air from the Schneefernerhaus inlet was measured simultaneously with all

analysers for 15 days in July 2020. The average difference between the hourly averaged CO, CO₂ and CH₄ mole fractions measured with the G2301/EP30 and G2401 analysers are -0.04 ± 0.58 ppb (mean \pm sd) in CO, 0.04 ± 0.02 ppm in CO₂ and 0.3 ± 0.2 ppb in CH₄.

Two corrections were applied to the CO data. First, a large difference of roughly 5 ppb between the Schneefernerhaus and mountain ridge data was observed in the first few months of the comparison measurement. The quality control measurements described in section 2.3 show that this difference is not caused by the analysers or the calibration. The main difference between the two setups for measuring the ambient air at Schneefernerhaus and at the mountain ridge, leaving aside the different intake lines, is that the air from the mountain ridge was not dried. This observed offset disappeared after the air from the mountain ridge has also been dried since 5 February 2019. To determine the influence on the CO data prior to 5 February 2019, the same air from the Schneefernerhaus intake was measured with all three analysers on 12 December 2018, where the air for the CRDS G2301 and OA-ICOS EP30 analysers were dried as usual, but the air for CRDS G2401 analyser was not. No significant difference was found for CO₂ and CH₄ mole fractions. However, the CO mole fractions were on average 5.2 ± 1.5 ppb lower for device G2401. Therefore, a correction factor of 5.2 ppm was applied to the CO mole fractions from the mountain ridge for the time without drying. Secondly, an influence of the pumps used in the flow path to the analysers could be detected and therefore the setup was modified on 16 July 2019. Test measurements of quality control gases were performed to determine this impact. In three measurements the pump was placed within the flow path, as usual, and in two cases the pump was excluded. The CO measurement showed a difference of 7.5 ppb. For CH₄ and CO₂, no significant influence by the pump could be observed. Thus, the CO mole fractions for both analysers are corrected by -7.5 ppb until 16 July 2019. Since the CO measurements of ambient air of Schneefernerhaus and of the mountain ridge were equally affected, the difference between both measurements was not influenced.

2.4 Data flagging

The station manager from UBA manually applied different flags to the Schneefernerhaus CO, CO₂ and CH₄ time series to document invalid values, such as artefacts and outliers, due to technical problems or work at the setup, as well as local pollution events. Additional data from other trace gas measurements, meteorological data, and station logbooks were used for this purpose. These data from Schneefernerhaus were called “ZSF QC” in this study. The half-hourly, hourly, daily and monthly averages of CO, CO₂ and CH₄ mole fractions at Schneefernerhaus reported to national and international databases such as the World Data Centre for Greenhouse Gases (WDCGG) are also based on one-minute averaged data where local pollution events have been flagged as invalid in addition to artefacts and outliers. However, since this study also aims to assess the impact of local pollution on the Schneefernerhaus time series, a dataset that excludes only instrumental and technical problems, but not local pollution events, was also used in this study. This dataset was called “ZFS with local pollution”. Since the mountain ridge is more elevated than Schneefernerhaus, no strong pollution events in the measured time series were assumed. Thus, only technical artefacts and outliers were flagged in the time series from the mountain ridge, called “ZGR”.

3 Results and Discussion

3.1 CO, CO₂ and CH₄ mole fractions measured at Schneefernerhaus and the mountain ridge

160 Since October 2018, CO, CO₂, and CH₄ mole fractions have been measured in ambient air of Schneefernerhaus and the mountain ridge. Figure 2 shows the time series of 1 min averaged values at both sites. The measurements from the mountain ridge (ZGR) are displayed in red, the Schneefernerhaus ZSF QC time series are shown in black, and the one “with local pollution” in blue. In contrast to the mountain ridge measurement, strong pollution events can be observed at Schneefernerhaus (ZSF time series “with local pollution”). These are mainly visible in the CO record, but also in CO₂ and partly in CH₄. High
165 CO pollution events of more than 400ppb can be noticed, especially during the snow season. These spikes are caused by snow clearing in front of the station with gasoline snow blowers or in preparation of the nearby ski area with snow groomers. In January 2019, heavy snowfall and an avalanche that passed over the station led to intensive use of gasoline snow blowers at the station. During this time, extremely high CO mole fractions of up to 28000ppb and corresponding CO₂ and CH₄ spikes were measured at Schneefernerhaus. In addition, CO₂ peaks can be caused by scientists working on the measurement terrace
170 near the inlet. These local pollution events at Schneefernerhaus were identified and flagged manually by the station manager to reduce and avoid their influence on the measurement.

The Schneefernerhaus ZSF QC time series had similar CO, CO₂ and CH₄ mole fractions than measurements from the mountain ridge and both time series followed the same seasonal variations. The CO mole fractions ranged from 48ppb to 342ppb. The CO₂ mole fractions showed a seasonal cycle with lower values in summer varying between 390ppm and 440ppm,
175 and the mole fractions of CH₄ ranged from 1872ppb to 2100ppb. As expected, the measurements of ambient air of the mountain ridge and Schneefernerhaus showed similar large scale patterns, but at the mountain ridge, they are much less influenced by local pollution.

3.2 Local pollution events of CO and CO₂ at Schneefernerhaus

In the CO and CO₂ Schneefernerhaus time series “with local pollution” that averaged over 1 min, large and narrow peaks can
180 occasionally be detected. Approximately 2% of CO or CO₂ data measured at Schneefernerhaus were strongly contaminated by local pollution and are therefore flagged manually by the station manager. This corresponds to 1 000 high CO events (12900 one-minute averaged values) and 2 100 high CO₂ events (25900 one-minute averaged values).

Figure 3 shows the number of local polluted and manually flagged CO and CO₂ data points against the time of day (a) and month (b). Due to their anthropogenic origin, strong diurnal and seasonal variations in the occurrence of pollution events are
185 observed. The polluted and manually marked CO and CO₂ data points occurred at 92% and 95%, respectively, during the day between 6 and 18 UTC. Furthermore, 94% and 92% of the polluted and manually flagged CO and CO₂ data occurred during the snow season at high altitude (October to May) and after snowfall events when snowblower and snow groomer are used more often.

In 2020, the COVID-19 pandemic led to temporary restrictions on tourism at Zugspitze and a reduction in work at Schneefernerhaus and the ski resort. Unfortunately, our CO analyser for ambient air from Schneefernerhaus was being validated at the
190

ICOS ATC Metrology Laboratory as part of the ICOS labelling process at that time. Instead, we analysed NO_2 measured at Schneefernerhaus. Especially for NO_2 , but also for CO_2 , a decrease in the occurrence of NO_2 and CO_2 peaks was observed in the measured time series at Schneefernerhaus, indicating a reduction of local pollution events.

3.3 Local pollution events of CH_4 at Schneefernerhaus

195 The CH_4 mole fraction measured at Schneefernerhaus is also occasionally influenced by local pollution events. The high CH_4 peaks in the Schneefernerhaus time series “with local pollution” coincide with high CO or CO_2 events and are manually excluded by the station manager. Furthermore, inverse CH_4 spikes of several ppb occurred in the Schneefernerhaus time series, which were not detected at the mountain ridge. Typically, these inverse spikes do not occur as single events, but during periods with multiple inverse spikes. These periods can last between some hours to multiple days and start and end abruptly (see
200 Fig. 4 a). Two tests were performed during such events with inverse CH_4 spikes. First, we have exchanged the analysers and second, we have used an independent intake line that receives ambient air from the same location on the terrace at Schneefernerhaus. The inverse CH_4 spikes were measured in air from both sampling lines to the Schneefernerhaus terrace and with both analysers when measuring ambient air from the Schneefernerhaus. However, they were not detected when measuring ambient air from the mountain ridge. Therefore, these spikes are neither an artefact nor caused by the analyser, the measurement setup
205 or the inlet. Thus, CH_4 depleted air must be transported occasionally to the measuring terrace.

From the Schneefernerhaus station, a rack railway tunnel leads downhill to the valley. An opening to this tunnel is located 45 m from the measurement terrace to direct the tunnel air with high radon concentration to the outside and to reduce the radon concentration inside the Schneefernerhaus station. For 15 days in November 2020, the CH_4 mole fraction was measured with a CRDS G1301 analyser (Picarro, Inc., Santa Clara, CA) inside the tunnel near the opening (see Fig. 4 b). The measured
210 mole fraction of CH_4 strongly depends on the direction of the tunnel air flow, which was measured in the tunnel with a 1D anemometer. As the tunnel air flows downhill, fresh air enters the tunnel through the opening, and the CH_4 mole fraction has only slightly lower values than those measured on the terrace. However, extremely low CH_4 mole fractions between 500 and 1000 ppb are measured when the tunnel air flows uphill. In this case, local wind pattern can transport the CH_4 -depleted tunnel air into the ambient air inlet line at Schneefernerhaus, resulting in inverse CH_4 spikes. Since the tunnel air flows downhill more
215 frequently in summer than in winter, the influence of the tunnel air, and thus of the inverse CH_4 spikes, seems to be larger in winter.

3.4 Comparison of local pollution events at Schneefernerhaus and the mountain ridge

Although strong local pollution events are noticeable in the CO and CO_2 mole fractions measured at Schneefernerhaus, such strong events did not occur in the mountain ridge time series. Only small peaks are sometimes noticeable in the mountain
220 ridge record, when the wind blows from southeast, from the station Schneefernerhaus to the mountain ridge (see Appendix B). However, these pollution events have much smaller amplitudes than those measured simultaneously at Schneefernerhaus. On approximately 83% of the days when local pollution events were visible in the CO or CO_2 measurements at Schneeferner-

haus, we did not find corresponding peaks at the mountain ridge. Even extremely high CO events of more than 1000ppb at Schneefernerhaus are usually not detectable at the mountain ridge.

225 Figure 5 shows frequency distributions of deviations between the Schneefernerhaus and mountain ridge time series averaged to minute values for CO (Fig. 5a) and CO₂ (Fig. 5b) mole fractions. On the left, the Schneefernerhaus time series “with local pollution” are used for the calculation and on the right the Schneefernerhaus ZSF QC dataset, where local pollution events were manually excluded by the station manager. The daytime data (red) show a high number of large positive differences when the Schneefernerhaus time series “with local pollution” are used. These strong positive deviations disappear on the right panels
230 when the Schneefernerhaus ZSF QC data are included in the calculation. This indicates that the large positive differences are caused by local pollution at Schneefernerhaus. Furthermore, it shows that strong local pollution events in the Schneefernerhaus time series are successfully excluded by the station manager.

3.5 Differences between Schneefernerhaus and the mountain ridge measurements

The above analyses of the Schneefernerhaus and the mountain ridge measurements have shown that the mountain ridge time series are less influenced by local pollution, and strong local pollution events at Schneefernerhaus are successfully flagged by
235 the station manager. To verify whether there are further influences from local sources in the Schneefernerhaus time series, which do not manifest themselves as spikes, the hourly mean values of the measurements from Schneefernerhaus were compared with those from the mountain ridge. These correspond to the QC datasets reported to national and international databases.

Figure 6 shows the hourly averaged mole fraction of CO, CO₂ and CH₄ measured at Schneefernerhaus (black) and at the
240 mountain ridge (red), as well as the difference between both locations. The hourly data for Schneefernerhaus used in this comparison are calculated out of the Schneefernerhaus ZSF QC time series in which local pollution events are manually excluded. The average difference between Schneefernerhaus and mountain ridge data is 0.3 ± 2.4 ppb for CO, 0.1 ± 0.4 ppm for CO₂ and -0.4 ± 3.4 ppb for CH₄ and meets the compatibility goal determined by the WMO (2020). The large standard deviations in CO, CO₂ and CH₄ differences cannot be explained by instrumental noise. A comparison of the CO, CO₂ and
245 CH₄ mole fractions measured for 15 days with OA-ICOS EP30 or CRDS G2401 and CRDS G2301 using exactly the same air shows a standard deviation of hourly averages of 0.6 ppb CO, 0.02 ppm CO₂ and 0.2 ppb CH₄. Thus, the large standard deviations of the hourly differences between the ambient air of Schneefernerhaus and the mountain ridge are probably caused by the different measuring locations. As local wind conditions can strongly vary between Schneefernerhaus and the mountain ridge (see Appendix B and Fig. B1), air masses often occur with a time delay between the two locations.

250 To analyse the distribution of the differences between the Schneefernerhaus ZSF QC and the mountain ridge measurement in more detail, the frequency of the one minute differences between both locations are shown on the right side in Fig. 6. In addition to the whole dataset (black), the distribution is shown for nighttime (blue, 18 to 6 UTC) and daytime (red) mole fractions. The differences in the CO mole fractions show a small shift of 0.4 ppb between nighttime and daytime and a symmetric distribution. However, the differences are still within the uncertainties of the analysers.

255 The CO₂ mole fractions show larger differences during the day, with an average of 0.13 ppm. This is approximately 0.07 ppm larger than that during the night and exceeds the WMO compatibility goal of 0.1 ppm (WMO, 2020). Furthermore, the distri-

bution is asymmetric to positive differences, which even increase during the day. This indicates elevated CO₂ mole fractions at Schneesfernerhaus at daytime values. One explanation for this may be that the influence of local sources is not completely excluded by manual flagging, but the different sites may also receive slightly different signals from regional pollution due to the
260 different altitudes and complex topography. The influence of local sources on the CO₂ time series at remote mountain station is noticed also at other stations. At the mountain station Jungfraujoch, the average difference between the usual location and another, less polluted one, is 0.01 ppm overnight and even 0.49 ppm during the day (Affolter et al., 2021).

In contrast to the CO₂ mole fractions, CH₄ time series from Schneesfernerhaus and the mountain ridge are comparable within the WMO compatibility goal of 2 ppb (WMO, 2020) overnight and during the day. However, the distribution is slightly
265 asymmetric towards negative values. This distribution is probably related to inverse CH₄ spikes visible at Schneesfernerhaus, which are caused by CH₄-depleted air from the rack railway tunnel (Sect. 3.3).

3.6 Annual and diurnal cycles

The annual and diurnal cycles of the mole fraction of CO, CO₂ and CH₄ measured in ambient air at Schneesfernerhaus and the mountain ridge are analysed and compared in Fig. 7. To describe the Schneesfernerhaus data the ZSF QC dataset (black)
270 and the ZSF time series “with local pollution” (blue) are used again. In the calculations, only data for which simultaneous measurements at both locations are available are taken into account. The diurnal cycles for the high alpine winter (October to March) and from April to September are determined by calculating the diurnal variations per month and then averaging them.

The general pattern of the annual and diurnal cycles of CO, CO₂ and CH₄ measured at Schneesfernerhaus and the mountain ridge are similar to those of other mountain stations in the Northern Hemisphere (Thoning et al., 1989; De Wekker et al., 2009).
275 The diurnal cycles are mainly determined by the planetary boundary layer height and convective upslope winds (Yuan et al., 2019). Depending on the season and the time of day, the air masses of the free troposphere or the planetary boundary layer are measured. Since air from the planetary boundary layer is typically more polluted in CO and CH₄, higher values are measured during the day especially in summer. Since CO₂ uptake by the biosphere reduces the CO₂ mole fraction in the boundary layer, the CO₂ mole fraction measured at Zugspitze is lower during the day in summer.

In the following, the difference between the Schneesfernerhaus and mountain ridge measurements is discussed in more detail. The monthly averaged CO and CH₄ mole fractions measured at Schneesfernerhaus (ZSF QC) and the mountain ridge (ZGR), as well as the diurnal cycles for both locations, do not show differences below the recommended compatibility of 2 ppb. For CH₄, a small annual cycle in the differences can be noticed, with slightly positive values in summer and negative values in winter. The especially low CH₄ mole fractions at Schneesfernerhaus between November 2018 and January 2019 compared to
285 the mountain ridge are caused by particularly frequent and strong inverse CH₄ spikes in the Schneesfernerhaus time series (see Sect. 3.3).

The monthly CO₂ mole fractions measured at Schneesfernerhaus are comparable to those at the mountain ridge in summer. However, stronger deviations are observed in winter. In addition, the diurnal CO₂ cycle at Schneesfernerhaus shows up to 0.24 ppm higher values in the afternoon than at the mountain ridge, although the strong local pollution events in the Schneesfernerhaus time series are already excluded. Otherwise, the difference would be up to 0.35 ppm in winter. The difference between
290

the diurnal cycles at Schneefernerhaus and at the mountain ridge in the afternoon decreases in summer compared to winter, but it does not disappear completely. Thus, the Schneefernerhaus ZSF QC time series still show influences from local or regional sources in winter and during the day, which are not visible at the mountain ridge due to the higher elevation, complex wind pattern (see Appendix B and Fig. B1), and the topography of the Zugspitze.

295 Other mountain stations also show local influences, which depend on the season and time of day. At Jungfraujoch, the diurnal cycle of CO₂ between measurements at the usual location and a less polluted site deviate up to 0.4 ppm in winter and up to 1.3 ppm in summer (Affolter et al., 2021). Thus, the CO₂ measurement at Jungfraujoch is more strongly influenced in summer, which is probably due to the larger number of tourists. At Schneefernerhaus, on the other hand, the strongest influence was observed in winter.

300 3.7 Working day dependency in the Schneefernerhaus time series

In this study, a more frequent occurrence of CO and CO₂ spikes during the day was found, which can be attributed to anthropogenic activities. In addition to the diurnal dependence, these local pollution events also have a working day dependence. Thus, 92% of the data points that were manually excluded due to local pollution occur on weekdays and not on weekends.

In a previous study by Yuan et al. (2019), which analysed the CO₂ time series at Schneefernerhaus between 2002 and
305 2016, a weekly cycle in the CO₂ measurement with lower CO₂ mole fractions on weekends and higher CO₂ mole fractions on weekdays is described. Additionally, they found stronger diurnal cycles on weekdays with higher values, especially in the morning, for the combined time period 2002 to 2016. They explained these findings by local anthropogenic working activities at Schneefernerhaus.

When analysing the Schneefernerhaus and mountain ridge measurements between October 2018 and October 2020 for the
310 weekly cycle and stronger diurnal cycles, no working day dependencies were detected for either the CO₂ or the CO and CH₄ data. Therefore, we investigated the CO₂ time record for Schneefernerhaus between 2002 and 2021, similar to Yuan et al. (2019). A statistical analysis was performed using the resampling technique described in Daniel et al. (2012) to determine if a weekly cycle showed a significant pattern. No significant weekend effect (mean weekend value minus mean weekday value) occurred until 2007, and after 2014. However, significantly higher values were evident during the week in the data between
315 2008 and 2014 (Fig. C1). Furthermore, the average diurnal CO₂ cycles on weekends were comparable for the three time periods 2002–2007, 2008–2014 and 2015–2021, which is not the case for weekdays. A strong difference of up to 1.5 ppm was noticed between the diurnal CO₂ cycles on weekends and weekdays in the time periods 2002–2007 and 2008–2014 (Fig. C2). Such a difference is not noticeable in the CO₂ data from 2015 to 2021.

A micro leakage in the inlet system could explain the noticed weekend effect and the higher CO₂ values during the day on
320 weekdays, as the strongest effect would be noticeable during the working hours when the CO₂ mole fraction in the laboratory is higher. Between 2014 and 2015, the entire measurement setup, including the intake system, was thoroughly checked for leakages, and old connections were renewed. In the process, the high temporal resolution CO₂ measurements of the CRDS analyser were also used. Presumably, the micro leakage was fixed at that time, as no working day dependency has been detectable since then.

325 4 Conclusions

Ambient air measured at Schneefernerhaus with high temporal resolution showed local pollution events caused by human activities, especially in CO and CO₂. These peaks occur mainly in the winter season and during the day. Therefore, approximately 2% of the measured data were flagged manually by the station manager.

330 To prevent the influence of this local pollution, additional measurements were taken at a more elevated location for comparison. This study showed that the atmospheric mole fractions of CO, CO₂ and CH₄ can be measured at the mountain ridge using a 290m long intake line. The time series determined in this way does not show a strong influence of local pollution events, unlike the measurements at Schneefernerhaus.

The comparison of measurements performed at Schneefernerhaus with baseline time series from the mountain ridge characterises the local influence on the CO, CO₂ and CH₄ time series. It is confirmed that strong local pollution events at Schneefernerhaus can be successfully removed from the Schneefernerhaus time series by the station manager. No significant difference in 335 the annual cycle of CO and CH₄ for the Schneefernerhaus time series with manually flagged pollution events and the mountain ridge could be noticed, and the diurnal cycles were similar at both locations, too. The annual and diurnal cycles in CO₂ measured at Schneefernerhaus and the mountain ridge also show the same pattern. In summer the difference between monthly values and diurnal cycles is negligible. In winter, however, higher CO₂ mole fractions occur at Schneefernerhaus especially 340 during the day, which indicates a small impact of local or regional pollution that is not yet excluded.

With the exception of slightly elevated CO₂ mole fractions during the day, the CO, CO₂, and CH₄ measurements at the Schneefernerhaus station are also consistent with background measured at the mountain ridge. However, as it is difficult to maintain the 290m intake line, especially in the winter months, and thus to ensure continuous series of measurements on the mountain ridge, it is important to continue measurements at both sites in the future.

345 *Data availability.* The high resolution (1 min) CO, CO₂ and CH₄ records of Zugspitze-Schneefernerhaus (ZSF) and the mountain ridge (ZGR) used in this study are available on request from the data owner (Cedric.Couret@uba.de). Hourly averaged CO, CO₂ and CH₄ records of Zugspitze-Schneefernerhaus (ZSF) are available from the World Data Centre for Greenhouse Gases (WDCGG) at <https://gaw.kishou.go.jp/>.

Appendix A: Experimental setup at Schneefernerhaus

Appendix B: Wind pattern

350 The wind direction and wind velocity at Schneefernerhaus and at Zugspitze summit are measured by DWD and at the mountain ridge by UFS. Figure B1 shows the wind roses computed from the available wind direction and the wind velocity at the three locations between October 2018 and October 2020. The colours indicate different wind speed ranges. The wind direction measured at Schneefernerhaus and at the mountain ridge is most likely influenced by the characteristic locations and thus by the topography of the mountain. The mountain ridge above Schneefernerhaus runs largely from west to east, so Schneefernerhaus

355 is on the southern slope of the mountain. Therefore, north and south winds are more frequently measured on the mountain ridge and the main wind directions at Schneesfernerhaus are east and west. At the Zugspitze summit, the inflow of air masses is less restricted, with more frequent winds from the northwest, southwest and southeast. Furthermore, the average wind velocity at the summit is 6.0 m s^{-1} and therefore larger than that at the mountain ridge (4.0 m s^{-1}) or Schneesfernerhaus (3.2 m s^{-1}).

Appendix C: Working day dependency in the Schneesfernerhaus time series between 2002 and 2021

360 To determine if the CO_2 mole fraction is generally higher on weekdays than on weekends, the data were detrended by subtracting a moving 31-day mean to remove the annual cycle. Since we are interested in the average variations within a week, weeks with missing days were excluded. A statistical analysis was performed using the resampling technique described in Daniel et al. (2012) to identify if a weekly cycle shows a significant pattern. For each year, we tested whether the weekend effect magnitude (mean weekend value minus mean weekday value) was significant. Until 2007, no significant ($p > 0.1$) weekend effect was noticeable. Between 2008 and 2014 every year except 2013 showed a significant ($p < 0.1$) weekend effect. No weekend effect has been visible since 2015. Therefore, we separate the CO_2 time series into three parts: 2002–2007, 2008–2014 and 2015–2021. Figure C1 shows the weekly cycles for all three time intervals. Even visually, the difference between the weekly cycles for 2008–2014 and the other two time periods is obvious. As expected, the weekly cycles for the time intervals 2002–2007 and 2015–2021 show no significant ($p < 0.05$) weekend effect, as does the weekly cycle for the time period 2008–2014.

370 Furthermore, we calculated the mean diurnal cycles for weekends and weekdays for each year. Before averaging, each single diurnal cycle is subtracted by the average nighttime (18 to 6 UTC) CO_2 mole fraction. For the years 2002 to 2014, a similar working day dependency can be noticed as reported by Yuan et al. (2019), with much stronger diurnal cycles on weekdays than on weekends. Since 2015, the difference in the diurnal cycles between weekdays and weekends disappeared. Figure C2 shows the diurnal cycles for weekdays and weekends for the winter (October to March) and summer (April to September). The weekday (Monday to Friday) diurnal cycles are shown in red and weekend cycles (Saturday and Sunday) in black. In addition, the difference between averaged weekday and weekend values are shown. For the first two time periods (2002–2007 and 2008–2014), stronger CO_2 peaks (5 to 15 UTC) were noticed during weekdays, which did not occur on weekends. The difference between weekdays and weekends was lower than 0.2 ppm during the night and reached values between 1 and 1.5 ppm during the day. In the time period 2015–2021 the diurnal cycles of weekends and weekdays showed no strong differences, with values lower than 0.2 ppm for the whole day. Comparing the three periods, it was further noticed that the diurnal cycles of weekends have the same amplitude and shape, especially in summer.

385 Yuan et al. (2019) explained the working day dependencies by local anthropogenic working activities at Schneesfernerhaus. This seems plausible, since the main increase in CO_2 occurred during working hours on weekdays. The tourist activities around Schneesfernerhaus are, on the other hand, normally stronger on weekends than on weekdays. However, the absence of the weekly cycle and of the difference in the diurnal cycles of weekday and weekend data since 2015 implicate a relevant change in the working activities at Schneesfernerhaus that did not occur. It is more likely that a micro leakage in the inlet system caused higher CO_2 values during the week until it was fixed between 2014 and 2015. The micro leakage affected the measured

CO₂ mole fractions only during working hours, when the CO₂ mole fraction inside the laboratory was much higher due to the presence of humans. The analysis has shown that nowadays there is no strong impact of local pollution on weekends or
390 weekdays.

Author contributions. AH and MS designed the study together with CC. AH evaluated the data and wrote the paper with the help of MS, CC and BH. CC was responsible for the GHG measurements at Scheefernerhaus and at the mountain ridge.

Competing interests. The authors declare that they have no conflict of interest.

395 *Acknowledgements.* The authors would like to thank the German Meteorological Service (DWD - Deutscher Wetterdienst) for providing the intake line to the mountain ridge. We also thank Gabriele Frank (DWD) and Josef Salvamoser (IGU - Institut für angewandte Isotopen-, Gas- und Umweltuntersuchungen) for making it possible to use the intake line to the mountain ridge for our comparison measurements. Our thanks also go to the UFS team for maintaining the intake lines and operating the weather station on the mountain ridge.

The data analysis of this study was funded by Projects 95297 and 167847 with the German Environment Agency (UBA). We wish to thank Frank Meinhardt and Ludwig Ries for their great support in these cooperation projects between the UBA and Heidelberg University.

400 References

- Affolter, S., Schibig, S., Berhanu, T., Bukowiecki, N., Steinbacher, M., Nyfeler, P., Hervo, M., Lauper, J. and Leuenberger, M.: Assessing local CO₂ contamination revealed by two near-by high altitude records at Jungfraujoch, Switzerland, *Environ. Res. Lett.*, 16, <https://doi.org/10.1088/1748-9326/abe74a>, 2021.
- Bousquet, P., Gaudry, A., Ciais, P., Kazan, V., Monfray, P., Simmonds, P.G., Jennings, S.G., O'Connor, T.C.: Atmospheric CO₂ concentration variations recorded at Mace Head, Ireland, from 1992 to 1994. *Phys. Chem. Earth* 21, pp. 477-481, 1996.
- 405 Ciattaglia, L.: Interpretation of atmospheric CO₂ measurements at Mt. Cimone (Italy) related to wind data, *J. Geophys. Res.*, 88, 1331, <https://doi.org/10.1029/JC088iC02p01331>, 1983.
- Daniel, J. S., Portmann, R. W., Solomon, S., and Murphy, D. M.: Identifying weekly cycles in meteorological variables: The importance of an appropriate statistical analysis, *J. Geophys. Res.-Atmos.*, 117, 14, <https://doi.org/10.1029/2012jd017574>, 2012.
- 410 De Wekker, S.F., Ameen, A., Song, G. et al.: A preliminary investigation of boundary layer effects on daytime atmospheric CO₂ concentrations at a mountaintop location in the Rocky Mountains, *Acta Geophys.* 57, 904–922, <https://doi.org/10.2478/s11600-009-0033-6>, 2009.
- Dlugokencky, E., NOAA/GML (https://gml.noaa.gov/ccgg/trends_ch4/), last access: 23 June 2022 ,2022.
- Dlugokencky, E. and Tans, P. , NOAA/GML (<https://gml.noaa.gov/ccgg/trends/>), last access: 23 June 2022 ,2022.
- 415 El Yazidi, A., Ramonet, M., Ciais, P., Broquet, G., Pison, I., Abbaris, A., Brunner, D., Conil, S., Delmotte, M., Gheusi, F., Guerin, F., Hazan, L., Kachroudi, N., Kouvarakis, G., Mihalopoulos, N., Rivier, L., and Serça, D.: Identification of spikes associated with local sources in continuous time series of atmospheric CO, CO₂ and CH₄, *Atmos. Meas. Tech.*, 11, 1599–1614, <https://doi.org/10.5194/amt-11-1599-2018>, 2018.
- Etheridge, D. M., Steele, L. P., Langenfelds, R. L., Francey, R. J., Barnola, J. M., and Morgan, V. I.: Natural and anthropogenic changes in atmospheric CO₂ over the last 1000 years from air in Antarctic ice and firn, *J. Geophys. Res.-Atmos.*, 101, 4115-4128, <https://doi.org/10.1029/95jd03410>, 1996.
- 420 Etheridge, D. M., Steele, L. P., Francey, R. J., and Langenfelds, R. L.: Atmospheric methane between 1000 AD and present: Evidence of anthropogenic emissions and climatic variability, *J. Geophys. Res.-Atmos.*, 103, 15979-15993, <https://doi.org/10.1029/98jd00923>, 1998.
- Frank, G., Salvamoser, J., Steinkopff, T. and Ries, L.: Radon-222 und Beryllium-7 als natürlicher tracer, UFS-Wissenschaftliche Resultate 425 2015/2016, Hrg. Umweltforschungsstation Schneefernerhaus GmbH, 2017.
- Ghasemifard, H., Yuan, Y., Luepke, M., Schunk, C., Chen, J., Ries, L., Leuchner, M., and Menzel, A.: Atmospheric CO₂ and delta C-13 Measurements from 2012 to 2014 at the Environmental Research Station Schneefernerhaus, Germany: Technical Corrections, Temporal Variations and Trajectory Clustering, *Aerosol Air Qual. Res.*, 19, 657-670, <https://doi.org/10.4209/aaqr.2018.01.0010>, 2019.
- Ghasemifard, H., Vogel, F. R., Yuan, Y., Luepke, M., Chen, J., Ries, L., Leuchner, M., Schunk, C., Vardag, S. N., and Menzel, A.: Pollution 430 Events at the High-Altitude Mountain Site Zugspitze-Schneefernerhaus (2670 m a.s.l.), Germany, *Atmosphere*, 10, 17, <https://doi.org/10.3390/atmos10060330>, 2019.
- Giemsa, E., Jacobeit, J., Ries, L., and Hachinger, S.: Investigating regional source and sink patterns of Alpine CO₂ and CH₄ concentrations based on a back trajectory receptor model, *Environ. Sci Eur.*, 31, 24, <https://doi.org/10.1186/s12302-019-0233-x>, 2019.
- Gomez-Pelaez, A. J., Ramos, R., Cuevas, E., Gomez-Trueba, V., and Reyes, E.: Atmospheric CO₂, CH₄, and CO with the CRDS technique 435 at the Izaña Global GAW station: instrumental tests, developments, and first measurement results, *Atmos. Meas. Tech.*, 12, 2043–2066, <https://doi.org/10.5194/amt-12-2043-2019>, 2019.

- Keeling, C. D., Bacastow, R. B., Bainbridge, A. E., Ekdahl, C. A., Guenther, P. R., Waterman, L. S., and Chin, J. F. S.: ATMOSPHERIC CARBON-DIOXIDE VARIATIONS AT MAUNA-LOA OBSERVATORY, HAWAII, *Tellus*, 28, 538-551, 1976.
- Navascues, B. and Rus, C.: Carbon dioxide observations at Izaña baseline station, Tenerife (Canary Islands): 1984–1988. *Tellus* 43B, 118–440 125, 1991.
- Reiter, R., Sladkovic, R., and Kanter, H.-J.: Concentration of trace gases in the lower troposphere, simultaneously recorded at neighboring mountain stations Part I: Carbon Dioxide, *Meteorol. Atmos. Phys.*, 35, 187–200, <https://doi.org/10.1007/BF01041811>, 1986.
- Schibig, M. F., Mahieu, E., Henne, S., Lejeune, B., and Leuenberger, M. C.: Intercomparison of in situ NDIR and column FTIR measurements of CO₂ at Jungfraujoch, *Atmos. Chem. Phys.*, 16, 9935–9949, <https://doi.org/10.5194/acp-16-9935-2016>, 2016.
- 445 Schmidt, M., Graul, R., Sartorius, H., and Levin, I.: Carbon dioxide and methane in continental Europe: A climatology, and (222)Radon-based emission estimates, *Tellus Ser. B-Chem. Phys. Meteorol.*, 48, 457-473, <https://doi.org/10.3402/tellusb.v48i4.15926>, 1996.
- Sturm, P., Leuenberger, M., and Schmidt, M.: Atmospheric O₂, CO₂ and $\delta^{13}\text{C}$ observations from the remote sites Jungfraujoch, Switzerland, and Puy de Dôme, France, *Geophys. Res. Lett.*, 32, 2467, <https://doi.org/10.1029/2005GL023304>, 2005.
- Thoning, K. W., Tans, P. P., and Komhyr, W. D.: ATMOSPHERIC CARBON-DIOXIDE AT MAUNA LOA OBSERVATORY .2. ANALYSIS 450 OF THE NOAA GMCC DATA, 1974-1985, *J. Geophys. Res.-Atmos.*, 94, 8549-8565, <https://doi.org/10.1029/JD094iD06p08549>, 1989.
- UFS 2020: Environmental Research Station Schneefernerhaus, <https://www.schneefernerhaus.de>, last access: 24 Sept 2020 ,2020.
- WMO 2020: 20th WMO/IAEA Meeting on Carbon Dioxide, Other Greenhouse Gases and Related Measurement Techniques (GGMT-2019), Jeju Island, South Korea 2–5 September 2019. GAW Report No. 255, 2020.
- Yuan, Y., Ries, L., Petermeier, H., Steinbacher, M., Gómez-Peláez, A. J., Leuenberger, M. C., Schumacher, M., Trickl, T., Couret, C., 455 Meinhardt, F., and Menzel, A.: Adaptive selection of diurnal minimum variation: a statistical strategy to obtain representative atmospheric CO₂ data and its application to European elevated mountain stations, *Atmos. Meas. Tech.*, 11, 1501–1514, <https://doi.org/10.5194/amt-11-1501-2018>, 2018.
- Yuan, Y., Ries, L., Petermeier, H., Trickl, T., Leuchner, M., Couret, C., Sohmer, R., Meinhardt, F., and Menzel, A.: On the diurnal, weekly, and seasonal cycles and annual trends in atmospheric CO₂ at Mount Zugspitze, Germany, during 1981–2016, *Atmos. Chem. Phys.*, 19, 460 999–1012, <https://doi.org/10.5194/acp-19-999-2019>, 2019.




Figure 1. Location of the Zugspitze-Schneefernerhaus measurement site. On the right side the two inlet lines at Schneefernerhaus (ZSF) and the mountain ridge (ZGR) are shown (Map data on left side: from © Google Earth).

Table 1. Calibrated target gas measurements which were performed with the CRDS G2301, G2401 and OA-ICOS EP30 analysers. The average and standard deviation of each analyser and the differences are shown. The CRDS G2301 and OA-ICOS EP30 analysers were usually used for the ambient air measurement at Schneefernerhaus and CRDS G2401 analyser for the measurements at the mountain ridge.

analyser	CO [ppb]	CO ₂ [ppm]	CH ₄ [ppb]
CRDS G2301	-	404.87 ± 0.03	1887.10 ± 0.23
CRDS G2401	124.6 ± 0.8	404.87 ± 0.02	1887.07 ± 0.03
OA-ICOS EP30	124.6 ± 0.1	-	-
difference	0.03 ± 0.87	-0.001 ± 0.017	0.03 ± 0.23




Figure 2. CO, CO₂ and CH₄ mole fractions in ambient air of Schneefernerhaus (ZSF QC: black, ZSF with local pollution: blue) and of the mountain ridge (ZGR: red). The shown data are averaged over 1 min.




Figure 3. Distribution of local polluted and manually flagged CO and CO₂ data at Schneefernerhaus as a function of month and time of day. The 1 min averaged dataset was used for this analysis.




Figure 4. (a): Example of inverse CH₄ spikes measured at Schneefernerhaus (ZSF: black), but not at the mountain ridge (ZGR: red). They are probably caused by CH₄ depleted air from the nearby rack railway tunnel.

(b): CH₄ mole fraction in ambient air at Schneefernerhaus (ZSF: black) and in tunnel air (blue) near the entrance of the rack railway tunnel. The beige-coloured time intervals correspond to a downhill wind-direction in the rack railway tunnel.




Figure 5. The frequency distribution of the difference between the 1 min averaged Schneefernerhaus (ZSF) and the mountain ridge (ZGR) measurements for CO (a) and CO₂ (b). The counts are displayed on a logarithmic scale. The blue colour corresponds to nighttime (18 to 6 UTC) and red to daytime (6 to 18 UTC) data. On the left side of the two figures, the Schneefernerhaus dataset with local pollution is used to calculate the differences, while on the right side the Schneefernerhaus ZSF QC dataset is used in which local pollution events have been manually removed by the station manager.




Figure 6. Left: Hourly averaged Schneefernerhaus (ZSF QC: black) and mountain ridge (ZGR: red) mole fractions of CO, CO₂ and CH₄, as well as the difference between both locations.

Right: The frequency of the differences between Schneefernerhaus and the mountain ridge calculated using the one-minute averaged data. All data are shown in black, whereas only the differences at night (18 to 6 UTC) and during the day are shown in blue and red, respectively.




Figure 7. Annual and diurnal cycles in the Schneefernerhaus (ZSF QC: black, ZSF with local pollution: blue) and mountain ridge (ZGR: red) measurements. In addition, the difference between the Schneefernerhaus and mountain ridge data is shown. The error bars are the standard errors of the averages. The beige coloured areas correspond to the WMO compatibility goals (WMO, 2020).




Figure A1. Experimental setup to measure CO, CO₂ and CH₄ in ambient air from Schneefernerhaus and from the mountain ridge.




Figure B1. Wind rose plots for the mountain ridge (left), Schneefernerhaus (middle) and Zugspitze summit (right) showing the wind directions and wind velocities between October 2018 and October 2020 provided by UFS (mountain ridge) and by DWD (Schneefernerhaus and Zugspitze summit). The colours correspond to different wind speed ranges.




Figure C1. Weekly cycles in CO₂ measured at Schneefernerhaus during three time periods: : 2002–2007 (blue), 2008–2014 (red), 2015–2021 (black). The CO₂ mole fraction is detrended by subtracting the moving 31-day average and weeks with missing days were excluded before averaging for each day of the week.




Figure C2. Mean diurnal cycles for Schneefernerhaus calculated from the weekend and weekday data between 2002 and 2021 for two seasons. Diurnal cycles calculated using weekday data are coloured red, and those using weekend data are in black. The top panels correspond to the seasons October to March, and the bottom panels correspond to April to September. The cycles in the lower part of the two panels show the difference between weekdays and weekends. The beige coloured areas correspond to the normal and extended WMO compatibility goals (WMO, 2020).