



Numerical Model Generation of Test Frames for Pre-launch Studies of EarthCARE's Retrieval Algorithms and Data Management System

Zhipeng Qu¹, David P. Donovan², Howard W. Barker¹, Jason N. S. Cole¹, Mark W. Shephard¹,
5 Vincent Huijnen²

¹ Environment and Climate Change Canada, Toronto, ON, Canada

² Royal Netherlands Meteorological Institute, De Bilt, Netherlands

Correspondence to: Howard W. Barker (howard.barker@canada.ca)

10 **Abstract.** The *Earth Cloud, Aerosol and Radiation Explorer* (EarthCARE) satellite consists of active and passive
sensors whose observations will be acted on by an array of retrieval algorithms. EarthCARE's retrieval algorithms
have undergone pre-launch verifications within a virtual observing system that consists of 3D atmosphere-surface
data produced by the Global Environmental Multi-scale (GEM) NWP model, and instrument simulators that when
applied to NWP data yield synthetic observations for EarthCARE's four sensors. Retrieval algorithms operate on the
15 synthetic observations and their estimates go into radiative transfer models that produce top-of-atmosphere solar and
thermal broadband radiative quantities, which are compared to synthetic broadband measurements thus mimicking
EarthCARE's radiative closure assessment. Three high-resolution test frames were simulated; each measures ~6,200
km along-track by 200 km across-track. Horizontal grid-spacing is 250 m and there are 57 atmospheric layers up to
10 mb. The frames span wide ranges of conditions and extend over: i) Greenland to The Caribbean crossing a cold
20 front off Nova Scotia; ii) Nunavut to Baja California crossing over Colorado's Rooky Mountains; and iii) central
equatorial Pacific Ocean that includes a mesoscale convective system. This report discusses how the test frames were
produced and presents their key geophysical features. All data are publicly available and, owing to their high-
resolution, could be used to simulate observations for other measurement systems.

1. Introduction

25 The *Earth Cloud, Aerosol and Radiation Explorer* (EarthCARE) satellite mission, which is scheduled for launch in
late-2023, is a joint venture funded by the European Space Agency (ESA) and Japanese Aerospace Exploration
Agency (JAXA) (Illingworth et al. 2015). The combination of Dopplerized cloud profiling radar (CPR), high-
spectral-resolution lidar (ATLID), and multi-spectral imager (MSI) will facilitate synergistic retrieves of profiles of
cloud, aerosol, and precipitation properties. Broadband top-of-atmosphere (TOA) radiances and fluxes calculated
30 using these profiles will be compared to near-coincidental observations made by EarthCARE's broadband radiometer



(BBR). This radiative closure assessment of retrievals will provide continuous feedback of performance to algorithm developers, as well as guidance to data users.

ESA's pre-launch phase of EarthCARE has relied much on *end-to-end simulation* of the measurement, retrieval, and data management systems. The primary objective was to build a virtual observing system in which retrieval algorithms, developed expressly for EarthCARE, get applied to synthetic observations that resemble closely those that will be made by all of EarthCARE's sensors. The initial step of this multi-stage process is definition of surface-atmosphere conditions. The obvious starting point was single homogeneous columns, but this quickly evolved into numerical simulation of realistic conditions for domains that span substantial portions of EarthCARE's planned orbit. These surface-atmosphere ensembles are then operated on by instrument simulators that yield synthetic observations suitable for ingestion by retrieval algorithms. One could stop here and assess performance by comparing retrieved geophysical quantities to their simulated counterparts (cf. Mason et al. 2023), but obviously the real mission lacks this luxury. As such, the next step in the *end-to-end* simulation chain is application of radiative transfer models to retrieved geophysical properties. This produces radiometric quantities that are commensurate with synthetic BBR observations that, in turn, come from application of similar radiative transfer models directly to the initially simulated atmosphere-surface fields. The comparison of these quantities defines EarthCARE's radiative closure assessment of its retrievals.

EarthCARE's data-handling system processes observations in "frames", of which there are six per orbit. As such, frames are ~6,500 km in the along-track direction. Their across-track width is 150 km as defined by the MSI's swath. Almost all measured and retrieved products are reported on the *Joint Standard Grid* (JSG), whose resolutions are ~1 km in both horizontal directions and 0.5 km in the vertical. It was established early on, by EarthCARE's science and engineering teams, that synthetic observations for end-to-end experiments need to cover entire frames and be resolved horizontally to better than 1 km. Environment and Climate Change Canada's numerical weather prediction (NWP) model, known as the Global Environment Multiscale (GEM) model (Côté et al., 1998, Girard et al., 2014), was used to produce three appropriately sized test frames with horizontal grid-spacings of 250 m and 57 atmospheric layers up to 10 mb. The frames include wide ranges of conditions and extend from: i) Greenland to The Caribbean, crossing a cold front off Nova Scotia; ii) Nunavut to Baja California, crossing over Colorado's Rooky Mountains; and iii) central equatorial Pacific Ocean, including a mesoscale convective system. The primary purposes of this



paper is to report on how these frames were constructed, their cloud, aerosol, and surface properties, as well as adjustments that were made to GEM's initial estimates of ice cloud particle sizes.

60 Full-frame datasets produced by GEM serve as inputs to the *EarthCARE simulator* (ECSIM) (Voors et al. 2007). ECSIM consists of radiative transfer and instrument models that are coupled to databases of optical and microwave scattering properties. Once bulk properties of atmospheric attenuators, such as GEM's 3D distributions of cloud water contents (CWC), are parsed into bin-resolved aerosol/cloud size distributions, they are used in ECSIM, which produces physically-consistent synthetic measurements to be made by EarthCARE's sensors. Production of simu-
65 ed L1 EarthCARE data using ECSIM is described by (Donovan et al., 2023).

The following section discusses how the test frames were defined. This is followed by descriptions of GEM and how it was configured and used for this study. Aerosols and surface optical properties were added to GEM's atmospheres, and these procedures are discussed in sections 4 and 5. Section 6 presents, and to a limited extent verifies, the simulated frames. Section 7 discusses issues with, and subsequent modifications to, GEM's simulated ice clouds. Con-
70 cluding remarks and information regarding acquisition of test frame data are provided in the final section.

2. Satellite orbit selection

Figure 1 shows several EarthCARE orbits, numbered 39316 through 39320. An orbit consists of eight frames; each frame's number having an appending letter from A to H. Frames are colour-coded and measure ~5,000 km along-track and 150 km across-track. Both ends of a frame overlap neighbouring frames by ~600 km for a total length of
75 ~6,200 km. All frames selected for testing correspond to local afternoon descending conditions (i.e., opposite to the A-Train). This is because night retrievals can be approximated by neglecting MSI solar channels and solar background for ATLID. Test frames should cover wide varieties of clouds, surface, meteorological, and solar illumination conditions. Locations and times needed to initialize simulations of test frames have been established by examining GOES satellite imagery and surface meteorological data. Also, A-Train's active sensor observations had to intersect
80 the frame.

As *Figure 1* shows, frame 39316D extends from southern Greenland, across extreme eastern Canada, and ends in the Atlantic Ocean roughly 500 km north of Dominican Republic. Because it passes close to the city of Halifax, Nova Scotia, it is referred to hereinafter as the *Halifax frame*. *Figure 2* shows GOES-13 reflectances and TOA brightness



temperatures for its 0.63 μm and 10.7 μm channels for 7-December-2014 at 18h00 UTC. *Table 1* lists surface condi-
85 tions reported at 18h00 UTC by several meteorological stations close to the ground-track (see dots on both *Figure 1*
and *Figure 2*). This frame includes no Sun over Greenland, cold surface air over eastern Canada, a cold-front with
deep clouds just off the coast of Nova Scotia, and scattered shallow clouds between Bermuda and Dominican Republic.

The second frame, 39318D, is referred to as the *Baja frame*. It stretches from the Canadian Arctic Archipelago, over
90 central North America's Great Plains and Rocky Mountains, and ends near Baja California Sur. *Figure 3* shows
GOES-15 imagery and EarthCARE's ground-track for 2-April-2015 at 21h00 UTC. *Table 2* lists surface conditions
reported at 21h00 UTC. At the north end of this frame surface conditions were cold with blowing snow and largely
cloudless. Through the Canadian Prairies there were low scattered clouds over snow-covered surfaces, while over the
Rocky Mountains skies were very cloudy. In the southern reaches, skies were clear with some cirrus, and surface
95 conditions were warm and very dry.

The third frame, 39320E, as shown in *Figure 1* and *Figure 4*, crosses the central Pacific Ocean, near Hawaii, on 24-
June-2015. It is referred to as the *Hawaii frame*. GOES-15 imagery at 00h00 UTC on 24-June-2015 indicates that the
central portion of the frame bisected a mesoscale convective system (MCS). North and south of the MCS, skies were
mostly cloudless with some broken cloud at variable altitudes. There was also a weak frontal system at its southern
100 extremity.

3. NWP model set-up

The NWP model used to produce EarthCARE's test frames was Environment and Climate Change Canada's GEM
model (Côté et al., 1998; Girard et al., 2014). GEM's dynamics are formulated in terms of the non-hydrostatic
primitive equations with a terrain-following hybrid vertical grid. It can be run as a global model or a limited-area
105 model and is capable of one-way self-nesting. For this work, GEM ran with four nested domains at horizontal grid-
spacings Δx of 10, 2.5, 1, and 0.25 km, with 79 hybrid levels for the 10 km outer-domain and 57 for the other three.

The simulations at Δx of 2.5, 1 and 0.25 km used Milbrandt and Yau's (2005a,b) double-moment bulk cloud micro-
physics scheme (referred to hereinafter as MY2), which predicts mass and number mixing ratio for each of six
hydrometeors classes: non-precipitating liquid droplets; ice crystals; rain; snow; graupel; and hail. For the $\Delta x = 10$



110 km domain, the Kain–Fritsch (KF) deep convection scheme (Kain and Fritsch 1990; 1993) was used. Its liquid and ice CWCs are passed later to MY2 as non-precipitating liquid droplet and ice crystal categories.

In addition to MY2 and KF, a planetary boundary-layer scheme can also produce liquid and ice clouds along with fractional cloudiness for cumulus and stratocumulus clouds (Bélair et al. 2005). Moreover, a shallow convection scheme (Bélair et al. 2005) also supplies estimates of liquid and ice CWCs and cloud fractions for cells with shallow
115 cumulus.

It was simplest to align GEM’s *computational equator* approximately along EarthCARE’s orbit, and divide 6,200 km long frames into 13 inner-domains ($\Delta x = 0.25$ km) and run them separately: 11 segments at 500 km along-track and both end segments at 350 km (all are 200 km wide). The downscaling transitional domains at Δx of 2.5 km and 1 km adapt themselves to the locations of the $\Delta x = 0.25$ km domains. A common $\Delta x = 10$ km domain was used for
120 all 13 segments. *Figure 5* illustrates this configuration, and *Table 3* summarizes domain sizes and Δx . Finally, the 13 inner-domains are simply concatenated to form 6,200 km frames. While this forms discontinuities, they are not a serious hindrance for the task at hand.

Simulations for the *Halifax frame* (39316D) were initialized at 12h00 UTC on 07-Dec-2014 and saved at 17h30 UTC. Likewise, the *Baja frame* (39318D) simulations were initialized at 12h00 UTC 02-Apr-2015 and saved at
125 21h00 UTC, while the *Hawaii frame* was initialized at 12:00 UTC 23-Jun-2015 and saved at 00h00 UTC on 24-Jun-2015. Data for all three frames are freely available. Files include CWC, number concentration, and effective radius R_{eff} for the six aforementioned hydrometeor types. Saved variables are listed in the Appendix.

4. Shortwave optical properties for land surfaces

GEM’s snow-free surface albedos were replaced by those based on MODIS’s MCD43GF 1 km resolution bidirec-
130 tional reflectance distribution function (BRDF) product for the period 2002 to 2013 (Schaaf et al. 2002). These data were interpolated, via nearest neighbour, to 0.25 km. Conditions for the *Baja frame* are shown here because it is primarily over land; the others are mostly over ocean.

Figure 6a illustrates the wide range of surface conditions for the *Baja frame*. The Rooky Mountains are crossed near 40°N, but this being mid-springtime only small amounts of mountain snow remain. *Figure 6b* highlights the distri-



135 bution of freshwater lakes in the Canadian Shield; *Figure 6c* shows that most are frozen and snow-covered. From
Figure 6c it is clear that shallow snow covers most of the Canadian Prairies with deeper snow north of the tree-line
(see 0.63 μm reflectances in *Figure 3*).

Following Schaaf et al. (2002), spectral-dependent black-sky albedos are defined as

$$\alpha_{bs}(\theta_0) = \alpha_1 + \alpha_2(-0.007574 - 0.070987\theta_0^2 + 0.307588\theta_0^3) + \alpha_3(-1.284909 - 0.166314\theta_0^2 + 0.041840\theta_0^3), \quad (1)$$

140 where θ_0 is solar zenith angle in radians, and α_1 , α_2 , and α_3 are separate sets of spectral-dependent BRDF kernel
weights for spectral ranges 300 - 700 nm and 700 - 50,000 nm. Corresponding white-sky albedos are defined as

$$\alpha_{ws} = \alpha_1 + 0.189184\alpha_2 - 1.377622\alpha_3. \quad (2)$$

Figure 7 shows α_{ws} for the *Baja frame*. Note that because these are snow-free values, they tend to be largest in the
southern areas; especially in the near-IR for forests of western Colorado and deserts of Arizona and Sonora. With
145 both variable surface elevation and surface albedos, this frame represents a stringent test for retrieval algorithms, as
opposed to the more straightforward ocean surfaces that dominate the other two frames.

For snow/ice covered areas, snow depth and ice fraction and/or surface land type should be used to determine albedo
of snow/ice covered surfaces. For ocean and lakes, wind speed is used to determine surface albedo. To determine
surface emissivity for different types of surface, an additional surface emissivity climatology is used. These are
150 discussed in another paper on broadband radiative quantities (ACM-COM and ACM-RT products) in this special
issue (Cole et al. 2023).

5. Aerosol properties

The ECSIM scene creation process requires 3D distributions of aerosol size distributions. As GEM lacks interactive
aerosol tracers and chemistry, aerosol fields were added to the test scenes using information from the Copernicus
155 Atmosphere Monitoring Service (CAMS) (Flemming et al. 2017; Inness et al. 2019). The aerosol scheme imple-
mented within ECSIM follows the *Hybrid End-To-End Aerosol Classification* (HETEAC) approach of defining a



certain set of basic aerosol types that, when weighted and summed, yield adequate representations of a wide range of observed aerosol optical properties (Wandinger et al. 2016; Wandinger et al. 2023). *Table 4* lists the CAMS aerosol fields, and the *Supplementary Material* section provides a detailed description of the mapping between CAMS fields and ECSIM scattering types. It also provides more details regarding aerosol representation.

6. Results: GEM simulations and verification

The purpose of this section is to show selected results that characterize the EarthCARE test frames produced directly by GEM. Post-simulation adjustments were made to ice microphysical properties as described and shown in both section 7 and *Supplementary Material*.

6.1. Halifax frame

Figure 8a and *c* show MODIS spectral fluxes (MYD02HKM product; MCST 2017a) for 0.459 - 0.479 μm and 10.8 - 11.3 μm for the *Halifax frame*. Key cloud-related features are a cold front between 40°N and 45°N, scattered clouds to its south, and mostly overcast conditions to its north. *Figure 8b* and *d* show TOA spectral fluxes for two wavebands, close to MODIS's bands, as simulated by the Rapid Radiative Transfer Model for GCMs (RRTMG - Mlawer et al. 1997; Iacono et al. 2000; 2008) using GEM data. At large-scales, GEM did well with respect to cloud occurrence. For the GEM scenes, discontinuities, stemming from the stitching together of the semi-independent high-resolution inner-domains, are clearly visible across the frontal system. While annoying to look at, they do not pose a serious problem for the task at hand.

Near 38°N GEM's longwave fluxes are significantly less than MODIS's. This is because GEM simulated widespread convection in this area whereas MODIS only observed isolated convective cells. This is also apparent in *Figure 9*, which shows cloudtop altitudes both inferred from MODIS radiances (Platnick et al. 2015) and computed by the MODIS simulator of the Cloud Feedback Model Intercomparison Project (CFMIP) Observation Simulator Package (Bodas-Salcedo et al. 2011; abbreviated as the *COSP simulator*). These differences cannot be explained by the slight time difference between MODIS observation (~17h20 UTC) and GEM simulation time (17h30 UTC). On the other hand, it is common for NWP models to simulate some characteristics of cloud systems quite well yet show temporal/spatial displacements relative to observations (e.g., Qu et al. 2018). The goal when simulating these test scenes



was, however, to produce large, well-resolved tracts of realistic clouds; emphasis on exactly what happened was secondary.

185 *Figure 10b* and *c* show cross-sections of ice CWC inferred from CloudSat radar reflectivities and simulated by GEM, respectively (for the transect indicated in *Figure 8*). While the cross-sections intersect only at latitude 43.6°N, the general forms of the fields agree well, and not just in the immediate vicinity of the intersection point. Unfortunately, CloudSat's retrieval of liquid CWC is problematic (e.g., Li et al. 2018) making a comparison to GEM useless. *Figure 10a* shows the ice water path (IWP) vertical integrals of values in *Figure 10b* and *c*. At and around the intersection point, CloudSat's IWP values are much larger than GEM's, but again, the forms of their curves are fairly similar. The
190 edge of GEM's inner-domain near 43°N is abundantly clear.

6.2. Baja frame

Figure 11 compares MODIS TOA fluxes to those computed by RRTMG acting on GEM data for the *Baja frame*. As with the *Halifax frame*, agreement is generally good, though GEM's fields exhibit some peculiarities. For instance, GEM's fluxes associated with clouds are less variable than MODIS's; especially between 40°N and 50°N. This could
195 be due to both GEM's clouds being simply too homogeneous or RRTMG's use of 1D radiative transfer models (see Barker et al. 2017). Also, the thin high clouds near latitude 32°N, which are also evident in *Figure 12* and positioned well in space, show an on-off pattern that is not seen in the observations. Furthermore, near latitude 55°N GEM failed to produce the very low and thin, but extensive, clouds. This is most apparent in *Figure 12*. Note too, that the discontinuities that stem from stitching together GEM's innermost domains are less apparent for this frame than they
200 are for the *Halifax frame*, though the discontinuity near 26°N is notably bad for it stands out in both visible and IR imagery.

Despite these discrepancies, *Figure 13* shows that in the vicinity of where the satellite tracks intersect, vertical realization of clouds by GEM is quite good though the smooth mid-level, yet very low density, cloud is much too extensive. The altitudes of GEM's clouds over the Rooky Mountains are also in fair agreement with CloudSat's.
205 Unlike the *Halifax frame*, the magnitudes of modelled and "observed" IWPs agree quite nicely, in general.



6.3. Hawaii frame

Figure 14 shows that for the *Hawaii frame*, GEM's positionings and approximate intensities of cloud systems near the Equator and $\sim 25^{\circ}\text{S}$ are very good. The harsh discontinuity in GEM's string of innermost domains near 2°S is due to a lack of high ice cloud, as seen in Figure 15, which likely stems from the lack of information, in the form of reduced outflow of high cirrus, coming into the sub-domain from the equatorial mesoscale system. Likewise, near 15°N the lack of upper-level cloud in GEM could be because this sub-domain was too disconnected from the mesoscale system to the south. Again, however, the point of this section is to show the gross verisimilitude of the test frames and hence their suitability for EarthCARE algorithm assessments.

As Figure 16 shows, despite CloudSat's track intersecting EarthCARE's well south of the mesoscale system situated near the centre of the *Hawaii frame*, the system was sufficiently large in the zonal direction that CloudSat's sampling of it can be compared to EarthCARE's sampling of GEM's simulation. The regions of high ice CWC values for the two samples match extremely well both vertically and horizontally. The distribution of ice CWC inferred from CloudSat reflectivities is very narrow while GEM's is much broader with many extremely small values (10^{-4} to 10^{-5} g m^{-3}) that are below the detection threshold of COSP. Aside from the huge spike in IWP for GEM near 3°N , which obviously included some precipitation, the magnitude and forms of the curves for CloudSat and GEM agree well.

What might appear to be a deficiency with GEM is the extreme lack of texture in the visible reflectance of cloud associated with the frontal system in the south of frame. As with the other frames, however, it is entirely likely that the smoothness GEM's field stems from application of a 1D radiative transfer model (see Barker et al. 2017). This is addressed explicitly in other papers in this special issue (Cole et al. 2023).

7. Alterations of GEM's ice crystal sizes

As GEM's scenes are to be input to ECSIM (Voors et al. 2007) to simulate synthetic L1-level measurements for ATLID, CPR, MSI, and BBR (Donovan et al. 2023), it is important that not only macrophysical cloud properties be realistic, for phenomena such as solar RT and CPR nonuniform beam filling (Tanelli et al. 2002), but cloud microphysical properties, such as size distributions and mass-diameter relationships, have to be, too. While the macrophysical cloud properties simulated by GEM were deemed satisfactory by algorithm development groups (see previous section), it was clear that there were shortcomings with its predicted ice cloud microphysical properties (cf. Qu et al.



2018). These deficiencies had a demonstrably negative impact on the realism of ECSIM's simulated measurements, and so adjustments to ice particle sizes were needed.

Basically, GEM predicts too many overly small ice crystals with $R_{eff} < 10 \mu\text{m}$. The cause of this appears to be overestimation of ice crystal number concentrations near cloudtops. Currently, secondary ice production (SIP) mechanisms are poorly understood (Field et al., 2017; Korolev and Leisner 2020), and while at least 6 SIP mechanisms are known (Korolev et al. 2020), only the Hallett-Mossop process (Hallett and Mossop 1974, Mossop and Hallett 1974) is parametrized in the MY2 scheme. It appears as though ice number concentrations in GEM's simulations are systematically underestimated near, or just above, the melting layer. Hence, cloud glaciation times will be too long, and an excess of liquid droplets will be sent too high by updrafts. In the current scheme, droplets will eventually be converted into ice crystals via homogenous freezing, and this will produce very high concentrations of small crystals at altitude. While ongoing studies aim to improve representations of SIP (e.g., Huang et al., 2021, Qu et al. 2022a), they are not yet ready for use in GEM. As such, more manual alterations to GEM's ice crystal sizes were needed.

To improve the realism of the synthetic observations, the following adjustments were made to the GEM's original fields:

1. Implicit liquid CWCs (see *Table 5*) at temperatures $< 273 \text{ K}$ were set to zero thereby reducing unrealistically large amounts of super-cooled droplets;
2. Implicit ice CWCs (see *Table 5*) were deleted because crystal R_{eff} were artificially fixed at $15 \mu\text{m}$;
3. The mass-dimension relationships used by GEM for ice and snow were replaced by those described in Erfani and Mitchell (2006) and the functional form of ice particle size D distribution was changed. Specifically, it was altered by multiplying by a factor of D^4 which had the effect of increasing mean D . Following these adjustments, particle number densities were recalculated subject to conservation of GEM's original total ice and snow water contents.

These alterations were found to produce significant, albeit from a qualitative perspective, improvements to GEM's simulated cloud properties. For example, considering the issue of too many, too small ice crystals, *Figure 17* shows



the relationship between cloud and ice particle R_{eff} before and after the adjustments listed above. It can be seen that the population of small crystals at temperatures above 245 K has been eliminated, whilst below 240 K, minimum
260 particle sizes after adjustments exceed 10 μm . Moreover, the distribution of R_{eff} after adjustments is more consistent with the phase-space indicated by real observations (e.g., Donovan and van Lammeren 2001; Wyser 1998). Lines in *Figure 17* are from parametrizations. While many observation-based parametrizations of ice crystal size distribution exist, they exhibit only moderate agreement, and so cannot be used to fully support the credibility of adjusted R_{eff} . It can be concluded, however, with some certainty, that the above adjustments removed unrealistically
265 small ice crystals and that the resulting temperature distribution of R_{eff} is in fair agreement with observations.

Censoring the implicit super-cooled liquid and ice water as well as the adjustment to ice and snow R_{eff} have important consequences for the vertical structure of optical extinction. This can be seen in *Figure 18* where nadir cross-sections of R_{eff} and extinction at 355 nm (the operating wavelength of ATLID) are shown both before and after adjustments were performed. Increases in R_{eff} and droplet extinction for clouds poleward of 50°N and at altitudes
270 below 5 km stem from a combination of removing super-cooled implicit water and increases to ice and snow R_{eff} . The reduction of cloud extinction, especially near cloudtops between 35°N and 45°N is mainly a consequence of increasing R_{eff} of ice particles.

In addition to these improvements in cloud optical properties, the same adjustments were found to improve the realism of cloud properties that are relevant to simulation of CPR observations. For example, after applying the
275 adjustments the relationship between ice CWC and simulated radar reflectivity now falls in phase-space that agrees well with real observations (e.g., Matrosov and Heymsfield 2017; Heymsfield et al. 2005). Various cross-sections of adjusted GEM+CAMS-derived fields can be found in the *Supplementary Material*.

8. Conclusions and data availability

In this day and age, it is difficult to see how a scientifically and technically advanced research satellite could be
280 launched without first having completed a pre-launch *end-to-end* numerical simulation programme that assesses myriad aspects of mission performance and demonstrates the likelihood of achieving the mission's science goals.



Such a programme would begin with simulation of surface-atmosphere conditions that ideally span, and resemble, much of what can be expected to be encountered during the mission. Virtual observations, to be made by the satellite's sensors, are then simulated for the mock surface-atmospheres, and they are, in turn, operated on by retrieval
285 algorithms. This exact end-to-end simulation programme has unfolded, over the past two decades, for the EarthCARE satellite mission (ESA 2001; Illingworth et al. 2015). The purpose of this paper was to summarize the surface-atmosphere test datasets.

The synthetic surface-atmosphere systems used for this study were produced by ECCO's Global Environment Multiscale model, which is abbreviated as "GEM" (Côté et al., 1998, Girard et al., 2014). This operational numerical
290 weather prediction model is well-known internationally (Leroy et al., 2014, 2022; Bélair et al., 2016; Milbrandt et al., 2016; Qu et al., 2018, 2020, 2022a; , McTaggart-Cowan et al. 2019). The end-to-end programme was intended initially to test retrieval algorithm performance, but was expanded to address ESA's communication and data handling segments, too. As such, large simulated domains were required. The fundamental data processing element is referred to as a *frame*. There are six frames per orbit, and so simulated "test frames" had to be ~6,200 km along-
295 track. The across-track swath of EarthCARE's multi-spectral imager (MSI) is 150 km, and so test frames had to be at least this wide, but to avoid edge-effects, their widths were extended to 200 km. Horizontal and vertical resolutions for GEM's simulations had to allow for at least some variability within the foot-prints of EarthCARE's sensors. Use of 57 vertical layers and horizontal grid-spacing Δx of 0.25 km were deemed adequate (see Qu et al. 2018).

Three test frames, that followed orbits provided by ESA, were identified via examination of satellite data and available
300 surface weather observations. Conditions that were captured include: a cold frontal system, broken shallow cumulus, a tropical mesoscale convective system, thin cirrus, multi-layer clouds; clear-sky conditions, with aerosols, over ocean, land (including mountains), and ice/snow surfaces. Surface bidirectional reflection distribution functions (BRDFs) and albedos from climatological data and aerosol properties were added to GEM's simulations. GEM's "computational equator" was oriented along each EarthCARE orbit and a set of nested simulations were performed
305 that culminated in 13, separately simulated, innermost domains at $\Delta x = 0.25$ km. These domains were concatenated to form the full 6,200 km test frames.

It was discovered that GEM's descriptions of some ice cloud properties lack the realism needed for adequate simulation of virtual observations and assessment of cloud and aerosol property retrieval algorithms. Hence, modifications



were made to the effective size of ice particles based on surface and *in-situ* observations. Most of the important
310 impacts were to particle sizes near cloudtops.

The test frames produced for this study have the potential to be used for other observation missions that could in-
clude platforms other than satellites; particularly those targeting cloud, aerosol, and radiation interactions. The
dataset is available publicly as summarized in the *Data availability* statement at the end. It includes atmospheric and
surface properties produced by GEM data, modified hydrometeor properties, climatological surface optical proper-
315 ties, and the properties of added aerosols (cf. tables in the Appendix).



Acknowledgements

This study was made possible by a series of contracts from the European Space Agency. The authors thank Jason Milbrandt, Sylvie Leroyer, Stéphane Bélair, and Manon Faucher for technical help and discussions.

320

Data availability

Halifax frame's data are available at Qu et al. 2022b,c. Baja frame's data are available at: Qu et al. 2022d,e. Hawaii frame's data are available at: Qu et al. 2022f,g,h.

325 Financial support

This study is supported by Clouds, Aerosol, Radiation - Development of INtegrated ALgorithms (CARDINAL) for the EarthCARE Mission.

Appendix: List of test frame variables

330 Please see Table 5 and 6 at the end of this manuscript.



References

- Barker, H. W., Z. Qu, S. Belair, S. Leroyer, J. A. Milbrandt, P. A. Vaillancourt, 2017: Scaling Properties of Observed and Simulated Satellite Visible Radiances. *J. Geophys. Res.*, **122**, 9413–9429, doi:10.1002/2017JD027146.
- 335 Belair, S., Mailhot, J., Girard, C., and Vaillancourt, A. P., 2005: Boundary layer and shallow cumulus clouds in a medium-range forecast of a large-scale weather system, *Mon. Weather Rev.*, **133**, 1938–1960.
- , Leroyer, S., Seino, N., Spacek, L., Souvranlasy, V., and Paquin-Ricard, D., 2017: Role and impact of the urban environment in the numerical forecast of an intense summertime precipitation event over Tokyo, *J. Meteorol. Soc. Jpn. II*, **96**, 77–94.
- 340 Bodas-Salcedo, A., Webb, M. J., Bony, S., Chepfer, H., Dufresne, J., Klein, S. A., Zhang, Y., Marchand, R., Haynes, J.M., Pincus, R. and John, V.O., 2011: COSP: satellite simulation software for model assessment. *Bull. Amer. Meteor. Soc.*, **92**, 1023–1043.
- Cole, J. N. S., H. W. Barker, Z. Qu, N. Villifranque, and M. W. Shephard, 2023: Broadband Radiative Quantities for the EarthCARE Mission: The ACM-COM and ACM-RT Products. *Atmos. Meas. Tech.*,... this issue.
- 345 Côté, J., Gravel, S., Méthot, A., Patoine, A., Roch, M., and Staniforth, A., 1998: The operational CMC–MRD global environmental multiscale (GEM) model. Part I: Design considerations and formulation, *Mon. Weather Rev.*, **126**, 1373–1395.
- Donovan, D. P., and van Lammeren, A. C. A. P., 2001: Cloud effective particle size and water content profile retrievals using combined lidar and radar observations: 1. Theory and examples, *J. Geophys. Res.*, **106**(D21), 27425–27448, doi:10.1029/2001JD900243.
- 350 —, and —, 2002: First ice cloud effective particle size parameterization based on combined lidar and radar data, *Geophys. Res. Lett.*, **29**(1), doi:10.1029/2001GL013731.
- , P. Kollias, and G-J van Zadelhoff, 2023: The Generation of EarthCARE L1 Test Data sets Using Atmospheric Model Data Sets, *This issue*.
- 355 Girard, C., Desgagné, M., McTaggart-Cowan, R., Côté, J., Charron, M., Gravel, S., Lee, V., Patoine, A., Qaddouri, A., Roch, M., Spacek, L., Tanguay, M., Vaillancourt, P. A., and Zadra, A., 2014: Staggered vertical discretization of the Canadian environmental multiscale (GEM) model using a coordinate of the log-hydrostatic-pressure type, *Mon. Weather Rev.*, **142**, 1183–1196.
- Erfani, E. and Mitchell, D. L., 2016: Developing and bounding ice particle mass- and area-dimension expressions for use in atmospheric models and remote sensing, *Atmos. Chem. Phys.*, **16**, 4379–4400, https://doi.org/10.5194/acp-16-4379-2016.
- 360 ESA., 2001: *The Five Candidate Earth Explorer Missions: EarthCARE –Earth Clouds, Aerosols and Radiation Explorer*, ESA SP-1257(1), September 2001. ESA Publications Division: Noordwijk, The Netherlands.
- Flemming, J., Benedetti, A., Inness, A., Engelen, R. J., Jones, L., Huijnen, V., Remy, S., Parrington, M., Suttie, M., Bozzo, A., Peuch, V.-H., Akritidis, D., and Katragkou, E., 2017: The CAMS interim Reanalysis of Carbon Monoxide, Ozone and Aerosol for 2003–2015, *Atmos. Chem. Phys.*, **17**, 1945–1983, https://doi.org/10.5194/acp-17-1945-2017.
- 365 Hallett, J. and Mossop, S., 1974: Production of secondary ice particles during the riming process, *Nature*, **249**, 26–28.
- Heymsfield, A. J., Wang, Z., and Matrosov, S., 2005: Improved Radar Ice Water Content Retrieval Algorithms Using Coincident Microphysical and Radar Measurements, *J. Appl. Meteorol.*, **44**(9), 1391–1412.
- 370 Huang, Y., Wu, W., McFarquhar, G. M., Wang, X., Morrison, H., Ryzhkov, A., Hu, Y., Wolde, M., Nguyen, C., Schwarzenboeck, A., Milbrandt, J., Korolev, A. V., and Heckman, I., 2021b: Microphysical processes producing high ice water contents (HIWCs) in tropical convective clouds during the HAIC-HIWC field campaign: evaluation of simulations using bulk microphysical schemes, *Atmos. Chem. Phys.*, **21**, 6919–6944, https://doi.org/10.5194/acp-21-6919-2021.



- Iacono, M. J., Mlawer, E. J., Clough, S. A., and Morcrette, J.-J., 2000: Impact of an improved longwave radiation model, RRTM. On the energy budget and thermodynamic properties of the NCAR community climate mode, CCM3, *J. Geophys. Res.*, **105**, 14873–14890.
- 380 —, Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D., 2008: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res.*, **113**, D13103, <https://doi.org/10.1029/2008JD009944>.
- Illingworth, A. J., Barker, H. W., Beljaars, A., Ceccaldi, M., Chepfer, H., Clerbaux, N., et al., 2015: The EarthCARE satellite: The next step forward in global measurements of clouds, aerosols, precipitation, and radiation. *Bull. Am. Met. Soc.*, **96**, 1311–1332.
- 385 Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., and Suttie, M., 2019: The CAMS reanalysis of atmospheric composition, *Atmos. Chem. Phys.*, **19**, 3515–3556, <https://doi.org/10.5194/acp-19-3515-2019>.
- 390 Kain, J. S. and Fritsch, J. M., 1990: A one-dimensional entraining/detraining plume model and its application in convective parameterization, *J. Atmos. Sci.*, **47**, 2784–2802.
- and —, 1993: Convective parameterization for mesoscale models: The Kain–Fritsch scheme. The Representation of Cumulus Convection in *Numerical Models*, *Meteor. Monogr.*, **24**, Amer. Meteor. Soc., 165–170.
- Korolev, A., Heckman, I., Wolde, M., Ackerman, A. S., Fridlind, A. M., Ladino, L. A., Lawson, R. P., Milbrandt, J., and Williams, E., 2020: A new look at the environmental conditions favorable to secondary ice production, *Atmos. Chem. Phys.*, **20**, 1391–1429.
- 395 — and Leisner, T., 2020: Review of experimental studies of secondary ice production, *Atmos. Chem. Phys.*, **20**, 11 767–11 797.
- Leroyer, S., Bélair, S., Husain, S., and Mailhot, J., 2014: Subkilometer numerical weather prediction in an urban coastal area: a case study over the Vancouver metropolitan area, *J. Appl. Meteorol. Clim.*, **53**, 1433–1453, <https://doi.org/10.1175/JAMC-D-13-0202>.
- 400 —, —, Souvanlasy, V., Vallée, M., Pellerin, S. and Sills, D., 2022: Summertime Assessment of an Urban-Scale Numerical Weather Prediction System for Toronto. *Atmosphere*, **13**, 1030. <https://doi.org/10.3390/atmos13071030>.
- Matrosov, S. Y., and A. J. Heymsfield, 2017: Empirical relations between size parameters of ice hydrometeor populations and radar reflectivity. *J. Appl. Meteorol. Clim.*, **56**, 2479–2488. doi:10.1175/JAMC-D-17-0076.1
- 405 Li, J. F., S. Lee, H.-Y. Ma, G. L. Stephens, and B. Guan, 2018: Assessment of the cloud liquid water from climate models and reanalysis using satellite observations. *Terr. Atmos. Ocean. Sci.*, **29**, 653–678, doi: 10.3319/TAO.2018.07.04.01
- McTaggart-Cowan, R., Vaillancourt, P. A., Zadra, A., Chamberland, S., Charron, M., Corvec, S., Milbrandt, J. A., Paquin-Ricard, D., Patoine, A., Roch, M., Separovic, L., and Yang, J., 2019: Modernization of atmospheric physics parameterization in Canadian NWP. *J. Adv. Model. Earth Syst.*, **11**, 3593–3635, <https://doi.org/10.1029/2019MS001781>.
- Milbrandt, J. A. and Yau, M. K., 2005a: A multi-moment bulk microphysics parameterization. Part I: Analysis of the role of the spectral shape parameter, *J. Atmos. Sci.*, **62**, 3051–3064, <https://doi.org/10.1175/JAS3534.1>.
- 415 — and Yau, M. K., 2005b: A multi-moment bulk microphysics parameterization. Part II: A proposed three-moment closure and scheme description, *J. Atmos. Sci.*, **62**, 3065–3081, <https://doi.org/10.1175/JAS3535.1>.
- , Bélair, S., Faucher, M., Vallée, M., Carrera, M. L., and Glazer, A., 2016: The pan-Canadian high resolution (2.5km) deterministic prediction system, *Weather Forecast.*, **31**, 1791–1816, <https://doi.org/10.1175/WAF-D-16-0035.1>.
- 420 Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A., 1997: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.*, **102**, 16663–16682.



- MODIS Characterization Support Team (MCST), 2017a. MODIS 500m Calibrated Radiance Product. NASA MODIS Adaptive Processing System, Goddard Space Flight Center, USA: <http://dx.doi.org/10.5067/MODIS/MYD0HKM.061>
- 425 MODIS Characterization Support Team (MCST), 2017b. MODIS 1km Calibrated Radiances Product. NASA MODIS Adaptive Processing System, Goddard Space Flight Center, USA: <http://dx.doi.org/10.5067/MODIS/MYD021KM.061>
- Mossop, S. C. and J. Hallett, 1974: Ice Crystal Concentration in Cumulus Clouds: Influence of the Drop Spectrum, *Science*, **186**, 632–634.
- 430 Platnick, S., Ackerman, S., King, M., et al., 2015. MODIS Atmosphere L2 Cloud Product (06_L2). NASA MODIS Adaptive Processing System, Goddard Space Flight Center, USA: http://dx.doi.org/10.5067/MODIS/MYD06_L2.061
- Qu, Z., H. W. Barker, A. V. Korolev, J. A. Milbrandt, M. Wolde, A. Schwarzenböck, D. Leroy, J. W. Strapp, J. N. S. Cole, L. Nguyen, and A. Heidinger, 2018: Evaluation of a high-resolution NWP model's simulated clouds using observations from CloudSat and in situ aircraft. *Q. J. R. Meteorol. Soc.*, **144**, 1681–1694. doi: 10.1002/qj.3318.
- 435 —, Huang, Y., Vaillancourt, P. A., Cole, J. N. S., Milbrandt, J. A., Yau, M.-K., Walker, K., and de Grandpré, J., 2020a: Simulation of convective moistening of the extratropical lower stratosphere using a numerical weather prediction model, *Atmos. Chem. Phys.*, **20**, 2143–2159, <https://doi.org/10.5194/acp-20-2143-2020>.
- , Korolev, A., Milbrandt, J. A., Heckman, I., Huang, Y., McFarquhar, G. M., Morrison, H., Wolde, M., and Nguyen, C., 2022a: The impacts of secondary ice production on microphysics and dynamics in tropical convection, *Atmos. Chem. Phys.*, **22**, 12287–12310, <https://doi.org/10.5194/acp-22-12287-2022>.
- 440 —, Donovan, D. P., Barker, H. W., Cole, J. N. S., Shephard, M. W., Huijnen, V., 2022b: Numerical Model Generated Halifax Test Scenes for EarthCARE Pre-launch Studies - Part 1: Atmospheric and Surface Properties, *Zenodo*, <https://doi.org/10.5281/zenodo.7258361>.
- , —, —, —, —, and —, 2022c: Numerical Model Generated Halifax Test Scenes for EarthCARE Pre-launch Studies - Part 2: Hydrometeor and Aerosol Properties, *Zenodo*, <https://doi.org/10.5281/zenodo.7254610>.
- 445 —, —, —, —, —, and —, 2022d: Numerical Model Generated Baja Test Scenes for EarthCARE Pre-launch Studies - Part 1: Atmospheric and Surface Properties, *Zenodo*, <https://doi.org/10.5281/zenodo.719623>.
- , —, —, —, —, and —, 2022e: Numerical Model Generated Baja Test Scenes for EarthCARE Pre-launch Studies - Part 2: Hydrometeor and Aerosol Properties, *Zenodo*, <https://doi.org/10.5281/zenodo.7255530>.
- 450 —, —, —, —, —, and —, 2022f: Numerical Model Generated Hawaii Test Scenes for EarthCARE Pre-launch Studies - Part 1: Atmospheric and Surface Properties, *Zenodo*, <https://doi.org/10.5281/zenodo.7196690>.
- , —, —, —, —, and —, 2022g: Numerical Model Generated Hawaii Test Scenes for EarthCARE Pre-launch Studies - Part 2: Hydrometeor and Aerosol Properties, *Zenodo*, <https://doi.org/10.5281/zenodo.7255758>.
- 455 —, —, —, —, —, and —, 2022h: Numerical Model Generated Hawaii Test Scenes for EarthCARE Pre-launch Studies - Part 3: Additional Data, *Zenodo*, <https://doi.org/10.5281/zenodo.7255800>.
- Schaaf, CB., Gao, F., Strahler, AH., Lucht, W., Li, XW., Tsang, T., Strugnell, NC., Zhang, XY., Jin, YF., Muller, JP., Lewis, P., Barnsley, M., Hobson, P., Disney, M., Roberts, G., Dunderdale, M., Doll, C., d'Entremont, RP., Hu, BX., Liang, SL., Privette, JL., Roy, D., 2002: First operational BRDF, albedo nadir reflectance products from MODIS. *Remote Sensing of Environment*, **83** (1-2), Article PII S0034-4257(02)00091-3. 10.1016/S0034-4257(02)00091-3.
- 460 Tanelli, S., E. Im, S. L. Durden, L. Facheris, and D. Giuli, 2002: The effects of nonuniform beam filling on vertical rainfall velocity measurements with a spaceborne Doppler radar. *J. Atmos. Ocean. Tech.*, **19**, 1019–1034.
- Voors, R., D. P. Donovan, J. Acarreta, M. Eisinger, R. Franco, D. Lajas, R. Moyano, F. Pirondini, J. Ramos, and T. Wehr, 2007: ECSIM: The simulator framework for EarthCARE, *Proc. SPIE 6744, Sensors, Systems, and Next-Generation Satellites XI*, 67441Y (26 October 2007); <https://doi.org/10.1117/12.737738>
- 465 Wandinger, U., H. Baars, R. Engelmann, A. Hünerbein, S. Horn, T. Kanitz, D. Donovan, G.-J. van Zadelhoff, D. Daou, J. Fischer, J. von Bismarck, F. Filipitsch, N. Docter, M. Eisinger, D. Lajas, and T. Wehr, 2016:



“HETEAC: The Aerosol Classification Model for EarthCARE”, *EPJ Web of Conferences* **119** 01004 (2016), DOI: 10.1051/epjconf/201611901004

470 Wyser, K., 1998: The Effective Radius in Ice Clouds, *J. Climate*, **11**(7), 1793-1802.

Table 1: Surface conditions (observed / modelled) near the *Halifax frame* (see orbit 39316D in *Figure 1*) for 7-December-2014 at 18h00 UTC

station	Temperature (°C)	Dew Point (°C)	Pressure (hPa)	Visibility (km)	Wind Direction	Wind Speed (km/h)	Conditions
Kangerlussaq	-15.0 / -16.3	-20.0 / -18.9	1006 / 1006	-	E	18.5	Mostly cloudy
Nuuk	-4.0 / -2.6	-13.0 / -6.1	1000 / 1000	-	ESE	33.3	Low Drifting Snow + Snow
Hopedale	-17.6 / -15.6	-23.9 / -21.1	1017 / 1022	-	SW	31.0	-
Goose Bay	-16.0 / -15.5	-24.0 / -18.6	1021 / 1024	24.1	WNW	22.2	Scattered Clouds
Charlottetown	-6.0 / -4.3	-11.0 / -8.8	1029 / 1029	24.1	NW	27.8	Mostly Cloudy
Halifax	-3.0 / -2.7	-6.0 / -7.2	1026 / 1028	24.1	NNW	31.5	Overcast
Bermuda	24.0 / 22.8	18.0 / 18.5	1009 / 1010	-	N	24.1	Scattered Clouds
Punta Cana	29.0 / 26.6	22.0 / 20.7	1012 / 1014	-	NE	18.5	Scattered Clouds



475 **Table 2:** As in *Table 1* but these are for the *Baja frame* (see orbit 39318D in *Figure 1*) for 2-April-2015 at 21h00 UTC

station	Temperature (°C)	Dew Point (°C)	Pressure (hPa)	Visibility (km)	Wind Direction	Wind Speed (km/h)	Conditions
Gjoa Haven	-26.0 / -24.6	-29.0 / -27.9	1024 / 1025	24.1	NNW	18.5	Ice Crystals
Baker Lake	-27.0 / -25.3	-31.0 / 29.0	1018 / 1022	4.8	N	40.7	SnowBlowing + Snow
Ennadai	-28.2 / -27.2	-32.2 / 31.6	1025 / 1028	-	NNW	50	Blowing
Key Lake	-11.0 / -12.1	-22.0 / 18.3	1025 / 1026	14.5	N	13	Clear
Saskatoon	0.0 / 1.0	-10.0 / -4.6	1024 / 1022	24.1	NE	11.1	Mostly Cloudy
Billings	7.2 / 8.2	-12.8 / 13.4	1021 / 1020	16.1	NW	25.9	Scattered Clouds
Big Piney	1.7 / 1.3	-16.1 / 11.1	1017 / 1022	16.1	NW	13	Overcast
Provo	8.0 / 0.0	-6.0 / -18.6	1019 / 1024	24.1	WNW	18.5	Mostly Cloudy
Page	18.9 / 18.6	-15.6 / -7.7	1008 / 1028	16.1	W	35.2	Clear
Phoenix	28.9 / 28.2	0.6 / 1.8	1010 / 1014	16.1	WNW	22.2	Mostly Cloudy
Hermosillo	34.0 / 30.5	2.0 / 3.3	1012 / 1011	16.1	SSW	29.6	Scattered Clouds
La Paz	31.0 / 21.6	9.0 / 15.9	1013 / 1014	16.1	W	11.1	Mostly Cloudy

Table 3: Sizes of GEM's downscaling domains (see *Figure 5*) and their horizontal resolutions

description	domain size (km) along-track x across-track	number of vertical layers	horizontal grid-spacing (km)
downscaling domain 1	8,600 x 3,600	79	10
downscaling domain 2	2,250 x 1350	57	2.5
downscaling domain 3	1000 x 600	57	1
innermost domains 1 and 13	350 x 200	57	0.25
innermost domains 2 to 12	500 x 200	57	0.25



480

Table 4: Aerosol classes from the Copernicus Atmosphere Monitoring Service (CAMS)

aerosol class	description
DD1-DD3	Dust (in different size intervals)
SS1-SS3	Sea Salt (in different size intervals)
SO4	Sulphate aerosol
BCB	Fine mode strongly absorbing aerosol
OMB	Weakly absorbing aerosol



Table 5: Variables for the original GEM simulations

Variables name	Units	Dimension	Notes
water_content_cloud	g m^{-3}	3	Explicit Cloud: From MY2 double-moment (MY2) scheme. It's an explicit scheme. The cloud optical properties could be calculated for each of the six species. There also could be properties from the implicit schemes such as the Planetary Boundary Layer (PBL) and shallow convection (SC) scheme (see below). Their optical properties could be calculated in a similar way to those for MY2 species. The final optical properties for use should be the combination of those of the concerned MY2 species and of the implicit clouds.
water_content_ice	g m^{-3}	3	
water_content_rain	g m^{-3}	3	
water_content_snow	g m^{-3}	3	
water_content_graupel	g m^{-3}	3	
water_content_hail	g m^{-3}	3	
number_concentration_cloud	$\# \text{ m}^{-3}$	3	
number_concentration_ice	$\# \text{ m}^{-3}$	3	
number_concentration_rain	$\# \text{ m}^{-3}$	3	
number_concentration_snow	$\# \text{ m}^{-3}$	3	
number_concentration_graupel	$\# \text{ m}^{-3}$	3	
number_concentration_hail	$\# \text{ m}^{-3}$	3	
effective_radius_cloud	m	3	
effective_radius_ice	m	3	
effective_radius_rain	m	3	
effective_radius_snow	m	3	
effective_radius_graupel	m	3	
effective_radius_hail	m	3	
implicit_cloud_solid_water_content	g m^{-3}	3	Implicit Cloud: Cloud condensates from implicit schemes (PBL+SC). R_{eff} for solid condensate is assumed to be 15 microns.
implicit_cloud_liquid_water_content	g m^{-3}	3	
implicit_cloud_liquid_effective_radius	m	3	
BRDF_iso	[0 1]	2 (8 bands)	Snow-free ground albedo climatology from MCD43GF. Ross-Li model (see Eq. 1 and 2). Add information of snow (X-MET, snow_depth etc.).
BRDF_vol	[0 1]	2 (8 bands)	
BRDF_geo	[0 1]	2 (8 bands)	
height_thermodynamic	m	3	Levels for all the 3D variables except those concerning the horizontal wind.
pressure_thermodynamic	Pa	3	
height_momentum	m	3	Levels for the variables concerning the horizontal wind (speed and direction).
pressure_momentum	Pa	3	
temperature	K	3	
specific_humidity	g m^{-3}	3	
relative_humidity	[0 1]	3	
wind_horizontal_speed	m s^{-1}	3	On thermodynamic levels
wind_horizontal_direction	deg	3	0: north, clockwise, on thermodynamic levels
wind_vertical_speed	m s^{-1}	3	
cloud_mask_3d	0/1	3	
cloud_mask_2d	0/1	2	
orography	m	2	
solar_zenithal_angle	deg	2	
surface_pressure	Pa	2	
water_land_fraction	[0 1]	2	1: 100% lake/sea/ocean, 0: 100% land
surface_temperature	K	2	Sea ice not included
ice_fraction	[0 1]	2	Includes sea ice + land ice. To distinguish, use <i>water_land_fraction</i> . Use <i>ice_fraction</i> to apply <i>ice_temperature</i> .
ice_temperature	K	2	
total_water_path	g m^{-2}	2	Liquid cloud water path + ice cloud water path
ice_water_path	g m^{-2}	2	Ice cloud water path
vertical_integrated_water_vapeur	g m^{-2}	2	




snow_depth	m	2	
total_precipitation_rate	m s ⁻¹	2	
liquid_precipitation_rate	m s ⁻¹	2	
longitude	deg	2	
latitude	deg	2	


485

Table 6: Variables archived for the test frames


Variables name		Units	Dimension	Notes
BRDF_iso		[0 1]	2 (8 bands)	Snow-free ground albedo climatology from MCD43GF. Ross-Li model (see Eqs. 1 and 2). NB. Add information of snow (X-MET, snow_depth etc.).
BRDF_vol		[0 1]	2 (8 bands)	
BRDF_geo		[0 1]	2 (8 bands)	
ice_fraction		[0 1]	2	Includes sea ice + land ice. To distinguish, use <i>water_land_fraction</i> . Use <i>ice_fraction</i> to apply <i>ice_temperature</i> .
ice_temperature		K	2	
total_precipitation_rate		m s ⁻¹	2	
liquid_precipitation_rate		m s ⁻¹	2	
orography		m	2	
snow_depth		m	2	
solar_zenithal_angle		deg	2	
surface_pressure		Pa	2	
water_land_fraction		[0 1]	2	
surface_temperature		K	2	
surface_wind_speed		m s ⁻¹	2	
temperature		K	3	
specific_humidity		g m ⁻³	3	
pressure		Pa	3	
wind_horizontal_speed		m s ⁻¹	3	
wind_vertical_speed		m s ⁻¹	3	
mass_content		g m ⁻³	3	For 6 types of hydrometeors: liquid cloud, ice cloud, rain, snow, graupel and hail, and for 4 types of aerosols: coarse dust, coarse salt, fine mode weakly absorbing and fine mode strongly absorbing.
effective_radius		micron	3	
number_concentration		cm ⁻³	3	




490 **Figure 1:** Examples of several successively numbered EarthCARE orbits as provided by ESA. Frames are colour-coded. The test frames are indicated by letters D and E. Frames 39316D, 39318D, and 39320E are referred to as “Halifax”, “Baja”, and “Hawaii”, respectively.




495 **Figure 2:** GOES-13 TOA reflectance at 0.63 μm and brightness temperature at 10.7 μm on 7-December-2014 at 18h00 UTC. Red lines indicate EarthCARE’s track for the *Halifax* frame (see orbit 39316D in Figure 1). Yellow dots mark locations of nearby surface meteorological stations. Blue shaded areas indicate GOES-13’s northern limit of observation.




500 **Figure 3:** As in *Figure 1* but this is GOES-15 imagery for 2-April-2015 at 21h00 UTC. Red lines indicate the *Baja* frame (see orbit 39318D in *Figure 1*).




505 **Figure 4:** As in *Figure 1* but this is GOES-15 imagery for 24-June-2015 at 0h00 UTC. Red lines indicate the *Hawaii* frame (see orbit 39320E in *Figure 1*).




510 **Figure 5:** Downscaling domains for the *Halifax frame* (39316D). Blue rectangles delineate successive downscaling domains that culminate in the 7th innermost domain whose horizontal grid-spacing is $\Delta x = 0.25$ km. Red rectangles are the 12 other innermost domains for this frame. EarthCARE's ground-track is indicated by the purple line, which is 60 km west of centre.




515 **Figure 6:** All panels are for the *Baja frame* (see *Figure 1* and *Figure 3*) and each panel's title is self explanatory. For (b) and (c), blue corresponds to land and yellow to either water or ice.




520 **Figure 7:** (a) Visible white-sky albedos, as defined in (2), for snow-free land for the *Baja frame*. (b) Same as (a) except these are near-IR values. EarthCARE's nadir-track is shown by red lines.




525 **Figure 8:** (a) MODIS TOA flux (approximated by radiance* π) for band 3 (459 - 479 nm) between 17h15 and 17h30 UTC on 7-Dec-2014. (b) RRTMG simulated upward TOA flux for 441.5 - 625 nm for GEM's simulation of the *Halifax frame*. (c) as in (a) but for band 31 (10.8-11.3 μm). (d) in as but for wavelengths 10.2 - 12.2 μm . Solid and dashed yellow lines indicate EarthCARE's and CloudSat's nadir-tracks. Blank areas are outside MODIS's field-of-view.



530 **Figure 9:** (a) MODIS cloudtop pressure (MYD06_L2 product) between 17h15 and 17h30 UTC on 7-Dec-2014. Blank areas are outside MODIS’s field-of-view. (b) GEM’s cloudtop pressure for the *Halifax frame* based on COSP’s MODIS simulator. Grey area in the northern portion has $\theta_0 > 90^\circ$ and so no COSP values.




535 **Figure 10:** (a) Ice water path (IWP) inferred from CloudSat observations at 17h21 UTC on 7-Dec-2014 as it crossed the *Halifax frame* between latitudes 41°N - 44°N (dashed yellow line in *Figure 8*), and as simulated by GEM along the nadir-track (solid yellow lines in *Figure 8*). (b) and (c) are ice CWC for CloudSat and GEM, respectively. Dashed white line indicates where CloudSat’s and EarthCARE’s tracks intersected.



540

Figure 11: As in *Figure 8* except these are for the *Baja frame*.



545

Figure 12: As in *Figure 9* except these are for the *Baja frame*.





Figure 13: As in *Figure 10* except these are for the *Baja frame*. CloudSat’s track indicated by the dashed yellow line in *Figure 11* is between latitudes 45°N - 49°N.



550

Figure 14: As in *Figure 8* except these are for the *Hawaii frame*.




Figure 15: As in Figure 9 except these are for the *Hawaii* frame.

555




Figure 16: As in Figure 10 except these are for the *Hawaii* frame. CloudSat’s track indicated by the dashed yellow line in Figure 14 is between latitudes 1.5°S - 6.5°S.

560




Figure 17: Effective radius R_{eff} as functions of temperature for GEM’s results both before and after application of adjustments discussed in the text. R_{eff} is defined in terms of the mass and cross-sectional area of crystals following



565

Donovan and van Lammeren (2001). Solid lines in the “before” panel correspond to the parametrization described in Wyser (1998), while dotted-lines follow Donovan and van Lammeren (2002). Colours of dots correspond to different ranges of IWC (g m^{-2}): red \rightarrow $\text{IWC} < 0.0001$, green $\rightarrow 0.0001 < \text{IWC} < 0.001$, blue $\rightarrow 0.001 < \text{IWC} < 0.01$, yellow $\rightarrow 0.01 < \text{IWC} < 0.1$, light-blue $\rightarrow 0.1 < \text{IWC}$.




Figure 18: Nadir cross-sections of R_{eff} and 355 nm extinction before and after making adjustments described in the text.

570