

Evaluating the effects of columnar NO₂ on the accuracy of aerosol optical properties retrievals

Theano Drosoglou¹, Ioannis-Panagiotis Raptis^{1,2}, Massimo Valeri³, Stefano Casadio³, Francesca Barnaba⁴, Marcos Herreras-Giralda⁵, Anton Lopatin⁵, Oleg Dubovik⁶, Gabriele Brizzi^{3,7}, Fabrizio Niro⁷,
5 Monica Campanelli⁴, Stelios Kazadzis⁸

¹Institute for Environmental Research and Sustainable Development, National Observatory of Athens (IERSD/NOA), 15236 Athens, Greece

²Laboratory of Climatology and Atmospheric Environment, Sector of Geography and Climatology, Department of Geology and Environment, National and Kapodistrian University of Athens, Athens, GR-15784, Greece

10 ³Serco Italia S.p.A., Frascati, Rome, Italy

⁴National Research Council, Institute of Atmospheric Sciences and Climate, CNR- ISAC, ~~National Research Council~~, Rome, Italy

⁵GRASP SAS, Remote Sensing Developments, 59260 Lezennes, France

⁶Univ. Lille, CNRS, UMR 8518 - LOA - Laboratoire d'Optique Atmosphérique, Lille, France

15 ⁷ESA-ESRIN, Frascati, Rome, Italy

⁸Physicalisch-Meteorologisches Observatorium Davos, World Radiation Center, CH-7260 Davos, Switzerland

Correspondence to: Theano Drosoglou (tdroso@noa.gr)

Abstract. We aim to evaluate the NO₂ absorption effect in aerosol columnar properties, namely the Aerosol Optical Depth (AOD), Ångström Exponent (AE) and Single Scattering Albedo (SSA), derived from sun-sky radiometers as well as the possible retrieval algorithm improvements by using more accurate characterization of NO₂ optical depth from co-located or satellite-based real-time measurements. For this purpose, we employ multiannual (2017-2022) records of Aerosol Optical Depth (AOD), Ångström Exponent (AE) and Single Scattering Albedo (SSA) collected by sun photometers at an urban and a suburban site in the Rome area (Italy) in the framework of both the AERONET and SKYNET networks. The uncertainties introduced in the aerosol retrievals by the NO₂ absorption are investigated using high-frequency observations of total NO₂ derived from co-located Pandora spectroradiometer systems as well as space-borne NO₂ products from the Tropospheric Monitoring Instrument (TROPOMI). For both AERONET and SKYNET, the standard network products were found to systematically overestimate AOD and AE. The average AOD bias found for Rome is relatively low for AERONET (~0.002 at 440 nm and ~0.003 at 380 nm) compared to the retrieval uncertainties, but quite higher for SKYNET (~0.007). On average, an AE bias of ~0.02 and ~0.05 was estimated for AERONET and SKYNET, respectively. In general, the correction seems to be low for areas with low columnar NO₂ concentrations, but it is still useful for lower AODs (< 0.3), where the majority of observations is-are found, especially under high NO₂ pollution events. For the cases of relatively high NO₂ levels (> ~0.7 DU), the mean AOD bias was found within the range 0.009 – 0.012 for AERONET, depending on wavelength and location, and about 0.018 for SKYNET. The analysis does not reveal any significant impact of the NO₂ correction on the derived aerosol temporal trends for the very limited data sets used in this study. However, the effect is expected to become more evident for trends derived from larger data sets as well as in the case of an important NO₂ trend. In addition, the comparisons of the NO₂-

20
25
30
35

modified ground-based AOD data with satellite retrievals from the Deep Blue (DB) algorithm of the NASA Moderate Resolution Imaging Spectroradiometer (MODIS) resulted in a slight improvement in the agreement of about 0.003 and 0.006 for AERONET and SKYNET, respectively. Finally, the uncertainty in assumptions on NO_2 seems to have a non-negligible impact on the retrieved values of SSA at 440 nm leading to an average positive bias of about 0.02 (2.5%) in both locations for high NO_2 loadings (> 0.79 DU).

1 Introduction

Atmospheric particles possess both direct and indirect effects on Earth's radiation budget and climate (IPCC, 2014). Direct radiative forcing arises from aerosols interaction with solar radiation through absorption and scattering processes (Hobbs, 1993). As an indirect impact, aerosols play an important role in cloud formation and properties by acting as cloud condensation nuclei on which water vapour condenses and by influencing the cloud albedo and lifetime (Rosenfeld et al., 2014). Moreover, heterogeneous chemical reactions can take place on the surfaces of atmospheric particles having a crucial effect on atmospheric chemistry and composition. Examples of such aerosol-driven reactions are those that lead to stratospheric ozone depletion in the polar regions (Solomon et al., 1986). In addition to their footprint on radiative forcing and climate, aerosols adversely affect human health and have been associated with a wide variety of health issues such as respiratory and neurological diseases, cancer, diabetes, cardiovascular diseases and hypertension (e.g., Lelieveld et al., 2015; Molina et al., 2020 and references therein).

The above effects of airborne particulate matter on Earth's climate and human health strongly depend on the intra-annual variations in its loading and properties. The most widely used variable for the estimation of columnar aerosol concentration in the atmosphere is the multi-wavelength Aerosol Optical Depth (AOD). Aerosol optical properties are monitored globally by satellite, e.g., the Moderate Resolution Imaging Spectroradiometer (MODIS) and ground-based networks of sun-photometers like the Aerosol Robotic Network (AERONET) (Holben et al., 1998), SKYNET (Nakajima et al., 2020) or the Global Atmosphere Watch Precision Filter Radiometer (GAW-PFR) network (Kazadzis et al., 2018a). Ground-based remote sensing allows accurate AOD retrievals, i.e. in the order of 0.01 – 0.02 depending on the AOD wavelength (Kazadzis et al., 2018b), which are in fact widely used as a validation reference for satellite or model-based AOD products (e.g., Chu et al., 2002; Remer et al., 2005; Green et al., 2009; Levy et al., 2010; Li et al., 2015; Sherman et al., 2016; Gkikas et al., 2021; Di Tomaso et al., 2022) and used as inputs on various modelling initiatives (e.g., Benedetti et al., 2018).

However, AOD retrieval from sun-photometers includes some assumptions in order to take into account all the non-aerosol effects in the retrieval spectral range. In particular, AOD retrievals are sensitive to the assumptions on the concentration of atmospheric trace gases, absorbing in the instrument spectral bands considered, among which are ozone (O_3) and nitrogen dioxide (NO_2). The exact effect of trace gases in the retrieval at a particular bandwidth depends also on their absorption cross-section. For the case of NO_2 , as filter radiometers retrieve the AOD in certain wavelength bands, based on their filter responsivity, such retrievals, especially in the standard wavelengths of 380 and 440 nm (AERONET), have to be corrected for

the NO₂ optical depth. Currently, some AOD retrievals do not take NO₂, ~~i.e. NO₂~~ optical depth, into consideration when deriving AOD (e.g., SKYNET, [Nakajima et al., 2020](#); GAW-PFR, [Kazadzis et al., 2018a](#)) while others use satellite-based climatological NO₂ data sets for estimating it (e.g., AERONET, [Giles et al., 2019](#)). In the case of the GAW-PFR network, the error introduced in AOD retrievals by NO₂ absorption can be assumed to be negligible due to the low NO₂ concentrations observed in the GAW remote stations (the annual mean values of NO₂ optical depth are in general < 0.001) ([Kazadzis et al., 2018a](#)). However, especially over polluted areas, NO₂ is characterized by rather short lifetime and high spatiotemporal variations, due to inhomogeneous local emission patterns and photochemical destruction (e.g., [Richter et al., 2005](#); [Boersma et al., 2008](#); [Tzortziou et al., 2014, 2015](#); [Drosoglou et al., 2017](#); [Fan et al., 2021](#)). Although the stratospheric component of NO₂ is quite stable spatially, the tropospheric NO₂ is highly variable in space and time and can bias the calculation of AOD if neglected ([Arola and Koskela, 2004](#); [Boersma et al., 2004](#)). Hence, areas with high tropospheric NO₂ emission will tend to have greater proclivity for deviating from climatological mean values which might not be representative of the actual NO₂ loading and spatial distribution in the atmosphere, introducing potential errors in AOD calculations in those spectral regions with significant NO₂ absorption footprint.

Satellite observations with improved spatial and temporal resolution, e.g., the Sentinel-5Precursor Tropospheric Monitoring Instrument (S5P/TROPOMI), models, or co-location with surface-based Pandora instruments from Pandora Global Network (PGN) spectroradiometers ([Cede et al., 2020](#)) measuring total column of NO₂ may assist in reducing the uncertainty of the NO₂ optical depth contribution in later versions of AOD retrieval algorithms. In the present study, we aim at evaluating if and how much AOD as well as its spectral variability, i.e., the associated Ångström Exponent (AE), and Single Scattering Albedo (SSA) retrievals could be improved by applying a specific correction using synchronous and co-located measurements of total NO₂ column from Pandora network spectroradiometers. To this purpose, we exploit the unique configuration of twin observational sites in the Rome area (Italy), where multiannual (2017-2022) records of both multispectral AOD observations and columnar NO₂ measurements are available both in the city centre and in a suburban location. High-frequency measurements of total NO₂ performed by co-located Pandora spectroradiometer systems ~~along with relatively high spatially resolved NO₂ observations from the S5P/TROPOMI satellite sensor~~ were used to evaluate the current uncertainty in the retrievals of aerosol properties. Aerosol retrieval modifications based on Pandora NO₂ measurements are proposed for both AERONET and SKYNET. In addition, relatively high spatially resolved NO₂ observations from the S5P/TROPOMI satellite sensor were used to demonstrate the possibility of applying the corrections globally. A first attempt to investigate the impact of those corrections on AOD and AE annual trends is also conducted.

2 Instrumentation, data and methodology

2.1 The target area and relevant observational sites

Rome is the capital and the most populous city of Italy with almost 3 million inhabitants and one of the most densely populated cities in the European Union (ISTAT 2021). It is located about 24 km east of the Tyrrhenian Sea, surrounded by an extensive

100 undulating plain and crossed by the Tiber and Aniene rivers. The city is part of the Lazio administrative region in the central part of the Italian peninsula. The economic activities in the metropolitan area are characterized by the absence of heavy industrial facilities and are related mainly to the services and high-technology sectors, as well as commercial activities and tourism. The city air quality is strongly affected by local emission sources, such as transportation and domestic heating, but it is also markedly affected by local circulation and mid-to-long range transport events of sea salt, wildfires and Saharan dust (e.g., Ciardini et al., 2012; Gobbi et al., 2013; Barnaba et al., 2017; Valentini et al., 2020; Di Bernardino et al., 2021).

105 Rome air quality is monitored on a regular basis by standard in situ instrumentation. These measurements are complemented by multi-platform, long-term observations of aerosol and trace gases performed by a variety of ground-based remote sensing instruments such as sun-sky radiometers, Raman and elastic lidars, automated lidar-ceilometers, Pandora, Brewer and DOAS spectrophotometers (e.g., Di Ianni et al., 2018; Iannarelli et al., 2021; Diemoz et al., 2021; Bellini et al., 2023). In this study,

110 we used remote sensing measurements of columnar NO₂ and aerosol properties performed in two stations located in the greater area of Rome. More specifically, observations were obtained from an urban station (APL-SAP hereafter) located at the Atmospheric Physics Laboratory of the Physics Department of ‘La Sapienza’ University in the city centre (41.90° N, 12.52° E; altitude 75 m a.s.l.) and a suburban site at the southern east edge of the city in the CNR-ISAC Rome Atmospheric Supersite, CIRAS, in Rome-Tor Vergata (41.84° N, 12.65° E; altitude 117 m a.s.l.). These two observational sites along with the rural

115 station of CNR-IIA in Montelibretti contribute to the Boundary-layer Air Quality-analysis Using Network of Instruments (BAQUNIN) supersite (Iannarelli et al., 2021) as well as to several national and international observing networks.

2.2 Aerosol data sets

2.2.1 AERONET

The Aerosol Robotic Network (AERONET) is a ground-based passive remote sensing aerosol monitoring network initiated by NASA and expanded by several national and international networks and collaborators (Holben et al., 1998). For more than

120 two decades, AERONET has been delivering continuous, long-term data sets of aerosol optical, microphysical and radiative properties to support aerosol studies and the validation of space-borne retrievals. The network uses the Cimel CE318-T Sun-Sky-Lunar multispectral photometers and provides standardization of instrument calibration and data acquisition as well as centralized data processing and distribution. The AERONET public domain database provides retrievals of spectral AOD, inversion products and precipitable water at a global scale (<https://aeronet.gsfc.nasa.gov/>, last access: 21 October 2022).

125 In this study, we employed Level 1.5 quality-assured retrievals of AOD at 380, 440, 500, 675 and 870 nm along with AE at 440-870 nm from Version 3 processing algorithm (Giles et al., 2019; Sinyuk et al., 2020). Level 1.5 data are cloud-screened and quality-assured, but final calibration may not have not been applied to them. However, they represent a good trade-off between quality and readiness, considering that our approach aims to perform a near-real-time improvement on aerosol

130 products. In the standard AERONET For AOD retrieval, the NO₂ optical depth is estimated from monthly climatological values of total NO₂ from the Ozone Monitoring Instrument (OMI/Aura) Level-3 retrievals during the 2004-2013 period at 0.25° by

0.25° spatial resolution and the NO₂ absorption coefficients from Burrows et al. (1998). The observations over the CNR-ISAC station used in this work cover the period March 2017 – mid-August 2022 (Fig. 1), in which synchronous data from the co-located Pandora instrument are also available. The respective period for APL-SAP is from April 2017 through early September 2022 (Fig. 1). The aerosol data sets for both locations are presented in Fig. 1. The average AE is 1.23 ± 0.4 and 1.31 ± 0.5 at APL-SAP and CNR-ISAC, respectively, while the average AOD is about 0.18 ± 0.1 at both stations. AOD has a quite marked yearly cycle, with higher AOD values recorded during summer months, i.e., about 0.22 ± 0.1 and 0.21 ± 0.1 at APL-SAP and CNR-ISAC, respectively. AE is also higher during summer with a mean value of 1.26 ± 0.4 for APL-SAP and 1.38 ± 0.5 for CNR-ISAC.

~~In addition, we employed Level 1.5 SSA retrievals at 440 nm from almucantar measurements performed at both AERONET stations. Almucantar scans (circle on the celestial sphere parallel to the horizon) are measurements of radiance as a function of scattering angle performed by CIMEL instruments in order to be used for the retrieval of inversion products such as optical absorption (SSA), aerosol scattering phase function and properties related to aerosol typing, e.g., aerosol volume size distribution and complex refractive index (Dubovik and King, 2000; Dubovik et al., 2000). SSA is a measure of the scattering/absorbing nature of aerosol mixture and theoretically ranges between 0 and 1. In Earth's atmosphere, its value is usually very close to 1, as most aerosols types mainly scatter solar light, and it is rarely found lower than 0.7 (Dubovik et al., 2002; Raptis et al., 2020). The complete SSA data set used for this study is formed by 1478 retrievals spanning from March 2017 to November 2020 in the case of APL-SAP station, and 2463 spanning from April 2017 to September 2021 for CNR-ISAC.~~

2.2.2 SKYNET

The SKYNET network, established at the beginning of the 2000s, is a ground-based radiation observation network dedicated to aerosol, cloud and solar radiation interaction researches using the Prede POM sun-sky radiometers (Takamura and Nakajima, 2004; Nakajima et al., 2020). It is based on the collaboration and maintenance by several universities and research institutes around the world. This network imposes the standardization of instrument calibration, data acquisition and data processing and implements two data analysis flows (SR-CEReS & ESR-MRI) mainly based on the SKYRAD.pack, a software package implemented for the POM sky radiometer (e.g., Nakajima et al., 1996) (<https://www.skynet-isdc.org/methodology.php>, last access: 21 October 2022). In contrast to AERONET AOD retrieval methodologies, no correction for NO₂ optical depth is applied in the calculation of SKYNET AOD (e.g., Campanelli et al., 2004; Estellés et al., 2012). Here, we used the ESR-MRI/SUNRAD processor version 0.9 Level 2 AOD at 400, 500, 675, 870, and 1020 nm and AE at 400-1020 nm data sets over APL-SAP from late September 2017 to May 2022 (Fig. 1), which are open-accessed online (<https://www.skynet-isdc.org/data.php>, last access: 21 October 2022). The SKYNET time series used in our analysis is also illustrated in Fig. 1. The calculated mean AOD and AE are 0.18 ± 0.1 and 1.23 ± 0.4 , respectively. These values are similar to AERONET APL-SAP averages mentioned in Sect. 2.2.1, though they correspond to slightly different wavelengths. SKYNET also reports higher values on average during summer, i.e., 0.22 ± 0.1 and 1.38 ± 0.5 for AOD and AE, respectively.

165 2.2.3 MODIS Deep Blue data

The Moderate Resolution Imaging Spectroradiometer (MODIS) is a key sensor onboard the NASA Terra and Aqua satellites flying respectively since 2000 and 2002. Terra MODIS (descending node, about 10:30 a.m. UTC) and Aqua MODIS (ascending node, about 1:30 p.m. UTC) are observing the entire Earth's surface every 1 to 2 days, acquiring data in 36 spectral bands ranging in wavelength from 0.4 μm to 14.4 μm , with a spatial resolution of 1 km at nadir (except for a few bands with
170 higher spatial resolution).

Inversion of MODIS observations allows retrievals of several geophysical quantities. Here, we used the aerosol AOD products retrieved using the MODIS Deep Blue (DB) algorithm (Hsu et al., 2004, 2006, 2013). The basic principle of DB algorithms is to utilize the pre-calculated land surface reflectance database in deep blue bands (0.412 μm), in which surface reflectance is relatively lower than those in longer bands. In particular, we used the Collection 6.1 DB AOD products for both Aqua and
175 Terra satellites. More details about the DB algorithm are in Hsu et al. (2013) and references therein. The spatial resolution of this product is 10 km. Wei et al. (2019) highlighted that the DB algorithm is relatively more stable and less affected by changes in atmospheric and surface conditions with respect to the Dark Target algorithm (Levy et al., 2013), showing better performances in urban areas for slightly polluted cases, such as the area of Rome. They also highlighted that Collection 6.1 AOD products perform better than the previous collections, especially in Europe and North America. The MODIS DB products
180 used in this study are available at the Level-1 and Atmosphere Archive and Distribution System Distributed Active Archive Center (<http://ladsweb.nascom.nasa.gov>, last access: 21 October 2022).

2.3 Total NO₂ observations

2.3.1 Pandora spectroradiometers

Pandora instruments are compact spectrometers that perform spectral measurements with high temporal resolution of direct
185 solar irradiance and scattered radiance for the retrieval of total and tropospheric column densities of atmospheric trace gases (e.g., NO₂, O₃ and HCHO) that affect air quality, as well as their near-surface concentrations and vertical profiles (e.g., Herman et al., 2009; Tzortziou et al., 2012, 2015). The total NO₂ vertical column data sets used in the present study were obtained from the Pandora spectrometer #115 operating at CNR-ISAC since March 2017 and the Pandora systems #117 and #138 both
deployed at APL-SAP since April 2016 and within the period August 2019 – October 2020, respectively. The above time
190 series have been affected by the COVID-19 lockdown period, February – May 2020 (Campanelli et al., 2021). The monthly averaged values from both stations are presented in Fig. 2 and inter-compared in the scatterplot of Fig. 3. On average, the Pandora total NO₂ column over APL-SAP is about 0.07% higher compared to the CNR-ISAC NO₂.

Pandora total NO₂ column product is derived from the direct-sun measurements in the UV-VIS spectral range 280-530 nm with an average resolution of 0.6 nm by means of the Blick software and the algorithm implemented there as described by
195 Cede (2021). The data sets employed for this work were obtained with the direct-sun retrieval code “nvs3” and the Blick processor version 1.8. Pandora instruments are part of the Pandonia Global Network (PGN) (Cede et al., 2020) and have been

200 fully characterized following the calibration procedures presented by Müller et al. (2020). The recorded raw spectrally resolved radiation measurements are centrally processed for the retrieval of atmospheric trace gas products, which are all publicly available online (<https://www.pandonia-global-network.org/>, last access: 21 October 2022). In the current study, high (flags 0 and 10) and medium (flags 1 and 11) quality data are employed. Information on the quality control of Pandora products can be found in Cede et al. (2021). Pandora NO₂ retrievals have been compared and validated with other ground-based and spaceborne observations during several field campaigns (e.g., Flynn et al., 2014; Martins et al., 2016; Lamsal et al., 2017; Herman et al., 2018; Kreher et al., 2020). Total NO₂ data from the Pandora instrument #117 located at APL-SAP have been compared with NO₂ observations retrieved by the co-located MkIV Brewer spectrophotometer with serial number #067, revealing a correlation coefficient above 0.96 and a negligible absolute median bias of 0.002 DU (Diémoz et al., 2021). According to Herman et al. (2009), the Pandora direct-sun total NO₂ has a clear-sky precision of 0.01 DU in the slant column and a nominal estimated accuracy of 0.1 DU in the vertical column. In the same study, a systematic difference of less than 1% was found between the relative slant columns of Pandora and a MultiFunction Differential Optical Absorption Spectroscopy (MFDOAS) instrument. In the current study, high (flags 0 and 10) and medium (flags 1 and 11) quality data are employed.

205

210 As already mentioned in Sect. 2.2.1, AERONET uses climatological values from OMI L3 products for the estimation of NO₂ optical depth in AOD retrievals. The corresponding OMI total NO₂ ranges between about 0.2 and 0.3 DU, with an average value of 0.26 (± 0.02) DU. ~~The bivariate relative distribution of the percentage time series of the deviation of Pandora columnar NO₂ differences from the AERONET climatological values against the PGN product for both urban (APL-SAP) and suburban (CNR-ISAC) locations is illustrated in the upper panel of Fig. 4. The Pandora NO₂ data are time-interpolated to AERONET measurements. The percentage frequency distributions of absolute Pandora-OMI deviation for both locations is are also presented (Fig. 4, lower panel). A systematic OMI climatology underestimation (positive deviations) is observed for about 89% and 87% of the data pairs for APL-SAP and CNR-ISAC, respectively (lower panel of Fig. 4). About 89% of the APL-SAP and 87% of the CNR-ISAC data pairs show an OMI climatology systematic underestimation of NO₂ (positive deviations in Fig. 4).~~ AERONET aerosol retrievals seem to significantly underestimate the NO₂ abundance over urban and suburban

215

220 locations with an average absolute difference between the actual Pandora measurements and the estimations from satellite climatology of about 0.15 ± 0.19 DU ($61.5 \pm 71.5\%$) and 0.16 ± 0.18 DU ($61.5 \pm 67.2\%$) for APL-SAP and CNR-ISAC, respectively. This underestimation of the NO₂ levels over urban locations, characterized by strong spatial gradients, can be attributed to the fact that OMI climatology cannot capture the temporal and spatial NO₂ variability within an urban context (e.g., Drosoglou et al., 2017; Herman et al., 2019). As expected Thus, since OMI climatology cannot capture the temporal NO₂ variability, both the absolute (not shown here) and percentage (Fig. 4) the derived differences of in total NO₂ are highly correlated to the Pandora measurements. Most of the data pairs lie within 0-200% deviation. The majority of PGN – OMI biases lie within 0 - 0.5 DU data from AERONET climatology corresponding to Pandora values lower than 1 DU. More specifically, ~~the~~ 90% of the PGN NO₂ data over APL-SAP differ within -0.14 DU (-50%) and 0.44 DU (150%) from OMI climatology, while the respective deviation range between -0.14 and 0.51 DU (-50% – 170%) for CNR-ISAC. However, there are quite a

225

230 few cases (~9.5% and ~8.8% for APL-SAP and CNR-ISAC, respectively) of higher PGN values (< 2 DU) leading to larger deviations (up to ~~600%~~, i.e. ~1.6 DU for APL-SAP and ~1.5 DU for CNR-ISAC).

2.3.2 TROPOMI

235 The Tropospheric Monitoring Instrument (TROPOMI) is a nadir-viewing spectrometer on board Sentinel-5 Precursor (S5P) satellite, launched on 13 October 2017. Since August 2019, TROPOMI has a pixel size of 5.5 km × 3.5 km (the initial resolution was 7 km × 3.5 km). NO₂ columns are retrieved using the backscatter solar radiation detected in the spectral window of 405-465 nm (van Geffen et al., 2015) by applying the DOAS technique (Platt, 1994; Platt and Stutz, 2008). The operational TROPOMI NO₂ products are generated using the algorithm described by van Geffen et al. (2022), which is an improvement of the NO₂ DOMINO algorithm (Boersma et al., 2011) developed by the Royal Netherlands Meteorological Institute (KNMI) for the OMI satellite sensor measurements. Both near-real-time (NRTI) and off-line (OFFL) NO₂ data sets are retrieved using 240 the KNMI standard algorithm (Eskes et al., 2022; Eskes and Eichmann, 2022). NRTI data files are available within 3 hours from the measurement, whereas the OFFL data are processed in off-line mode and the respective files are generated a few days after the sensing time (van Geffen et al., 2022).

In this study, the OFFL NO₂ retrievals are employed, which are the main S5P/TROPOMI product. The extracted NO₂ data set covers the period October 2018 – August 2022 and includes observations obtained from several processor versions; beginning 245 with version 01.02.00 before March 2019 and going up to version 02.04.00 after July 2022. The total NO₂ column was calculated from the sum of the tropospheric and stratospheric components, which is preferred over the TROPOMI total NO₂ product for comparisons with ground-based data, because the latter suffers from retrieval uncertainties due to its significant dependence on the ratio of the a-priori tropospheric and stratospheric columnar data (van Geffen et al., 2022). Additionally, the satellite pixels have been filtered to keep only those with QA value > 0.75, corresponding to cloud radiance fractions < 0.5 250 (Eskes and Eichmann, 2022). The S5P/TROPOMI NO₂ products have been downloaded from the Sentinel-5P Pre-Operations Data Hub of the Copernicus Open Access Hub (<https://scihub.copernicus.eu/>, last access: 21 October 2022).

For visualization purposes, averages of the summed NO₂ column re-gridded on a 500m grid are plotted for the greater Rome area (Fig. 5). The data used in Fig. 5 cover the period from 2018 to 2021, excluding the COVID-19 lockdown period (February – May 2020) in order to prevent the average NO₂ values from being affected by the low values observed during that period.

255 2.4 Methodology AOD and AE corrections for NO₂ absorption

2.4.1 AOD retrievals

The methodology to derive AOD (also referred to as τ) from photometric measurements is based on the Lambert-Beer law (Eq. 1), which describes light attenuation by atmospheric components. $I_0(\lambda)$ is the intensity of the incident light and $I(\lambda)$ denotes the radiation intensity after traversing through the atmosphere at a specific wavelength λ .

$$260 \quad I(\lambda) = I_0(\lambda) \cdot e^{-(m_\tau(\lambda)\tau(\lambda) + m_R(\lambda)\tau_R(\lambda) + \sum_j m_j(\lambda)\tau_j(\lambda))}, \quad (1)$$

$$\frac{\ln I(\lambda)}{\ln I_0(\lambda)} = -\left(m_\tau(\lambda)\tau(\lambda) + m_R(\lambda)\tau_R(\lambda) + \sum_j m_j(\lambda)\tau_j(\lambda)\right), \quad (2)$$

The quantities τ and τ_R describe the optical depth of radiation extinction due to aerosols (Mie scattering) and atmospheric molecules (Rayleigh scattering), whereas m_τ and m_R are the respective air mass factors. $\sum_j m_j\tau_j$ represents the sum of the extinction due to absorption from atmospheric gases (Eq. 3), this depending on wavelength.

$$265 \quad \sum_j m_j(\lambda)\tau_j(\lambda) = m_{NO_2}(\lambda)\tau_{NO_2}(\lambda) + m_{O_3}(\lambda)\tau_{O_3}(\lambda) + m_{H_2O}(\lambda)\tau_{H_2O}(\lambda) + \dots, \quad (3)$$

In our study we investigate the effects of using an independent, direct measurement of $\tau_{NO_2}(\lambda)$ rather than the climatological value used in the AERONET inversion in determining the AOD (τ). Thus, by combining Eq. (2) with Eq. (3), assuming that the air mass factor in direct-sun measurements is equal to $\sec(\theta)$ for both aerosol and NO_2 , where θ is the solar zenith angle, and absorption from all the other gaseous components keeps the same, the difference in AOD due to the different estimation of NO_2 optical depth is obtained by Eq. (4):

$$\Delta\tau(\lambda) = \tau_{NO_2PGN}(\lambda) - \tau_{NO_2AER}(\lambda), \quad (4)$$

where τ_{NO_2AER} is the NO_2 absorption optical depth climatology used by AERONET and τ_{NO_2PGN} is the optical depth calculated from Pandora NO_2 measurements. The latter is derived using Eq. (5):

$$\tau_{NO_2PGN}(\lambda) = \sigma_{NO_2}(\lambda) \cdot c_{NO_2PGN}, \quad (5)$$

275 The quantity $\sigma_{NO_2}(\lambda)$ in Eq. (5) refers to the absorption cross-section of NO_2 at wavelength λ (Burrows et al., 1998) and c_{NO_2PGN} is the total NO_2 column from Pandora instrument. The modified AOD values (τ_{AER_mod}) are obtained from the standard AERONET AOD (τ_{AER}) by applying the following equation:

$$\tau_{AER_mod}(\lambda) = \tau_{AER}(\lambda) - \left(\left(\sigma_{NO_2}(\lambda) \cdot c_{NO_2PGN}\right) - \tau_{NO_2AER}(\lambda)\right), \quad (6)$$

The same approach was also applied to the SKYNET AOD data. However, since the SKYNET retrievals assume $\tau_{NO_2SKYNET} = 0$, Eq. (4) and (6) are modified as:

$$\Delta\tau(\lambda) = \sigma_{NO_2}(\lambda) \cdot c_{NO_2PGN}, \quad (7)$$

$$\tau_{SKYNET_mod}(\lambda) = \tau_{SKYNET}(\lambda) - \left(\sigma_{NO_2}(\lambda) \cdot c_{NO_2PGN}\right), \quad (8)$$

where $\tau_{SKYNET}(\lambda)$ denotes the standard SKYNET AOD at spectral channel λ and $\tau_{SKYNET_mod}(\lambda)$ is the modified AOD at wavelength λ .

285 2.4.2 AE retrievals

The spectral variability of AOD is generally expressed as:

$$\tau = \beta \cdot \lambda^{-\alpha}, \quad (9)$$

$$\ln \tau = \ln \beta - \alpha \cdot \ln \lambda, \quad (10)$$

where α stands for the Ångström Exponent (AE).

290 ~~In~~ The AERONET AE product (Eck et al., 1999), ~~AE~~ is calculated by applying a least squares regression fit on Eq. (10) using the AOD and wavelength logarithms for each non-polarized wavelength channels in different spectral ranges, i.e. 340-440, 380-500, 440-675, 440-870 and 500-870 nm. The negative slope of this linear fit is the Ångström exponent α (Eq. 11).

$$\alpha = - \frac{N \sum \ln \tau_i \ln \lambda_i - \sum \ln \lambda_i \sum \ln \tau_i}{N \sum (\ln \lambda_i)^2 - (\sum \ln \lambda_i)^2}, \quad (11)$$

Here, we also investigate the impact of using synchronous Pandora total NO₂ data in AOD algorithm as described in Sect. 295 2.4.1 on AE retrievals. For this, the AERONET AE product in the range 440-870 nm was used along with the AOD of non-polarized channels included in this range, i.e., 440, 500, 675 and 870 nm. AE was recalculated based on Eq. (11) using the modified AOD at wavelengths 440 and 500 nm obtained from Eq. (6). For the other channels (675 and 870 nm), in which NO₂ absorption is negligible, the standard AOD data from AERONET were employed.

For SKYNET, AE is calculated by applying a least squares regression fit on Eq. (10) using the AOD and wavelength logarithms 300 at all wavelengths 400, 500, 675, 870, and 1020 nm. Again, AOD was recalculated using Eq. (8) only at wavelengths 400 and 500 nm, where the impact of NO₂ absorption is significant.

The difference in AE due to the different estimation of NO₂ optical depth in AOD retrievals is expressed as:

~~$$\Delta \alpha(\lambda) = \alpha_{mod}(\lambda) - \alpha(\lambda), \quad (12)$$~~

~~$$\Delta \alpha(\lambda) = \alpha(\lambda) - \alpha_{mod}(\lambda), \quad (12)$$~~

305 where $\alpha_{mod}(\lambda)$ represents the modified AE data and $\alpha(\lambda)$ denotes the AE standard product from AERONET or SKYNET network.

2.5 Trend calculations

~~2.4.3 Trend calculations~~

In this study we also evaluate the impact of modified AOD and AE ~~modified~~ retrievals, as described in Sect. 2.4.1 and 2.4.2, 310 on aerosol temporal trends. This is only a first attempt to investigate the possible effect of NO₂ absorption on the AOD and AE trends since the data sets used here are quite short for statistically meaningful calculations.

The annual trends in AOD and AE were estimated by applying the weighted least squares fitting technique introduced by Weatherhead et al. (1998), previously adopted in several aerosol trend analysis studies from space and the ground (e.g., Zhang and Reid, 2010; Yoon et al., 2012; Logothetis et al., 2021). The applied linear trend model is based on the following formula:

$$315 \quad Y_t = \mu + \omega X_t + \varepsilon_t, \quad t = 1, \dots, T, \quad (13)$$

where Y_t is the monthly average aerosol property of interest, μ is a constant term representing the linear fit offset at the start of the time series, ω stands for the magnitude of the trend per year and ε_t is the monthly average noise not represented by the linear fit. $X_t = t/12$ is the decimal number of years since the first month of the time series, t is the month index, T denotes the total number of months and $T/12$ is the total number of years in the time series.

320 In order to account for data variability due to severe aerosol events and cloud disturbance, we introduced a monthly weighting factor w_t in the linear fitting procedure (Eq. 14) (Yoon et al., 2012). This weighting factor is defined as the square root of the number of observations available each month n_t divided by the monthly standard deviation σ_t (Eq. 15).

$$\chi^2(\mu, \omega) = \sum_{t=1}^T (w_t \cdot (Y_t - \mu - \omega X_t))^2, \quad (14)$$

$$w_t = \frac{\sqrt{n_t}}{\sigma_t}, \quad (15)$$

325 In order to derive statistically significant monthly mean values, a minimum number of 10 observations in a daily basis was ensured. In addition, qualified monthly averages require the availability of measurements from at least 10 days per month. Data were filtered based on the above criteria and days and/or months that did not fulfil them were excluded from the data sample used in the trend calculations. It should be noted that the data sets employed in this study are quite short for statistically meaningful aerosol trend analysis. However, this is a first attempt to investigate the impact of modified AOD and AE modified calculations on the derived temporal trends.

330

2.6 GRASP algorithm

The Generalized Retrieval of Atmosphere and Surface Properties (GRASP) (Dubovik et al., 2021) is a state-of-the-art inversion algorithm based on a statistically optimized multi-term Least Square Method (LSM) proposed by Dubovik et al. (2004). GRASP has been applied to numerous applications covering a vast variety of instruments and, which is more interesting, to

335 very different combinations between them. Among the different applications of GRASP, it is possible to find: GRASP/POLDER-3 (Chen et al. 2020), GRASP/AOD (Torres et al., 2017), OLCI/GRASP (Chen et al., 2022), the combination of active lidar measurements and ground-based radiometry (Lopatin et al., 2013; 2021; Román et al., 2018; Herreras et al., 2019), the retrieval of all-sky cameras (Román et al., 2017; 2022) or for example applications to in situ measurements including polar nephelometers (Espinosa et al., 2017; 2019; Schuster et al., 2019).

340 GRASP scientific core was borne from the heritage of the AERONET inversion algorithm (Dubovik and King, 2000; Dubovik et al., 2000; 2004; King and Dubovik, 2013). At the same time, as discussed above and by Dubovik et al. (2011; 2021), the possibilities of GRASP have been extended due to the totally generalized nature of the inversion module and the continuous developments of the forward model.

345 For this study, GRASP has been used to mimic AERONET standard retrieval in order to understand the effects of the NO₂ concentration on the retrieved SSA at 440 nm. In this case, two different approaches were followed for the GRASP algorithm.

350 First of all, GRASP has been used as close as possible to the standard AERONET retrieval, which means that the input measurements of the algorithm are the Total Optical Depth (TOD) and the almucantar sky measurement routine at 440, 675, 870 and 1020 nm. In the first approach (GRASP/AERONET NO₂ hereafter), the NO₂ absorption is taken into account using OMI climatology, exactly as in AERONET. On the other hand, GRASP flexibility allows the use of different assumptions of the gaseous properties. Therefore, in addition to the standard approach, the aerosol retrieval has been done also using the total columnar NO₂ concentrations provided by the Pandora spectrometers co-located with AERONET instruments at the two stations selected for this study. This methodology will be hereafter referred to as GRASP/Pandora NO₂. Thus, in addition to the standard AERONET retrieval products, GRASP has provided aerosol retrieval using these more accurate NO₂ concentrations. The NO₂ absorption features were calculated more precisely from those concentrations by using a K-Distribution approach; the “kbin” code (Doppler et al., 2014a; 2014b) to speed up the calculations.

3 Results and discussion

3.1 Differences in AOD and AE retrievals using Pandora NO₂ data

360 The differences in AOD ($\Delta\tau$) at 440 nm and, thus, of its spectral variability through the AE ($\Delta\alpha$ at 440-870 nm) correcting for measured NO₂ effects with respect to the standard AERONET retrievals are illustrated in Fig. 6 for both the Rome CNR-ISAC and APL-SAP stations. The frequency distributions of AOD, $\Delta\tau$ and $\Delta\alpha$ are also included in Fig. 6. $\Delta\tau$ is defined as the standard minus the modified AOD ($\tau_{\text{AER}} - \tau_{\text{AER_mod}}$, see Eq. 4-6). Similarly, whereas $\Delta\alpha$ is defined as $\alpha_{\text{AER_mod}} - \alpha_{\text{AER_mod}}$ (Eq. 12). The derived values are presented versus the AOD at 440 nm and are color-coded with respect to the Pandora NO₂ retrievals. The dependency of $\Delta\tau$ on NO₂ is quite clear. As expected, higher $\Delta\tau$ absolute values are obtained for higher NO₂ concentrations, regardless of the initial measured AOD. Also, the absolute percentage of $\Delta\tau$ with respect to the AOD is higher for lower aerosol loadings, which means that the impact of the NO₂ correction is more significant on lower AODs. This fact is also clear from $\Delta\alpha$, which is higher not only for higher NO₂, but for lower AOD values as well. Interestingly, based on Fig. 6, the highest Pandora NO₂ retrievals (reddish colors) are not associated to the highest AOD values, indicating that in Rome the high AOD loadings are not strictly associated to high NO₂ pollution events. In fact, high AODs are frequently related to long-range transport of elevated layers of desert dust, fires plumes or a combination of both (e.g., Barnaba et al., 2011; Gobbi et al., 2019; Campanelli et al., 2021; Andrés Hernandez et al., 2022). Hence, it might be worth to modify aerosol retrievals for high NO₂ in those pollution-related events with low to medium and middle AOD pollution-related events levels. More about AOD and aerosol type climatology for the Rome area can be found in Di Ianni et al., (2018) and in Campanelli et al. (2022).

375 In general, considering the climatological value chosen for Rome in AERONET retrievals, the use of actual, coincident NO₂ measurements on the calculations of aerosol properties seems to be still useful for AOD < 0.3, while quite negligible-low (less than 10%) for AOD > 0.5 and almost negligible for AOD > 0.8. In most cases AERONET retrievals seem to overestimate AOD and AE. However, there are cases of underestimation, especially in AE retrievals, which seems to be higher for lower AODs. Those underestimations correspond to overestimation of NO₂ from satellite monthly climatological values used in

AERONET retrievals. The estimated AOD and AE deviations are below 0.01 and 0.1, respectively, for the majority of observations, i.e., about 96 - 98% of occurrences for both CNR-ISAC and APL-SAP (see also distributions in Fig. 6). The average AOD bias is between 0.002 ± 0.003 and 0.003 ± 0.003 (with the higher values observed at 380nm), while the average AE bias is $\sim 0.02 \pm 0.03$. Overall, the mean AOD bias is low compared to the estimated uncertainties for the standard AERONET product, i.e., 0.01 - 0.02 (with the higher errors observed in the UV) (Sinyuk et al., 2020). However, the mean AOD bias for the cases of high NO_2 levels ($> \sim 0.7$ DU) is $\sim 0.011 \pm 0.003$ at 440 nm and $\sim 0.012 \pm 0.003$ at 380 nm for APL-SAP and $\sim 0.009 \pm 0.003$ at 440 nm and $\sim 0.010 \pm 0.003$ at 380 nm for CNR-ISAC, which is comparable to the AERONET reported uncertainties. The estimated mean bias of AE retrievals for the cases with high NO_2 ($> \sim 0.7$ DU) is $\sim 0.08 \pm 0.04$ for both Rome sites. The threshold for NO_2 has been selected as the average Pandora NO_2 (~ 0.4) calculated from the whole data set plus two times the standard deviation.

The results for SKYNET observations are similar (Fig. 7), but only positive $\Delta\tau$ and $\Delta\alpha$ values are derived, indicating overestimation of the aerosol properties, since the NO_2 optical depth is not considered in the standard retrieval processes (see Eq. 7-8). $\Delta\tau$ is defined as $\tau_{\text{SKYNET}} - \tau_{\text{SKYNET_mod}}$ (see Eq. 7-8), whereas $\Delta\alpha$ stands for $\alpha_{\text{SKYNET_mod}} - \alpha_{\text{SKYNET_mod}}$ (Eq. 12). In addition, the derived deviations in aerosol properties reach higher values compared to AERONET. Especially AE differences extend up to a value of about 0.7, which is more than double compared to AERONET results. Interestingly, these quite large $\Delta\alpha$ values (> 0.3) correspond to relatively low NO_2 loadings (< 1.2 DU). The differences observed between the two networks can be partly attributed to the different wavelength channels used for AOD and AE retrievals. Similarly to AERONET, the derived AOD and AE biases for SKYNET are below 0.01 and 0.1, respectively, for the majority of observations, i.e., about 85% of occurrences for AOD and about 90% for AE (see also distributions in Fig. 7). The overall average AOD bias is $\sim 0.007 \pm 0.003$, which can be assumed low considering that Nakajima et al. (2020) have estimated a root-mean-square difference (RMSD) of about 0.03 for wavelengths < 500 nm in city areas in AOD comparisons with other networks. However, the mean AOD bias for the cases with high NO_2 levels ($> \sim 0.7$ DU) is found to be about 0.018 ± 0.003 , which is comparable to the RMSD value reported by Nakajima et al. (2020). The overall average AE bias calculated in this study is $\sim 0.05 \pm 0.04$, whereas the AE bias averaged over the high NO_2 cases is about 0.10 ± 0.05 .

WMO (2005) states that, when comparing AOD retrieved from sun-photometers, 95% of the AOD differences should lie within $\pm (0.005 + 0.01/m)$ of AOD, where m is the optical air mass. The first term of the equation (0.005) represents the maximum tolerance for the uncertainty due to the atmospheric parameters used for the AOD calculation (additional atmospheric trace gas corrections, i.e., Ozone and NO_2 , and Rayleigh scattering), while the second term (0.01/m) describes the calibration-related relative uncertainties, for which WMO recommends an upper limit of 1% (e.g., Cuevas et al., 2019; Kazadzis et al., 2018a). Based on the above, although the average deviations found in this study are low compared to the retrieval uncertainties, they cannot be considered negligible, especially the average systematic underestimation of AOD of about 0.007 from SKYNET, having also in mind that there are locations with much higher average NO_2 compared to the city of Rome.

The statistics showing mean differences in AOD and AE AERONET and SKYNET retrievals using actual, coincident NO₂ measurements are presented in Table 1. AERONET AOD retrievals at 380 nm are also included in the table. In addition, deviations of AOD and AE using daily or monthly averages of NO₂ in AERONET and SKYNET observations are also investigated. Table 1 shows that the average deviations of AOD and AE values do not change significantly whether the actual Pandora NO₂ measurements or the daily or monthly mean values are used for the retrievals. The percentage differences for AOD lie within the range ~~1.2~~ $1.2 - 1.9\%$ for AERONET, while they are more than doubled (~~-5.3~~ $-5.3 - 5.7\%$) for SKYNET. For the standard aerosol products of the latter, NO₂ optical depth is not considered. The estimated percentage differences for AE are ~~within~~ $1.2 - 1.7\%$ and ~~> 2.5~~ $2.6 - 2.9\%$ for AERONET CNR-ISAC and APL-SAP, respectively, and ~~about 7.8~~ $7 - 7.9\%$ for SKYNET APL-SAP. It should be noted that the spectral channels used in AERONET retrievals are 380 and 440 nm for AOD and 440-870 nm for AE, whereas SKYNET data refer to 400 nm and 400-1020 nm for AOD and AE, respectively.

3.2 AOD and AE retrievals based on TROPOMI NO₂ data

Satellite sensors perform measurements globally and provide information on the air quality even over regions that lack ground-based observations. However, as already mentioned for OMI in Sect. 2.3.1, the spatial resolution of the satellite retrievals is limited by the pixel size. Co-located S5P/TROPOMI observations, characterized by improved spatial and temporal resolution compared to previous satellite missions (e.g., OMI), were also employed to investigate whether the ground-based retrievals of aerosol properties could be improved on a global scale. Again, the approach described in Sect. 2.4.1 and 2.4.2 was applied by replacing the Pandora total NO₂ (c_{NO_2PGN}) with corresponding columnar retrievals from TROPOMI. Based on the current satellite footprint (5.5 km × 3.5 km), a radius of 5 km around each ground-based station was selected for the spatial co-location. The TROPOMI NO₂ data were time-interpolated to AERONET and SKYNET measurements. Despite the improved spatial resolution of TROPOMI, the NO₂ corrections using TROPOMI data are expected to be less accurate than those performed with the Pandora product. For example, Lambert et al. (2021) showed a bias between TROPOMI and Pandora total NO₂ column ranging from -23% over polluted stations to +4.1% over clean areas with a median bias of -7.1%, in the frame of the standard validation process of TROPOMI Level 2 NO₂ products. Other studies have concluded similar results. For example, Zhao et al. (2020) showed a negative bias for the standard TROPOMI total NO₂ product in the range 23 - 28% over urban and suburban environments and a positive bias of 8 - 11% at a rural site, while Park et al. (2022) showed 26 - 29% negative bias and R² within 0.73-0.76 over the Seoul Metropolitan Area in Korea.

The statistical metrics of the averaged deviations of the modified AERONET and SKYNET AOD and AE retrievals using actual, co-located TROPOMI NO₂ measurements from the network standard products are presented in Table 2. Similarly to Sect. 3.1 and Table 1, deviations of AOD and AE retrievals derived by employing daily or monthly mean TROPOMI total NO₂ were also investigated. The average deviations of AOD and AE values do not change significantly whether the actual TROPOMI NO₂ measurements or the daily mean values are used for the retrievals. This behaviour is expected considering that TROPOMI overpasses occur once or twice per day and, hence, they do not capture daily variations of NO₂. In the case of monthly averaged TROPOMI NO₂ data, the estimated differences between the standard and modified aerosol products drop

notably for AERONET. However, there are still differences compared to OMI NO₂ climatology due to the improved spatial resolution of [the TROPOMI pixel](#). [The average AOD bias is \$\sim 0.001 \pm 0.001\$ \(with the higher values observed at 380nm\), while the average AE bias is \$\sim 0.01 \pm 0.01\$ for both AERONET stations. For the cases of high NO₂ levels \(\$> \sim 0.7\$ DU\), the mean AOD bias is \$\sim 0.004 \pm 0.001\$ at 440 nm and \$\sim 0.005 \pm 0.002\$ at 380 nm for APL-SAP and \$\sim 0.003 \pm 0.001\$ at both 440 nm and 380 nm for CNR-ISAC. The estimated mean bias of AE retrievals for the cases with high NO₂ \(\$> \sim 0.7\$ DU\) is \$\sim 0.05 \pm 0.04\$ and \$\sim 0.02 \pm 0.01\$ for APL-SAP and CNR-ISAC, respectively. In the case of SKYNET, the overall average AOD bias is \$\sim 0.005 \pm 0.002\$ for AOD and \$\sim 0.04 \pm 0.03\$ for AE. For the high NO₂ cases, a mean AOD bias of about \$0.011 \pm 0.002\$ and an average AE bias of \$\sim 0.07 \pm 0.04\$ were calculated.](#) Interestingly, the deviations of SKYNET retrievals using monthly TROPOMI data are very similar to those derived using the actual overpasses or daily averaged TROPOMI NO₂, probably due to the fact that the NO₂ optical depth is not included in the standard network AOD retrieval processes.

The percentage differences for AOD lie within the range $0.2\% - 0.9\%$ for AERONET and are about $3.8 - 3.9\%$ for SKYNET, which are much lower compared to those derived using Pandora NO₂ (see Table 1). The estimated percentage differences for AE are $\sim 0.8 - 0.9\%$ and $\sim 1.6 - 1.7\%$ for AERONET CNR-ISAC and APL-SAP, respectively, and about 4% for SKYNET APL-SAP using actual or daily TROPOMI data. It should be noted again that the spectral channels used in AERONET retrievals are 380 and 440 nm for AOD and 440-870 nm for AE, whereas SKYNET data refer to 400 nm and 400-1020 nm for AOD and AE, respectively.

460 3.3 Case study: Impact of high Pandora NO₂ on low AOD

In order to investigate further the impact of high NO₂ during pollution events on the retrieval of relatively low levels of AOD, we used measurements performed at APL-SAP on June 25th 2020, in the morning of which there was a high NO₂ event. In the upper panels of [Fig. 8](#) and [Fig. 9](#), the total NO₂ measured from Pandora during that day is illustrated. For AERONET ([left panels of Fig. 8](#)), the satellite climatological values used in the retrieval of standard AOD product and their deviations from Pandora NO₂ are also displayed. The standard and [NO₂-modified AOD and AE data from both AERONET and SKYNET \(see also Sect. 2.4.1 and 2.4.2\)](#), as well as the magnitude of the respective differences ($\Delta\tau$ and $\Delta\alpha$), are presented in the middle and lower panels of [both figures Fig. 8](#).

The differences in AOD and AE retrievals from both networks are significant only within a time span of about 3 hours around the high NO₂ event ($\sim 7:00-10:00$ UT) and can be assumed negligible for the rest of the day when the NO₂ levels remain quite low. [The median AOD bias for AERONET is about 0.003 with a maximum of about 0.02 at the peak of the event. The median and maximum AE biases are 0.014 and 0.11, respectively.](#) It can be also noted that in the case of SKYNET both AOD ([median value of \$\sim 0.008\$ with a maximum of \$\sim 0.03\$](#)) and AE deviations ([median and maximum values of \$\sim 0.03\$ and 0.10, respectively](#)) are a bit higher compared to the respective AERONET deviations of synchronous data. This can be mainly attributed to the fact that SKYNET standard AOD retrieval processes do not account for the NO₂ absorption and partly explained by the different channels used in the detectors of the two networks.

3.4 Impact on AOD and AE trends

In this section, a first attempt is conducted to investigate the effect of the modified AOD and AE retrievals based on the Pandora total NO₂ observations on the annual trends of those aerosol properties. The annual trends of AERONET/SKYNET AOD and AE over both APL-SAP and CNR-ISAC sites, calculated by applying the approach described in paragraph 2.5.4.3, as well as their uncertainties (standard errors of the regression slope) are presented in Table 3. ~~In general, the calculated trends from AERONET over the suburban area (CNR-ISAC) are quite higher compared to the AERONET derived trends for the urban location (APL-SAP), but comparable with those from SKYNET.~~

It should be noted here that the aerosol data sets from the two networks correspond to slightly different time periods. In addition, there are significant gaps in the time series from CNR-ISAC due to instrumental problems and ~~the~~ COVID-19 lockdown period (February – May 2020) has been excluded from the data ~~sets~~ analysis.

~~Therefore, the results in Table 3 are mainly intended to highlight how a different NO₂ correction may affect the aerosol trends and should be interpreted separately for each individual site. Interpretation of the trend significance for the Rome area is not possible using this short period of time (~5.5 years), considering that the estimated trends are quite small and the uncertainties introduced by linear regression are relatively high. do not reveal any notable impact of NO₂-modified retrievals on AOD trends. It should be noted that the aerosol time series employed in this study are quite short (~5.5 years) for statistically meaningful trend analysis. In addition, the estimated trends are quite small and the uncertainties introduced by linear fit are relatively high, especially for AOD. Thus, we have not conducted any study on the significance of these trends. However, the NO₂ effect on the aerosol retrievals, although small, is comparable with the calculated trends and may be significant for trends estimated over larger data sets.~~

~~One aspect shown here is that the difference in the AOD and AE trends for the two data sets (original and NO₂-modified) is comparable with the calculated trends. As expected, AE trends with and without NO₂ correction show relatively higher differences, as AE is much more sensitive to spectral AOD changes. However, the linear fitting uncertainty on AE is also high. In addition, NO₂ effects on AOD trends would be more obvious in the case of a significant NO₂ trend during ~~the a certain~~ period. A thorough long-term trend analysis is out of the scope of this work and could be the topic for a future study. AE trends with and without NO₂ correction show relatively higher differences as AE is much more sensitive to spectral AOD changes. However, the linear fitting uncertainty on AE is also high.~~

3.5 ~~Impact on the inter-comparison of AERONET and SKYNET ground-based AOD products against and satellite~~ MODIS-DB AOD data

In this section, ~~in addition to the analysis of ground-based data,~~ we have analysed a potential effect of considered NO₂ corrections on the agreement of AERONET and SKYNET AOD products with relevant satellite data. Indeed, it is well known that most satellite retrievals are validated against ground-based measurements of AOD that are considered as a ground-truth. Moreover, most satellite retrieval algorithms are substantially tuned to closely match AERONET observations. For example, all MODIS algorithms, including DB, rely, in one way or another, on AERONET dynamic aerosol models and climatologies

of AERONET retrievals. Nonetheless, since MODIS retrievals fundamentally rely on MODIS radiances that are fully
510 independent of AERONET data, some inaccuracies in assumptions, such as those on NO₂ amount, can cause some additional
biases between AERONET and MODIS AOD results.

To evaluate the effects of the proposed correction, we have compared AERONET and SKYNET AOD products against
MODIS DB AOD products at 470 nm for the 2017-2022 period. In the inter-comparison, we considered only MODIS DB
AOD products for which the distance between the center of the pixel and the AERONET site location (APL-SAP or CNR-
515 ISAC) does not exceed 5 km. Furthermore, we considered all the AERONET (or SKYNET) AOD data within ± 30 minutes
from the MODIS satellite overpasses. In order to guarantee the quality of the data, we used MODIS DB AOD with QA index
 ≥ 2 , which corresponds to good and very good products (Wei et al., 2019).

The inter-comparison has been performed using MODIS DB AOD at 470 nm. Consequently, we computed the AERONET
and SKYNET AOD at 470 nm exploiting the AE. The AERONET AOD at 470 nm was calculated using the standard
520 AERONET AOD at 440 nm and AE at 440-870 nm. Similarly, the SKYNET AOD at 470 nm was computed using the standard
SKYNET AOD at 400 nm and AE at 400-1020 nm. The NO₂-modified AERONET and SKYNET AOD at 470 nm were also
computed with using the same approach and the ~~modified~~-AOD and AE retrievals that have been modified using the Pandora
NO₂ data.

We observe a generally satisfactory agreement between the ground-based (both AERONET and SKYNET) and MODIS DB
525 AOD data with a Pearson correlation (r) higher than 0.7. In general, MODIS DB AOD slightly overestimates the AOD
observed by the sun-photometers. The bias (calculated as satellite minus sun-photometer AOD) between MODIS DB and the
different ground-based data sets before the correction (upper panels of Fig. ~~109~~) varies from -0.009 for SKYNET APL-SAP
data (-0.008 considering AERONET) to 0.027 for AERONET CNR-ISAC. AERONET data, available for both sites, highlight
a lower agreement for the CNR-ISAC site, with a bias about 3 times larger with respect to the APL-SAP site. The correction
530 introduces a slight change of about 0.003 in the agreement between MODIS DB and AERONET AOD products and of 0.006
between MODIS and SKYNET data (lower panels of Fig. ~~109~~). Figure ~~10-9~~ also shows an improvement in the percentage of
MODIS AOD data falling within the expected error (EE) of $\pm (0.05 + 20 \%)$ (Hsu et al., 2013) for APL-SAP adopting the
correction on both AERONET and SKYNET.

In Fig. ~~11-10~~, we show the absolute correction (computed as the difference between original AERONET/SKYNET AOD data
535 at 470 nm and modified ones) as a function of the MODIS DB AOD and the NO₂ column retrieved by the Pandora instruments
located at APL-SAP and CNR-ISAC sites (upper panels). As already highlighted, we observe that the correction only depends
on the NO₂ amount and not on the AOD. Figure ~~11-10~~ also highlights that, although the improvement is relatively slight-low
on average, the correction can be larger than 10/15% in many cases.

This inter-comparison exercise demonstrated that the proposed correction slightly improves the agreement between MODIS
540 DB AOD data and AERONET and SKYNET AOD products, even if, on average, it is not statistically significant. Nevertheless,
as shown in Fig. 10, the improvement becomes significant when the differences between the NO₂ values observed by Pandora
and the OMI NO₂ climatology are also significant (lower panels of Fig. 10). Furthermore, since the proposed correction

depends on the amount of NO₂, the improvement is more evident in the correspondence of high values of NO₂ (upper panels of Fig. 10), typical of highly polluted areas such as the urban area of Rome (APL-SAP). Also, a slight improvement is also achieved in the suburban area of Rome (CNR-ISAC). Finally, in the case of SKYNET AOD products, the systematic overestimation, due to neglected NO₂ extinction in the official retrieval chain, is eliminated. Since the proposed correction depends on the amount of NO₂, the improvement is more significant in the correspondence of high values of NO₂, typical of highly polluted areas such as the urban area of Rome (APL-SAP). Nevertheless, a slight improvement is achieved also in the suburban area of Rome (CNR-ISAC).

550 3.6 Impact on SSA

One of the main impacts of accurate characterization of the columnar NO₂ concentration is certainly expected on the retrieved values of SSA in spectral ranges coinciding with NO₂ absorptions. In order to quantify this effect, the sensitivity of AERONET retrieval of SSA at 440 nm has been tested. For that purpose, the Generalized Retrieval of Atmosphere and Surface Properties (GRASP) algorithm (Dubovik et al., 2021) was applied to the standard AERONET aerosol retrieval measurements (TOD and almucantar at 440, 675, 870 and 1020 nm) following two different approaches. First, the reference for these comparisons was established by using GRASP retrieval configured as closely as possible to mimic the AERONET operational retrieval. Specifically, the NO₂ absorption values were taken from the same source as done in the AERONET processing (OMI climatology). On the other hand, GRASP is very flexible and allows the use of different assumptions of the gaseous properties. Therefore, in addition to the standard approach, the accurate aerosol retrieval has been done by using the total columnar NO₂ concentrations provided by the Pandora spectrometers co-located with AERONET instruments at the two stations selected for this study. Thus, in addition to the standard AERONET retrieval, GRASP has provided aerosol retrieval using these more accurate NO₂ concentrations. The NO₂ absorption features were estimated more precisely from those concentrations by using a K-Distribution approach; the “k bin” code (Doppler et al., 2014a; Doppler et al., 2014b) has been used to speed up the calculations. As previously explained (Sect. 2.6), two different GRASP approaches have been applied to this purpose: the GRASP/AERONET NO₂ and the GRASP/Pandora NO₂. Despite the close methodological basis between GRASP and AERONET retrievals, the divergence in the development of both algorithms has led to some differences in the retrieved products. Thus, in order to assure that the difference in the retrieved SSA at 440 nm is produced exclusively by the changes in the description of NO₂ absorption and to avoid the inclusion of any other sources of discrepancy, GRASP code has been used in both approaches instead of the standard AERONET SSA product.

The comparisons of the SSA at 440 nm obtained with both methodologies GRASP/AERONET NO₂ and GRASP/Pandora NO₂ improved absorption features for the two stations for the complete data set (not shown) do not show a clear influence of the change of NO₂ concentration. High correlations ($r^2-R > 0.98$) and a Mean Bias Error (MBE < 0.002) very close to zero are obtained. The mean NO₂ column concentration for the retrievals presented here is 0.4 DU. Thus, in general the analysed improvements are not expected to produce an important change in the retrieved parameters at 440 nm in the conditions with relatively low NO₂ absorption. However, in the cases where NO₂ concentration is elevated compared to the climatological

580 expected range, significant changes in the SSA at 440 nm retrievals can be appreciated. Figure ~~12-11~~ shows the comparisons of the SSA at 440 nm obtained with GRASP following an AERONET-like approach (X-axis) and the approach with the new NO₂ concentrations provided by Pandora (Y-axis), filtered for NO₂ concentrations higher than 0.79 DU, which corresponds to the average NO₂ plus two times the standard deviation. The two stations are correspondingly represented in the left and right panels. As it can be noted, for both stations in conditions of high NO₂ concentrations there is a consistent positive bias of ~ 0.02 (~~~ 2.5~~ %). However, a high correlation (~~r^2~~ $R > 0.965$) and Root Mean Square Errors (RMSE < 0.035) are also observed. Previous studies, ~~regarding found SSA retrieval uncertainty, found it~~ in the range of 0.02-0.03 (Eck et al., 2003; Corr et al., 2009; Jethva et al., 2014; Kazadzis et al., 2016), whereas the correction, when high NO₂ is recorded, is usually higher. Thus, it is clear that in conditions of high NO₂ concentrations an accurate characterization of this gas is necessary in order to avoid
585 noticeable bias in the affected AERONET channel around 440 nm.

4 Summary and conclusions

The retrievals of aerosol properties from sun-photometers may be ~~significantly~~ affected by NO₂ absorption in the observed spectral range and, thus, ~~the~~ accurate assumptions on NO₂ concentrations are highly desirable. Currently, some ground-based aerosol networks, such as SKYNET, do not take NO₂ optical depth into consideration in AOD retrieval processes, while others
590 (e.g., AERONET) use satellite-based NO₂ climatology for estimating it. However, ~~quite~~ significant errors ~~may~~ could be introduced in the AOD retrievals, especially over urban areas, where NO₂ variability can be high and also the occurrence of high NO₂ events is more frequent. Such errors may occur only in the cases where NO₂ is not taken into account or the used NO₂ climatology underestimates such high-NO₂ events. ~~is characterized by rather short lifetime and high spatial and temporal variations, due to inhomogeneous local emission patterns and photochemical destruction.~~

595 Actual co-located surface-based NO₂ measurements (e.g., from Pandora instruments) or space-borne observations with improved spatial and temporal resolution (e.g., S5P/TROPOMI) may be helpful for reducing the uncertainty of the NO₂ optical depth contribution in later versions of AOD retrieval algorithms. In this study, we evaluated the possible improvements of AOD and AE retrievals by applying a specific correction using synchronous and co-located measurements of the total NO₂ column from Pandora spectroradiometers and the TROPOMI satellite sensor. For this purpose, we used multiannual (2017-
600 2022) observations from both AERONET and SKYNET multispectral AOD observations co-located with Pandora instruments and collected over two locations in Rome (Italy) with different anthropic pressure, one in the city centre and the other in a suburban area.

The deviations of the NO₂-modified AOD retrievals from the network standard products were investigated. AERONET-used NO₂ climatology was found to systematically underestimate Pandora-measured NO₂ over both sites. The impact of the
605 correction is higher in the case of SKYNET since the NO₂ optical depth is not considered at all in the standard retrieval processes of that network. At the same time, the observed differences in the results between the two networks can be partly explained also by the different channels used for the retrievals. For both AERONET and SKYNET, a low but systematic AOD

overestimation was found. Although in most of the cases the differences are lower than 0.01 for AOD and lower than 0.1 for AE retrievals, the correction can still be ~~still~~-useful for lower AODs (< 0.3), where the majority of observations ~~is~~-are found, especially under high NO_2 pollution events. The mean AOD bias derived for the high NO_2 cases ($> \sim 0.7$ DU) is $\sim 0.011 \pm 0.003$ at 440 nm and $\sim 0.012 \pm 0.003$ at 380 nm for AERONET APL-SAP and $\sim 0.009 \pm 0.003$ at 440 nm and $\sim 0.010 \pm 0.003$ at 380 nm for AERONET CNR-ISAC. The mean AE bias for the high NO_2 is $\sim 0.08 \pm 0.04$ for both Rome AERONET sites. In the case of SKYNET, the mean bias for the cases with high NO_2 levels ($> \sim 0.7$ DU) is $\sim 0.018 \pm 0.003$ and $\sim 0.10 \pm 0.05$ for AOD and AE, respectively. Overall, the average biases in AOD retrievals are systematic but within the reported AOD uncertainties.

615 However, they are important to be reported here, as AOD retrieval uncertainties not linked with instrument calibration (e.g. Rayleigh, ozone and NO_2 related optical depths) are considered to have an upper limit of 0.005 as a goal for sun-photometers according to WMO (WMO, 2005). As expected, the effect of improved NO_2 assumption in the retrievals is more evident in both AOD and AE when the actual synchronous ground-based Pandora NO_2 measurements are employed compared to the situations when the used correction was based on daily or monthly averaged Pandora data or TROPOMI NO_2 retrievals. The use of TROPOMI NO_2 data is a demonstration of the possibility for corrections on a global scale. However, the underestimation of NO_2 concentrations by TROPOMI compared to Pandora NO_2 data for Rome leads to lower AOD corrections. In addition, AERONET used NO_2 climatology was found to systematically underestimate Pandora measured NO_2 over both sites.

In addition, a first attempt to evaluate the impact of those corrections ~~to~~-on AOD and AE annual trends was conducted. However, the aerosol data sets employed in this trend analysis are quite short for a statistically meaningful robust trend analysis.

625 Here, only quantitative comparisons are performed for each individual data set, i.e., corresponding to specific instrument and site, before and after the NO_2 -based correction. Although the effect of NO_2 on the derived trends seems to be insignificant and the linear fit trend calculations introduce uncertainties similar or higher to the NO_2 effects on AOD, the more pronounced impact may be expected for trends derived from larger data sets as well as in the case of a significant NO_2 trend.

We also investigated the possible effects of the proposed NO_2 optical depth correction on the agreement between ground-based and space-borne AOD retrievals. In particular, we compared ~~of~~-MODIS DB AOD retrievals at 470 nm with AERONET and SKYNET AOD products. In general, the agreement between ground-based (both AERONET and SKYNET) and MODIS DB AOD is quite good, revealing a correlation coefficient (r) higher than 0.7. The use of Pandora NO_2 in the sun-photometers retrievals introduces a slight improvement in absolute values of ~ 0.003 in the agreement between MODIS DB and AERONET AOD and an improvement of ~ 0.006 between MODIS and SKYNET observations. Although the impact on the comparisons

635 between space-borne and ground-based observations of AOD is quite small, it can be quite useful for eliminating or decreasing possible biases in the inter-comparisons of satellite and ground-based data in situations with NO_2 concentrations typical for highly polluted areas.

Moreover ~~Finally~~, we investigated the impact of using a precise characterization of the total NO_2 concentration on the SSA retrieval at 440 nm from AERONET measurements. For this, the GRASP algorithm was used to evaluate the effect of NO_2 correction on AERONET aerosol retrievals obtained by inverting TOD and almucantar radiances at 440, 675, 870 and 1020 nm. GRASP aerosol retrieval using the actual total NO_2 concentration provided by the co-located Pandora, over both stations

640

selected for this study, were compared with GRASP retrievals mimicking AERONET operational retrievals. The results showed that, in general, the effect in the retrieved parameters at 440 nm under low NO₂ absorption conditions was not significant. At the same time, for the cases with high NO₂ loadings (> 0.79 DU), important changes in the retrieved SSA were observed, with an average positive bias of 0.02 (2.5 %) for both locations.

In general, the effect of NO₂ absorption can be relatively important in the retrievals of aerosol properties, especially AE, AOD and SSA at 440 nm and 380nm, when NO₂ is not included in the retrieval algorithms or in cases where NO₂ absorption is significantly higher than the NO₂ climatology used. ~~is very important to account in the retrieval of aerosol properties especially AE, AOD and SSA at 440 nm.~~ Moreover, ~~if~~ NO₂ absorption is accounted from climatological data, the accuracy of such approach may not be sufficient at locations where NO₂ has high diurnal variability during high NO₂ concentration episodes that cannot be captured by the satellite climatology. In such situations, the use of accurate co-located NO₂ observations, e.g., by Pandora instruments, is highly desirable. Thus, based on the results of this study, the effect of NO₂ correction could be considered relatively small for large fraction of the observations, nonetheless the correction has certainly contributed towards lowering the uncertainty of AOD and, especially, aerosol SSA provided by sun-photometers.

In future studies, the effect of NO₂ correction on Absorption Ångström Exponent (AAE) could be explored. AAE is an aerosol optical property that describes the absorption variation with respect to wavelength and is significantly influenced by particle size, shape, and chemical composition used for aerosol characterization and apportionment studies (e.g., Schuster et al., 2006). Since AAE is a function of spectral AOD and SSA, the NO₂ correction for certain AOD wavelengths and SSAs, shown in this study, is expected to impact the AAE calculations towards lower values (as the NO₂-corrected AOD is systematically lower, and the corrected SSA is higher).

Finally, the improved technology including real-time NO₂ monitoring (e.g., the Pandonia network), real-time satellite-based products at high spatial resolution (e.g., TROPOMI) and the foreseen more precise NO₂ products (e.g., from Sentinel 4) tend to positively contribute towards improving retrieved aerosol properties in the spectral range (~380 – 440 nm) affected by NO₂ absorption.

Data availability. The AOD and AE products from the ~~CIMEL-Cimel~~ sunphotometer measurements as well as the NO₂ optical depth used in the retrievals are available in the AERONET data server (<http://aeronet.gsfc.nasa.gov/>). The SKYNET AOD and AE data sets were downloaded from the international SKYNET data center (<https://www.skynet-isdc.org/data.php>). The Pandora total NO₂ columns are available in the Pandonia Global Network website (<https://www.pandonia-global-network.org/>). The S5P/TROPOMI NO₂ products were obtained from the Sentinel-5P Pre-Operations Data Hub of the Copernicus Open Access Hub (<https://scihub.copernicus.eu/>). The MODIS DB products are available at the Level-1 and Atmosphere Archive and Distribution System Distributed Active Archive Center (<http://ladsweb.nascom.nasa.gov>). The SSA retrievals from the ~~CIMEL-Cimel~~ almucantar measurements can be accessed by contacting the author.

675 *Author contribution.* The manuscript was prepared by TD, I-PR, and MV. TD and I-PR developed and implemented the
correction algorithm for AOD and AE retrievals and conducted the trend analysis. MV and SC conducted the inter-comparison
of ground-based AOD with MODIS DB products and performed the S5P/TROPOMI data extraction and visualization. MH-G
and AL developed the SSA retrieval algorithm and conducted the analysis on SSA results. SC, FB and MC supervised the
maintenance and operation of ground-based instruments as well as the acquisition and curation of the respective data sets. OD
680 contributed in the discussions on the SSA analysis and the inter-comparisons with MODIS DB. I-PR, SK, GB and FN
supervised the investigation and contributed towards methodological ideas and their presentation. All authors reviewed and
edited the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

685

Acknowledgements. The work has been supported by the European Space Agency (ESA) in the frame of the Instrument Data
quality Evaluation and Assessment Service - Quality Assurance for Earth Observation (IDEAS-QA4EO) project. OD and SK
acknowledge the European Metrology Program for Innovation and Research (EMPIR) within the joint research project EMPIR
MAPP: “Metrology for aerosol optical properties”. The EMPIR is jointly funded by the EMPIR participating countries within
690 EURAMET and the European Union. SK would like to acknowledge the ACTRIS Switzerland project funded by the Swiss
State Secretariat for Education Research and Innovation.

References

- 695 Andrés Hernández, M. D., Hilboll, A., Ziereis, H., Förster, E., Krüger, O. O., Kaiser, K., Schneider, J., Barnaba, F., Vrekoussis, M., Schmidt, J., Huntrieser, H., Blechschmidt, A.-M., George, M., Nenakhov, V., Harlass, T., Holanda, B. A., Wolf, J., Eirenschmalz, L., Krebsbach, M., Pöhlker, M. L., Kalisz Hedegaard, A. B., Mei, L., Pfeilsticker, K., Liu, Y., Koppmann, R., Schlager, H., Bohn, B., Schumann, U., Richter, A., Schreiner, B., Sauer, D., Baumann, R., Mertens, M., Jöckel, P., Kilian, M., Stratmann, G., Pöhlker, C., Campanelli, M., Pandolfi, M., Sicard, M., Gómez-Amo, J. L., Pujadas, M., Bigge, K., Kluge, F., Schwarz, A., Daskalakis, N., Walter, D., Zahn, A., Pöschl, U., Bönisch, H., Borrmann, S., Platt, U., and
- 700 Burrows, J. P.: Overview: On the transport and transformation of pollutants in the outflow of major population centres – observational data from the EMERGe European intensive operational period in summer 2017, Atmos. Chem. Phys., 22, 5877–5924, <https://doi.org/10.5194/acp-22-5877-2022>, 2022.
- Arola, A., and Koskela, T.: On the sources of bias in aerosol optical depth retrieval in the UV range, *J. Geophys. Res.*, 109, D08209, doi:10.1029/2003JD004375, 2004.
- 705 Barnaba, F., Angelini, F., Curci, G., and Gobbi, G. P.: An important fingerprint of wildfires on the European aerosol load, Atmos. Chem. Phys., 11, 10487–10501, [10.5194/acp-11-10487-2011](https://doi.org/10.5194/acp-11-10487-2011), 2011.
- Barnaba, F., Bolignano, A., Di Liberto, L., Morelli, M., Lucarelli, F., Nava, S., Perrino, C., Canepari, S., Basart, S., Costabile, F., Dionisi, D., Ciampichetti, S., Sozzi, R., Gobbi, G.P.: Desert dust contribution to PM10 loads in Italy: Methods and recommendations addressing the relevant European Commission Guidelines in support to the Air Quality Directive
- 710 2008/50, Atmos. Environ., 161, 288–305, 2017.
- Bellini, A., Diémoz, H., Di Liberto, L., Gobbi, G.P., and Barnaba, F.: Monitoring the aerosol vertical profiles within the Italian Automated Lidar Ceilometer network (Alicenet): retrievals and applications for air quality and multi risk warning systems, in preparation, 2023.
- Benedetti, A., Reid, J. S., Knippertz, P., Marsham, J. H., Di Giuseppe, F., Rémy, S., Basart, S., Boucher, O., Brooks, I. M.,
- 715 Menut, L., Mona, L., Laj, P., Pappalardo, G., Wiedensohler, A., Baklanov, A., Brooks, M., Colarco, P. R., Cuevas, E., da Silva, A., Escribano, J., Flemming, J., Huneus, N., Jorba, O., Kazadzis, S., Kinne, S., Popp, T., Quinn, P. K., Sekiyama, T. T., Tanaka, T., and Terradellas, E.: Status and future of numerical atmospheric aerosol prediction with a focus on data requirements, *Atmos. Chem. Phys.*, 18, 10615–10643, <https://doi.org/10.5194/acp-18-10615-2018>, 2018.
- Boersma, K. F., Eskes, H. J., and Brinkma, E. J.: Error analysis for tropospheric NO₂ retrieval from space, *J. Geophys. Res.*,
- 720 109, D04311, doi:10.1029/2003JD003962, 2004.
- Boersma, K. F., Jacob, D. J., Eskes, H. J., Pinder, R. W., Wang, J., and van der A, R. J.: Inter-comparison of SCIAMACHY and OMI tropospheric NO₂ columns: Observing the diurnal evolution of chemistry and emissions from space, *J. Geophys. Res.*, 113, 1–14, doi:10.1029/2007JD008816, 2008.

- Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veeffkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitão, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO₂ column retrieval algorithm for the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 4, 1905–1928, <https://doi.org/10.5194/amt-4-1905-2011>, 2011.
- Burrows, J.P., Dehn, A., Deters, B., Himmelmann, S., Richter, A., Voigt, S., and Orphal, J.: Atmospheric remote-sensing reference data from GOME: Part 1. Temperature-dependent absorption cross-sections of NO₂ in the 231-794 nm range" *J. Quant. Spectrosc. Radiat. Transfer*, 60, 1025-1031, 1998.
- Campanelli, M., Nakajima, T., and Olivieri, B.: Determination of the solar calibration constant for a sun-sky radiometer: proposal of an in-situ procedure, *Appl. Opt.* 43, 651-659, 2004.
- Campanelli, M., Iannarelli, A.M., Mevi, G., Casadio, S., Diémoz, H., Finardi, S., Dinoi, A., Castelli, E., di Sarra, A., Di Bernardino, A., Casasanta, G., Bassani, C., Siani, A.M., Cacciani, M., Barnaba, F., Di Liberto, L., Argentini, S.: A wide-ranging investigation of the COVID-19 lockdown effects on the atmospheric composition in various Italian urban sites (AER – LOCUS), *Urban Climate*, Volume 39, 100954, ISSN 2212-0955, <https://doi.org/10.1016/j.uclim.2021.100954>, 2021.
- [Campanelli, M., Diémoz, H., Siani, A. M., di Sarra, A., Iannarelli, A. M., Kudo, R., Fasano, G., Casasanta, G., Tofful, L., Cacciani, M., Sanò, P., and Dietrich, S.: Aerosol optical characteristics in the urban area of Rome, Italy, and their impact on the UV index, *Atmos. Meas. Tech.*, 15, 1171–1183, <https://doi.org/10.5194/amt-15-1171-2022>, 2022.](#)
- Cede, A., Tiefengraber, M., Dehn, A., Lefer, B., von Bismarck, J., Casadio, S., Abuhassan, N., Swap, R., and Valin, L.: Operational satellite validation with data from the Pandonia Global Network (PGN), EGU General Assembly 2020, Online, 4–8 May 2020, EGU2020-13850, <https://doi.org/10.5194/egusphere-egu2020-13850>, 2020.
- Cede, A.: Manual for Blick Software Suite 1.8, 10 Sep 2021, Issue 1.8-4, available at: https://www.pandonia-global-network.org/wp-content/uploads/2021/09/BlickSoftwareSuite_Manual_v1-8-4.pdf (last access: 22 April 2022), 2021.
- [Chen, C., Dubovik, O., Fuertes, D., Litvinov, P., Lapyonok, T., Lopatin, A., et al.: Validation of GRASP Algorithm Product from POLDER/PARASOL Data and Assessment of Multi-Angular Polarimetry Potential for Aerosol Monitoring, *Earth Syst. Sci. Data* 12, 3573–3620, \[doi:10.5194/essd-12-3573-2020\]\(https://doi.org/10.5194/essd-12-3573-2020\), 2020.](#)
- [Chen, C., Dubovik, O., Litvinov, P., Fuertes, D., Lopatin, A., Lapyonok, T., Matar, C., Karol, Y., Fischer, J., Preusker, R., Hangler, A., Aspetsberger, M., Bindreiter, L., Marth, D., Chimot, J., Fougnie, B., Marbach, T., Bojkov, B.: Properties of aerosol and surface derived from OLCI/Sentinel-3A using GRASP approach: Retrieval development and preliminary validation, *Remote Sensing of Environment*, 280, 113142, \[Doi:10.1016/j.rse.2022.113142\]\(https://doi.org/10.1016/j.rse.2022.113142\), 2022.](#)
- Chu, D., Kaufman, Y. J., Ichoku, C., Remer, L., Tanre, D., and Holben, B. N.: Validation of MODIS aerosol optical depth retrieval over land, *Geophys. Res. Lett.*, 29(12), 1617, [doi:10.1029/2001GL013205](https://doi.org/10.1029/2001GL013205), 2002.
- Ciardini, V., Di Iorio, T., Di Liberto, L., Tirelli, C., Casasanta, G., di Sarra, A., Fiocco, G., Fuà, D., and Cacciani, M.: Seasonal variability of tropospheric aerosols in Rome, *Atmos. Res.*, 118, 205–214, <https://doi.org/10.1016/j.atmosres.2012.06.026>, 2012.

- Corr, C. A., Krotkov, N., Madronich, S., Slusser, J. R., Holben, B., Gao, W., Flynn, J., Lefer, B., and Kreidenweis, S. M.: Retrieval of aerosol single scattering albedo at ultraviolet wavelengths at the T1 site during MILAGRO, *Atmos. Chem. Phys.*, 9, 5813–5827, doi:10.5194/acp-9-5813-2009, 2009.
- 760 [Cuevas, E., Romero-Campos, P. M., Kouremeti, N., Kazadzis, S., Räisänen, P., García, R. D., Barreto, A., Guirado-Fuentes, C., Ramos, R., Toledano, C., Almansa, F., and Gröbner, J.: Aerosol optical depth comparison between GAW-PFR and AERONET-Cimel radiometers from long-term \(2005–2015\) 1 min synchronous measurements, *Atmos. Meas. Tech.*, 12, 4309–4337, <https://doi.org/10.5194/amt-12-4309-2019>, 2019.](#)
- Di Bernardino, A., Iannarelli, A. M., Casadio, S., Mevi, G., Campanelli, M., Casasanta, G., Cede, A., Tiefengraber, M., Siani, A. M., Spinei, E., and Cacciani, M.: On the effect of sea breeze regime on aerosols and gases properties in the urban area of Rome, Italy, *Urban Climate*, 37, 100842, <https://doi.org/10.1016/j.uclim.2021.100842>, 2021.
- 765 Di Ianni A, Costabile F, Barnaba F, Di Liberto L, Weinhold K, Wiedensohler A, Struckmeier C, Drewnick F, Gobbi GP.: Black Carbon Aerosol in Rome (Italy): Inference of a Long-Term (2001–2017) Record and Related Trends from AERONET Sun-Photometry Data. *Atmosphere*. 9(3), 81, <https://doi.org/10.3390/atmos9030081>, 2018.
- 770 Di Tomaso, E., Escribano, J., Basart, S., Ginoux, P., Macchia, F., Barnaba, F., Benincasa, F., Bretonnière, P.-A., Buñuel, A., Castrillo, M., Cuevas, E., Formenti, P., Gonçalves, M., Jorba, O., Klose, M., Mona, L., Montané Pinto, G., Mytilinaios, M., Obiso, V., Olid, M., Schutgens, N., Votsis, A., Werner, E., and Pérez García-Pando, C.: The MONARCH high-resolution reanalysis of desert dust aerosol over Northern Africa, the Middle East and Europe (2007–2016), *Earth Syst. Sci. Data*, 14, 2785–2816, <https://doi.org/10.5194/essd-14-2785-2022>, 2022.
- 775 Diémoz, H., Siani, A. M., Casadio, S., Iannarelli, A. M., Casale, G. R., Savastiouk, V., Cede, A., Tiefengraber, M., and Müller, M.: Advanced NO₂ retrieval technique for the Brewer spectrophotometer applied to the 20-year record in Rome, Italy, *Earth Syst. Sci. Data*, 13, 4929–4950, <https://doi.org/10.5194/essd-13-4929-2021>, 2021.
- Doppler, L., Carbajal-Henken, C., Pelon, J., Ravetta, F., and Fischer, J.: Extension of radiative transfer code MOMO, matrix-operator model to the thermal infrared—Clear air validation by comparison to RTTOV and application to CALIPSO-IIR. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 144, 49-67, 2014a.
- 780 Doppler, L., Preusker, R., Bennartz, R., and Fischer, J.: k-bin and k-IR: k-distribution methods without correlation approximation for non-fixed instrument response function and extension to the thermal infrared—Applications to satellite remote sensing. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 133, 382-395, 2014b.
- Drosoglou, T., Bais, A. F., Zyrichidou, I., Kouremeti, N., Poupkou, A., Liora, N., Giannaros, C., Koukouli, M. E., Balis, D., and Melas, D.: Comparisons of ground-based tropospheric NO₂ MAX-DOAS measurements to satellite observations with the aid of an air quality model over the Thessaloniki area, Greece, *Atmos. Chem. Phys.*, 17, 5829–5849, <https://doi.org/10.5194/acp-17-5829-2017>, 2017.
- 785 Dubovik, O. and King, M. D.: A flexible inversion algorithm for retrieval of aerosol optical properties from Sun and sky radiance measurements, *J. Geophys. Res.*, 105, 20,673-20,696, 2000.

- 790 Dubovik, O., Smirnov, A., Holben, B. N., King, M. D., Kaufman, Y. J., Eck, T. F., and Slutsker, I.: Accuracy assessments of aerosol optical properties retrieved from AERONET Sun and sky-radiance measurements, *J. Geophys. Res.*, 105, 9791–9806, 2000.
- ~~Dubovik, O., Holben, B. N., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanré, D., and Slutsker, I.: Variability of absorption and optical properties of key aerosol types observed in worldwide locations, *J. Atmos. Sci.*, 59, 590–608, 2002.~~
- 795 Dubovik, O.: Optimization of Numerical Inversion in Photopolarimetric Remote Sensing, in: Photopolarimetry in Remote Sensing, edited by: Videen, G., Yatskiv, Y., and Mishchenko, M., Kluwer Academic Publishers, Dordrecht, Netherlands, 65–106, 2004.
- Dubovik, O., Herman, M., Holdak, A., Lapyonok, T., Tanré, D., Deuzé, J. L., Ducos, F., Sinyuk, A., and Lopatin, A.: Statistically optimized inversion algorithm for enhanced retrieval of aerosol properties from spectral multi-angle polarimetric satellite observations, *Atmos. Meas. Tech.*, 4, 975–1018, <https://doi.org/10.5194/amt-4-975-2011>, 2011.
- 800 Dubovik, O., Fuertes, D., Litvinov, P., Lopatin, A., Lapyonok, T., Dubovik, I., et al.: A comprehensive description of multi-term LSM for applying multiple a priori constraints in problems of atmospheric remote sensing: GRASP algorithm, concept, and applications. *Front. Remote Sens*, 2021.
- Eck T. F., Holben, B. N., Reid, J. S., Dubovik, O., Kinne, S., Smirnov, A., O'Neill, N. T., and Slutsker, I.: The wavelength dependence of the optical depth of biomass burning, urban and desert dust aerosols, *J. Geophys. Res.*, 104, 31,333–31,350, 805 1999.
- Eck, T. F., Holben, B. N., Ward, D. E., Mukelabai, M. M., Dubovik, O., Smirnov, A., Schafer, J. S., Hsu, N. C., Piketh, S. J., Queface, A., and Roux, J. L.: Variability of biomass burning aerosol optical characteristics in southern Africa during the SAFARI 2000 dry season campaign and a comparison of single scattering albedo estimates from radiometric 810 measurements, *J. Geophys. Res.-Atmos.*, 108, 2156–2202, doi:10.1029/2002JD002321, 2003.
- Eskes, H., van Geffen, J., Boersma, F., Eichmann, K.-U., Apituley, A., Pedergnana, M., Sneep, M., Veeffkind, J. P., and Loyola, D.: Sentinel-5 precursor/TROPOMI Level 2 Product User Manual Nitrogen dioxide, Tech. Rep. S5PKNMI-L2-0021-MA, Koninklijk Nederlands Meteorologisch Instituut (KNMI), CI-7570-PUM, issue 4.1.0, available at: <https://sentinel.esa.int/documents/247904/2474726/Sentinel-5P-Level-2-Product-User-Manual-Nitrogen-Dioxide.pdf> 815 (last access: 21 October 2022), 2022.
- Eskes, H. J. and Eichmann, K.-U.: S5P Mission Performance Centre Nitrogen Dioxide [L2_NO2_] Readme, issue 2.2, version 02.04.00, 21 pp., 17 November 2021, available at: <https://sentinel.esa.int/documents/247904/3541451/Sentinel-5P-Nitrogen-Dioxide-Level-2-Product-Readme-File> (last access: 21 October 2022), 2022.
- Espinosa, W. R., Remer, L. A., Dubovik, O., Ziemba, L., Beyersdorf, A., Orozco, D., Schuster, G., Lapyonok, T., Fuertes, D., and Martins, J. V.: Retrievals of aerosol optical and microphysical properties from Imaging Polar Nephelometer scattering measurements, *Atmos. Meas. Tech.*, 10, 811–824, <https://doi.org/10.5194/amt-10-811-2017>, 2017.
- 820

Espinosa, W. R., Martins, J. V., Remer, L. A., Dubovik, O., Lapyonok, T., Fuertes, D., et al.: Retrievals of Aerosol Size Distribution, Spherical Fraction, and Complex Refractive Index from Airborne In Situ Angular Light Scattering and Absorption Measurements, *J. Geophys. Res. Atmos.* 124, 7997–8024, doi:10.1029/2018JD030009, 2019.

825 Estellés, V., Campanelli, M., Smyth, T. J., Utrillas, M. P., and Martínez-Lozano, J. A.: Evaluation of the new ESR network software for the retrieval of direct sun products from CIMEL CE318 and PREDE POM01 sun-sky radiometers, *Atmos. Chem. Phys.*, 12, 11619–11630, <https://doi.org/10.5194/acp-12-11619-2012>, 2012.

Fan, C., Li, Z., Li, Y., Dong, J., van der A, R., and de Leeuw, G.: Variability of NO₂ concentrations over China and effect on air quality derived from satellite and ground-based observations, *Atmos. Chem. Phys.*, 21, 7723–7748, 830 <https://doi.org/10.5194/acp-21-7723-2021>, 2021.

Flynn, C. M., Pickering, K. E., Crawford, J. H., Lamsal, L., Krotkov, N., Herman, J., Weinheimer, A., Chen, G., Liu, X., Szykman, J., Tsay, S.-C., Loughner, C., Hains, J., Lee, P., Dickerson, R. R., Stehr, J. W., and Brent, L.: Relationship between column-density and surface mixing ratio: Statistical analysis of O₃ and NO₂ data from the July 2011 Maryland DISCOVER-AQ mission, *Atmos. Environ.*, 92, 429–441, <https://doi.org/10.1016/j.atmosenv.2014.04.041>, 2014.

835 Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korkin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 database – automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements, *Atmos. Meas. Tech.*, 12, 169–209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.

840 Gkikas, A., Proestakis, E., Amiridis, V., Kazadzis, S., Di Tomaso, E., Tsekeri, A., Marinou, E., Hatzianastassiou, N., and Pérez García-Pando, C.: ModIs Dust AeroSol (MIDAS): a global fine-resolution dust optical depth data set, *Atmos. Meas. Tech.*, 14, 309–334, <https://doi.org/10.5194/amt-14-309-2021>, 2021.

Gobbi, G.P., Angelini F., Barnaba F., Costabile F., Baldasano J.M., Basart S., Sozzi R., Bolignano A.: Changes in particulate matter physical properties during Saharan advections over Rome (Italy): a four-year study, 2001–2004. *Atmos Chem Phys* 13:7395–7404. <https://doi.org/10.5194/acp-13-7395-2013>, 2017.

845 Gobbi, G.P., Barnaba, F., Di Liberto, L., Bolignano, A., Lucarelli, F., Nava, S., Perrino, C., Pietrodangelo, A., Basart, S., Costabile, F., Dionisi, D., Rizza, U., Canepari, S., Sozzi, R., Morelli, M., Manigrasso, M., Drewnick, F., Struckmeier, C., Poenitz, K., Wille, H.: An inclusive view of Saharan dust advections to Italy and the Central Mediterranean, *Atmospheric Environment*, 201, 242-256, 10.1016/j.atmosenv.2019.01.002, 2019.

850 Green, M, Kondragunta, S, Ciren, P, Xu, C.: Comparison of GOES and MODIS aerosol optical depth (AOD) to aerosol robotic network (AERONET) AOD and IMPROVE PM_{2.5} mass at Bondville, Illinois, *J Air Waste Manag Assoc.*, 59(9):1082-91, doi: 10.3155/1047-3289.59.9.1082, 2009.

Herman, J., Cede, A., Spinei, E., Mount, G., Tzortziou, M., and Abuhassan, N.: NO₂ column amounts from ground-based Pandora and MFDOAS spectrometers using the direct-sun DOAS technique: Intercomparisons and application to OMI 855 validation, *J. Geophys. Res.*, 114, D13307, doi:10.1029/2009JD011848, 2009.

- Herman, J., Spinei, E., Fried, A., Kim, J., Kim, J., Kim, W., Cede, A., Abuhassan, N., and Segal-Rozenhaimer, M.: NO₂ and HCHO measurements in Korea from 2012 to 2016 from Pandora spectrometer instruments compared with OMI retrievals and with aircraft measurements during the KORUS-AQ campaign, *Atmos. Meas. Tech.*, 11, 4583–4603, <https://doi.org/10.5194/amt-11-4583-2018>, 2018.
- 860 [Herman, J., Abuhassan, N., Kim, J., Kim, J., Dubey, M., Raponi, M., and Tzortziou, M.: Underestimation of column NO₂ amounts from the OMI satellite compared to diurnally varying ground-based retrievals from multiple PANDORA spectrometer instruments, *Atmos. Meas. Tech.*, 12, 5593–5612, <https://doi.org/10.5194/amt-12-5593-2019>, 2019.](#)
- [Herrerias, M., Román, R., Cazorla, A., Toledano, C., Lyamani, H., Torres, B., et al.: Evaluation of Retrieved Aerosol Extinction Profiles Using as Reference the Aerosol Optical Depth Differences between Various Heights, *Atmos. Res.* 230, 104625, \[doi:10.1016/j.atmosres.2019.104625\]\(https://doi.org/10.1016/j.atmosres.2019.104625\), 2019.](#)
- 865
- Hobbs, P. V.: Aerosol-cloud interactions, in: *Aerosol-Cloud-Climate Interactions*, Academic, San Diego, California, 1993.
- Holben, B.N., Eck, T.F., Slutsker, I.A., Tanre, D., Buis, J.P., Setzer, A., Vermote, E., et al.: AERONET—A federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.* 66 (1), 1–16, 1998.
- Hsu, N. C., Tsay, S.-C., King, M. D., and Herman, J. R.: Aerosol Properties Over Bright-Reflecting Source Regions, *IEEE T. Geosci. Remote Sens.*, 42, 557–569, <https://doi.org/10.1109/TGRS.2004.824067>, 2004.
- 870
- Hsu, N. C., Tsay, S.-C., King, M. D., and Herman, J. R.: Deep Blue Retrievals of Asian Aerosol Properties During ACE-Asia, *IEEE T. Geosci. Remote S.*, 44, 3180–3195, <https://doi.org/10.1109/TGRS.2006.879540>, 2006.
- Hsu, N. C., Jeong, M.-J., Bettenhausen, C., Sayer, A. M., Hansell, R., Seftor, C. S., Huang, J., and Tsay, S.-C.: Enhanced Deep Blue aerosol retrieval algorithm: The second generation, *J. Geophys. Res.-Atmos.*, 118, 9296–9315, <https://doi.org/10.1002/jgrd.50712>, 2013.
- 875
- Iannarelli, A. M., Di Bernardino, A., Casadio, S., Bassani, C., Cacciani, M., Campanelli, M., Casasanta, G., Cadau, E., Diémoz, H., Mevi, G., Siani, A. M., Cardaci, M., Dehn, A., and Goryl, P.: The Boundary Layer Air Quality-Analysis Using Network of Instruments (BAQUNIN) Supersite for Atmospheric Research and Satellite Validation over Rome Area, *B. Am. Meteorol. Soc.*, E599–E618, <https://doi.org/10.1175/BAMS-D-21-0099.1>, 2021.
- 880 [IPCC, 2021: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* \[Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou \(eds.\)\]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, In press, \[doi:10.1017/9781009157896\]\(https://doi.org/10.1017/9781009157896\).](#)
- 885 [IPCC: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.](#)

- ISTAT: Annual demographic balance 2021,
 890 <http://demo.istat.it/bilmens/query.php?anno=2021&lingua=ita&Rip=S3&Reg=R12&Pro=P058&Com=91&submit=Tavo>
 la, (last access: 21 August 2022), 2021.
- Jethva, H., Torres, O., and Ahn, C.: Global assessment of OMI aerosol single-scattering albedo using ground-based AERONET
 inversion, *J. Geophys. Res.-Atmos.*, 119, 9020–9040, doi:10.1002/2014JD021672, 2014.
- Kazadzis, S., Raptis, P., Kouremeti, N., Amiridis, V., Arola, A., Gerasopoulos, E., and Schuster, G. L.: Aerosol absorption
 895 retrieval at ultraviolet wavelengths in a complex environment, *Atmos. Meas. Tech.*, 9, 5997–6011,
<https://doi.org/10.5194/amt-9-5997-2016>, 2016.
- Kazadzis, S., Kouremeti, N., Nyeki, S., Gröbner, J., and Wehrli, C.: The World Optical Depth Research and Calibration Center
 (WORCC) quality assurance and quality control of GAW-PFR AOD measurements, *Geosci. Instrum. Method. Data Syst.*,
 7, 39–53, <https://doi.org/10.5194/gi-7-39-2018>, 2018a.
- 900 Kazadzis, S., Kouremeti, N., Diémoz, H., Gröbner, J., Forgan, B. W., Campanelli, M., Estellés, V., Lantz, K., Michalsky, J.,
 Carlund, T., Cuevas, E., Toledano, C., Becker, R., Nyeki, S., Kosmopoulos, P. G., Tatsiankou, V., Vuilleumier, L., Denn,
 F. M., Ohkawara, N., Ijima, O., Goloub, P., Raptis, P. I., Milner, M., Behrens, K., Barreto, A., Martucci, G., Hall, E.,
 Wendell, J., Fabbri, B. E., and Wehrli, C.: Results from the Fourth WMO Filter Radiometer Comparison for aerosol optical
 depth measurements, *Atmos. Chem. Phys.*, 18, 3185–3201, <https://doi.org/10.5194/acp-18-3185-2018>, 2018b.
- 905 [King, M. D., and Dubovik, O.: Determination of Aerosol Optical Properties from Inverse Methods, *Aerosol Remote Sensing*,
 Editors J. Lenoble, L. Remer, and D. Tanré \(Chischester, UK: Springer, Praxis Publishers\), 101–136. doi:10.1007/978-3-
 642-17725-5_5, 2013.](#)
- Kreher, K., Van Roozendaal, M., Hendrick, F., Apituley, A., Dimitropoulou, E., Frieß, U., Richter, A., Wagner, T., Lampel,
 J., Abuhassan, N., Ang, L., Anguas, M., Bais, A., Benavent, N., Bösch, T., Bogner, K., Borovski, A., Bruchkouski, I.,
 910 Cede, A., Chan, K. L., Donner, S., Drosoglou, T., Fayt, C., Finkenzeller, H., Garcia-Nieto, D., Gielen, C., Gómez-Martín,
 L., Hao, N., Henzing, B., Herman, J. R., Hermans, C., Hoque, S., Irie, H., Jin, J., Johnston, P., Khayyam Butt, J., Khokhar,
 F., Koenig, T. K., Kuhn, J., Kumar, V., Liu, C., Ma, J., Merlaud, A., Mishra, A. K., Müller, M., Navarro-Comas, M.,
 Ostendorf, M., Pazmino, A., Peters, E., Pinardi, G., Pinharanda, M., Piter, A., Platt, U., Postlyakov, O., Prados-Roman,
 C., Puentedura, O., Querel, R., Saiz-Lopez, A., Schönhardt, A., Schreier, S. F., Seyler, A., Sinha, V., Spinei, E., Strong,
 915 K., Tack, F., Tian, X., Tiefengraber, M., Tirpitz, J.-L., van Gent, J., Volkamer, R., Vrekoussis, M., Wang, S., Wang, Z.,
 Wenig, M., Wittrock, F., Xie, P. H., Xu, J., Yela, M., Zhang, C., and Zhao, X.: Intercomparison of NO₂, O₄, O₃ and HCHO
 slant column measurements by MAX-DOAS and zenith-sky UV–visible spectrometers during CINDI-2, *Atmos. Meas.*
Tech., 13, 2169–2208, <https://doi.org/10.5194/amt-13-2169-2020>, 2020.
- [Lambert, J.-C., A. Keppens, S. Compernelle, K.-U. Eichmann, M. de Graaf, D. Hubert, Q. Kleipool, B. Langerock, M.K. Sha,
 920 T. Verhoelst, T. Wagner, C. Ahn, A. Argyrouli, D. Balis, K.L. Chan, I. De Smedt, H. Eskes, A.M. Fjæraa, K. Garane, J.F.
 Gleason, F. Goutail, J. Granville, P. Hedelt, K.-P. Heue, G. Jaross, M.L. Koukouli, J. Landgraf, R. Lutz, S. Nanda, S.
 Niemeijer, A. Pazmiño, G. Pinardi, J.-P. Pommereau, A. Richter, N. Rozemeijer, M. Sneep, D. Stein Zweers, N. Theys, G.](#)

- 925 [Tilstra, O. Torres, P. Valks, J. van Geffen, C. Vigouroux, P. Wang, and M. Weber: Quarterly Validation Report of the Copernicus Sentinel-5 Precursor Operational Data Products #13: April 2018 – December 2021, S5P MPC Routine Operations Consolidated Validation Report series, issue #13, version 13.01.00, 189 pp., 17 December 2021, available at: \[https://mpc-vdaf.tropomi.eu/ProjectDir/reports/pdf/S5P-MPC-IASB-ROCVR-13.00.10-20211217_signed.pdf\]\(https://mpc-vdaf.tropomi.eu/ProjectDir/reports/pdf/S5P-MPC-IASB-ROCVR-13.00.10-20211217_signed.pdf\) \(last access: 12 February 2023\), 2021.](https://mpc-vdaf.tropomi.eu/ProjectDir/reports/pdf/S5P-MPC-IASB-ROCVR-13.00.10-20211217_signed.pdf)
- Lamsal, L. N., Janz, S. J., Krotkov, N. A., Pickering, K. E., Spurr, R. J. D., Kowalewski, M. G., Loughner, C. P., Crawford, J. H., Swartz, W. H., and Herman, J. R.: High-resolution NO₂ observations from the Airborne Compact Atmospheric Mapper: Retrieval and validation, *J. Geophys. Res.*, 122, 1953–1970, <https://doi.org/10.1002/2016JD025483>, 2017.
- 930 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367–371, <https://doi.org/10.1038/nature15371>, 2015.
- Levy, R. C., Remer, L. A., Kleidman, R. G., Mattoo, S., Ichoku, C., Kahn, R., and Eck, T. F.: Global evaluation of the Collection 5 MODIS dark-target aerosol products over land, *Atmos. Chem. Phys.*, 10, 10399–10420, doi:10.5194/acp-10-10399-2010, 2010.
- 935 Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmos. Meas. Tech.*, 6, 2989–3034, <https://doi.org/10.5194/amt-6-2989-2013>, 2013.
- Li, J., Carlson, B. E., Lacis, A. A.: How well do satellite AOD observations represent the spatial and temporal variability of PM_{2.5} concentration for the United States?, *Atmospheric Environment*, Volume 102, Pages 260-273, ISSN 1352-2310, <https://doi.org/10.1016/j.atmosenv.2014.12.010>, 2015.
- 940 Logothetis, S.-A., Salamalikis, V., Gkikas, A., Kazadzis, S., Amiridis, V., and Kazantzidis, A.: 15-year variability of desert dust optical depth on global and regional scales, *Atmos. Chem. Phys.*, 21, 16499–16529, <https://doi.org/10.5194/acp-21-16499-2021>, 2021.
- 945 [Lopatin, A., Dubovik, O., Chaikovskiy, A., Goloub, P., Lapyonok, T., Tanré, D., and Litvinov, P.: Enhancement of aerosol characterization using synergy of lidar and sun-photometer coincident observations: the GARRLiC algorithm, *Atmos. Meas. Tech.*, 6, 2065–2088, <https://doi.org/10.5194/amt-6-2065-2013>, 2013.](https://doi.org/10.5194/amt-6-2065-2013)
- 950 [Lopatin, A., Dubovik, O., Fuertes, D., Stenchikov, G., Lapyonok, T., Veselovskii, I., Wienhold, F. G., Shevchenko, I., Hu, Q., and Parajuli, S.: Synergy processing of diverse ground-based remote sensing and in situ data using the GRASP algorithm: applications to radiometer, lidar and radiosonde observations, *Atmos. Meas. Tech.*, 14, 2575–2614, <https://doi.org/10.5194/amt-14-2575-2021>, 2021.](https://doi.org/10.5194/amt-14-2575-2021)
- Martins, D. K., Najjar, R. G., Tzortziou, M., Abuhassan, N., Thompson, A. M., and Kollonige, D. E.: Spatial and temporal variability of ground and satellite column measurements of NO₂ and O₃ over the Atlantic Ocean during the Deposition of Atmospheric Nitrogen to Coastal Ecosystems Experiment, *J. Geophys. Res.*, 121, 14175–14187, <https://doi.org/10.1002/2016JD024998>, 2016.

- 955 Molina, C., Toro A., R., Manzano, C.A., Canepari, S., Massimi, L., Leiva-Guzmán, M.A. Airborne Aerosols and Human Health: Leapfrogging from Mass Concentration to Oxidative Potential. *Atmosphere*, 11, 917. <https://doi.org/10.3390/atmos11090917>, 2020.
- Müller, M., Gebetsberger, M., Tiefengraber, M., and Cede, A.: LuftBlick Report 2019003, Fiducial Reference Measurements for Air Quality, Calibration Procedures Document, Tech. rep., LuftBlick, https://www.pandonia-global-network.org/wp-content/uploads/2021/01/LuftBlick_FRM4AQ_CPD_RP_2019003_v4.0.pdf (last access: 22 April 2022), 960 2020.
- Nakajima, T., Tonna, G. and Rao, R., Boi, P., Kaufman, Y., Holben, B.: Use of sky brightness measurements from ground for remote sensing of particulate polydispersions, *Appl. Opt.*, 35, pp. 2672-2686, 1996.
- Nakajima, T., Campanelli, M., Che, H., Estellés, V., Irie, H., Kim, S.-W., Kim, J., Liu, D., Nishizawa, T., Pandithurai, G., 965 Soni, V. K., Thana, B., Tugjurn, N.-U., Aoki, K., Go, S., Hashimoto, M., Higurashi, A., Kazadzis, S., Khatri, P., Kouremeti, N., Kudo, R., Marengo, F., Momoi, M., Ningombam, S. S., Ryder, C. L., Uchiyama, A., and Yamazaki, A.: An overview of and issues with sky radiometer technology and SKYNET, *AMT*, 13, 4195-4218, 2020.
- [Park, J.-U., Park J.-S., Santana Diaz D., Gebetsberger M., Müller M., Shalaby L., Tiefengraber M., Kim H.-J., Park S.-S., Song C.-K., and Kim S.-W.: Spatiotemporal inhomogeneity of total column NO₂ in a polluted urban area inferred from TROPOMI and Pandora intercomparisons, *GIScience & Remote Sensing*, 59:1, 354-373, DOI: 10.1080/15481603.2022.2026640, 2022.](#) 970
- Platt, U.: Differential optical absorption spectroscopy (DOAS), in: *Air Monitoring by Spectroscopic Techniques*, Chem. Anal. Ser. 127, John Wiley, New York, 27–84, 1994.
- Platt, U. and Stutz, J.: *Differential Optical Absorption Spectroscopy: Principles and Applications*, Springer, Germany, Berlin, 975 2008.
- ~~Raptis, I.P., Kazadzis, S., Amiridis, V., Gkikas, A., Gerasopoulos, E., and Mihalopoulos, N.: A decade of aerosol optical properties measurements over Athens, Greece. *Atmosphere*, 11(2), p.154, 2020.~~
- Remer, L., Kaufman, Y., Tanre, D., Mattoo, S., Chu, D., Martins, J., et al.: The MODIS aerosol algorithm, products, and validation, *J. Atmos. Sci.*, 62(4), 947–973, 2005.
- 980 Richter, A., Burrows, J. P., Nüszlig, H., Granier, C., and Niemeier, U.: Increase in tropospheric nitrogen dioxide over China observed from space (and supplementary discussion on: Error estimates for changes in tropospheric NO₂ columns as derived from satellite measurements), *Nature*, 437, 129–132, doi:10.1038/nature04092, 2005.
- [Román, R., Torres, B., Fuertes, D., Cachorro, V., Dubovik, O., Toledano, C., Cazorla, A., Barreto, A., Bosch, J., Lapyonok, T., González, R., Goloub, P., Perrone, M., Olmo, F., de Frutos, A., and Alados-Arboledas, L.: Remote sensing of lunar aureole with a sky camera: Adding information in the nocturnal retrieval of aerosol properties with GRASP code, *Remote Sens. Environ.*, 196, 238–252, <https://doi.org/10.1016/j.rse.2017.05.013>, 2017.](#) 985
- [Román, R., Benavent-Oltra, J. A., Casquero-Vera, J. A., Lopatin, A., Cazorla, A., Lyamani, H., Denjean, C., Fuertes, D., Pérez-Ramírez, D., Torres, B., Toledano, C., Dubovik, O., Cachorro, V. E., de Frutos, A., Olmo, F. J., and Alados-](#)

- 990 [Arboledas, L.: Retrieval of aerosol profiles combining sunphotometer and ceilometer measurements in GRASP code, Atmos. Res., 204, 161–177, 2018.](#)
- [Román, R., Antuña-Sánchez, J. C., Cachorro, V. E., Toledano, C., Torres, B., Mateos, D., Fuertes, D., López, C., González, R., Lapyonok, T., Herreras-Giralda, M., Dubovik, O., and de Frutos, Á. M.: Retrieval of aerosol properties using relative radiance measurements from an all-sky camera, Atmos. Meas. Tech., 15, 407–433, <https://doi.org/10.5194/amt-15-407-2022>, 2022.](#)
- 995 Rosenfeld, D., Andreae, M.O., Asmi, A., Chin, M., de Leeuw, G., Donovan D.P., Kahn, R., Kinne, S., Kivekäs, N., Kulmala, M., Lau, W., Schmidt, K.S., Suni, T., Wagner, T., Wild, M., and Quaas, J.: Global observations of aerosol-cloud-precipitation-climate interactions, Rev. Geophys., 52, 750–808, <https://doi.org/10.1002/2013RG000441>, 2014.
- [Schuster, G. L., Dubovik, O., and Holben, B. N.: Angstrom exponent and bimodal aerosol size distributions, J. Geophys. Res., 111, D07207, <https://doi.org/10.1029/2005JD006328>, 2006.](#)
- 1000 [Schuster, G., Espinosa, W., Ziemba, L., Beyersdorf, A., Rocha-Lima, A., Anderson, B., et al.: A Laboratory Experiment for the Statistical Evaluation of Aerosol Retrieval \(STEAR\) Algorithms, Remote Sensing 11, 498, doi:10.3390/rs11050498, 2019.](#)
- Sherman, J.P., Gupta, P., Levy, R.C. and Sherman, P.J.: An Evaluation of MODIS-Retrieved Aerosol Optical Depth over a Mountainous AERONET Site in the Southeastern US. Aerosol Air Qual. Res. 16: 3243-3255. <https://doi.org/10.4209/aaqr.2015.09.0568>, 2016.
- 1005 Sinyuk, A., Holben, B. N., Eck, T. F., Giles, D. M., Slutsker, I., Korkin, S., Schafer, J. S., Smirnov, A., Sorokin, M., and Lyapustin, A.: The AERONET Version 3 aerosol retrieval algorithm, associated uncertainties and comparisons to Version 2, Atmos. Meas. Tech., 13, 3375–3411, <https://doi.org/10.5194/amt-13-3375-2020>, 2020.
- Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone, Nature, 321, 755–758, <https://doi.org/10.1038/321755a0>, 1986.
- 1010 Takamura T. and Nakajima T., Overview of SKYNET and its activities, Opt. Pura Apl. 37, 3303-3308, 2004.
- [Torres, B., Dubovik, O., Fuertes, D., Schuster, G., Cachorro, V. E., Lapyonok, T., Goloub, P., Blarel, L., Barreto, A., Mallet, M., Toledano, C., and Tanré, D.: Advanced characterisation of aerosol size properties from measurements of spectral optical depth using the GRASP algorithm, Atmos. Meas. Tech., 10, 3743–3781, <https://doi.org/10.5194/amt-10-3743-2017>, 2017.](#)
- 1015 Tzortziou, M., Herman, J. R., Cede, A., and Abuhassan, N.: High precision, absolute total column ozone measurements from the Pandora spectrometer system: Comparisons with data from a Brewer double monochromator and Aura OMI. Journal of Geophysical Research: Atmospheres, 117(D16), 2012.
- Tzortziou, M., Herman, J. R., Ahmad, Z., Loughner, C. P., Abuhassan, N., and Cede, A.: Atmospheric NO₂ dynamics and impact on ocean color retrievals in urban nearshore regions, J. Geophys. Res.-Oceans, 119, 3834–3854, <https://doi.org/10.1002/2014JC009803>, 2014.
- 1020

- Tzortziou, M., Herman, J. R., Cede, A., Loughner, C. P., Abuhassan, N., and Naik, S.: Spatial and temporal variability of ozone and nitrogen dioxide over a major urban estuarine ecosystem, *J. Atmos. Chem.*, 72, 287–20 309, <https://doi.org/10.1007/s10874-013-9255-8>, <https://doi.org/10.1007/s10874-013-9255-8>, 2015.
- 1025 Valentini, S., Barnaba, F., Bernardoni, V., Calzolari, G., Costabile, F., Di Liberto, L., Forello A.C., Gobbi, G.P., Gualtieri, M., Lucarelli F., Nava, S., Petralia, E., Valli, G., Wiedensohler, A., and Vecchi, R.: Classifying aerosol particles through the combination of optical and physical-chemical properties: Results from a wintertime campaign in Rome (Italy). *Atmospheric Research*, 235, 104799, 2020.
- van Geffen, J. H. G. M., Boersma, K. F., Van Roozendaal, M., Hendrick, F., Mahieu, E., De Smedt, I., Sneep, M., and Veefkind, J. P.: Improved spectral fitting of nitrogen dioxide from OMI in the 405–465 nm window, *Atmos. Meas. Tech.*, 8, 1685–1699, <https://doi.org/10.5194/amt-8-1685-2015>, 2015.
- 1030 van Geffen, J.H.G.M.; Eskes, H.J.; Boersma, K.F.; Veefkind, J.P.: TROPOMI ATBD of the Total and Tropospheric NO₂ Data Products, available at: <https://sentinel.esa.int/documents/247904/2476257/Sentinel-5P-TROPOMI-ATBD-NO2-data-products> (last access: 21 October 2022), 2022.
- 1035 Weatherhead, E. C., Reinsel, G.C., Tiao, G.C., Meng, X.-L., Choi, D., Cheang, W.-K., Keller, T., DeLuisi, J., Wuebbles, D.J., Kerr, J.B., Miller, A.J., Oltmans, S.J. and Frederick, J.E.: Factors affecting the detection of trends: Statistical considerations and applications to environmental data, *J. Geophys. Res.*, 103(D14), 17149– 17161, doi:10.1029/98JD00995, 1998.
- Wei, J., Li, Z., Peng, Y., Sun, L.: MODIS Collection 6.1 aerosol optical depth products over land and ocean: validation and comparison, *Atmos. Environ.*, 201, 428–440, <https://doi.org/10.1016/j.atmosenv.2018.12.004>, 2019.
- 1040 [WMO: WMO/GAW Experts Workshop on a Global Surface-Based Network for Long Term Observations of Column Aerosol Optical Properties, GAW Report No. 162, WMO TD No. 1287, available at: https://library.wmo.int/index.php?lvl=notice_display&id=11094, 2005.](https://library.wmo.int/index.php?lvl=notice_display&id=11094)
- Yoon, J., von Hoyningen-Huene, W., Kokhanovsky, A. A., Vountas, M., and Burrows, J. P.: Trend analysis of aerosol optical thickness and Ångström exponent derived from the global AERONET spectral observations, *Atmos. Meas. Tech.*, 5, 1271–1299, <https://doi.org/10.5194/amt-5-1271-2012>, 2012.
- Zhang, J. and Reid, J. S.: A decadal regional and global trend analysis of the aerosol optical depth using a data-assimilation grade over-water MODIS and Level 2 MISR aerosol products, *Atmos. Chem. Phys.*, 10, 10949–10963, <https://doi.org/10.5194/acp-10-10949-2010>, 2010.
- 1050 [Zhao, X., Griffin, D., Fioletov, V., McLinden, C., Cede, A., Tiefengraber, M., Müller, M., Bognar, K., Strong, K., Boersma, F., Eskes, H., Davies, J., Ogyu, A., and Lee, S. C.: Assessment of the quality of TROPOMI high-spatial-resolution NO₂ data products in the Greater Toronto Area, Atmos. Meas. Tech., 13, 2131–2159, https://doi.org/10.5194/amt-13-2131-2020, 2020.](https://doi.org/10.5194/amt-13-2131-2020)

1055 Table 1. Deviation of Pandora total NO₂ column from satellite climatology used for AERONET retrievals and differences in modified AERONET and SKYNET AOD and AE from the standard products over CNR-ISAC and APL-SAP calculated using actual Pandora total NO₂ observations, as well as daily and monthly averaged values of NO₂. Note that the spectral channels used in AERONET retrievals are 380 and 440nm for AOD and 440-870nm for AE, whereas for SKYNET the wavelength channels are 400nm and 400-1020nm for AOD and AE, respectively.

		PGN NO ₂ Actual Measurements					PGN NO ₂ Daily Mean					PGN NO ₂ Monthly Mean				
		AERONET		AERONET	SKYNET	AERONET		AERONET	SKYNET	AERONET		AERONET	SKYNET			
		CNR-ISAC		APL-SAP	APL-SAP	CNR-ISAC		APL-SAP	APL-SAP	CNR-ISAC		APL-SAP	APL-SAP			
NO ₂	Channel [nm]	380	440	380	440	400	380	440	380	440	400	380	440	380	440	400
	% Mean Deviation	61.2 3	64.8 4	63.5 4	64.5 4	-	53.2 2	57.2 4	65.3 0	66.3 74	-	45.4 5	49.6 3	59.4 4	60.7 46	-
	Mean Deviation [DU]	0.163	0.168	0.162	0.163	-	0.141	0.147	0.167	0.169	-	0.120	0.127	0.151	0.153	-
	STD [DU]	0.170	0.171	0.182	0.182	-	0.099	0.101	0.118	0.119	-	0.033	0.035	0.062	0.062	-
	Minimum Deviation [DU]	1.3×10 ⁻⁵	0.3×10 ⁻⁵	2.83×10 ⁻⁶	0.6×10 ⁻⁶	-	0.6×10 ⁻⁵	0.7×10 ⁻⁵	0.1×10 ⁻⁵	0.5×10 ⁻⁵	-	0.6×10 ⁻⁶	0.004	0.010	0.012	-
	Maximum Deviation [DU]	2.066	2.080	2.406	2.410	-	0.803	0.815	0.773	0.777	-	0.297	0.311	0.291	0.293	-
AOD	Channel [nm]	380	440	380	440	400	380	440	380	440	400	380	440	380	440	400
	% Mean Deviation	1.5 42	1.7 695	1.7 677	1.7 682	5.3 275	1.3 30	1.5 40	1.9 886	1.9 897	5.6 597	1.2 164	1.3 44	1.8 44	1.8 14	5.7 684
	Mean Deviation	0.00 327	0.0024	0.00 327	0.0024	0.0074	0.002 3	0.002 4	0.00 328	0.0024	0.00 876	0.002 0	0.00 218	0.00 325	0.002 2	0.0074
	STD	0.00 328	0.0025	0.003 0	0.00 326	0.0034	0.00 216	0.00 215	0.002 0	0.00 217	0.002 2	0.0005	0.0004	0.001 0	0.0009	0.001 4
	Minimum Deviation	0.02×10 ⁻⁶	0.05×10 ⁻⁶	0.04×10 ⁻⁶	0.09×10 ⁻⁷	0.0022	0.01×10 ⁻⁵	0.01×10 ⁻⁵	0.02×10 ⁻⁶	0.07×10 ⁻⁶	0.0029	0.01×10 ⁻⁶	0.06×10 ⁻³	0.0002	0.0002	0.0052
	Maximum Deviation	0.034	0.030	0.040	0.035	0.043	0.013	0.012	0.013	0.011	0.018	0.005	0.004	0.005	0.004	0.010
AE	Spectral Range [nm]	440-870		440-870		400-1020	440-870		440-870		400-1020	440-870		440-870		400-1020
	% Mean Deviation	1.7 34		2.6 552		6.9 607.0	1.4 15		2.7 696		7.6 559	1.2 178		2.9 895		7.9 38
	Mean Deviation	0.019		0.021		0.053	0.016		0.022		0.057	0.012		0.021		0.058
	STD	0.027		0.026		0.036	0.019		0.023		0.041	0.011		0.017		0.044
	Minimum Deviation	0.02×10 ⁻⁶		0.01×10 ⁻⁶		0.002	0.08×10 ⁻⁵		0.01×10 ⁻⁴		0.002	0.12×10 ⁻⁵		0.04×10 ⁻⁵		0.003
	Maximum Deviation	0.309		0.291		0.701	0.215		0.322		0.621	0.139		0.248		0.640

1060


Table 2. Similar to Table 1, using TROPOMI measurements instead of Pandora total NO₂ for the estimation of NO₂ abundance in AERONET and SKYNET aerosol retrievals.

		TROPOMI NO ₂ Actual Measurements					TROPOMI NO ₂ Daily Mean					TROPOMI NO ₂ Monthly Mean				
		AERONET CNR-ISAC		AERONET APL-SAP		SKYNET APL-SAP	AERONET CNR-ISAC		AERONET APL-SAP		SKYNET APL-SAP	AERONET CNR-ISAC		AERONET APL-SAP		SKYNET APL-SAP
NO ₂	Channel [nm]	380	440	380	440	400	380	440	380	440	400	380	440	380	440	400
	% Mean Deviation	19.549	19.81	24.10	24.329	-	18.73	19.01	23.41	23.64	-	6.327	6.63	12.90	13.24	-
	Mean Deviation [DU]	0.053	0.053	0.064	0.064	-	0.051	0.051	0.062	0.062	-	0.017	0.018	0.034	0.035	-
	STD [DU]	0.048	0.048	0.066	0.067	-	0.046	0.046	0.064	0.064	-	0.019	0.018	0.025	0.025	-
	Minimum Deviation [DU]	6×10 ⁻⁵	8×10 ⁻⁶	2×10 ⁻⁵	2×10 ⁻⁵	-	3×10 ⁻⁶	4×10 ⁻⁶	2×10 ⁻⁶	3×10 ⁻⁶	-	2×10 ⁻⁶	3×10 ⁻⁵	5×10 ⁻⁸	9×10 ⁻⁷	-
	Maximum Deviation [DU]	0.408	0.422	0.565	0.567	-	0.398	0.412	0.564	0.566	-	0.103	0.089	0.149	0.151	-
AOD	Channel [nm]	380	440	380	440	400	380	440	380	440	400	380	440	380	440	400
	% Mean Deviation	0.536	0.6574	0.9855	0.845	3.8784	0.517	0.550	0.839	0.814	3.812	0.2169	0.2182	0.538	0.5451	3.9865
	Mean Deviation	0.0009	0.0008	0.0011	0.0009	0.0051	0.0008	0.0007	0.0010	0.0009	0.0051	0.0003	0.0003	0.0006	0.0005	0.0051
	STD	0.0008	0.0007	0.0011	0.0010	0.0017	0.0008	0.0007	0.0011	0.0009	0.0017	0.0003	0.0003	0.0004	0.0004	0.0008
	Minimum Deviation	1×10 ⁻⁶	1×10 ⁻⁷	3×10 ⁻⁷	4×10 ⁻⁷	0.0023	5×10 ⁻⁸	6×10 ⁻⁸	4×10 ⁻⁸	5×10 ⁻⁸	0.0024	3×10 ⁻⁸	5×10 ⁻⁷	7×10 ⁻¹⁰	1×10 ⁻⁸	0.0038
	Maximum Deviation	0.007	0.006	0.009	0.008	0.017	0.007	0.006	0.009	0.008	0.015	0.002	0.001	0.002	0.002	0.008
AE	Spectral Range [nm]	440-870		440-870		400-1020	440-870		440-870		400-1020	440-870		440-870		400-1020
	% Mean Deviation	0.9853		1.645		4.03-963	0.843		1.7674		4.000	0.524		0.616		4.227
	Mean Deviation	0.009		0.012		0.038	0.009		0.011		0.038	0.006		0.005		0.039
	STD	0.011		0.017		0.026	0.010		0.016		0.027	0.004		0.006		0.027
	Minimum Deviation	1×10 ⁻⁷		1×10 ⁻⁷		0.001	2×10 ⁻⁷		7×10 ⁻⁸		0.001	1×10 ⁻⁷		7×10 ⁻⁸		0.002
	Maximum Deviation	0.116		0.239		0.286	0.116		0.238		0.393	0.036		0.090		0.252

1065 **Table 3. AOD and AE trends and their uncertainties for both standard and modified AERONET and SKYNET products over CNR-ISAC and APL-SAP. Note that the spectral channels used in AERONET retrievals are 440 nm for AOD and 440-870 nm for AE, whereas for SKYNET are 400 nm and 400-1020 nm for AOD and AE, respectively. The trend uncertainties refer to the standard error of the regression slope. The differences are calculated on the absolute trend values.**

		AERONET CNR-ISAC		AERONET APL-SAP		SKYNET APL-SAP	
		standard	modified	standard	modified	standard	modified
Number of years		5.7	5.7	5.7	5.7	5.4	5.4
AOD	Trend [/year]	0.00 437	0.0024	0.000 219	0.0005 2	0.00 215	0.00 218
	% trend [/year]	2.04	1. 438	0.14	0.3 0	0.8 2	1.00 99
	Uncertainty	0.0054	0.0054	0.00 549	0.00 549	0.006 2	0.006 2
	Modified – standard		-0.001 2 (-33.1%)		0.0003 3 (174.3%)		0.00024 (15.8%)
AE	Trend [/year]	0.0474	0.04 216	-0.022 1	-0.0181	-0.061 1	-0.05 769
	% trend [/year]	3.8 0	3. 437	-1.8 5	-1.54	-5. 656	-5. 546
	Uncertainty	0.025 2	0.02 657	0.0184	0.01 989	0.02 656	0.0264
	Modified – standard		-0.00 659 (-12.4%)		-0.004 0 (-18.1%)		-0.004 2 (-6.9%)

1070



1075 **Figure 1: Time series of monthly averaged AOD (upper panel) and AE (lower panel) measurements over APL-SAP (AERONET and SKYNET) and CNR-ISAC (AERONET). Note that AERONET AOD and AE correspond to the wavelength channels of 440nm and 440-870nm, respectively, whereas SKYNET AOD and AE refer to 400nm and 400-1020nm, respectively. The shaded areas correspond to the monthly 1-sigma standard deviation.**




Figure 2: Time series of monthly NO₂ total column from Pandora instruments over APL-SAP (blue line) and CNR-ISAC (yellow line). The shaded areas correspond to the 1-sigma standard deviation of the monthly averaged values. The NO₂ concentration is clearly affected by the COVID-19 lockdown during February – May 2020.






Figure 3: Monthly NO₂ total column from Pandora over CNR-ISAC against synchronous APL-SAP observations. The grey shaded area corresponds to the 95% confidence interval of the linear regression fit (red line).


1090






1095

Figure 4: ~~Percentage Time series of the deviation of Pandora total NO₂ data-deviation from AERONET NO₂ climatological values (OMI) for both APL-SAP (left upper panel) and CNR-ISAC (right upper panel) versus Pandora NO₂. The data pairs are grouped in color-sealed bins of bivariate percentage distribution. The corresponding relative frequency distributions of absolute Pandora-OMI deviation for both locations are illustrated in the lower panel. Note that the distribution bars for the two locations are stacked.~~



1100

Figure 5: S5P/TROPOMI summed total NO₂ column averaged for the period 2018-2021, excluding the COVID-19 lockdown period. The data are gridded on a 500m grid. The locations of the two observational sites used in this study are also reported for reference.



1105

Figure 6: The differences of modified AERONET AOD at 440 nm (upper panels) and AE at 440-870 nm (lower panels) over CNR-ISAC (left panels) and APL-SAP (right panels) from the standard products illustrated with respect to the standard AERONET AOD measurements at 440 nm and the actual NO₂ observed by Pandora (color scale). The corresponding distributions of all variables are also included.

1110





Figure 7: Similar to Fig. 6, but for SKYNET retrievals over APL-SAP. Note that the spectral channels for the retrievals are different compared to AERONET, i.e. 400nm for AOD and 400-1020nm for AE. Also the axis scales are different.






Figure 8: Case study over APL-SAP for 25th June 2020 **for both AERONET (left panels) and SKYNET (right panels)**. Upper panels: Pandora total NO₂ column and its deviation from climatology. Middle panels: **AERONET** AOD (solid blue line), its improvement using Pandora NO₂ (dashed blue line) and the magnitude of improvement (light orange line and right y-axis). Lower panels: Similar to middle panels, but for AE retrievals. **Note that the spectral channels for the retrievals are different for the two networks.**

120





Figure 9: Similar to Fig. 8, but for SKYNET retrievals over APL-SAP location. Note that the spectral channels for the retrievals are different compared to AERONET.


125







1130

Figure 109: Inter-comparison of MODIS DB with standard (upper panels) and modified (lower panels) ground-based AOD at 470 nm for CNR- ISAC (left panels) and APL-SAP (middle and right panels) sites. In the left and middle panels, against AERONET AOD products, whereas in the right panels against SKYNET AOD. The y=x lines and MODIS DB EE envelopes $\pm (0.05 + 20\%)$ are plotted as dashed lines. The inter-comparison was performed considering a maximum distance between the center of the MODIS DB pixel and the site location of 5 km and Δt_{\max} (time between MODIS and AERONET/SKYNET observations) of ± 30 minutes.

1135



140 **Figure 4-10: Upper row:** Absolute correction as a function of the corresponding MODIS DB AOD data and PGN NO₂ data for CNR-
 145 **ISAC** (left panel) and **APL-SAP** (middle and right panels) sites. In the left and middle panels the inter-comparison was performed
 using **AERONET AOD** products, in the right panel **SKYNET AOD** was used. The color scale represents the **PGN NO₂** retrieved
 in correspondence of the **AERONET/SKYNET AOD** products. The analysis was performed considering a maximum distance
 between the center of the MODIS DB pixel and the site location of 5 km and Δt_{max} of ± 30 minutes. **Lower row:** As in the upper
 row, but here the color scale represents the absolute difference between **PGN and OMI climatological NO₂** data in correspondence
 with the **AERONET AOD** products.






Figure 1211: Comparisons of SSA at 440 nm obtained with GRASP following the standard AERONET procedure (X axis) and a similar approach but precisely accounting for NO₂ concentration (Y axis) from the co-located Pandora instruments in two different stations: APL-SAP (right upper panel) from March 2017 to November 2020, and CNR-ISAC (left upper panel) from April 2017 to September 2021. The data has been filtered to show retrievals corresponding to NO₂ concentration higher than 0.79 DU. The color of the circles is an indicator of the density of points, i.e., colors closer to red, indicate higher number of points close together. The absolute Mean Bias Error (MBE) (percent in parenthesis), the Root Mean Square Error (RMSE) and the correlation coefficient of the linear fit are also shown in the figures. The probability density functions of the difference between both methodologies (GRASP/Pandora NO₂ - GRASP/AERONET NO₂) can be found in the lower panels correspondingly for each station. The probability density functions for SSA values higher or lower than 0.9 are also included.