

Response to Referee #2's Comments: amt-2022-341

We appreciate the referee for his/her review and helpful comments. We have considered those comments very carefully and substantially revised our manuscript. A point-by-point response to the referee's comments is as follows.

General comments

The manuscript describes the impact of Aeolus observations on the forecast of Hurricane Ida (2021) and a mesoscale convective system that occurred during NASA's CPEX-AW field campaign. The authors claim that the novelty of this investigation is that it evaluates the potential impact of the "near-real-time" assimilation of this data set.

As mentioned in the paper, previous work has focused on observing system experiments (OSEs). As a matter of fact, OSEs mimic near-real-time impacts as they use observations that were available at the time of the experiment. Thus, I don't understand what the limitations of earlier investigations are. In fact, these earlier studies used all the observations that were available for assimilation – not just conventional data.

The previous observing system experiments did not examine how assimilating the Aeolus HLOS winds influences the dynamic and thermodynamic structures of the hurricane in detail, such as the vorticity and divergence in the inner core regions of the hurricane. Moreover, they did not examine the impacts on the mesoscale convective system. Our study attempted to fill these gaps.

In fact, the "conventional data" we referred to in the paper are all routinely assimilated data in the NCEP GDAS operational data assimilation system, including both conventional types of data and satellite data products. We have made clarifications in the text of the paper by mentioning them as "GDAS data".

The findings of this study are limited to one single hurricane and one single mesoscale system, which makes it impossible to draw any general conclusions. Therefore, the evaluation of the impact of Aeolus is over-emphasized.

Even if we have one main hurricane case in this paper, we did very comprehensive experiments. We performed the cycling data assimilations for hurricane Ida (2021), which can produce the forecasts initialized at different data assimilation cycles. We also carried out the sensitivity study of the initial time, horizontal, and vertical ensemble length scales for the experiments of hurricane Ida (2021). The results of horizontal and vertical were added to the revised version based on Referee #1's comments.

Meanwhile, we agree with you that more cases should be used in future studies. Therefore, we have mentioned these at the end of the revised paper to address your concern. Please see the sentences in the last paragraph of the revised manuscript.

In addition, some of the technical elements are not well described. An example are the observations errors. Those are critical in data assimilation as reported in previous Aeolus data impact studies.

We agree with you that the observation errors are critical for assimilating the Aeolus data. We considered the observational errors very carefully when performing our experiments. We did not use any default numbers for the observational errors. Instead, we used the error characteristics from the Aeolus data samples. We did assign different errors for Mie and Rayleigh winds (see updated Fig.2 and related text).

The paper doesn't show any statistics of background/analysis departures either that are critical to evaluate the behavior of the assimilation algorithms used here.

We added the statistics of background/analysis departures into the revised version of the paper (See Figure 3 and new Section 3.2).

More detailed comments

L74: What is the meaning of near-real-time HLOS here – and what is different from the methodology used in previous studies (OSEs)? What is new in this study since only focuses on one single hurricane?

"Near-real-time HLOS" refer to the type of Aeolus data we used. We have made the clarification in the text now.

As mentioned above, even if we have one main hurricane case in this paper, we did very comprehensive experiments. We performed the cycling data assimilations for hurricane Ida (2021), which can produce the forecasts initialized at different data assimilation cycles. We also carried out the sensitivity study of the initial time, horizontal, and vertical ensemble length scales for the experiments of hurricane Ida (2021).

L99: Is the assimilated done in the parent domain, the inner domain or both?

Data assimilation was performed in both domains. We clarified this in the revision.

Fig.2: Are the Aeolus values observations or background simulations? Also, is the y-axis the instrument error or the observation error? It is not clear to this reviewer how the observation errors are estimated.

The Aeolus values are observations. The y-axis is the instrument error provided by the Aeolus L2B product. We have revised this figure and the associated paragraph to explain how we estimated the observation errors.

In a real-time environment, all available observations would be assimilated – not just conventional data. Why is it not done here? Are the reconnaissance data been assimilated?

We agree. In fact, the "conventional data" we referred to in the paper include all conventional data available in the NCEP operational system, including both conventional types of data and satellite-derived data products. We have made clarifications in the text of the paper by mentioning them as "GDAS data". The reconnaissance data should be part of operational data; thus, they should be assimilated.

The impacts of assimilating Aeolus horizontal line-of-sight winds on numerical predictions of Hurricane Ida (2021) and a mesoscale convective system over the Atlantic Ocean

Chengfeng Feng¹, Zhaoxia Pu¹

5 ¹Department of Atmospheric Sciences, University of Utah, Salt Lake City, 84112, United States

Correspondence to: Zhaoxia Pu (Zhaoxia.Pu@utah.edu)

Abstract. On 22 August 2018, the European Space Agency (ESA) launched the first spaceborne wind lidar, the Aeolus satellite, measuring horizontal line-of-sight (HLOS) winds globally. The assimilation of Aeolus HLOS winds has been proven to improve numerical weather predictions (NWP). Still, its influences on forecasts of tropical cyclones (TCs) and tropical convective systems have yet to be examined in detail. This study investigates the impacts of assimilating Aeolus HLOS winds on the analysis and forecasts of Hurricane Ida (2021) and a mesoscale convective system (MCS) embedded in an African easterly wave (AEW) over the Atlantic Ocean (AO) with the mesoscale community Weather Research and Forecasting (WRF) model and the NCEP-GSI based three-dimensional ensemble-variational (3DnVAR) hybrid data assimilation (DA) system. Mie-cloudy and Rayleigh-clear winds are assimilated. The results for Ida (2021) show that assimilating Aeolus HLOS winds leads to better track predictions. The intensity forecasts are improved in some cases, even with limited coverage of Aeolus HLOS winds within the inner core region of Ida (2021). In addition, the structure of heavy precipitation associated with Ida (2021) is refined after assimilation of Aeolus HLOS winds. Further diagnosis demonstrates that the improved intensity and precipitation forecasts result from enhanced divergence in the upper level of the troposphere after assimilation of Aeolus HLOS winds. Additional results from the MCS associated with an AEW indicate that assimilating Aeolus HLOS winds enhances forecasts of its precipitation structure and the associated low-level divergence. In short, this study demonstrates the potential of assimilation of Aeolus HLOS winds to improve forecasts for TCs and tropical convective systems.

Deleted:

1 Introduction

Measuring three-dimensional wind profiles in the Global Observing System (GOS), especially over the oceans and remote land areas, is essential for exploring atmospheric dynamics, understanding critical issues of climate change, improving the estimation of air pollutant dispersion, and creating better initial conditions (ICs) for numerical weather predictions (NWPs; WMO, 2017; Baker et al., 2014; Pu et al., 2017; Zhang and Pu, 2010; Pu et al. 2010; Rennie et al., 2021b). Large areas of the tropical atmosphere lack measurement of wind profiles, which suggests the potential for significantly improving forecasting skills for various tropical convective systems with additional wind observations (Baker et al., 2014). To provide high spatial

30 and temporal near-vertical wind profiles globally, the Aeolus satellite, the first spaceborne wind lidar, was launched by the European Space Agency (ESA) on 22 August 2018 (Reitebuch et al., 2020; ESA, 2022).

After successfully launching into a sun-synchronous orbit at about 320 km altitude with a weekly cycle, the Aeolus satellite now passes the equator at 18:00 (6:00) local time (LT) during ascending (descending) orbits (Andersson et al., 2008; Krisch et al., 2022; ESA, 2022). The azimuth angle of the Aeolus satellite closely approaches 270° (90°) for ascending (descending) orbits near the equator, and the viewing angle toward the atmosphere is 35° off-nadir. The horizontal line-of-sight (HLOS) wind component, derived from the measured wind along the laser beam line-of-sight (LOS), is approximately east-west oriented for most of the orbits (Krisch et al., 2022).

40 The Atmospheric Laser Doppler INstrument (ALADIN), on board the Aeolus mission, measures Doppler-shifted backscattered light scattering by atmospheric molecules and particulates with two separate interferometers: the Fizeau (Mie channel) for large particles, cloud droplets, ice crystals, dust, and aerosols, and the Fabry-Perot (Rayleigh channel) for moving molecules, including oxygen and nitrogen (Andersson et al., 2008; Reitebuch et al., 2009; Ingmann and Straume, 2016). Based on the signal-to-noise ratio, four types of HLOS winds are available, including Mie-clear, Mie-cloudy, Rayleigh-clear, and Rayleigh-
45 cloudy (Jos de Kloe et al., 2022). Rayleigh-clear winds perform better than Rayleigh-cloudy winds due to little or no contamination from Mie scattering. Mie-cloudy winds are better than Mie-clear winds because measurements from the Mie channel require strong backscattering from aerosols, water droplets, or ice crystals (Rennie et al., 2021a). The horizontal resolution of the Mie channel is typically about 10-15 km along the ground track and about 90 km for the Rayleigh channel. Vertically, the sizes of 24 range bins vary from 250 m to 2 km (Krisch et al., 2022).

50 Aeolus HLOS winds have been validated with different reference observations over many regions since launching, such as ground-based radar measurements (Zuo et al., 2022), ground-based coherent Doppler wind lidars (Wu et al., 2022), airborne Doppler wind lidars (Witschas et al., 2020; Witschas et al., 2022), in situ Atmospheric Motion Vectors (AMVs; Rani et al., 2022; Lukens et al., 2022), NWP equivalents (Rani et al., 2022; Martin et al., 2021), and radiosonde observations (Martin et al., 2021; Baars et al., 2020). Validation shows that the quality of Mie-cloudy winds is better than that of Rayleigh-clear winds (Zuo et al., 2022; Rani et al., 2022). Furthermore, Aeolus HLOS winds can capture atmospheric dynamic structures well, such as the Low-Level Jet (LLJ), Tropical Easterly Jet (TEJ; Rani et al., 2022), atmospheric gravity waves (GWs; Banyard et al., 2021). As the Aeolus products are continuously calibrated and validated, the product processor is updated and the performance of the Aeolus Level-2B (L2B) wind product improves (Wu et al., 2022). Thus, the current Aeolus products are suitable for
60 data assimilation (DA) in the global forecast system (GFS; Pourret et al., 2022; Guo et al., 2021).

Assimilation of Aeolus HLOS winds has already been shown to improve analyses and forecasts in many global NWP models, including the Météo-France global NWP model ARPEGE (Pourret et al., 2022), the Canadian Global Deterministic Prediction

System (GDPS; Laroche and St-James, 2022), and ECMWF's Integrated Forecasting System (IFS; Rennie et al., 2021a).
65 Several other studies have assessed the impacts of assimilating Aeolus HLOS winds on tropical cyclones (TCs) by Observing
System Experiments (OSEs). Rani et al. (2022) conducted OSEs to examine the impacts of assimilation of Aeolus HLOS
winds on simulations of the location, intensity, and vertical extent of North Indian Ocean (NIO) cyclones, and the
improvements due to Aeolus winds were marginal. Marinescu et al. (2022) carried out OSEs with the operational Hurricane
Weather and Research Forecasting (HWRF) model to assess the impacts of assimilating Aeolus HLOS winds on forecasting
70 TCs, and their results suggest that the most significant potential impacts from assimilation of Aeolus observations are likely
to occur in the upper troposphere and lower stratosphere and within about 500 km of the TC center. The OSEs performed with
the NOAA Finite-Volume Cubed-Sphere Global Forecast System (FV3GFS) suggested that assimilation of Aeolus HLOS
winds can reduce track forecast error in the Eastern Pacific basin and Atlantic basin (Garrett et al., 2022). However, the
potential impacts of assimilating [these so-called "Aeolus near-real-time \(NRT\)"](#) HLOS winds on simulations of TCs and
75 tropical convective systems have not been investigated. From August to September 2021, the NASA Convective Processes
Experiment - Aerosols & Winds (CPEX-AW) field campaign, in collaboration with ESA, conducted post-launch calibration
and validation activities for the Aeolus satellite near St. Croix over the Atlantic Ocean (AO), marking notable TCs and other
tropical convective systems observed by the Aeolus satellite. In this study, we use Hurricane Ida (2021), a major hurricane,
and a mesoscale convective system (MCS) embedded in an African easterly wave (AEW) during NASA's CPEX-AW
80 campaign to assess the impacts of assimilating Aeolus HLOS winds.

The paper is organized as follows. Section 2 introduces the mesoscale community Weather Research and Forecasting (WRF)
model, the NCEP Gridpoint Statistical Interpolation (GSI)-based three-dimensional ensemble-variational (3DnVAR) hybrid
DA system, the Integrated Multi-satellitE Retrievals for GPM (IMERG) precipitation dataset, and the statistical evaluation
85 metrics. Section 3 discusses the impacts of assimilating Aeolus HLOS winds on forecasts of Hurricane Ida (2021). Section 4
diagnoses the influence of [the](#) assimilation of Aeolus HLOS winds on the analysis of atmospheric conditions associated with
Hurricane Ida (2021). Section 5 evaluates the influence of [the](#) assimilation of Aeolus HLOS winds on an MCS embedded in
an AEW. Finally, Sect. 6 summarizes the results and assesses the need for future work.

90

Deleted: ¶

Deleted: ¶

2 WRF Model, DA system, IMERG, and statistical evaluation metrics

2.1 WRF Model

100 An advanced research version of WRF model (WRF-ARW) V4.4 (Skamarock et al., 2019) is applied in this study, with 61 terrain-following levels and the model top set at 10 hPa, as indicated by Table 1. Two domains are employed in all experiments. Their sizes differ in the experiments of Hurricane Ida (2021) and the MCS within an AEW, as illustrated in Fig. 1. The horizontal resolutions of the parent and inner domains are 12 and 4 km, and the time steps of the parent and inner domains are 30 and 10 s, respectively. The boundary conditions of the parent domain are obtained from the $0.25^\circ \times 0.25^\circ$ horizontal resolution NCEP GFS final analyses (FNL). Details of the parameterization scheme options are listed in Table 1.

105

Table 1: Configuration of the WRF-GSI cycling DA [system](#).

Model	Resolution	12 km (domain 1, d01, or parent domain), and 4 km (domain 2, d02, or inner domain) 61 vertical levels with model top at 10 hPa
	Time steps	30 s (d01) and 10 s (d02)
	Physics	Longwave radiation: RRTM (Mlawer et al., 1997) Shortwave radiation: Dudhia (Dudhia, 1989) Microphysics: WSM6 (Hong and Lim, 2006) Cumulus: Kain-Fritsch (Activated only in the parent domain; Kain, 2004) PBL: YSU (Hong et al., 2006) Surface layer: Monin-Obukhov Similarity (Jiménez et al., 2012) Land surface: Unified Noah LSM (Tewari et al., 2004)
	Boundary condition	Spin up and cycling DA period: NCEP GFS FNL Forecast period: NCEP GFS Forecasts
Analysis	DA system	NCEP GSI-based 3DnVAR hybrid DA system V3.7
	State vector	\mathbf{u} (Zonal wind), \mathbf{v} (Meridional wind), \mathbf{t} (Virtual temperature), \mathbf{q} (Specific humidity), \mathbf{prse} (Pressure), \mathbf{ps} (Surface pressure), \mathbf{sst} (Sea surface temperature)
	Control vector	\mathbf{sf} (Stream function; 0.60), \mathbf{vp}_{ub} (Unbalanced velocity potential; 0.60), \mathbf{ps}_{ub} (Unbalanced surface pressure; 0.75), \mathbf{t}_{ub} (Unbalanced temperature; 0.75), \mathbf{rh} (Normalized RH; 0.75), \mathbf{sst} (Sea surface temperature; 1.00) The numbers indicate normalized scale factors for their background error variances.
	Assimilation window	± 3 h
	Background error covariance	Static part (0.2), and flow-dependent part (0.8, estimated from 80-member 6-h ensemble forecasts from the NCEP GSI 4DnVAR system)
	Ensemble localization scale	Horizontal: 110 km Vertical: 3 grid units

Deleted: system





Figure 1: Two nested domains (d01: 12 km, and d02: 4 km) in experiments 2418_L2B (a) and 1918_L2B (b). The Aeolus measurement swaths (colored line) are from 24 August 2021, 12:00 UTC to 27 August 2021, 00:00 UTC (a) and from 20 August 2021, 00:00 UTC to 21 August 2021, 00:00 UTC (b). The track and maximum surface wind speeds (MWSS, colored dot) of Hurricane Ida (2021) in (a) are adopted from the NHC best-track data, and the days are illustrated below the open markers indicating 00:00 UTC. Two classifications (TD: tropical depression; TS: tropical storm) and different categories (Cat1: Category 1; Cat2: Category 2; Cat3: Category 3; Cat4: Category 4) based on the Saffir-Simpson Hurricane wind scale are given below the color bar of MWS. The AEW in (b) is tracked manually using the GFS-analyzed relative vorticity maxima at 700 hPa (purple line with black edges). The white dots with the numbers below along the AEW track indicate 00:00 UTC of a day. The geopotential height at 700 hPa (white line) and relative humidity at 850 hPa (RH; yellow line), obtained from GFS analysis, are overlaid atop GOES-16 channel 8 brightness temperatures (BTs; shaded) on 25 August 2021, 12:00 UTC (a), and 21 August 2021, 00:00 UTC (b). Only RH less than 35% is shown in (a) and (b).



Deleted:

Deleted: 12 UTC 24 Aug to 00 UTC 27 Aug 2021...(a) and from 20 August 2021, 00:00 UTC to 21 August 2021, 00:00 UTC 00 UTC 20 Aug to 00 UTC 21 Aug 2021...(b). The track maximum surface wind speeds (MWSS, colored dot) of Hurricane Ida (2021) in (a) are adopted from the NHC best-track data, and the days are illustrated below the open markers indicating 00:00 UTC. Two classifications (TD: tropical depression; TS: tropical storm) and different categories (Cat1: Category 1; Cat2: Category 2; Cat3: Category 3; Cat4: Category 4) based on the Saffir-Simpson Hurricane wind scale are given below the color bar of MWS. The AEW in (b) is tracked manually using the GFS-analyzed 700-hPa...relative vorticity maxima at 700 hPa (purple line with black edges). The white dots with the numbers below along the AEW track indicate 00:00 UTC of a day. The 700-hPa ...geopotential height at 700 hPa (white line) and 850-hPa ...relative humidity at 850 hPa (RH; yellow line), obtained from GFS analysis, are overlaid atop GOES-16 channel 8 brightness temperatures (BTs; shaded) on 25 August 2021, 12:00 UTC 25 Aug...(a), and 21 August 2021, 00:00 UTC 00 UTC 21 Aug 2021... (1)

2.2 NCEP GSI-based 3DnVAR hybrid DA system

The DA system used in this study is the NCEP GSI-based 3DnVAR hybrid DA system V3.7, the details of which are listed in Table 1. The cost function of the GSI system has two terms: the background and observational error terms.

$$J(\mathbf{x}, \boldsymbol{\beta}) = \frac{1}{2} (\mathbf{x}_b - \mathbf{x})^T (\alpha_1 \mathbf{B}_1 + \alpha_2 \mathbf{B}_2)^{-1} (\mathbf{x}_b - \mathbf{x}) + \frac{1}{2} [\mathbf{y}_0 - H(\mathbf{x})]^T \mathbf{R}^{-1} [\mathbf{y}_0 - H(\mathbf{x})] \quad (1)$$

In the background error term (the first term in Eq. (1)), \mathbf{x} is the analysis, and \mathbf{x}_b is the first guess, namely a 6-h WRF model simulation. The hybrid background error covariance matrix consists of a prescribed static part \mathbf{B}_1 , and a flow-dependent part \mathbf{B}_2 . The weighting factors of these two portions are 0.2 (α_1) and 0.8 (α_2), respectively. The flow-dependent part is estimated by 80-member 6-h ensemble forecasts from the NCEP GSI 4DnVAR system. The default homogeneous isotropic horizontal ensemble localization scale is 110 km, and the default vertical localization scale is 3 grid units (see Table 1). In the observational error term (the second term in Eq. (1)), \mathbf{y}_0 is the observation, H is the forward model, and \mathbf{R} is the observation error covariance. Thus, the innovation Observation-Minus-Background (OmB) is defined as $\mathbf{y}_0 - H(\mathbf{x})$.

The observations \mathbf{y}_0 in this study are the Aeolus L2B baseline 12-NRT HLOS winds (ESA, 2021). The quality control (QC) steps used in this study are the following:

1. Mie-clear and Rayleigh-cloudy products are not used in this study due to their poor quality (Rennie et al., 2021a; Pourret et al., 2022; Laroche and St-James, 2022).
2. Aeolus HLOS winds marked as invalid retrievals by the L2B processor are eliminated (Rennie et al., 2021a).
3. The GSI system rejects Mie-cloudy and Rayleigh-clear HLOS winds when their L2B estimated instrumental errors (before scaling) are over 12 and 10 m s⁻¹, respectively (Rennie et al., 2021a).
4. Aeolus HLOS winds are discarded when their absolute value is larger than 1000 m s⁻¹.
5. A gross check is adopted to remove observations with normalized OmBs (OmB normalized by σ) larger than 4 σ , while σ is the standard deviation of the observation errors (similar to QC Step 6 in Pourret et al. (2022)).

The forward model $H(\mathbf{x})$ of the Mie-cloudy and Rayleigh-clear HLOS winds (Pourret et al., 2022; Laroche and St-James, 2022; Rennie et al., 2021a) is defined as:

$$H(\mathbf{x}) = -\mathbf{u} \sin \theta - \mathbf{v} \cos \theta \quad (2)$$

The forward model $H(\mathbf{x})$ interpolates the WRF model winds (zonal wind component \mathbf{u} and meridional wind component \mathbf{v}) according to the geolocation of the Aeolus observations and projects the interpolated model winds to the HLOS winds with

Deleted: -

Deleted: 6-h

Deleted: near-real-time

the horizontal azimuth angle θ . The vertical velocity is assumed to be zero in the forward model because it seems to be a second-order problem even for tropical convective systems (Rennie and Isaksen, 2020).

The observation error covariance matrix \mathbf{R} can be divided into $\mathbf{R} = \Sigma^{\frac{1}{2}} \mathbf{C} \Sigma^{\frac{1}{2}}$. \mathbf{C} is the correlation matrix. Since we assume that the observations are uncorrelated in this study, \mathbf{C} is an identity matrix. $\Sigma^{\frac{1}{2}}$ is a diagonal matrix of the standard deviation of the observation error σ . The observation error consists of the instrument and representation error, while the representation error includes the observation-operator error and the error due to unresolved scales and processes (Janjić et al., 2018). Since the representation error is unknown and the primary goal of this study is to investigate the impacts of assimilation of Aeolus HLOS winds on Hurricane Ida (2021) and an MCS embedded in an AEW, we simply use the L2B dynamic estimated instrumental errors to estimate the standard deviation of the observation error σ . Figure 2 shows scatter plots of the valid Mie-cloudy (a) and Rayleigh-clear (b) HLOS winds against their estimated instrumental errors from 25 August 2021, 00:00 UTC to 26 August 2021, 18:00 UTC (cycling DA period of experiment 2418 L2B). The Mie-cloudy HLOS winds are clustered primarily between -20 and 20 m s^{-1} , and their estimated instrumental errors lie mostly between 1 and 4 m s^{-1} . Based on the statistics of the data, the standard deviation of the observation errors for the Mie-cloudy winds is assigned to 2.5 m s^{-1} in this study, as indicated by the black line in Fig. 2 (a). The Rayleigh-clear HLOS winds are mainly between -30 and 30 m s^{-1} , and their estimated instrumental errors vary primarily from 3 to 6 m s^{-1} . The standard deviation of the observation errors for the Rayleigh-clear winds is set at 4.5 m s^{-1} based on the statistics of data samples. The Rayleigh-clear winds have a higher standard deviation of observation errors because the quality of the Mie-cloudy winds is better than that of the Rayleigh-clear winds (Zuo et al., 2022; Rani et al., 2022).





Figure 2: Scatter plots (black circle) of valid observed Aeolus HLOS winds against their estimated instrumental errors from 25 August 2021, 06:00 UTC to 27 August 2021, 00:00 UTC (cycling DA period of experiment 2500 L2B) for Mie-cloudy winds (a) and Rayleigh-clear winds (b). The black lines in (a) and (b) indicate the estimated instrumental errors for the greatest population of valid


Deleted: :00...UTC to 26 August 2021, 18:00:00 ... [2]

Deleted: , so...the standard deviation of the observation errors for the Mie-cloudy winds is simply ... [3]

Deleted: , so...the standard deviation of the observation errors for the Rayleigh-clear winds is simply ... [4]



Deleted:



Deleted:

Deleted: Figure 2: Scatter plots (black circle) of valid Aeolus HLOS winds against their estimated instrumental errors from 25 August 2021, 00:00 UTC to 26 August 2021, 18:00 UTC (cycling DA period of experiment 2418 L2B) for Mie-cloudy winds (a) and Rayleigh-clear winds (b). The black lines in (a) and (b) indicate the estimated observation errors for the greatest population of valid Aeolus HLOS winds, which are 2.5 and 4.5 m s^{-1} , respectively. The results of ... [5]

observed Aeolus HLOS winds, which are 2.5 and 4.5 m s⁻¹, respectively. The results of experiments 2406 L2B, 2412 L2B, and 2418 L2B are similar to those of experiment 2500 L2B.

255

2.3 IMERG Data

The precipitation dataset used for verifying rainfall forecasts in this study is IMERG Version 6B-Final. IMERG precipitation estimates combine various satellite passive microwave sensors related to precipitation, including the GPM constellation and microwave precipitation-calibrated geo-IR fields. IMERG precipitation estimates are half-hourly and on a 0.1° × 0.1° grid over the globe. This dataset is available from June 2000 to the present (Huffman et al., 2019).

260

2.4 Statistical evaluation metrics

2.4.1 RMSE

Root-mean-square error (RMSE) can measure errors of the track, MWS, and minimum sea-level pressure (MSLP) between the forecasts and NHC best-track data. RMSE is defined as follows:

265

$$RMSE = \sqrt{(x - x_{ref})^2} \quad (3)$$

where x is the location, MWS, or MSLP forecast of a TC, while x_{ref} is adopted from the NHC best-track data.

2.4.2 ETS

An Equitable Threat Score (ETS) is a corrected ratio of the number of correctly predicted events to the total number of predicted or observed events. The definition of ETS is:

270

$$ETS = \frac{N_H - ref}{N_H + N_{FA} + N_M - ref} \quad (4)$$

where ref is the chance forecast:

275

$$ref = \frac{(N_H + N_{FA})(N_H + N_M)}{N_H + N_{FA} + N_H + N_M} \quad (5)$$

Other variables are computed by a contingency table (Table 2), which defines Hit (N_H), False Alarm (N_{FA}), Miss (N_M), and Correct Negative (N_{CN}).

280

Deleted: ¶

Table 2: Contingency table

		Observation	
		Yes	No
Forecast	Yes	Hit (N_H)	False Alarm (N_{FA})
	No	Miss (N_M)	Correct Negative (N_{CN})

3 Impacts of Aeolus data on numerical simulations of Hurricane Ida (2021)

3.1 Case description and experiment design

285 Ida (2021) originated from an AEW on 14 August 2021. On 24 August 2021, the AEW moved into the Caribbean Sea, reached
the area near Aruba, Bonaire, and Curaçao, and interacted with MCSs along the northern coast of South America. On 25
290 August 2021, 12:00 UTC, as shown by Fig. 1 (a), the convection, indicated by the GOES-R channel 8 BTs, was concentrated
in the eastern area of a broad low-pressure system south of the subtropical ridge over the southwestern Caribbean Sea, which
is shown by geopotential height at 700 hPa and RH at 850 hPa. Steered by the subtropical ridge, the convective disturbance
295 began to turn north to northwestward on 26 August 2021, as demonstrated by the track of Ida (2021) in Fig. 1 (a). A TD formed
on 26 August 2021, 12:00 UTC, and then intensified into a TS 6 hours after cyclogenesis. Early on 27 August 2021, the first
rapid intensification (RI) period occurred, and Ida (2021) strengthened into a hurricane on 27 August 2021, 18:00 UTC. After
moving across the Isle of Youth, Ida (2021) made landfall in Cuba around 28 August 2021, 00:00 UTC. After passing Cuba,
Ida (2021) experienced the second RI period from 28 August 2021, 12:00 UTC to 29 August 2021, 12:00 UTC (Beven II et
al., 2022).

As shown by Table 3, 8 experiments, including experiments 2406, 2406_L2B, 2412, 2412_L2B, 2418, 2418_L2B, 2500, and
2500_L2B, are performed to investigate the impacts of assimilating Aeolus Mie-cloudy and Rayleigh-clear HLOS winds on
the analysis and subsequent forecasts of Hurricane Ida (2021). [data assimilation was performed for both model domains.](#)

300 Experiment 2406 is initialized by the NCEP GFS FNL on 24 August 2021, 06:00 UTC. [The first two digits of the experiment
name indicate the day of the initial time, and the last two digits represent the hour of the initial time.](#) After a 6 h spin-up, a
cycling DA period from 24 August 2021, 12:00 UTC to 26 August 2021, 06:00 UTC (8 DA cycles in total) is performed in
experiment 2406. [The data that were assimilated routinely into the NCEP Global Data Assimilation System \(GDAS\), including
global upper air and surface weather observations, as well as satellite data,](#) are assimilated. The subsequent 48 h forecasts are
305 initialized from the analysis of DA cycles 5, 6, 7, and 8. Experiment 2406_L2B is similar to experiment 2406, but it also
assimilates Aeolus HLOS winds during cycling DA. The forecasts are not initialized from the analysis of DA cycles 1, 2, 3,
and 4 in this study because the impacts of assimilating Aeolus HLOS winds are tiny if the number of DA cycles is not large
enough.

Deleted: :00

Deleted: brightness temperatures (BTs)

Deleted: 700-hPa

Deleted: 850-hPa

Deleted: 0:00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: , so

Deleted: t

Deleted: 6-h

Deleted: :00

Deleted: :00

Deleted: ,

Deleted: and. Only NCEP

Deleted: ADP

Deleted:

Deleted: 48-hour

330 Since Ida (2021) was a broad low-pressure system interacting with MCSs along the northern coast of South America on 24 August 2021, the forecasts of Ida (2021) are sensitive to its ICs. Consequently, except for experiments 2406 and 2406_L2B, we carry out three more sets of experiments with different initial times: 2412 and 2412_L2B (initialized on 24 August 2021, 12:00 UTC), 2418 and 2418_L2B (initialized on 24 August 2021, 18:00 UTC), and 2500 and 2500_L2B (initialized on 25 August 2021, 00:00 UTC). The cycling DA periods of all experiments are before the first RI of Ida (2021), enabling us to

335 investigate the impacts of assimilating Aeolus HLOS winds on the intensity forecasts of the first and second rapid RI periods of Ida (2021).

Since data assimilation results are sensitive to the homogeneous isotropic horizontal and vertical ensemble localization scales, we performed additional sensitivity experiments. For instance, corresponding to experiment 2500_L2B, the horizontal ensemble localization scale of experiment 2500_L2B_H1 reduces to 55 km, half of the default value for experiment 2500_L2B. The horizontal localization scale of experiments 2500_L2B_H2 increases to 220 km, double the default value, when the vertical localization scale is fixed at 3 grid units. Experiments 2500_L2B_V1 and 2500_L2B_V2 are similar to experiments 2500_L2B_H1 and 2500_L2B_H2, but for testing the sensitivity to vertical localization scale (see details in Table 3).

Table 3: List of experiment configurations

Experiment	Case	Initial Time	Cycling DA Period	Horizontal Scale (km)	Vertical Scale (grid units)	DA Observation	Forecast
2406	Ida	24 August 2021, 06:00 UTC	From 24 August 2021, 12:00 UTC to 26 August 2021, 06:00 UTC (3DEnVAR)	110	3	NCEP GDAS data	48 h forecasts initialized from DA Cycle 5, 6, 7, and 8
2406_L2B	Ida	24 August 2021, 06:00 UTC	From 24 August 2021, 12:00 UTC to 26 August 2021, 06:00 UTC (3DEnVAR)	110	3	NCEP GDAS data Aeolus L2B HLOS winds	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2412	Ida	24 August 2021, 12:00 UTC	From 24 August 2021, 18:00 UTC to 26 August 2021, 12:00 UTC (3DEnVAR)	110	3	NCEP GDAS data	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2412_L2B	Ida	24 August 2021, 12:00 UTC	From 24 August 2021, 18:00 UTC to 26 August 2021, 12:00 UTC (3DEnVAR)	110	3	NCEP GDAS data Aeolus L2B HLOS winds	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2418	Ida	24 August 2021, 18:00 UTC	From 25 August 2021, 00:00 UTC to 26 August 2021, 18:00 UTC (3DEnVAR)	110	3	NCEP GDAS data	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2418_L2B	Ida	24 August 2021, 18:00 UTC	From 25 August 2021, 00:00 UTC to 26 August 2021, 18:00 UTC (3DEnVAR)	110	3	NCEP observations Aeolus L2B HLOS winds	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2500	Ida	25 August 2021, 00:00 UTC	From 25 August 2021, 06:00 UTC to 27 August 2021, 00:00 UTC (3DEnVAR)	110	3	NCEP GDAS data	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2500_L2B	Ida	25 August 2021, 00:00 UTC	From 25 August 2021, 06:00 UTC to 27 August 2021, 00:00 UTC (3DEnVAR)	110	3	NCEP GDAS data Aeolus L2B HLOS winds	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2500_L2B_H1	Ida	25 August 2021, 00:00 UTC	From 25 August 2021, 06:00 UTC to 27 August 2021, 00:00 UTC (3DEnVAR)	55	3	NCEP GDAS data Aeolus L2B HLOS winds	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2500_L2B_H2	Ida	25 August 2021, 00:00 UTC	From 25 August 2021, 06:00 UTC to 27 August 2021, 00:00 UTC (3DEnVAR)	220	3	NCEP GDAS data Aeolus L2B HLOS winds	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2500_L2B_V1	Ida	25 August 2021, 00:00 UTC	From 25 August 2021, 06:00 UTC to 27 August 2021, 00:00 UTC (3DEnVAR)	110	1.5	NCEP GDAS data Aeolus L2B HLOS winds	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
2500_L2B_V2	Ida	25 August 2021, 00:00 UTC	From 25 August 2021, 06:00 UTC to 27 August 2021, 00:00 UTC (3DEnVAR)	110	6	NCEP GDAS data Aeolus L2B HLOS winds	48 h forecasts initialized from DA cycles 5, 6, 7, and 8
1918	MCS	19 August 2021, 18:00 UTC	From 20 August 2021, 00:00 UTC to 21 August 2021, 00:00 UTC (3DVAR)	1	1	NCEP GDAS data	30 h forecasts initialized from DA cycles 5

Deleted: ...ince Ida (2021) was just ... broad low-pressure... [6]

Deleted: In addition to ICs, the forecasts of Ida (2021) may... [7]

Formatted ... [8]

Formatted ... [10]

Formatted Table ... [9]

Deleted: (2021) [1]

Deleted: :00 ... [44]

Deleted: - ... [11]

Formatted ... [43]

Deleted: :00...UTC [20]

Deleted: :00...UTC (3DEnVAR8 cycles) ... [34]

Deleted: (2021) [1]

Deleted: :00 ... [45]

Deleted: and ... [21]

Deleted: :00...UTC [35]

Deleted: C ... [22]

Deleted: :00...UTC (3DEnVAR8 cycles) ... [36]

Deleted: C ... [23]

Deleted: :00...UTC (3DEnVAR8 cycles) ... [37]

Deleted: C ... [12]

Deleted: :00 ... [13]

Deleted: :00...UTC [24]

Deleted: :00...UTC (3DEnVAR8 cycles) ... [38]

Deleted: C ... [25]

Formatted ... [14]

Deleted: :00...UTC (3DEnVAR8 cycles) ... [39]

Deleted: C ... [26]

Deleted: :00 ... [27]

Deleted: conventional observations ... [28]

Deleted: and ... [29]

Deleted: - ... [30]

Deleted: :00...UTC [31]

Formatted ... [32]

Deleted: :00...UTC (3DEnVAR8 cycles) ... [33]

Deleted: C ... [34]

Deleted: :00 ... [35]

Deleted: conventional observations ... [36]

Deleted: - ... [37]

Deleted: [38]

3.2. Distribution of O-B and O-A

To reveal the influence of data assimilation on initial analysis, Figure 3 shows the distribution of O-B (observation minus background) and O-A (observation minus analysis) for data assimilation experiments in both domains. Compared to observations, the analyses with data assimilation enhanced the model fields as O-A are smaller than the O-B, and the O-A is closer (relative to O-B) to the zero line. The decreased O-A relative to O-B is associated with better hurricane forecasts in these experiments (see next sections).




Figure 3: Probability density functions (PDFs) of normalized Observation-Minus-Background (OmB, pink) and normalized Observation-Minus-Analysis (OmA, green) in domain 1 (d01, solid line), and domain 2 (d02, dashed line) from 25 August 2021, 06:00 UTC to 27 August 2021, 00:00 UTC (cycling DA period of experiment 2500_L2B) for Mie-cloudy winds (a) and Rayleigh-clear winds (b). The PDFs are computed in 0.2 bins of normalized OmB or OmA.


3.3. Track forecasts

Figure 4(a) compares the 48 h track forecasts between experiments 2406 (without assimilation of Aeolus HLOS winds) and 2406_L2B (with assimilation of Aeolus HLOS winds) during the last four cycles of the cycling DA period (from 25 August 2021, 12:00 UTC to 26 August 2021, 06:00 UTC). The 48 h track forecasts of experiment 2406 initialized from the last four DA cycles show that all track forecasts, where Ida generally moves northwestward, are southwest of the NHC best track. As shown by Fig. 4(b), (c), and (d), the 48 h track forecasts of experiments 2412, 2418, and 2500 (as in experiment 2406) also have systematic biases toward the southwest of the NHC best track of Ida (2021). In addition, the track forecast errors are reduced in experiments without assimilation of Aeolus HLOS winds (see the numbers in Fig. 4(e)-(h)) when more NCEP GDAS data are assimilated, in addition to the forecasts initialized from DA cycles 5 and 6 of experiment 2406. Compared to experiments without assimilation of Aeolus HLOS winds, the 48 h track forecasts become closer to the NHC best track of Ida (see Fig. 4(a)-(d)), and the averaged track forecast errors are reduced consistently in the experiments with assimilation of

- Deleted: :00
- Deleted:
- Deleted: conventional observations
- Deleted: s and
- Deleted: -
- Formatted: Centered
- Deleted: :00
- Deleted: ¶
- Deleted: :00
- Deleted: 5 cycles
- Deleted: C
- Formatted: Font: (Default) +Body (Times New Roman), Bold, Not Italic
- Formatted: Font: (Default) +Body (Times New Roman)
- Formatted: Font: (Default) +Body (Times New Roman)
- Formatted: Normal, Indent: First line: 0.33"
- Formatted: Font: (Default) +Body (Times New Roman)
- Formatted: Font: (Default) +Body (Times New Roman), Font color: Custom Color(RGB(34,34,34))
- Formatted: Centered
- Formatted: Normal
- Formatted: Font: 9 pt, Bold, Font color: Black
- Deleted: 2
- Deleted: 3
- Deleted: 48-hour
- Deleted: :00
- Deleted: :00
- Deleted: 48-hour
- Deleted: 3
- Deleted: 48-hour
- Deleted: 3
- Deleted: conventional
- Deleted: 48-hour
- Deleted: 3

Aeolus HLOS winds, as shown by Fig. 4 (e)-(h). The reductions of the averaged track forecast errors range from 10 to around 60 km after assimilation of Aeolus HLOS winds.

Deleted: 3



620 | **Figure 4:** Comparison of 48 h track forecasts of Hurricane Ida (2021) initialized from the analysis of DA cycles 5 (blue), 6 (green), 7
 | (brown), and 8 (pink) between experiments 2406 (dashed lines) and 2406_L2B (solid lines) (a), 2412 and 2412_L2B (b), 2418 and
 | 2418_L2B (c), and 2500 and 2500_L2B (d). Tracks of Ida (2021) adopted from the NHC best-track data are shown by solid black
 | lines in (a)-(d), and the days are illustrated above the open markers indicating 00:00 UTC. The forecasts of Ida (2021) are tracked
 | by the GFDL vortex tracker. Compared with the experiments without assimilation of Aeolus HLOS winds, the averaged RMSE
 | reductions of the 48 h track forecasts initialized from the analysis of DA cycles 5 (blue), 6 (green), 7 (brown), and 8 (pink) for
 | experiments 2406_L2B, 2412_L2B, 2418_L2B, and 2500_L2B are shown in (e)-(h), respectively. The numbers indicate the averaged
 | RMSEs of track forecasts of the reference experiments. Positive values of RMSE reductions mean improvement, while negative
 | values indicate degradation.

Deleted: 3
 Deleted: -h

Deleted: -

3.4 Intensity forecasts

630 | As introduced in Sect. 3a, Ida (2021) made cyclogenesis on 26 August 2021, 12:00 UTC and strengthened into a TS 6 hours
 | later. Ida (2021) experienced its first RI period from 00:00 UTC to 18:00 UTC on 27 August 2021. After making landfall and
 | passing Cuba, Ida (2021) experienced the second RI period from 28 August 2021, 12:00 UTC to 29 August 2021, 12:00 UTC.
 | Figure 5(a) compares the 48 h MSLP forecasts initialized from the last four cycles (from 25 August 2021, 12:00 UTC to 26
 | August 2021, 06:00 UTC) between experiments 2406 (without assimilation of Aeolus HLOS winds) and 2406_L2B (with
 | assimilation of Aeolus HLOS winds). The 48 h MSLP forecasts of experiments 2406 and 2406_L2B capture the intensification
 | processes well, although they cannot capture the platform between the two RI periods (from 27 August 2021, 18:00 UTC to
 | 28 August 2021, 12:00 UTC) because their forecasts fail to predict the landfall of Ida (2021) in Cuba (see Fig. 4(a)). Figure 5
 | (e) indicates a neutral impact of assimilation of Aeolus HLOS winds on the MSLP forecasts for Ida (2021), compared to
 | experiment 2406. Figure 6(a) and (e) compare the 48 h MWS forecasts initialized from the last four cycles between
 | experiments 2406 and 2406_L2B, and the conclusions are consistent with the MSLP forecasts of experiments 2406 and
 | 2406_L2B. Similarly, the 48 h forecasts of experiments 2412 and 2412_L2B also predict the MSLP (see Fig. 5(b)) and the
 | MWS (see Fig. 6(b)) well, expect for the forecasts initialized from DA cycle 5, and the impacts of assimilating Aeolus HLOS
 | winds are neutral as well (see Fig. 5(f)).

Deleted: 3

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: 4

Deleted: -hour

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: 3

Deleted: 4

Deleted: 5

Deleted: -hour

Deleted: -h

Deleted: 4

Deleted: 5

Deleted: 4





Figure 5: As in Fig. 4 but for MSLP forecasts.

Deleted: 4
Deleted: 3



670 Figure 6: As in Fig. 4, but for MWS forecasts.

Nevertheless, Fig. 5 and Fig. 6 (g) and (h) indicate the positive impacts of assimilating Aeolus HLOS winds on the MSLP and MWS forecasts, compared to experiments 2418 and 2500. The averaged improvements of the MSLP forecasts are up to 4 hPa, while the averaged improvements of the MWS forecasts reach 5 Knots, after assimilation of Aeolus HLOS winds. As shown

Deleted: 5
 Deleted: 3
 Deleted: 4
 Deleted: 5

by Fig. 5 and Fig. 6 (c) and (d), the 48 h MSLP and MWS forecasts of experiments 2418 and 2500 underestimate the intensity of Ida (2021) from the first to second RI period. The corresponding 48 h MSLP and MWS forecasts from experiments 2418_L2B and 2500_L2B also underestimate the intensity of Ida, but they are stronger than those of the reference experiments owing to deeper MSLPs and higher MWSs.

- Deleted: 4
- Deleted: 5
- Deleted: -h
- Deleted: -h

3.5 Precipitation forecasts

- Deleted: 4

To evaluate the impacts of assimilating Aeolus HLOS winds on the rainfall structures of Ida (2021), we calculate ETS scores using the forecasts of 6 h accumulated precipitation against the IMERG precipitation data within a $10^\circ \times 10^\circ$ box centered on Hurricane Ida (2021). Figure 7 (a) exhibits the improvements of the 48 h averaged ETS scores of experiment 2406_L2B (with assimilation of Aeolus HLOS winds) for precipitation thresholds 10 and 15 mm when compared to experiment 2406 (without assimilation of Aeolus HLOS winds). The improvements of experiment 2406_L2B are evident for 15 mm precipitation thresholds, compared with experiment 2406. Positive impacts of assimilation of Aeolus HLOS winds on the 6 h accumulated precipitation forecasts for 10 and 15 mm precipitation thresholds can also be found in experiments 2412_L2B, 2418_L2B, and 2500_L2B (see Fig. 7 (b)-(d)) when compared to their corresponding reference experiments. These results imply that assimilating Aeolus HLOS winds can improve simulations of the rainfall structure of strong convection for Ida (2021).

- Deleted: -h
- Deleted: 6
- Deleted: reductions
- Deleted: -
- Deleted: different
- Deleted: -
- Deleted: mm
- Deleted: 6




Figure 7: Improvements of the 48 h averaged ETSs of forecasts initialized from the analysis DA cycles 5 (blue), 6 (green), 7 (brown), and 8 (pink) for different precipitation thresholds: 10 mm (without white lines) and 15 mm (with white lines) in experiments 2406 L2B (a), 2412 L2B (b), 2418 L2B (c), and 2500 L2B (d), compared with the corresponding experiments without assimilation of Aeolus HLOS winds. The ETS scores are calculated using the forecasts of 6 h accumulated precipitation against the IMERG precipitation data within a $10^\circ \times 10^\circ$ box centered on Hurricane Ida (2021).

3.6 Sensitivity of Aeolus data assimilation to horizontal and vertical localization scales

The data assimilation results are affected by the homogeneous isotropic horizontal and vertical ensemble localization scales. In experiment 2500 L2B H1, Ida (2021) remains a disturbance and then dissipates 12 hours after the initial time (results not shown), implying that the reduction of the horizontal localization scale to 55 km degrades the forecasts of Ida (2021). The reductions of the 48 h averaged track forecast errors, MSLP forecast errors, MWS forecast errors, and the improvements of ETSs for experiments 2500 L2B H2, 2500 L2B V1, and 2500 L2B V2, compared with experiment 2500 L2B, can be found in Figure 8. Specifically, Figures 8 (e)-(h) demonstrate that reducing the vertical localization scale to 1.5 grid points also leads to the degradation of the forecasts of Ida (2021). Consequently, reducing the horizontal and vertical localization

Deleted: ¶ [53]

Deleted: 6

Deleted: Reductions

Deleted: -h

Deleted: green

Deleted: pink

Deleted: -

Deleted: Positive values of ETS score reductions mean improvement, while negative values indicate degradation.

Deleted: ¶

Formatted: English (US)

Deleted: 5

Deleted: study

Deleted: Since the forecasts of Ida (2021) are

Deleted: ,

Deleted: we performed the sensitivity study of the localization scales for experiment 2500 L2B.

Deleted: so

Deleted: ing

Deleted: s

Deleted: .

Deleted: 7

Deleted: .

Deleted: 7

scales to half of their default values (in 2500_LB2) degraded the forecasts of Ida (2021) when compared with experiment 2500_L2B. Figures 8(a)-(d), and (i)-(l) show that the track forecasts are improved continuously in experiments 2500_L2B_H2 and 2500_L2B_V2 when compared with experiment 2500_L2B. However, the impacts on intensity and precipitation forecasts are small and mixed. Overall, the default horizontal localization scale (110 km) and vertical localization scale (3 grid units) are the best configurations for investigating the impacts of assimilating Aeolus HLOS winds on the analysis and forecasts of Hurricane Ida (2021); thus, they are used for other experiments in this study.




Figure 8: The reductions of the 48 h averaged track forecast errors (a), MSLP forecast errors (b), MWS forecast errors (c), and the improvements of ETSs (d) for DA cycles 5 (blue), 6 (green), 7 (brown), and 8 (pink) in experiments 2500_L2B_H2, compared with

Deleted: s

Deleted: 7

Deleted: tiny

Deleted: Thus

Deleted: the intensity forecasts of Ida (2021) are not sensitive to the horizontal and vertical localization scales when they are larger than the default values in the GSI system, which means that

Deleted: reasonable

Deleted: .

Formatted: Centered

Deleted: 7

Formatted: Caption

[experiment 2500_L2B. Positive values of reductions of track, MSLP, and MWS forecast errors mean improvement, while negative values indicate degradation. \(e\)-\(h\), and \(i\)-\(l\) as in \(a\)-\(d\), but for experiments 2500_L2B_V1, and 2500_L2B_V2.](#)

Formatted: English (US)

765 4 Diagnosis of influence of Aeolus DA on analysis of Hurricane Ida (2021)

4.1 Analysis increments of Aeolus HLOS winds

As mentioned in Sect. 3c, assimilating Aeolus HLOS winds leads to better intensity forecasts in experiments 2418_L2B and 2500_L2B from the first to second RI periods of Ida (2021). Thus, it is necessary to understand how assimilation of Aeolus HLOS winds influences Ida's dynamic and thermodynamic structure and then improves the intensity forecasts in experiments 770 2418_L2B and 2500_L2B. Commonly, improvements in a hurricane's inner core structure lead to positive impacts on hurricane intensity forecasts. Therefore, we choose experiments that assimilate Aeolus measurement swaths close to the center of Ida (2021) to examine their analysis increments over the hurricane's inner core region. As demonstrated by Fig. 1 (a), the Aeolus descending measurement swath on 25 August 2021, 12:00 UTC, and another on 26 August 2021, 12:00 UTC, are close to the center of Ida (2021) during the cycling DA period of experiments 2418_L2B and 2500_L2B and are suitable for 775 investigating the analysis increments of Aeolus HLOS winds. Figure 9 (a) and (c) show vertical cross sections of the analysis increments of Mie-cloudy winds and Rayleigh-clear winds for the selected Aeolus measurement swath on 25 August 2021, 12:00 UTC in experiment 2418_L2B. Since the HLOS winds of the Mie channel are derived by the Doppler-shifted backscattered light from the Fizeau interferometer, which detects aerosols and small hydrometeors, the analysis increments of Mie-cloudy winds are located primarily near the cloud top (between 10 and 16 km), as revealed by Fig. 9 (a). The analysis 780 increments of Mie-cloudy winds are mostly positive near the center of Ida (2021). Figure 9 (b) shows that the Rayleigh-clear winds, whose range bin thickness is 750 m between 12 and 15 km, are measured from the surface up to over 20 km in the tropical region (0 – 30° N). In comparison, the maximum measurement height in the extratropical region (30 – 60° N) is approximately 17.5 km, and the vertical resolution between 5 and 10 km (500 m) is higher due to the detection of the jet stream (ESA, 2020). However, ALADIN is totally attenuated by optically thick clouds or aerosols, so there are no Rayleigh-clear 785 winds under the cloud top near the center of Ida (2021). The resolution of the Mie-cloudy winds is much finer than that of the Rayleigh-clear winds, but the Rayleigh-clear winds have more extensive coverage. The analysis increments of Rayleigh-clear winds are roughly consistent with those of Mie-cloudy winds where they overlap. Figure 9 (b)-(d) and (f)-(h) are similar to Fig. 9 (a) and (c), respectively, but for different experiments or times. Note that the analysis increments of Mie-cloudy winds are mainly negative near the center of Ida (2021) on 25 August 2021, 12:00 UTC in experiment 2500_L2B.

Deleted: :00

Deleted: :00

Deleted: 8

Deleted: 7

Deleted: :00

Deleted: 8

Deleted: 7

Deleted: 8

Deleted: 7


Deleted: 8

Deleted: 7


Deleted: 8

Deleted: 7

Deleted: :00



805 **Figure 9:** Vertical cross sections of analysis increments of Mie-cloudy winds for the Aeolus descending measurement swath close to the center of Ida (2021) in experiments 2418_L2B (a) and 2500_L2B (b) on 25 August 2021, 12:00 UTC. The triangles indicate the locations of Ida (2021) in the simulations on 25 August 2021, 12:00 UTC. (c)-(d) As in (a)-(b) but for 26 August 2021, 12:00 UTC. (e)-(h) As in (a)-(d), but for Rayleigh-clear winds.




Deleted:

Deleted: 8

Deleted: 7... Vertical cross sections of analysis increments of Mie-cloudy winds for the Aeolus descending measurement swath close to the center of Ida (2021) in experiments 2418_L2B (a) and 2500_L2B (b) on 25 August 2021, 12:00...UTC. The triangles indicate the locations of Ida (2021) in the simulations on 25 August 2021, 12:00:00...UTC. (c)-(d) As in (a)-(b) but for 12 UTC on ...


4.2 Vertical cross-section of analysis increments of zonal winds and RH

825 The Aeolus HLOS winds are perpendicular to the Aeolus orbit, about 10 degrees off the zonal direction (Krisch et al., 2022). Thus, assimilating Aeolus HLOS winds generally impact the zonal wind component more than the meridional wind component. Figure 10, (c) compares vertical cross sections of zonal wind increments along the selected Aeolus descending measurement swath on 25 August 2021, 12:00 UTC, between experiments 2418 and 2418 L2B. The significant impacts of assimilating Aeolus HLOS winds on the analysis increments of zonal winds are located at the upper troposphere (near the cloud top), especially near the center of Ida (2021), which is consistent with Marinescu et al. (2022), and Garrett et al. (2022). As shown by Fig. 10, (c), the difference in the analysis increments of the zonal winds is negative at the cloud tops near the center of Ida (2021) due to the positive analysis increments of Mie-cloudy and Rayleigh-clear winds (descending orbit). Figure 10, (f) is similar to Fig. 10, (c), but for experiments initialized on 25 August 2021, 00:00 UTC. Figure 10, (f) demonstrates that the difference in the zonal wind increments is primarily positive at the cloud tops near the center of Ida (2021), owing to the negative analysis increments of Mie-cloudy and Rayleigh-clear winds. Figure 11, examines vertical cross sections of analysis increments of RH along the selected Aeolus descending measurement swath on 25 August 2021, 12:00 UTC, for experiments initialized on 24 August 2021, 18:00 UTC, and 25 August 2021, 00:00 UTC. As with the zonal wind increments, the impacts of assimilating Aeolus HLOS winds on RH increments are also located primarily in the upper troposphere (between 10 and 15 km).



840 **Figure 10:** Vertical cross section of zonal wind increments along the Aeolus descending measurement swath close to the center of Ida (2021) on 25 August 2021, 12:00 UTC in experiments 2418 (a) and 2418 L2B (b). (c) shows the differences in the zonal wind

Deleted: 9
Deleted: 8... (c) compares vertical cross sections of zonal wind increments along the selected Aeolus descending measurement swath on 25 August 2021, 12:00:00 (... [55])
Deleted: 9
Deleted: 8
Deleted: 9
Deleted: 8
Deleted: 9
Deleted: 8... (c), but for experiments initialized on 25 August 2021, 00:00:00 (... [56])
Deleted: 9
Deleted: 8
Deleted: 10
Deleted: 9... examines vertical cross sections of analysis increments of RH along the selected Aeolus descending measurement swath on 25 August 2021, 12:00:00... UTC, for (... [57])



Deleted:
Deleted: 9
Deleted: 8... Vertical cross section of zonal wind increments along the Aeolus descending measurement swath close to the center of Ida (2021) on 25 August 2021, 12:00:00 (... [58])

increments between experiments 2418_L2B and 2418. The triangles indicate the locations of Ida (2021) in the simulations. (d)-(f) As in (a)-(c), but for experiments initialized on 25 August 2021, 00:00 UTC.





Figure 11: As in Fig. 10, but for RH.

895 4.3 Vertical profiles of averaged analysis increments of divergence and RH

To further investigate the impacts of assimilating Aeolus HLOS winds on the dynamic and thermodynamic structure of Hurricane Ida (2021), we calculate the averaged analysis increments of divergence and RH within a 300 km radius of the center of Ida (2021). Figure 12 (a) shows that after assimilation of Aeolus HLOS winds on 25 August 2021, 12:00 UTC, the analysis increments of divergence slightly increase in the upper level of the troposphere (near 200 hPa), which may help the intensification processes of Ida (2021). The difference in the analysis increments of divergence is tiny in the middle and lower troposphere between experiments 2418 and 2418_L2B because the major impacts of assimilation of Aeolus HLOS winds are located at the cloud top near the center of Ida, as mentioned in Sect. 3f. On 26 August 2021, 12:00 UTC, the analysis increments of divergence decrease in the lower and middle troposphere (below 400 hPa), which may be caused by assimilation of NCEP GDAS data and different center locations of Ida (2021) in experiments 2418 and 2418_L2B. Figure 12 (b) demonstrates that the analysis increments of divergence near 200 hPa become stronger on 25 August 2021, 12:00 UTC, and the analysis increments of convergence in the upper troposphere become weaker on 26 August 2021, 12:00 UTC after assimilation of Aeolus HLOS winds. Figure 12 (c) and (d) show that assimilation of Aeolus HLOS winds has only tiny impacts on the moisture

Deleted: :00



Deleted:

Deleted: 0

Deleted: 9

Deleted: 9

Deleted: 8

Deleted: 1

Deleted: 0

Deleted: :00

Deleted: differences

Deleted: are

Deleted: :00

Deleted: conventional observations

Deleted: 1

Deleted: 0



Deleted: :00

Deleted: :00

Deleted: 1

Deleted: 0

structure of Ida (2021). In short, assimilation of Aeolus HLOS winds leads to stronger divergence in the upper level of the troposphere, which could be one reason for the improved intensity forecasts of experiments 2418_L2B and 2500_L2B.



930 **Figure 12:** Vertical profiles of averaged analysis increments of divergence (a) of experiments 2418 (dashed line) and 2418_L2B (solid line) on 25 August 2021, 12:00 UTC (green) and 26 August 2021, 12:00 UTC (pink). The analysis increments of divergence are averaged within a 300 km radius of the center of Ida (2021). (b) As in (a), but for experiments 2500 and 2500_L2B. (c)-(d) As in (a)-(b), but for averaged analysis increments of RH.

5 Impacts of Aeolus data on numerical simulations of an MCS associated with an AEW

940 5.1 Case description and experiment design

As indicated by the GOES-R channel 8 (see Fig. 1 (b)), there was a strong subtropical high over the AO on 21 August 2021, 00:00 UTC. TS Henri, moving northeastward, was west of the subtropical high, and MCSs could be found on the north coast of South America. As shown by the track of GFS-analyzed relative vorticity maxima at 700 hPa, an AEW, propagating westward on the south edge of the subtropical high, moved into the Caribbean Sea around 19 August 2021, 12:00 UTC. Steered by the subtropical ridge, the AEW, with scattered convection embedded inside, continued to move westward to northwestward and reached the western Caribbean Sea around 22 August 2021, 00:00 UTC. Experiments 1918 and 1918_L2B are carried out to assess the impacts of the assimilation of Aeolus HLOS winds on the analysis and forecasts of the MCS associated with the AEW. Experiment 1918 is initialized by the NCEP GFS FNL on 19 August 2021, 18:00 UTC, and assimilates only the NCEP GDAS data with 3DVAR during the cycling DA period from 20 August 2021, 00:00 UTC to 21 August 2021, 00:00 UTC (5 DA cycles in total). The subsequent forecast is initialized from the analysis of DA cycle 5. Experiment 1918_L2B is similar to experiment 1918 but also assimilates Aeolus HLOS winds.

5.2 Results

Figure 13 shows the 6 h accumulated precipitation from 21 August 2021, 18:00 UTC to 22 August 2021, 00:00 UTC, and the divergence over $5 \times 10^{-5} \text{ s}^{-1}$ at 850 hPa on 22 August 2021, 00:00 UTC. As shown by Fig. 13 (a), a large area of 6 h accumulated rainfall over 10 mm from the IMERG precipitation dataset is located within a 150 km radius and west of the center of the AEW. Due to the downward airflow created by the heavy rainfall, divergence over $5 \times 10^{-5} \text{ s}^{-1}$ from the GFS analysis can also be found west of the center of the AEW on 22 August 2021, 00:00 UTC. Figure 13 (b) shows only scattered convection and rainfall-induced low-level divergence near the center of the AEW in experiment 1918. As for experiment 1918_L2B (see Fig. 13 (c)), a large area of 6 h accumulated rainfall over 10 mm and rainfall-induced low-level divergence are located at the center of the AEW within a 150 km radius, which is more consistent with the IMERG precipitation and the low-level divergence from the GFS analysis, implying a positive impact on numerical simulations of the MCS by assimilating Aeolus HLOS winds.

Deleted: :00

Deleted: 700-hPa

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: conventional observations

Deleted: :00

Deleted: :00

Deleted: 2

Deleted: 1

Deleted: -

Deleted: :00

Deleted: :00

Deleted: 850-hPa

Deleted: :00

Deleted: 2

Deleted: 1

Deleted: 6-h

Deleted: :00

Deleted: 2

Deleted: 1

Deleted: 2

Deleted: 1

Deleted: 6-h





Figure 13: (a) 6 h accumulated IMERG precipitation (color shading) from 21 August 2021, 18:00 UTC to 22 August 2021, 00:00 UTC and divergence over $5 \times 10^{-5} s^{-1}$ at 850 hPa (black contour) from the GFS analysis on 22 August 2021, 00:00 UTC. (b)-(c) As in (a), but the precipitation and divergence are calculated by the WRF model simulations of experiment 1918 (b) and experiment 1918_L2B (c). The gray circles indicate radii of 150, 300, and 450 km, respectively.

6 Conclusions

Measurement of three-dimensional wind profiles in the GOS is essential for improving the ICs of global NWP, but it is insufficient over the oceans and remote land areas (WMO, 2017; Baker et al., 2014; Pu et al., 2017; Zhang and Pu, 2010; Pu et al. 2010; Rennie et al., 2021b). To fill the gap, the Aeolus satellite was launched by the ESA on 22 August 2018 and became the first spaceborne wind lidar (Reitebuch et al., 2020; ESA, 2022). Four types of Aeolus HLOS winds are available: Mie-clear, Mie-cloudy, Rayleigh-clear, and Rayleigh-cloudy (Jos de Kloe et al., 2022), but Mie-cloudy and Rayleigh-clear winds have better quality than the others (Zuo et al., 2022; Rani et al., 2022). The influences of assimilating Aeolus Mie-cloudy and Rayleigh-clear NRT HLOS winds on the forecasts of TCs and tropical convective systems have yet to be investigated. Thus, this study assesses the impacts of assimilating Aeolus Mie-cloudy and Rayleigh-clear HLOS winds on the analysis and forecasts of Hurricane Ida (2021) and an MCS embedded in an AEW during the CPEX-AW field campaign (2021). The WRF model and NCEP GSI-based 3DnVAR hybrid DA system are applied in this study.

Ida (2021) originated from an AEW on 14 August 2021 and became a TD on 26 August 2021, 12:00 UTC. Six hours after cyclogenesis, it intensified into a TS. Ida (2021) experienced two RI periods: from 00 UTC to 18 UTC on 27 August 2021 (RI1); and from 28 August 2021, 12:00 UTC to 29 August 2021, 12:00 UTC (RI2). Between these two RI periods, Ida (2021) made landfall in Cuba around 28 August 2021, 00:00 UTC (Beven II et al., 2022). Eight cycling DA experiments, including experiments 2406, 2406_L2B, 2412, 2412_L2B, 2418, 2418_L2B, 2500, and 2500_L2B, are carried out to investigate the impacts of assimilating Aeolus HLOS winds on the track and intensity forecasts of Ida (2021), especially during the two RI periods. The horizontal and vertical ensemble localization scales of these eight cycling DA experiments are set to 110 km and



Deleted:

Deleted: 2

Deleted: 1

Deleted: 6-h

Deleted: :00

Deleted: :00

Deleted: 850-hPa

Deleted: :00

Deleted: near-real-time

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: cycling

1025 [3 grid units, respectively, which has been proved reasonable for the data assimilation experiments, by a sensitivity study.](#)

Deleted: Ida (2021)

1030 Compared with the experiments without assimilation of Aeolus HLOS winds (experiments 2406, 2412, 2418, and 2500), the track forecasts are improved after assimilation of Aeolus HLOS winds (experiments 2406_L2B, 2412_L2B, 2418_L2B, 2500_L2B). The impacts on intensity forecasts are neutral in experiments 2406_L2B and 2412_L2B, while small but continuous improvements in intensity forecasts can be found in experiments 2418_L2B, and 2500_L2B. In addition, the ETS scores against the IMERG precipitation data near the center of Ida (2021) show that assimilating Aeolus HLOS winds can improve the [6 h](#) accumulated precipitation forecasts for strong convection (10 mm and 15 mm). One reason for the improved intensity and precipitation forecasts after assimilation of Aeolus HLOS winds is the stronger divergence in the upper level of the troposphere, as indicated by the averaged analysis increments of divergence within a 300 km radius of the center of Ida (2021) on 25 August 2021, 12:00 UTC and 26 August 2021, 12:00 UTC in experiments 2418_L2B and 2500_L2B. In addition

Deleted: 6-h

1035 to Hurricane Ida (2021), we also perform experiments 1918 and 1918_L2B to examine the impacts of assimilating Aeolus HLOS winds on an MCS embedded in an AEW, which was steered by a subtropical high and propagated westward from 19 August 2021, 12:00 UTC to 20 August 2021, 00:00 UTC in the Caribbean Sea. The results of the MCS demonstrate that assimilating Aeolus HLOS winds leads to better structure of the [6 h](#) accumulated precipitation from 21 August 2021, 18:00 UTC to 22 August 2021, 00:00 UTC, and rainfall-induced low-level divergence on 22 August 2021, 00:00 UTC.

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: 6-h

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: :00

Deleted: Although t

Deleted: and the MCS associated with an AEW, the assimilation techniques require further investigation and enhancement

Deleted: will

Deleted: improve

Deleted: -hourly

1040 [This study demonstrates the positive impacts of assimilating Aeolus Mie-cloudy and Rayleigh-clear winds on the forecasts of Hurricane Ida \(2021\) with comprehensive experiments.](#) More case studies are needed if the operational assimilation of Aeolus HLOS winds is required. Future studies [should emphasize improving](#) the technique of Aeolus HLOS wind assimilation with more cases of [hurricanes and](#) tropical convective systems.

1045 **Data availability.** The NCEP operational Global Forecast System analysis and forecasts at a [6 h](#) interval are obtained from <https://rda.ucar.edu/datasets/ds084.1> (last access: 31 August 2022, National Center for Atmospheric Research). The Aeolus L2B scientific wind products are accessed via <https://aeolus-ds.eo.esa.int/oads/access/collection> (last access: 05 September 2022, European Space Agency). The NHC best-track data can be accessed via <https://www.nhc.noaa.gov/data> (last access: 14 September 2022, National Hurricane Center and Central Pacific Hurricane Center). IMERG precipitation data are downloaded from https://disc.gsfc.nasa.gov/datasets/GPM_3IMERGHH_06/summary (last access: 12 April 2022, Goddard Earth Sciences Data and Information Services Center).

1055 **Author contribution.** Chengfeng Feng collected the data, performed the experiments, analyzed the experiment results, and drafted the manuscript. Zhaoxia Pu evolved the overall research goals and aims, acquired the financial support for the project leading to this publication, and revised the manuscript critically.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We appreciate the NCAR WRF model development group for providing the WRF model, the UCAR Developmental Tested Center (DTC) for the GSI DA system source code, and the Center for High-Performance Computing (CHPC) at the University of Utah for computer support. The NASA Award 80NSSC20K0900 is acknowledged for its support.

Comments from two anonymous reviewers are helpful and greatly appreciated.

Deleted: also

References

Andersson, E., Dabas, A., Endemann, M., Ingmann, P., Källén, E., Offiler, D., and Stoffelen, A.: ADM-Aeolus science report, ESA, 130 pp., <https://esamultimedia.esa.int/multimedia/publications/SP-1311/SP-1311.pdf>, 2008.

1085 Baars, H., Herzog, A., Heese, B., Ohneiser, K., Hanbuch, K., Hofer, J., Yin, Z., Engelmann, R., and Wandinger, U.: Validation of Aeolus wind products above the Atlantic Ocean, *Atmos. Meas. Tech.*, 13, 60070-6024, <https://doi.org/10.5194/amt13-6007-2020>, 2020.

1090 Baker, W. E., Atlas, R., Cardinali, C., Clement, A., Emmitt, G. D., Gentry, B. M., Hardesty, R. M., Källén, E., Kavaya, M., J., Langland, R., and Ma, Z.: Lidar-measured wind profiles: The missing link in the global observing system, *Bull. Amer. Meteor.*, 95, 543-564, <https://doi.org/10.1175/BAMS-D-12-00164.1>, 2014.

1095 Banyard, T. P., Wright, C. J., Hindley, N. P., Halloran, G., Krisch, I., Kaifler, B., and Hoffmann, L.: Atmospheric gravity waves in Aeolus wind lidar observations, *Geophys. Res. Lett.*, 48, e2021GL092756, <https://doi.org/10.1029/2021GL092756>, 2021.

Beven II, J. L., Hagen, A., and Berg, R.: Hurricane Ida (AL092021), National Hurricane Center, 163 pp., https://www.nhc.noaa.gov/data/tcr/AL092021_Ida.pdf, 2022.

1100 Dudhia, J.: Numerical study of convection observed during the winter monsoon experiment using a mesoscale two-dimensional model, *J. Atmos. Sci.*, 46, 3077-3107, [https://doi.org/10.1175/1520-0469\(1989\)046<3077:NSOCOD>2.0.CO;2](https://doi.org/10.1175/1520-0469(1989)046<3077:NSOCOD>2.0.CO;2), 1989.

1105 ESA-A guide to Aeolus range bin settings: <https://earth.esa.int/eogateway/news/a-guide-to-aeolus-range-bin-settings>, last access: 24 December 2022, 2020.

- ESA-Facts and figures: https://www.esa.int/Applications/Observing_the_Earth/Aeolus/Facts_and_figures, last access: 24 December 2022.
- 1110
ESA: ESA Aeolus online dissemination system, ESA [data set], <https://aeolus-ds.eo.esa.int/oads/access/>, 2021.
- Garrett, K., Liu, H., Ide, K., Hoffman, R. N., and Lukens, K. E.: Optimization and Impact Assessment of Aeolus HLOS Wind Data Assimilation in NOAA's Global Forecast System, *Q. J. R. Meteorol. Soc.*, 148, 2703-2716,
1115 <https://doi.org/10.1002/qj.4331>, 2022.
- Guo, J., Liu, B., Gong, W., Shi, L., Zhang, Y., Ma, Y., Zhang, J., Chen, T., Bai, K., Stoffelen, A., and de Leeuw, G.: First comparison of wind observations from ESA's satellite mission Aeolus and ground-based radar wind profiler network of China, *Atmos. Chem. Phys.*, 21, 2945-2958, <https://doi.org/10.5194/acp-21-2945-2021>, 2021.
- 1120
Hong, S. Y., and Lim, J. O. J.: The WRF single-moment 6-class microphysics scheme (WSM6), *J. Korean Meteor. Soc.*, 42, 129-151, 2006.
- Hong, S. Y., Noh, Y., and Dudhia, J.: A new vertical diffusion package with an explicit treatment of entrainment processes,
1125 *Mon. Wea. Rev.*, 134, 2318-2341, <https://doi.org/10.1175/MWR3199.1>, 2006.
- Huffman, G. J., Stocker E. F., Bolvin D. T., Nelkin E. J., and Tan, J.: GPM IMERG Final Precipitation L3 Half Hourly 0.1 degree x 0.1 degree V06, Goddard Earth Sciences Data and Information Services Center [data set],
<https://doi.org/10.5067/GPM/IMERG/3B-HH/06>, 2019.
- 1130
Ingmann, P., and Straume, A. G.: ADM-AEOLUS mission requirements document, ESA, 57 pp.,
https://esamultimedia.esa.int/docs/EarthObservation/ADM-Aeolus_MRD.pdf, 2016.
- Janjić, T., Bormann, N., Bocquet, M., Carton, J. A., Cohn, S. E., Dance, S. L., Losa, S. N., Nichols, N. K., Pothast, R.,
1135 Waller, J. A., and Weston, P.: On the representation error in data assimilation, *Q. J. R. Meteorol. Soc.*, 144, 1257-1278,
<https://doi.org/10.1002/qj.3130>, 2018.
- Jiménez, P. A., Dudhia, J., González-Rouco, J. F., Navarro, J., Montávez, J. P., and García-Bustamante, E.: A revised scheme for the WRF surface layer formulation, *Mon. Wea. Rev.*, 140, 898-918, <https://doi.org/10.1175/MWR-D-11-00056.1>, 2012.
- 1140

- Jos de Kloe, K. N. M. I., and Rennie, M.: Aeolus data innovation science cluster DISC ADM-Aeolus Level-2B/2C processor input/output data definitions interface control document, ECMWF, 128 pp.,
<https://earth.esa.int/eogateway/documents/20142/37627/Aeolus-L2B-2C-Input-Output-DD-ICD.pdf>, 2022.
- 1145
- Kain, J. S.: The Kain–Fritsch convective parameterization: An update, *J. Appl. Meteor. Climatol.*, 43, 170-181,
[https://doi.org/10.1175/1520-0450\(2004\)043<0170:TKCPAU>2.0.CO;2](https://doi.org/10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2), 2004.
- Krisch, I., Hindley, N. P., Reitebuch, O., and Wright, C. J.: On the derivation of zonal and meridional wind components
1150 from Aeolus horizontal line-of-sight wind, *Atmos. Meas. Tech.*, 15, 3465-3479, <https://doi.org/10.5194/amt-15-3465-2022>,
2022.
- Laroche, S., and St-James, J.: Impact of the Aeolus Level-2B horizontal line-of-sight winds in the Environment and Climate
Change Canada global forecast system, *Q. J. R. Meteorol. Soc.*, 148, 2047-2062, <https://doi.org/10.1002/qj.4300>, 2022.
- 1155
- Lukens, K. E., Ide, K., Garrett, K., Liu, H., Santek, D., Hoover, B., and Hoffman, R. N.: Exploiting Aeolus level-2b winds to
better characterize atmospheric motion vector bias and uncertainty, *Atmos. Meas. Tech.*, 15, 2719-2743,
<https://doi.org/10.5194/amt-15-2719-2022>, 2022.
- 1160
- Marinescu, P. J., Cucurull, L., Apodaca, K., Bucci, L., and Genkova, I.: The characterization and impact of Aeolus wind
profile observations in NOAA's regional tropical cyclone model (HWRF), *Q. J. R. Meteorol. Soc.*, 148, 3491-3508,
<https://doi.org/10.1002/qj.4370>, 2022.
- Martin, A., Weissmann, M., Reitebuch, O., Rennie, M., Geiß, A., and Cress, A.: Validation of Aeolus winds using
1165 radiosonde observations and numerical weather prediction model equivalents, *Atmos. Meas. Tech.*, 14, 2167-2183,
<https://doi.org/10.5194/amt-14-2167-2021>, 2021.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous
atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.*, 102, 16663-16682,
1170 <https://doi.org/10.1029/97JD00237>, 1997.
- Pourret, V., Šavli, M., Mahfouf, J. F., Raspaud, D., Doerenbecher, A., Bénichou, H., and Payan, C.: Operational assimilation
of Aeolus winds in the Météo-France global NWP model ARPEGE, *Q. J. R. Meteorol. Soc.*, 148, 2652-2671,
<https://doi.org/10.1002/qj.4329>, 2022.
- 1175

- Pu, Z., Zhang, L., and Emmitt, G. D.: Impact of airborne Doppler Wind Lidar data on numerical simulation of a tropical cyclone, *Geophys. Res. Lett.*, 37, L05801, <https://doi.org/10.1029/2009GL041765>, 2010.
- 1180 Pu, Z., Zhang, L., Zhang, S., Gentry, B., Emmitt, D., Demoz, B., and Atlas, R.: The impact of Doppler wind lidar
measurements on high-impact weather forecasting: Regional OSSE and data assimilation studies in: *Data Assimilation for
Atmospheric, Oceanic and Hydrologic Applications (Vol. III)*, edited by: Park, S. K., and Xu, L., Springer, Berlin,
Heidelberg, Germany, 259-283, https://doi.org/10.1007/978-3-319-43415-5_12, 2017.
- 1185 Rani, S. I., Jangid, B. P., Kumar, S., Bushair, M. T., Sharma, P., George, J. P., George, G., and Das Gupta, M.: Assessing the
quality of novel Aeolus winds for NWP applications at NCMRWF, *Q. J. R. Meteorol. Soc.*, 148, 1344-1367,
<https://doi.org/10.1002/qj.4264>, 2022.
- 1190 Reitebuch, O., Lemmerz, C., Lux, O., Marksteiner, U., Rahm, S., Weiler, F., Witschas, B., Meringer, M., Schmidt, K.,
Huber, D., Nikolaus, I., Geiss, A., Vaughan, M., Dabas, A., Flament, T., Stieglitz, H., Isaksen, L., Rennie, M., de Kloe, J.,
Marseille, G.-J., Stoffelen, A., Wernham, D., Kanitz, T., Straume, A.-G., Fehr, T., von Bismark, J., Floberghagen, R., and
Parrinello, T.: Initial assessment of the performance of the first Wind Lidar in space on Aeolus, *The 29th International Laser
Radar Conference*, Heifei, China, 24-28 June 2019, 01010, <https://doi.org/10.1051/epjconf/202023701010>, 2020.
- 1195 Reitebuch, O., Lemmerz, C., Nagel, E., Paffrath, U., Durand, Y., Endemann, M., Fabre, F., and Chaloupy, M.: The airborne
demonstrator for the direct-detection Doppler wind lidar ALADIN on ADM-Aeolus. Part I: Instrument design and
comparison to satellite instrument, *J. Atmos. Ocean Technol.*, 26, 2501-2515, <https://doi.org/10.1175/2009JTECHA1309.1>,
2009.
- 1200 Rennie, M. P., and Isaksen, L.: The NWP impact of Aeolus level-2B winds at ECMWF, ECMWF, 110 pp.,
<https://doi.org/10.21957/alif7mhr>, 2020.
- Rennie, M. P., Isaksen, L., Weiler, F., de Kloe, J., Kanitz, T., and Reitebuch, O.: The impact of Aeolus wind retrievals on
ECMWF global weather forecasts, *Q. J. R. Meteorol. Soc.*, 147, 3555-3586, <https://doi.org/10.1002/qj.4142>, 2021a.
- 1205 Rennie, M., Stoffelen, A., Khaykin, S., Osprey, S., Wright, C., Banyard, T., Straume, A. G., Reitebuch, O., Krisch, I.,
Parrinello, T., and Von Bismarck, J.: Demonstrated Aeolus benefits in atmospheric sciences, in: *2021 IEEE International
Geoscience and Remote Sensing Symposium IGARSS*, Brussels, Belgium, 11-16 July 2021, 763-766,
<https://doi.org/10.1109/IGARSS47720.2021.9554267>, 2021b.

- 1210 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., and Huang, X. Y.: A description of the advanced research WRF model version 4, NCAR, 145 pp., <https://doi.org/10.5065/1dfh-6p97>, 2019.
- Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M. A., Mitchell, K., Ek, M., Gayno, G., Wegiel, J., and Cuenca, R.
- 1215 H.: Implementation and verification of the unified NOAA land surface model in the WRF model, 20th Conf. on Weather Analysis and Forecasting/16th Conf. on Numerical Weather Prediction, Seattle, WA, U.S., January 2004, 14.2a, https://ams.confex.com/ams/84Annual/techprogram/paper_69061.htm, 2004.
- WMO: Guide to meteorological instruments and methods of observation, 2014 edition, WMO, Geneva, Switzerland, 1177
- 1220 pp., <http://dx.doi.org/10.25607/OBP-432>, 2017.
- Witschas, B., Lemmerz, C., Geiß, A., Lux, O., Marksteiner, U., Rahm, S., Reitebuch, O., and Weiler, F.: First validation of Aeolus wind observations by airborne Doppler wind lidar measurements, *Atmos. Meas. Tech.*, 13, 2381-2396, <https://doi.org/10.5194/amt-13-2381-2020>, 2020.
- 1225 Witschas, B., Lemmerz, C., Geiß, A., Lux, O., Marksteiner, U., Rahm, S., Reitebuch, O., Schäfler, A., and Weiler, F.: Validation of the Aeolus L2B wind product with airborne wind lidar measurements in the polar North Atlantic region and in the tropics, *Atmos. Meas. Tech.*, 15, 7049-7070, <https://doi.org/10.5194/amt-15-7049-2022>, 2022.
- 1230 Wu, S., Sun, K., Dai, G., Wang, X., Liu, X., Liu, B., Song, X., Reitebuch, O., Li, R., Yin, J., and Wang, X.: Inter-comparison of wind measurements in the atmospheric boundary layer and the lower troposphere with Aeolus and a ground-based coherent Doppler lidar network over China, *Atmos. Meas. Tech.*, 15, 131-148, <https://doi.org/10.5194/amt-15-131-2022>, 2022.
- 1235 Zhang, L., Pu, Z.: An Observing System Simulation Experiment (OSSE) to assess the impact of Doppler wind lidar (DWL) measurements on the numerical simulation of a tropical cyclone, *Adv. Meteorol.*, 2010, Article ID 743863, <https://doi.org/10.1155/2010/743863>, 2010.
- Zuo, H., Hasager, C. B., Karagali, I., Stoffelen, A., Marseille, G. J., and De Kloe, J.: Evaluation of Aeolus L2B wind product
- 1240 with wind profiling radar measurements and numerical weather prediction model equivalents over Australia, *Atmos. Meas. Tech.*, 15, 4107-4124, <https://doi.org/10.5194/amt-15-4107-2022>, 2022.