

# Observing Low Altitude Features in Ozone Concentrations in a Shoreline Environment via Uncrewed Aerial Systems

Josie K. Radtke<sup>1</sup>, Benjamin N. Kies<sup>1</sup>, Whitney A. Mottishaw<sup>1</sup>, Sydney M. Zeuli<sup>1</sup>, Aidan T.H. Voon<sup>1</sup>,  
Kelly L. Koerber<sup>1</sup>, Grant W. Petty<sup>2</sup>, Michael P. Vermeuel<sup>3†</sup>, Timothy H. Bertram<sup>3</sup>, Ankur R. Desai<sup>2</sup>,  
5 Joseph P. Hupy<sup>4</sup>, R. Bradley Pierce<sup>2,5</sup>, Timothy J. Wagner<sup>5</sup>, Patricia A. Cleary<sup>1</sup>

<sup>1</sup>Department of Chemistry, University of Wisconsin-Eau Claire, 105 Garfield Ave, Eau Claire, WI

<sup>2</sup>Department of Atmospheric and Oceanic Sciences, University of Wisconsin-Madison, 1225 W Dayton St, Madison, WI

<sup>3</sup>Department of Chemistry, University of Wisconsin-Madison, 500 Lincoln Dr, Madison, WI

<sup>4</sup>Aviation and Transportation Technology, Purdue Polytechnic Institute, Purdue University, West Lafayette, IN

10 <sup>5</sup>Space Sciences and Engineering Center, University of Wisconsin-Madison, Madison WI

<sup>†</sup>Now at Department of Soil, Water, and Climate, University of Minnesota, St. Paul, MN

*Correspondence to:* Patricia A. Cleary (clearypa@uwec.edu)

**Abstract.** Ozone is a pollutant formed in the atmosphere by photochemical processes involving nitrogen oxides (NO<sub>x</sub>) and volatile organic compounds (VOCs) when exposed to sunlight. Tropospheric boundary layer ozone is regularly measured at  
15 ground stations and sampled infrequently through balloon, lidar, and crewed aircraft platforms, which have demonstrated characteristic patterns with altitude. Here, to better resolve vertical profiles of ozone within the atmospheric boundary layer, we developed and evaluated an uncrewed aircraft system (UAS) platform for measuring ozone and meteorological parameters of temperature, pressure, and humidity. To evaluate this approach, an UAS was flown with a portable ozone monitor and a meteorological temperature and humidity sensor to compare to tall tower measurements in northern Wisconsin. In June 2020,  
20 as a part of the WiscoDISCO20 campaign, a DJI M600 hexacopter UAS was flown with the same sensors to measure Lake Michigan shoreline ozone concentrations. This latter UAS experiment revealed low-altitude structure in ozone concentrations in a shoreline environment showing highest ozone at altitudes from 20-100 mAGL. These first such measurements of low-altitude ozone via UAS in the Great Lakes Region revealed a very shallow layer of ozone rich air lying above the surface.

## 25 1 Introduction

Ozone at elevated concentrations near the surface is a pollutant that causes respiratory irritation in humans (Bell et al., 2006; Brauner et al., 2016 ) and oxidative stress on photosynthesizing organisms in many ecosystems (Fuhrer, 2002). In the troposphere, ozone is generated by reactions of nitrogen oxides (NO<sub>x</sub> = NO + NO<sub>2</sub>) and volatile organic compounds (VOCs) exposed to sunlight (Sillman, 1999). NO<sub>x</sub> compounds are emitted from combustion sources and VOCs are emitted by biogenic  
30 processes and anthropogenic industrial sources such as transportation and evaporated solvents (benzene, formaldehyde, ethyl alcohol, etc.). While ozone is monitored at the surface to meet various air quality monitoring standards, or to understand ozone

depositional losses, ozone gradients aloft have been measured in various ways over the years using sondes that reach the stratosphere (Beekmann et al., 1995; Witte et al., 2018), instrumented towers (Crawford et al., 1996; Desjardins et al., 1995), tethered balloons (Chandrasekar et al., 2003; Li et al., 2018; Mazzuca et al., 2017; Zhang et al., 2019; Tang et al., 2021; Demuer et al., 1997; Greenberg et al., 2009; Knapp et al., 1998), and crewed aircraft (e.g. (Kaser et al., 2017; Crawford et al., 1996; Tanimoto et al., 2015; Tarasick et al., 2019; Desjardins et al., 1995). Because ozone is generated by chemical reactions, the confinement of primary pollutants near the surface via atmospheric inversions tends to produce higher ozone concentration events at the surface. Understanding the volume of air in and above an inversion at a shoreline location prone to high ozone events can help elucidate the chemical evolution processes in this environment (Chai et al., 2013; Tang et al., 2021; Tang et al., 2009).

Recently there have been an expansion of efforts for Uncrewed Aircraft Systems (UAS) to be used for atmospheric profiling (Telg et al., 2017; Chilson et al., 2019; De Boer et al., 2021; Hemingway et al., 2017; Jacob et al., 2018; Koch et al., 2018; Wainwright et al., 2015; Li et al., 2018 ). Tethered balloons have been used to study vertical ozone (Demuer et al., 1997; Peng et al., 2008; Knapp et al., 1998; Zhang et al., 2019; Greenberg et al., 2009), and meteorological conditions (Chandrasekar et al., 2003) gathering data at heights ranging from ground level to 1500 meters above ground level, which included evaluations of episodes of biomass burning (Xu et al., 2018) and mesoscale modeling of ozone in the upper troposphere (Peng et al., 2008). UAS platforms measuring atmospheric properties have deployed at heights ranging from ground level to 4000 meters above ground level (Adkins and Sescu, 2017; Chilson et al., 2019; Cook et al., 2013; Greatwood et al., 2017; Hemingway et al., 2017). The portable Personal Ozone Monitor (2B Tech POM) mounted on a UAS performed consistently in comparison to a larger ozone photoanalyzer equipped to a tethered airship in the lower troposphere (Li et al., 2018) but with some significant discrepancies between platforms within the planetary boundary layer. Through modeling efforts using Generalized Additive Models (GAMs) Li et al. (2018) attributed these discrepancies to horizontal separation of platforms and vertical variations in atmospheric structure including temperature and relative humidity.

The effect of lake breeze or sea breeze on regional ozone in shoreline environments has been a point of interest in several studies. The association of sea breezes and lake breezes with elevated ozone at shoreline locations have been documented in Houston (Banta et al., 2005), Toronto (Levy et al., 2010; Sills et al., 2011), New York City during LISTOS (Zhang et al., 2020) , and near Chesapeake Bay (Gronoff et al., 2019), but few studies have explored vertical profiles within the marine layer structure. The lake and sea breeze meteorology develop from colder air parcels moving over land underneath buoyant warmer air which can create capping inversion which can trap pollutants (Lu and Turco, 1994; Gaza, 1998; Levy et al., 2010; Sills et al., 2011). Multiple groups have found there to be a notable difference in ozone levels during a sea or lake breeze including OWLETS (The Ozone Water-Land Transition Study) in the Chesapeake Bay region (Sullivan et al., 2019), ABLE (Amazon Boundary Layer Experiment) over Manaus Brazil (Guimaras et al., 2020), and a research team in the Salt Lake City region (Blaylock et al., 2017). OWLETS analyzed ozone pollution using ozone sensors mounted onto ships and UAS. These measurements showed that ozone builds up over the bay due to the effect of sea breeze up to 2000 m above sea level (Sullivan et al., 2019). With these observations, Sullivan et al. (2019) attempted to forecast chemical emissions based upon emissions


from ships and other emission sources in the bay. During ABLE, Guimaras et al. (2020) used UAS to study the urban nighttime boundary layer over Manaus, Brazil in both the dry and wet seasons. They conducted flights from the center of the city from ground level up to 500 m to quantify the effect humidity has on ozone pollution over Manaus at night (Guimaras et al., 2020). Crewed aircraft were used over the Great Salt Lake in Utah to study ozone levels up to 4000 m above ground level and  
70 demonstrated complicating factor of lake breeze transporting contrasting air masses into the region (Blaylock et al., 2017; Crosman et al., 2017; Horel et al., 2016).

The relationship between ambient ozone and coastal environments has been investigated by aircraft, mobile platforms for the 2017 Lake Michigan Ozone Study (LMOS) (Cleary et al., 2022b; Doak et al., 2021; Stanier et al., 2021) and UAS for the OWLETs campaign (Gronoff et al., 2019) and multi-UAS strategies for WiscoDISCO-21 (Tirado et al., 2023; Cleary et al.,  
75 2022a). Ozone concentrations have been shown to vary with altitude sharply in low-altitude crewed aircraft flights over Lake Michigan (Cleary et al., 2022b; Stanier et al., 2021). During the OWLETs campaign, the high-over-water ozone was investigated by UAS and ship-based platforms, including low ozone titration events. In these transitional environments, model and observation agreement can be improved with the capture of small gradients and modelling marine inversions over water (Abdi-Oskouei et al., 2020; Mcnider et al., 2018; Cleary et al., 2015). Recent observations over riverine environments have  
80 demonstrated the viability of UAS for detecting low altitude variations in ozone and plume chemistry (Li et al., 2021; Guimaras et al., 2020; Ye et al., 2022). The horizontal extent of lake breeze has also been documented at the shoreline to Lake Michigan where horizontal gradients close to the shoreline were observed during 2017 LMOS (Cleary et al., 2022b; Stanier et al., 2021).

The goal of this study is to develop a technique for investigating the vertical profiles of ozone at a shoreline location impacted by high ozone episodes. Chiwaukee Prairie, WI hosts a regulatory site at a shoreline state natural area, which is one of the few  
85 in Wisconsin which regularly exceed federal ozone standards and is regularly impacted by lake breeze. The large sources of emissions for ozone precursors are mainly concentrated in the Chicago metro area and the presence of Lake Michigan provides an inverted atmosphere at times in which to trap said pollutants. The role of the inversion over Lake Michigan, the advection of pollutants over Lake Michigan and then back on land during the meso-scale meteorological phenomenon of the Lake Breeze is the focus of the WiscoDISCO field campaigns. We first outline how the instrumentation was tested in a non-lake shore  
90 environment during CHEESEHEAD19 and then describe improvements to instrumentation performance for the first WiscoDISCO field campaign in 2020. Here, the UAS-based observations of ozone and meteorological variables were compared to tower observations in a forested environment in 2019 and then ground observations at a Lake Michigan shoreline in 2020 demonstrating improved performance and viability of a UAS atmospheric profiler to investigate lower atmospheric variability at a site impacted by lake breeze and poor air quality.

**2.1 CHEESEHEAD19 and PEcorINO Measurement Campaigns**

The University of Wisconsin-Eau Claire team joined the Chequamegon Heterogenous Ecosystem Energy-Balance Study Enabled by a High-density Extensive Array of Detectors 2019 (CHEESEHEAD19) campaign (Butterworth et al., 2021) in July 2019 in order to compare UAS-based observations with tower observations made during the first 7-day intensive observation period of the field campaign. CHEESEHEAD19 was the multi-institute campaign that sought to give insight into atmosphere-land exchanges in a temperate mixed forest (Butterworth et al., 2021). The CHEESEHEAD19 domain incorporated a swath of land in the Chequamegon National Forest near Park Falls, WI, where multiple tower, UAS, aircraft, ground, and remote sensing observations were conducted, focused around the 447 m instrumented tower operated by WLEF-TV (45.946 N, 90.273 W) and owned by the State of Wisconsin. Local vehicular traffic at the tall tower site was light and mixed trucking, forestry, and automobile traffic on WI Highway 182 (Figure 1). The tower has been in operation since 1995 as a National Oceanic and Atmospheric Administration (NOAA) greenhouse gas tall tower site (LEF) and since 1996 as an Ameriflux eddy covariance site (US-PFa), with sampling inlets and flux measurements currently at 30, 122 and 396 m above ground level. Ozone concentration observations were at two specific heights on the tower (30 m and 122 m) by two instruments: a chemical ionization time-of-flight mass spectrometer (CI-ToFMS, Tofwerk, and Aerodyne) using oxygen anion ( $O_2^-$ ) ionization chemistry (1 s L.O.D. ~10 ppt) (Novak et al., 2020) and an EPA standard photometric analyzer (L.O.D. 0.5 ppb ThermoScientific 49i) (Vermeuel et al., 2021). The fast observations of ozone by the CI-ToFMS instrument were used for flux measurements (Vermeuel et al., 2021). For the purposes of proving the viability of a UAS-mounted ozone measurement, the tower ozone measurements were compared to ozone gradient measurements from the UAS-mounted POM.



115 **Figure 1: During the CHEESEHEAD19 and PECorINO campaigns, measurements were taken at the WLEF TV ‘Very Tall Tower’ (northern Wisconsin). During the WISCODisco20 campaign, measurements were taken by the Chiwaukee Prairie Air Monitoring Station (southeastern Wisconsin). Map was made with ArcPro 2.8 using ESRI basemap data.**

120 A follow-up study Probing Ecosystem Responses Involving Notable Organics (PEcorINO) (Vermeuel et al., 2023) was conducted in September 2020 at the WLEF tower with observations of VOC and ozone at the 30 m inlet (Figure 1). A high-resolution proton-transfer reaction time-of-flight mass spectrometer (HR-PTR-ToFMS) (Vocus; Aerodyne Research Inc. and Tofwerk AG (Krechmer et al., 2018)), collected continuous 10-Hz measurements of VOCs and a photometric analyzer (ThermoScientific 49i) collected 1 Hz O<sub>3</sub> measurements at 30 m. Routine US-PFa site measurements of 10 Hz temperature and 1-Hz measurements of relative humidity (HMP-45C) were also collected during this period.

125 For CHEESEHEAD19, the Yuneec Typhoon H hexa-copter UAS was flown four days in July 2019 (July 8, 11, 12, and 16) during the campaign at the WLEF tower and September 13 and 14, 2020 during PEcorINO. The Typhoon H, owned by UW-Eau Claire, was chosen for this campaign because it was an inexpensive commercial UAS with capability of holding the payload of the POM. Flights in 2019 were conducted in the time window of 11 am - 3 pm local time (CDT) and at 11 am and 6 pm CDT in 2020. The Typhoon H was equipped with the POM for each of the flights at the tall tower, and an Internet Systems meteorological sensor, the iMet-XQ2, for the flights on July 16, 2019, and September flights from 2020 (See Figure 130 2a). The iMet-XQ2 sensor was placed on the landing gear of the Typhoon H to balance the payload. The days were chosen for

suitable flying conditions without strong winds ( $< 15$  mph gusts) or rainstorms or other precipitation. The Typhoon H was flown from a location roughly 100 ft from the tall tower in different patterns to hover for 5 min at 30 m, 60 m, 90 m and 122 m above ground level. Tower gradient uncertainties were determined from 1 standard deviation of the data from 30 and 122 m. The instruments sampling at the 122 m and 30 m heights from the tall tower were switched periodically (Vermeuel et al., 2021). The POM ozone data were collected at 10 s intervals and averaged to 5 minutes.

Before the CHEESEHEAD19 campaign, numerous test flights were necessary to work out payload distribution and to devise flight strategies. The Typhoon H had an approximate 15-minute flight time per battery with the payload. Each flight of the Typhoon H flights consisted of 2 hovers at different heights for 5 minutes. UAS flight log data were saved and was used as primary source for GPS data. All UAS flights were conducted under Federal Aviation Administration (FAA) Part 107 UAS regulations with a licensed UAS pilot.

## 2.2 The WiscoDISCO20 Campaign

The purpose of the Wisconsin's Dynamic Influence of Shoreline Circulations on Ozone (WiscoDISCO) campaign was to investigate the marine inversion influence on ozone measurements at the Lake Michigan shoreline by using an UAS at Chiwaukee Prairie Natural Area in Kenosha County, WI. A regulatory monitor at Chiwaukee Prairie managed by the Wisconsin Department of Natural Resources (WDNR) records some of the highest ozone in the state of Wisconsin and many Wisconsin shoreline Lake Michigan counties are in nonattainment of federal ozone standards (Stanier et al., 2021). Chiwaukee Prairie is located at the border between Wisconsin and Illinois and is situated between the coastal communities of Winthrop Harbor, Illinois, and Pleasant Prairie, Wisconsin. Suburban housing developments and mixed farmland surround the prairie (Figure 1). Local automobile traffic near to the monitor and UAS launch site was light and limited to neighborhood traffic and occasional train traffic.

The main goal of this campaign was to capture ozone exceedance days at this site where there was an influence of the lake breeze circulation. Ozone exceedance days are typically those in which the synoptic winds bring air from the south northward with high pressure systems over the Ohio Valley (Hanna and Chang, 1995), which are influenced heavily by Chicago pollution plumes. In this environment, temperature inversions commonly form when near-surface air is chilled by thermal exchange with the comparatively cold water of Lake Michigan and are exacerbated when lake breezes advect this dense but shallow layer of cold air inland (Wagner et al., 2022). The result is a shallow pool of colder, denser air overlain by warmer air aloft, with the inversion defined by the temperature increase with height at the boundary between the dissimilar air masses. Inversions act as a cap on the vertical mixing of air that would otherwise dilute and disperse  $\text{NO}_x$  and VOCs within these pollution plumes. Thus, these ozone precursors can accumulate in the near-surface air to relatively high concentrations.

During WiscoDISCO20 UAS were deployed on June 8, 9 and 15-19, 2020. The WiscoDISCO20 campaign was in collaboration with the Wisconsin Department of Natural Resources' (DNR)'s enhanced monitoring plan for the Chiwaukee Prairie site and included Pandora (Herman et al., 2009) (a ground-based differential optical absorption spectrometer which uses the sun as a light source to obtain total column trace gas measurements) and Doppler lidar observations at the site, provided

by the Space Science and Engineering Center at the University of Wisconsin-Madison. The Doppler lidar instruments were  
165 deployed on June 9, 2020 and operated continuously throughout the summer. The Pandora instrument is part of the Pandonia  
Global Network, (Verhoelst et al., 2021) which provides automated measurements of total column and tropospheric column  
NO<sub>2</sub>.

A DJI M600 hexa-copter was utilized in a collaborative research endeavor with Purdue University for the WiscoDISCO20  
170 campaign with an FAA compliant Part 107 UAS pilot, Joe Hupy. The DJI M600 had an increased payload capacity with its  
camera removed and the ability to place a top-mount for the sensor package, thus increasing the stability of the payload and  
providing a longer flight time than the Typhoon H (See Figure 2b). A 3D printed bracket to support the POM was mounted  
to the top of the vehicle. The inlet filter cartridge for the POM was held at a position with the least influence from propeller  
wash at the center of the top position of the UAS with a ~6 cm inlet tube. The iMet-XQ2 sensor was mounted to the bracket  
175 and secured with cable ties (See SI: Figure S2). During WiscoDISCO20, a series of flights were conducted to produce an  
atmospheric vertical profile with fixed altitudes where the UAS hovered for 5 minutes at each designated altitude. The flight  
times were approximately 15-20 minutes where the UAS would ascend for 15 m altitude increments where it would hover  
for 5 minutes. In an approximate 1.25-hour time window, 8 heights were sampled from 0-122 m AGL with 3 individual  
flights (See SI: Table S1). Flights were conducted from a gravel road inside of the Chiwaukee Prairie State Natural Area,  
with two focused vertical profile sampling periods: one in the morning at approximately 7-9 am local time (CDT) and one in  
180 the afternoon at approximately 2-4 pm local time (CDT).



**Figure 2.** a) Typhoon H UAS with mounted POM and iMet-XQ2 as flown during CHEESEHEAD-19 in 2019. POM was housed in foam for vibration dampening. b) Top mounted iMet-XQ2 and POM on a DJI M600 as flown in Sept. 2020 for WiscoDISCO20. The inlet to the POM is held up with a bracket to hold the inlet filter assembly (blue and white).

### 2.3 Personal Ozone Monitor, POM

The 2B Tech personal ozone monitor, POM, measured ozone concentrations via UV absorption spectroscopy, which is designed to account for a known interference with humidity in the atmosphere (Wilson and Birks, 2006). The POM measures ozone concentrations by calculating the difference in absorption between a whole air and a scrubbed ozone air sample in-series with one optical cell. The POM operates with an in-series duty cycle of measuring the whole air sample for 5 seconds and an ozone-scrubbed background air sample for another 5 seconds in the same optical cell (Andersen et al., 2010). This differs from current robust ground analysers, such as the ThermoScientific 49i, which use dual optical cells, one chamber of whole air and another chamber with ozone scrubbed out to measure a real-time background interference in the absorption signal (Ollison et al., 2013) (Wilson and Birks, 2006). This duty cycle must be considered when the POM is on a moving platform, as the air sampled in the first 5 seconds must be representative of the air sampled in the second 5 seconds for each measurement, therefore slow movement of the UAS during sampling was preferred. For all measurements described here, the UAS was held at one altitude for 5 minutes to collect representative data from that airmass. The absorption spectroscopy principle behind the POM with the active background humidity subtraction has a higher specificity to ozone than other light-weight electrochemical sensors (Kim et al., 2018). The POM was calibrated with the 2B Tech Model 309 transfer standard ozone generator within 24 hours of UAS flights during CHEESEHEAD19. The POM was placed in a foam case to dampen any vibrations associated with the UAS to which it was fastened. The filter on the POM was used for all flights to protect the optical cell from atmospheric particles and debris. The POM was independently powered by lithium-ion batteries. During WiscoDISCO20, the POM was calibrated with the Model 309 ozone generator within 2 hours of each atmospheric profile series of UAS flights. Zero drift of the POM was monitored by collecting scrubbed-ozone data using a cartridge ozone scrubber in between flights. The 2B Tech listed POM accuracy and precision are given as 1.5 ppb or 2% of observations whichever is highest. For the range of observations in this study, the accuracy and precision ranged from 1.5 ppb for many morning observations to up to 2.1 ppb for high ozone afternoon observations.

### 2.4 iMet-XQ2

The iMet-XQ2 sensor is lightweight and portable which allows it to measure temperature (bead thermistor), relative humidity (capacitive), and pressure (piezoresistive) along with recording GPS data with its own internal storage and power systems. The International Met Systems listed iMet-XQ2 accuracy and resolution of  $\pm 0.3$  °C and 0.01 °C for temperature,  $\pm 5\%$



and 0.1% for relative humidity,  $\pm 1.5$  hPa and 0.01 hPa for pressure, and an accuracy of 12 m for vertical GPS data. The data  
210 were extracted from the iMet-XQS after each flight.

Previous studies have evaluated the accuracy of the iMet-XQ2 on UAS platforms (Kimball et al., 2020; Inoue and  
Sato, 2022). Kimball et al. executed an exhaustive study on the performance of the iMet-XQ on a UAS in certain solar radiation  
and wind speed conditions. While they found that under low solar radiation, the accuracy and precision of the temperature  
measurement followed the listed accuracy and precision, with some direct solar radiation, higher wind speeds on the thermistor  
215 allowed for improved precision of the measurements. In cold conditions, shielding the thermistor from both solar radiation and  
heating from the UAS is important (Inoue and Sato, 2022). Sensor position on the UAS has been found to be important for  
preventing additional bias from motor heating and propeller wash if the sensor is placed too close to UAS motors (Greene et  
al., 2019). For this study, a lower accuracy of the iMet-XQ was considered reasonable to ascertain the vertical profile structure  
of the atmosphere at a shoreline location, if the solar radiation conditions and flight conditions were similar throughout the  
220 data collection flight.

### 3 Results and Discussion

#### 3.1 UAS to Tower Comparisons

During the CHEESEHEAD19 campaign, an intercomparison was conducted between the observations of ozone from the  
WLEF tall tower and UAS.. The tall tower ozone measurements were from either a ThermoFischer 49i photometric analyzer  
225 or a CI-ToFMS instrument. Each sampled air from either the 122 m or 30 m inlet simultaneously, and source inlets (i.e. sampled  
heights) were switched between instruments periodically. The absolute ozone concentrations at the 122 m and 30 m heights  
from the tall tower ranged from mid-day highs of 40-60 ppb. Tower ozone gradients were calculated as the difference between  
the ozone measured at 122 m and 30 m inlet heights. These tower observations were compared to the gradient ozone  
observations made by hovering the UAS at the 122 m and 30 m altitudes for 5 minutes each. The gradient ozone observations  
230 reproduced the reported ozone gradients on the tall tower within the considerable uncertainty (See Table 1). The absolute  
concentrations from the POM were as much as 20 ppb higher than the tower observations (See SI: Figure S3), with tower  
observations from both the 49i and TOF considered to be the more reliable measurements with consistent calibration  
procedures and low detection limits (Vermeuel et al., 2021; Vermeuel et al., 2023). Technically the overall comparison between  
tower gradients and UAS gradients show agreement; however, the considerable uncertainties make POM gradients from 8 and  
235 11 July indistinguishable from zero (See Table 1). This evaluation demonstrated a likely source of inaccuracy with POM ozone  
observations, with significant offset from the absolute tower observations. These inaccuracies have since been attributed to  
zero-point drift of the POM, which was substantiated by further laboratory experiments on calibration conditions of the POM.  
Those experiments showed differences in calibrations due to different sources of power to the POM (batteries versus wall-  
power). Large noise in the POM observations was attributed to disrupted airflow from propeller wash which was exacerbated  
240 by the bottom-mount of the POM on the UAS, as observed as higher noise during take-off and at the start of every hover.

Improvements to the UAS sensor package for the WiscoDISCO20 system were developed as a result of these findings as follows: a) the POM was mounted at the top of a larger UAS with the inlet positioned to the center of a larger, more robust platform, b) the POM was calibrated with the same independent POM battery source as the flights and calibrations were conducted within 2 hours of every flight and c) zero drift was monitored by placing an in-line ozone scrubber on the POM inlet directly after each flight for 5 minutes. The temperature and relative humidity measurements observed from the UAS using the iMet were found to vary from the tower measurements by no more than 1.7°C for temperature and 8% RH (See Table 2).

**Table 1: Comparison of ozone gradients made from Tall Tower at Park Falls and UAS-based POM during CHEESEHEAD-19. Ozone gradient,  $\Delta O_3$ , calculated as measured  $O_3$  at 122 m –  $O_3$  at 30 m. The tower measurements were selected as coincident with UAS-mounted POM measurements. Uncertainties reported are propagated from 1 standard deviation at each altitude.**

<b>Day-Month- Year of Flight</b>	<b>POM UAS</b>	<b>Tower</b>
	$\Delta O_3 \pm \sigma$ (ppb)	$\Delta O_3 \pm \sigma$ (ppb)
<b>08-Jul-19</b>	$-5.9 \pm 9.6$	$1.0 \pm 1.1$
<b>11-Jul-19</b>	$11.9 \pm 21.7$	$8.7 \pm 0.8$
<b>12-Jul-19</b>	$16.1 \pm 13.2$	$9.1 \pm 1.3$

255

**Table 2: Comparison of average air temperature and relative humidity made from Tall Tower at Park Falls and iMet-XQ during CHEESEHEAD-19 and PÉcorINO in 2020. The average Tower temperatures at the 30-meter inlet were computed at the time intervals when the UAS altitude was 30 meters AGL. The iMET and Tower data were evaluated at 1 Hz, therefore approximately n=300 for each 5-min hover period.**

<b>Day-Month- Year of Flight</b>	<b>Altitude</b>	<b>iMet UAS</b>	<b>Tower</b>	<b>iMet UAS</b>	<b>Tower</b>
	A (meters)	$T \pm \sigma$ (°C)	$T \pm \sigma$ (°C)	$RH \pm \sigma$ (%)	$RH \pm \sigma$ (%)
<b>16-Jul-19</b>	30	$25.0 \pm 0.4$	$24.43 \pm 0.07$	$61.2 \pm 1.3$	$66.8 \pm 0.4$
<b>13-Sep-20</b>	30	$15.5 \pm 0.3$	$13.8 \pm 0.9$	$68.9 \pm 0.8$	$76.7 \pm 5.5$


260

### 3.2 Observations at Lake Michigan Shoreline: WiscoDISCO20 UAS to Ground Comparisons

265 The viability for UAS-mounted ozone observations to capture low-altitude features in ozone is well-matched to the small-scale vertical structure of marine layer ozone concentrations in a near-shore environment. For the June 2020 observations, the UAS platform was the DJI M600 with a top-mounted bracket for positioning the filter cartridge for the POM in a space for minimal disruption of the air mass from propeller wash. The iMet-XQ2 sensor was mounted to this bracket to the side of the POM (See SI: Figure S2). The DJI M600 was flown at the Chiwaukee Prairie State Natural Area to capture shoreline airmasses impacted by lake breeze onshore flow during time of high ozone. The week of June 15-19, 2020, provided ideal conditions for high ozone and lake breeze conditions (as seen in Figure 3) where daytime winds shifted regularly to southeasterly and daily maximum temperatures increased throughout the week (See SI for identification of lake breeze from GOES-East satellite imagery). Most days during the week of June 15-19 had observable cumulous cloud suppression fronts over land near to the shoreline of Lake Michigan which is indicative of marine air incursion over land (see SI: Figures S4-S5). Particulate matter concentrations also increased during the week. The UAS was flown in a 2-hour window to capture morning and afternoon

270

vertical atmospheric profiles. A single battery flight of the UAS accounted for 3-4 hover heights and multiple sets of batteries were used to hover for 10 total points to get a vertical distribution.




275


**Figure 3: June 8-22, 2020 a) temperature in °C b) wind direction, c)  $PM_{2.5}$  in  $\mu g/m^3$  and d)  $O_3$  as measured at the WDNR ground station (black) and on the UAS via POM (red).**

The accuracy of the ozone concentration, temperature, and relative humidity (RH) observations made aloft on the UAS was evaluated by comparing the lowest altitude hover altitude at 9 meters above ground level (m AGL) to 1-minute data from the local air monitoring station in Chiwaukee Prairie (AIRSID# 55-059-0019). The uncertainty in the UAS-mounted POM was

determined to be the 1 standard deviation in the averaged 10 s data. A regression analysis of the two observations is given in Figure 4a; these data are strongly correlated as the  $R^2$  value is 0.939. The linear fit to the graph is weighted by the highest ozone data and the RMSD = 5.3 ppb. Some disagreement could be from the discrepancy in altitudes for the two observations (the DNR inlet is at 4.5 m in comparison to the first altitude for hovers at 9 m), or to accuracy issues with the zero drift as observed during CHEESEHEAD-19. A similar comparison was conducted for the iMet temperature measured at the lowest hovering altitude (approx. 9 m) with ground temperatures (Figure 4b) with an agreement at  $R^2= 0.944$ . With these added observations, the accuracy for the  $O_3$  concentrations via UAS-mounted POM are considered accurate within 10 ppb, and the iMet temperatures within 11 %. This figure has some similarities for the Li et al. (Li et al., 2020) figure 5a, where they saw a linear fit of  $0.7x - 7$  for a POM correlation to a regulatory ozone measurement instrument standard. The difference between our measurement and theirs is that we see more observations along the 1:1 line with higher ozone concentrations deviating the most from the center line, whereas Li et al. (2020) showed a consistent linear response at  $\sim 70\%$  of the regulatory  $O_3$  measurement.



a)



b)

295

Figure 4: a) Intercomparison  $O_3$  UAS POM measurements in comparison to  $O_3$  WiDNR Chiwaukee Prairie measurements on June 8, 9, 15-19 2020. The 5- minute average WiDNR and UAS POM data from the lowest hovering altitude is with uncertainties as  $1\sigma$  from the mean. The grey line demonstrates a 1:1 line and the red line depicts a linear regression fit ( $R^2 = 0.939$ ), with a fit of  $[O_3 \text{ POM}] = 0.861 (\pm 0.078) [O_3 \text{ WiDNR}] + 0.9 (\pm 4.8)$ . b) Intercomparison of temperature from lowest altitude reading from the UAS-mounted iMet-XQ2 and the WiDNR ground station. Red line indicates the linear regression ( $T_{\text{iMet-XQ2}} = 1.12 (\pm 0.11) T_{\text{WiDNR}} - 2.7 (\pm 2.5)$ ,  $R^2 = 0.944$ ) and the gray line is 1:1 fit.

300

### 3.3 Case study: Low Altitude Gradients at the Lake Michigan Shoreline

The week of June 15-19, 2020 had 4 days where  $O_3$  concentrations exceeded 70 ppb (Figure 3 d). The dominant winds were from the south and lake breezes were observed on all days that week (Figure 3 b), which are conditions conducive to higher ozone concentrations along the Lake Michigan shoreline due to Chicago emissions getting trapped over Lake Michigan during optimal conditions for photochemical production of ozone and then advecting ozone back on land at the shoreline (Vermeuel et al., 2019; Abdi-Oskouei et al., 2020; Cleary et al., 2022b; Baker et al., 2023). The conditions near Lake Michigan were consistently sunny at the shoreline with some evidence for cumulus cloud formation inland on June 19, 2020 often used as a identifying signature of lake breeze from satellite observations (Levy et al., 2010; Sills et al., 2011).

310

Vertical profiles for UAS flights were constructed using hovering altitudes of the UAS and time stamps for each altitude to determine observed average  $O_3$ , temperature, pressure and relative humidity (RH) for each altitude. Because of limited battery time, each vertical profile was constructed from 3-4 hovering altitudes during 3 separate 20-minute flights, covering a time window of approximately 1.25 hours (See SI Table S2). Figure 5 depicts vertical profiles of potential temperature overlaid

315 with ozone concentrations through the week of June 15-19, 2020. Every day shows an inverted stable atmosphere. Some days  
show a well-mixed buoyant internal boundary layer in the lowest 40-100 m AGL (Figure 5) where the potential temperature  
is close to a vertical line with respect to altitude, particularly in the June 18 and 19 afternoon flights. This discontinuity of most  
vertical profiles exists where the lowest 40-60 m AGL is closer to a more vertical potential temperature profile, which we refer  
320 a gradient of 5 K or more within 60-100 m AGL. The morning of June 18 (Fig 5-c) was the only day to show a steep inversion  
down to the surface with no discontinuity. Ozone concentrations in all ascents had maximum observations below the maximum  
altitude of the flight. Ozone concentrations tended to be highest near areas of steep inversion (June 15, June 17 am and pm,  
and June 18 pm flights) or near/within the internal boundary layer (June 16 pm, June 19) except on June 18 in the morning  
when ozone concentrations were highest at the lowest altitudes when the inversion extended to the surface. For all 5 days,  
325 observed afternoon maximum ozone concentrations were observed at higher altitudes than adjacent to the surface (Figures 5  
a-e). The higher ozone concentrations in the vertical profiles tended to be at or near the maximum inversion generally in the  
region of 40-60 m AGL.

The variation in height of the steep inversion layer is evident in the day-to-day differences, from as low as 40 m AGL (June  
15, 16, and 17) to as high as 100 m AGL on June 19. Morning to afternoon differences on July 16 and 17 show a steeper  
330 gradient in temperature later in the afternoon with not much change in the inversion height. By contrast, on the morning of  
July 18, the vertical profile of temperature shows an inversion starting at the surface (Figure 5d) and by the afternoon the  
inversion height starts at 60 m AGL. In comparison, turbulent kinetic energy (TKE) based boundary layer depths given by the  
High-Resolution Rapid Refresh (HRRR) (Dowell et al., 2022) atmospheric model outputs extend from 80 to 250 m AGL for  
this location, not as low as the data in Figure 5. HRRR boundary layer height is a metric which addresses how photochemical  
335 models are treating vertical profiles when computing photochemical ozone production. The use of the HRRR boundary layer  
height highlights the sub-grid scale of the vertical profiling, which indicates that UAS observations can sample important  
properties of the marine air incursion of a lake breeze. The lower boundary layer heights in the afternoon in comparison to the  
morning are proposed to arise from stronger synoptic winds degrading the inversion from above (Lyman and Tran, 2015).  
Doppler lidar measurements (which cannot make observations below 100 m AGL) show high aerosol loading in the afternoons  
340 at altitudes close to the ~100 m cut-off altitudes below which the instrument has a dead zone, which may correspond to  
continued inversion up to 200 m or more. The UAS observations give a complementary measurement to indicate the region of  
inversion and the compositional layering below, within, and above the inversion to demonstrate a more complicated picture of  
mixing and vertical stratification in the lower atmosphere. While these measurements may not adequately address exactly why  
models do not represent the shoreline effectively (See SI Figure S6), they can shed light on the required resolution and vertical  
345 structure that encompasses plume volume within an inverted atmosphere near Lake Michigan.


The UAS observations at Chiwaukee Prairie shown here demonstrate a very shallow internal boundary layer (40-100  
m AGL) developed over land underneath the temperature inversion (modeled boundary layer heights 80-250 m AGL), where  
ozone is found to be in highest abundance near the maximum inversion. On two days with the highest internal boundary layer

height, ozone concentrations were highest within the internal boundary layer, suggesting that an extended internal boundary layer height over land could possibly play a role in transport of pollutants in the marine layer. However, more observations of atmospheric profiles of ozone and meteorology are required over land and over water to better characterize this transitional environment.

The work by Li et al. (2020) described use of POM and particle observation on a fixed-wing UAS flying at a speed of 150 km/hr and compared measurements from those instruments to regulatory instruments on a tethered airship and addressed intercomparison with the POM and a regulatory ozone measurement instrument ( $O_3$ 42M from ESA). They used an insulated box for the POM and were able to show high correlation with a regulatory monitor, but with an offset. Their conclusions are that the POM measures atmospheric variability consistent with a regulatory monitor but demonstrates a negative bias. Here, we flew the POM at a much lower flight speed, and only averaged data from a single hovered point at which we stayed for 5 minutes each flight. This was to address the duty-cycle limitations of the POM with the on-off in series subtraction of the water vapor absorption. Li et al. address only that the regulatory monitor they used for comparison which employed an in-line heating method for removing water vapor interference, instead of a dual-cell active subtraction in parallel as is typical for other regulatory monitors. While Li et al. 2020 demonstrated some correlation between RH and variability between the UAS-mounted POM and tethered-airship-platform regulatory monitors, they do show that vertical gradients can be captured by UAS and tethered airship, but with discrepancies in location of planetary boundary layer. This is consistent with our observations that the gradient observations from UAS are in agreement (with high variability) with tower-based observations in the lowest 120 m AGL. What we cannot account for here is the difference in POM variability on a UAS which hovers for 5 minutes in comparison to a fixed-wing travelling at 150 km hr<sup>-1</sup>, which may also lead to additional variability in the measurement due to inlet pressure changes and optical cell vibrations. Additional improvements to the POM performance could arise from a) thermal insulation and b) a slow constant ascent instead of hovering and are intended for future studies. Additional improvements to the iMET-XQ2 performance could arise from a slow ascent (to assist in aspirating the thermistor) and shielding the iMET to account for solar irradiation of the sensors.

The feasibility of using UAS in shoreline environments depends on the vertical scale of the phenomenon, the UAS flight time and requisite battery life for such UAS observations and the legal flight conditions within each municipality. Some researchers have successfully used UAS for vertical ozone profiles up to 1000 m (Chen et al., 2022; Wu et al., 2021), tethered balloons (Li et al., 2020; Chen et al., 2022) and thermally insulated a UAS-mounted POM in the winter (Chen et al., 2020; Chen et al., 2019). The scales of sea breeze influence on vertical profiles have ranged from 400-600 m AGL at coastal locations in Nova Scotia (Gong et al., 2000), 600-800 m AGL at coastal locations in China (Wu et al., 2010) and 400-800 m AGL in lake breeze influenced locations in Saskatchewan (Sun et al., 1998). The lake breeze vertical dimensionality near Lake Michigan has been shown to have large effects at altitudes from 50-500 m AGL from crewed aircraft (Stanier et al., 2021; Cleary et al., 2022b), remote sensing (Wagner et al., 2022) and UAS studies (Tirado et al., 2023).





385 **Figure 5: Altitude (m AGL) versus potential temperature (K) with O<sub>3</sub> (ppb) for a) June 15 flights, June 16 b) morning, and c) afternoon flights, June 17 d) morning and e) afternoon flights, June 18 f) morning and g) afternoon flights, and h) June 19 afternoon flights. All times in CDT (2020). Grey and black bars indicate HRRR boundary layer heights for morning and afternoon, respectively.**

## 4 Conclusions

An UAS atmospheric profiler was tested in comparison to tower-based instrumental observations indicating a need for careful adjustments to operating parameters for the ozone measurements. Improvements including a top-mount for the sensor package, a larger UAS and higher frequency calibrations and zero-drift checks were shown to improve overall accuracy of the ozone observations near a ground air monitoring station. The improved vertical atmospheric profiler was shown to capture atmospheric variability in the lowest 120 m of the atmosphere at a Lake Michigan shoreline region, demonstrating a feasible use for UAS in atmospheric sampling to connect the scales of ground-based observations and tower or remote sensing aloft. These observations are the first UAS observations of ozone near Lake Michigan that document the over-land penetration of the marine layer and gradients in ozone within it. This work highlights the necessity for higher vertical resolution in observations in this shoreline location to inform improvements to how air quality models represent the stratification and mixing of air parcels at locations like these.

Suggestions for further improvements: in this study, the POM performance on UAS was improved by inlet positioning and slow flight parameters, top-mount placement on a robust UAS and increasing the rate of calibrations to pair each calibration with specific battery power source improved the precision and accuracy. However, added thermal insulation, as described by Li et al, appears another promising additional consideration for improved performance of the POM on UAS. The POM appears to be a robust enough instrument for course atmospheric measurements in the atmosphere (to 2 ppb precision or 2% of reading) but integration onto a UAS should be carefully considered. A wider variety of studies have been conducted on iMET performance on UAS (Kimball et al., 2020; Inoue and Sato, 2022; Greene et al., 2019).

## Appendices

Supplemental Info

### Data Availability

A data repository was generated for CHEESEHEAD19 at: [https://data.eol.ucar.edu/master\\_lists/generated/cheesehead/](https://data.eol.ucar.edu/master_lists/generated/cheesehead/) (last accessed 7/6/2023).

A data repository was generated for the WiscoDISCO 2020 field campaign at: <https://zenodo.org/communities/wiscodisco2020/> (last accessed 7/6/2023). Dataset available at DOI: 10.5281/zenodo.8118176. Each data file is in a .txt tab delimited structure with descriptive column titles. Any data file with a full suite of data from both iMET and POM instruments is given without a qualifier. On days when data was collected from one of the instruments, the file names indicate them as such.

## Author Contributions

JR, BK, MPV and KK contributed to data acquisition, data analysis and manuscript writing. WM, SZ, and AV contributed to data analysis. GP and TW contributed to data acquisition and manuscript writing and editing. AD, TB, and RBP contributed to field campaign planning and manuscript editing and writing. JPH contributed to field campaign planning, data analysis and manuscript writing and editing. PAC contributed to field campaign management, data acquisition, data analysis, manuscript writing and editing.

## Competing Interests

The contact author has declared that none of the authors has any competing interests.

## Acknowledgements

University of Wisconsin -Eau Claire team acknowledges funding from the Office of Research and Sponsored Programs Faculty/Student Research Collaboration Grants through Blugold differential tuition. This research is funded through the National Science Foundation Grant AGS-1918850. Ankur Desai coordinated the efforts for CHEESEHEAD19 and acknowledges funding from NSF AGS-1822420 and the Dept of Energy Ameriflux Network Management Program award to the ChEAS core site cluster. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation. Ancestral & Indigenous Lands Acknowledgement: We acknowledge that our research took place at the ancestral lands of the Ojéwéthi Šakówinj, Anishinabewaki, Ho-Chunk, Myaamia, Potawatomi, Kaskaskia, Peoria and Kiikaapoi people.

## References

- Abdi-Oskouei, M., Carmichael, G., Christiansen, M., Ferrada, G., Roozitalab, B., Sobhani, N., Wade, K., Czarnetzki, A., Pierce, R. B., Wagner, T., and Stanier, C.: Sensitivity of Meteorological Skill to Selection of WRF-Chem Physical Parameterizations and Impact on Ozone Prediction During the Lake Michigan Ozone Study (LMOS), *Journal of Geophysical Research-Atmospheres*, 125, 10.1029/2019jd031971, 2020.
- Adkins, K. A. and Sescu, A.: Observations of relative humidity in the near-wake of a wind turbine using an instrumented unmanned aerial system, *Int. J. Green Energy*, 14, 845-860, 10.1080/15435075.2017.1334661, 2017.
- Andersen, P. C., Williford, C. J., and Birks, J. W.: Miniature Personal Ozone Monitor Based on UV Absorbance, *Analytical Chemistry*, 82, 7924-7928, 10.1021/ac1013578, 2010.

- Baker, K. R., Liljegren, J., Valin, L., Judd, L. M., Szykman, J., Millet, D. B., Czarnetzki, A., Whitehill, A. R., Murphy, B. P., and Stanier, C.: Photochemical model representation of ozone and precursors during the 2017 Lake Michigan Ozone Study (LMOS), *Atmospheric Environment*, 293, 119465, <https://doi.org/10.1016/j.atmosenv.2022.119465>, 2023.
- 450 Banta, R. M., Senff, C. J., Nielsen-Gammon, J., Darby, L. S., Ryerson, T. B., Alvarez, R. J., Sandberg, S. R., Williams, E. J., and Trainer, M.: A bad air day in Houston, *Bulletin of the American Meteorological Society*, 86, 657-+, 10.1175/bams-86-5-657, 2005.
- Beekmann, M., Ancellet, G., Martin, D., Abonne, C., Duverneuil, G., Eidelman, F., Bessemoulin, P., Fritz, N., and Gizard, E.: INTERCOMPARISON OF TROPOSPHERIC OZONE PROFILES OBTAINED BY ELECTROCHEMICAL SONDES, A GROUND-BASED LIDAR AND AN AIRBORNE UV-PHOTOMETER, *Atmospheric Environment*, 29, 1027-1042, 10.1016/1352-2310(94)00336-j, 1995.
- 455 Bell, M. L., Peng, R. D., and Dominici, F.: The exposure-response curve for ozone and risk of mortality and the adequacy of current ozone regulations, *Environmental Health Perspectives*, 114, 532-536, 10.1289/ehp.8816, 2006.
- 460 Blaylock, B. K., Horel, J. D., and Crosman, E. T.: Impact of Lake Breezes on Summer Ozone Concentrations in the Salt Lake Valley, *Journal of Applied Meteorology and Climatology*, 56, 353-370, 10.1175/jamc-d-16-0216.1, 2017.
- 465 Brauner, E. V., Karotki, D. G., Frederiksen, M., Kolarik, B., Spilak, M., Andersen, Z. J., Vibenholt, A., Ellermann, T., Gunnarsen, L., and Loft, S.: Residential ozone and lung function in the elderly, *Indoor and Built Environment*, 25, 93-105, 10.1177/1420326x14539339, 2016.
- Butterworth, B. J., Desai, A. R., Metzger, S., Townsend, P. A., Schwartz, M. D., Petty, G. W., Mauder, M., Vogelmann, H., Andresen, C. G., Augustine, T. J., Bertram, T. H., Brown, W. O. J., Buban, M., Cleary, P., Durden, D. J., Florian, C. R., Iglinski, T. J., Kruger, E. L., Lantz, K., Lee, T. R., Meyers, T. P., Mineau, J. K., Olson, E. R., Oncley, S. P., Paleri, S., Pertzborn, R. A., Pettersen, C., Plummer, D. M., Riihimaki, L., Ruiz Guzman, E., Sedlar, J., Smith, E. N., Speidel, J., Wagner, T. J., Wang, Z., Wanner, L., White, L. D., Wilczak, J. M., Wright, D. B., and Zheng, T.: Connecting land-atmosphere interactions to surface heterogeneity in CHEESEHEAD19, *Bulletin of the American Meteorological Society*, <https://doi.org/10.1175/BAMS-D-19-0346.1>, 2021.
- 475 Chai, T., Kim, H. C., Lee, P., Tong, D., Pan, L., Tang, Y., Huang, J., McQueen, J., Tsidulko, M., and Stajner, I.: Evaluation of the United States National Air Quality Forecast Capability experimental real-time predictions in 2010 using Air Quality System ozone and NO<sub>2</sub> measurements, *Geoscientific Model Development*, 6, 1831-1850, 10.5194/gmd-6-1831-2013, 2013.
- 480 Chandrasekar, A., Philbrick, C. R., Doddridge, B., Clark, R., and Georgopoulos, P.: A comparison study of RAMS simulations with aircraft, wind profiler, lidar, tethered balloon and RASS data over Philadelphia during a 1999 summer episode, *Atmospheric Environment*, 37, 4973-4984, 10.1016/j.atmosenv.2003.08.030, 2003.
- Chen, L., Pang, X. B., Li, J. J., Xing, B., An, T. C., Yuan, K. B., Dai, S., Wu, Z. T., Wang, S. Q., Wang, Q., Mao, Y. P., and Chen, J. M.: Vertical profiles of O<sub>3</sub>, NO<sub>2</sub> and PM in a major fine chemical industry park in the Yangtze River Delta of China detected by a sensor package on an unmanned aerial vehicle, *Science of the Total Environment*, 845, 10.1016/j.scitotenv.2022.157113, 2022.

- 490 Chen, Q., Li, X. B., Song, R. F., Wang, H. W., Li, B., He, H. D., and Peng, Z. R.: Development and utilization of hexacopter unmanned aerial vehicle platform to characterize vertical distribution of boundary layer ozone in wintertime, *Atmos. Pollut. Res.*, 11, 1073-1083, 10.1016/j.apr.2020.04.002, 2020.
- Chen, Q., Wang, D. S., Li, X. B., Li, B., Song, R. F., He, H. D., and Peng, Z. R.: Vertical Characteristics of Winter Ozone Distribution within the Boundary Layer in Shanghai Based on Hexacopter Unmanned Aerial Vehicle Platform, *Sustainability*, 11, 10.3390/su11247026, 2019.
- 495 Chilson, P. B., Bell, T. M., Brewster, K. A., de Azevedo, G. B. H., Carr, F. H., Carson, K., Doyle, W., Fiebrich, C. A., Greene, B. R., Grimsley, J. L., Kanneganti, S. T., Martin, J., Moore, A., Palmer, R. D., Pillar-Little, E. A., Salazar-Cerreno, J. L., Segales, A. R., Weber, M. E., Yearly, M., and Droegemeier, K. K.: Moving towards a Network of Autonomous UAS Atmospheric Profiling Stations for Observations in the Earth's Lower Atmosphere: The 3D Mesonet Concept, *Sensors*, 19, 10.3390/s19122720, 2019.
- 500 Cleary, P. A., de Boer, G., Hupy, J. P., Borenstein, S., Hamilton, J., Kies, B., Lawrence, D., Pierce, R. B., Tirado, J., Voon, A., and Wagner, T. J.: Observations of the Lower Atmosphere From the 2021 WiscoDISCO campaign, *Earth System Science Data*, 14, 2129-2145, <https://doi.org/10.5194/essd-14-2129-2022>, 2022a.
- 505 Cleary, P. A., Fuhrman, N., Schulz, L., Schafer, J., Fillingham, J., Bootsma, H., McQueen, J., Tang, Y., Langel, T., McKeen, S., Williams, E. J., and Brown, S. S.: Ozone distributions over southern Lake Michigan: comparisons between ferry-based observations, shoreline-based DOAS observations and model forecasts, *Atmospheric Chemistry and Physics*, 15, 5109-5122, 10.5194/acp-15-5109-2015, 2015.
- 510 Cleary, P. A., Dickens, A. J., McIlquham, M., Sanchez, M., Geib, K., Hedberg, C., Hupy, J., Watson, M. W., Fuoco, M., Olson, E. R., Pierce, R. B., Stanier, C., Long, R., Valin, L., Conley, S., and Smith, M.: Impacts of lake breeze meteorology on ozone gradient observations along Lake Michigan Shorelines in Wisconsin, *Atmospheric Environment*, 269, <https://doi.org/10.1016/j.atmosenv.2021.118834>, 2022b.
- 515 Cook, D. E., Strong, P. A., Garrett, S. A., and Marshall, R. E.: A small unmanned aerial system (UAS) for coastal atmospheric research: preliminary results from New Zealand, *Journal of the Royal Society of New Zealand*, 43, 108-115, 10.1080/03036758.2012.695280, 2013.
- Crawford, T. L., Dobosy, R. J., McMillen, R. T., Vogel, C. A., and Hicks, B. B.: Air-surface exchange measurement in heterogeneous regions: Extending tower observations with spatial structure observed from small aircraft, *Global Change Biology*, 2, 275-285, 10.1111/j.1365-2486.1996.tb00079.x, 1996.
- 520 Crosman, E. T., Jacques, A. A., and Horel, J. D.: A novel approach for monitoring vertical profiles of boundary-layer pollutants: Utilizing routine news helicopter flights, *Atmos. Pollut. Res.*, 8, 828-835, 10.1016/j.apr.2017.01.013, 2017.
- 525 de Boer, G., Elston, J., Houston, A., Pillar-Little, E., Argrow, B., Bell, T., Chilson, P., Choate, C., Greene, B., Islam, A., Detweiler, C., Jacob, J., Natalie, V., Rhodes, M., Rico, D., Stachura, M., Lappin, F., Whyte, S., and Wilson, M.: Evaluation and Intercomparison of Small Uncrewed Aircraft Systems Used for Atmospheric Research, in preparation, *Journal of Atmospheric and Oceanic Technology*, 2021.

- 530 DeMuer, D., Heylen, R., VanLoey, M., and DeSadelaeer, G.: Photochemical ozone production in the convective mixed layer, studied with a tethered balloon sounding system, *Journal of Geophysical Research-Atmospheres*, 102, 15933-15947, 10.1029/97jd01211, 1997.
- Desjardins, R. L., Macpherson, J. I., Neumann, H., Denhartog, G., and Schuepp, P. H.: FLUX ESTIMATES OF LATENT AND SENSIBLE HEAT, CARBON-DIOXIDE, AND OZONE USING  
535 AN AIRCRAFT-TOWER COMBINATION, *Atmospheric Environment*, 29, 3147-3158, 10.1016/1352-2310(95)00007-1, 1995.
- Doak, A. G., Christiansen, M. B., D.A., H., Bertram, T. H., Carmichael, G., Cleary, P., Czarnetzki, A. C., Dickens, A. F., Janssen, M., Kenski, M., Millet, D. B., Novak, G., Pierce, B. R., Stone, E. A., Szykman, J., Vermeuel, M., Wagner, T. J., Valin, L., and Stanier, C. O.: Characterization of ground-  
540 based atmospheric pollution and meteorology sampling stations during the Lake Michigan Ozone Study 2017, *Journal of the Air & Waste Management Association*,  
<https://doi.org/10.1080/10962247.2021.1900000>, 2021.
- Dowell, D. C., Alexander, C. R., James, E. P., Weygandt, S. S., Benjamin, S. G., Manikin, G. S., Blake, B. T., Brown, J. M., Olson, J. B., Hu, M., Smirnova, T. G., Ladwig, T., Kenyon, J. S., Ahmadov, R.,  
545 Turner, D. D., Duda, J. D., and Alcott, T. I.: The High-Resolution Rapid Refresh (HRRR): An Hourly Updating Convection-Allowing Forecast Model. Part I: Motivation and System Description, *Weather and Forecasting*, 37, 1371-1395, 10.1175/waf-d-21-0151.1, 2022.
- Fuhrer, J.: Ozone impacts on vegetation, *Ozone-Science & Engineering*, 24, 69-74, 10.1080/01919510208901597, 2002.
- 550 Gaza, R. S.: Mesoscale meteorology and high ozone in the northeast United States, *Journal of Applied Meteorology*, 37, 961-977, 10.1175/1520-0450(1998)037<0961:Mmahoi>2.0.Co;2, 1998.
- Gong, W. M., Mickle, R. E., Bottenheim, J., Froude, F., Beauchamp, S., and Waugh, D.: Marine/coastal boundary layer and vertical structure of ozone observed at a coastal site in Nova Scotia during the 1996 NARSTO-CE field campaign, *Atmospheric Environment*, 34, 4139-4154, 10.1016/s1352-  
555 2310(00)00226-0, 2000.
- Greatwood, C., Richardson, T. S., Freer, J., Thomas, R. M., MacKenzie, A. R., Brownlow, R., Lowry, D., Fisher, R. E., and Nisbet, E. G.: Atmospheric Sampling on Ascension Island Using Multirotor UAVs, *Sensors*, 17, 24, 10.3390/s17061189, 2017.
- Greenberg, J. R., Guenther, A. B., and Turnipseed, A.: Tethered balloon-based soundings of ozone, aerosols, and solar radiation near Mexico City during MIRAGE-MEX, *Atmospheric Environment*, 43, 2672-2677, 10.1016/j.atmosenv.2009.02.019, 2009.
- 560 Greene, B. R., Segales, A. R., Bell, T. M., Pillar-Little, E. A., and Chilson, P. B.: Environmental and Sensor Integration Influences on Temperature Measurements by Rotary-Wing Unmanned Aircraft Systems, *Sensors*, 19, 10.3390/s19061470, 2019.
- 565 Gronoff, G., Robinson, J., Berkoff, T., Swap, R., Farris, B., Schroeder, J., Halliday, H. S., Knepp, T., Spinei, E., Carrion, W., Adcock, E. E., Johns, Z., Allen, D., and Pippin, M.: A method for quantifying near range point source induced O<sub>3</sub> titration events using Co-located Lidar and Pandora measurements, *Atmospheric Environment*, 204, 43-52, 10.1016/j.atmosenv.2019.01.052, 2019.
- Guimaras, P., Ye, J. H., Batista, C., Barbosa, R., Ribeiro, I., Medeiros, A., Zhao, T. N., Hwang, W. C.,  
570 Hung, H. M., Souza, R., and Martin, S. T.: Vertical Profiles of Atmospheric Species Concentrations and

- Nighttime Boundary Layer Structure in the Dry Season over an Urban Environment in Central Amazon Collected by an Unmanned Aerial Vehicle, *Atmosphere*, 11, 10.3390/atmos11121371, 2020.
- Hanna, S. R. and Chang, J. C.: Relations between meteorology and ozone in the Lake Michigan region, *Journal of Applied Meteorology*, 34, 670-678, 10.1175/1520-0450(1995)034<0670:rbmaoi>2.0.co;2, 1995.
- 575 Hemingway, B. L., Frazier, A. E., Elbing, B. R., and Jacob, J. D.: Vertical Sampling Scales for Atmospheric Boundary Layer Measurements from Small Unmanned Aircraft Systems (sUAS), *Atmosphere*, 8, 18, 10.3390/atmos8090176, 2017.
- Herman, J., Cede, A., Spinei, E., Mount, G., Tzortziou, M., and Abuhassan, N.: NO<sub>2</sub> column amounts from ground-based Pandora and MFDOAS spectrometers using the direct-sun DOAS technique: Intercomparisons and application to OMI validation, *Journal of Geophysical Research-Atmospheres*, 114, 10.1029/2009jd011848, 2009.
- 580 Horel, J., Crosman, E., Jacques, A., Blaylock, B., Arens, S., Long, A., Sohl, J., and Martin, R.: Summer ozone concentrations in the vicinity of the Great Salt Lake, *Atmospheric Science Letters*, 17, 480-486, 10.1002/asl.680, 2016.
- 585 Inoue, J. and Sato, K.: Toward sustainable meteorological profiling in polar regions: Case studies using an inexpensive UAS on measuring lower boundary layers with quality of radiosondes, *Environmental Research*, 205, 10.1016/j.envres.2021.112468, 2022.
- Jacob, J. D., Chilson, P. B., Houston, A. L., and Smith, S. W.: Considerations for Atmospheric Measurements with Small Unmanned Aircraft Systems, *Atmosphere*, 9, 16, 10.3390/atmos9070252, 2018.
- 590 Kaser, L., Patton, E. G., Pfister, G. G., Weinheimer, A. J., Montzka, D. D., Flocke, F., Thompson, A. M., Stauffer, R. M., and Halliday, H. S.: The effect of entrainment through atmospheric boundary layer growth on observed and modeled surface ozone in the Colorado Front Range, *Journal of Geophysical Research-Atmospheres*, 122, 6075-6093, 10.1002/2016jd026245, 2017.
- 595 Kim, J., Shusterman, A. A., Lieschke, K. J., Newman, C., and Cohen, R. C.: The BERkeley Atmospheric CO<sub>2</sub> Observation Network: field calibration and evaluation of low-cost air quality sensors, *Atmospheric Measurement Techniques*, 11, 1937-1946, 10.5194/amt-11-1937-2018, 2018.
- Kimball, S. K., Montalvo, C. J., and Mulekar, M. S.: Evaluating Temperature Measurements of the iMET-XQ, in the Field, under Varying Atmospheric Conditions, *Atmosphere*, 11, 10.3390/atmos11040335, 2020.
- 600 Knapp, K. G., Jensen, M. L., Balsley, B. B., Bognar, J. A., Oltmans, S. J., Smith, T. W., and Birks, J. W.: Vertical profiling using a complementary kite and tethered balloon platform at Ferryland Downs, Newfoundland, Canada: Observation of a dry, ozone-rich plume in the free troposphere, *Journal of Geophysical Research-Atmospheres*, 103, 13389-13397, 10.1029/97jd01831, 1998.
- 605 Koch, S. E., Fengler, M., Chilson, P. B., Elmore, K. L., Argrow, B., Andra, D. L., and Lindley, T.: On the Use of Unmanned Aircraft for Sampling Mesoscale Phenomena in the Preconvective Boundary Layer, *Journal of Atmospheric and Oceanic Technology*, 35, 2265-2288, 10.1175/jtech-d-18-0101.1, 2018.
- 610 Krechmer, J., Lopez-Hilfiker, F., Koss, A., Hutterli, M., Stoermer, C., Deming, B., Kimmel, J., Warneke, C., Holzinger, R., Jayne, J., Worsnop, D., Fuhrer, K., Gonin, M., and de Gouw, J.: Evaluation

- of a New Reagent-Ion Source and Focusing Ion-Molecule Reactor for Use in Proton-Transfer-Reaction Mass Spectrometry, *Analytical Chemistry*, 90, 12011-12018, 10.1021/acs.analchem.8b02641, 2018.
- 615 Levy, I., Makar, P. A., Sills, D., Zhang, J., Hayden, K. L., Mihele, C., Narayan, J., Moran, M. D., Sjostedt, S., and Brook, J.: Unraveling the complex local-scale flows influencing ozone patterns in the southern Great Lakes of North America, *Atmospheric Chemistry and Physics*, 10, 10895-10915, 10.5194/acp-10-10895-2010, 2010.
- 620 Li, X. B., Peng, Z. R., Lu, Q. C., Wang, D. F., Hu, X. M., Wang, D. S., Li, B., Fu, Q. Y., Xiu, G. L., and He, H. D.: Evaluation of unmanned aerial system in measuring lower tropospheric ozone and fine aerosol particles using portable monitors, *Atmospheric Environment*, 222, 10.1016/j.atmosenv.2019.117134, 2020.
- 625 Li, X. B., Wang, D. F., Lu, Q. C., Peng, Z. R., Fu, Q. Y., Hu, X. M., Huo, J. T., Xiu, G. L., Li, B., Li, C., Wang, D. S., and Wang, H. Y.: Three-dimensional analysis of ozone and PM<sub>2.5</sub> distributions obtained by observations of tethered balloon and unmanned aerial vehicle in Shanghai, China, *Stoch. Environ. Res. Risk Assess.*, 32, 1189-1203, 10.1007/s00477-018-1524-2, 2018.
- 630 Li, Y. W., Liu, B., Ye, J. H., Jia, T. J., Khuzestani, R. B., Jia, Y. S., Cheng, X., Zheng, Y., Li, X., Wu, C., Xin, J. Y., Wu, Z. H., Tomoto, M. A., McKinney, K. A., Martin, S. T., Yong, J. L., and Chen, Q.: Unmanned Aerial Vehicle Measurements of Volatile Organic Compounds over a Subtropical Forest in China and Implications for Emission Heterogeneity, *Acs Earth and Space Chemistry*, 5, 247-256, 10.1021/acsearthspacechem.0c00271, 2021.
- 635 Lu, R. and Turco, R. P.: AIR POLLUTANT TRANSPORT IN A COASTAL ENVIRONMENT .1. 2-DIMENSIONAL SIMULATIONS OF SEA-BREEZE AND MOUNTAIN EFFECTS, *Journal of the Atmospheric Sciences*, 51, 2285-2308, 10.1175/1520-0469(1994)051<2285:Aptiac>2.0.Co;2, 1994.
- 640 Lyman, S. and Tran, T.: Inversion structure and winter ozone distribution in the Uintah Basin, Utah, USA, *Atmospheric Environment*, 123, 156-165, 10.1016/j.atmosenv.2015.10.067, 2015.
- 645 Mazzuca, G. M., Pickering, K. E., Clark, R. D., Loughner, C. P., Fried, A., Zweers, D. C. S., Weinheimer, A. J., and Dickerson, R. R.: Use of tethered sonde and aircraft profiles to study the impact of mesoscale and microscale meteorology on air quality, *Atmospheric Environment*, 149, 55-69, 10.1016/j.atmosenv.2016.10.025, 2017.
- 650 McNider, R. T., Pour-Biazar, A., Doty, K., White, A., Wu, Y. L., Qin, M. M., Hu, Y. T., Odman, T., Cleary, P., Knipping, E., Dornblaser, B., Lee, P., Hain, C., and McKeen, S.: Examination of the Physical Atmosphere in the Great Lakes Region and Its Potential Impact on Air Quality Overwater Stability and Satellite Assimilation, *Journal of Applied Meteorology and Climatology*, 57, 2789-2816, 10.1175/jamc-d-17-0355.1, 2018.
- 655 Novak, G. A., Vermeuel, M. P., and Bertram, T. H.: Simultaneous detection of ozone and nitrogen dioxide by oxygen anion chemical ionization mass spectrometry: a fast-time-response sensor suitable for eddy covariance measurements, *Atmospheric Measurement Techniques*, 13, 1887-1907, 10.5194/amt-13-1887-2020, 2020.
- 660 Ollison, W. M., Crow, W., and Spicer, C. W.: Field testing of new-technology ambient air ozone monitors, *Journal of the Air & Waste Management Association*, 63, 855-863, 10.1080/10962247.2013.796898, 2013.



- Peng, Y. P., Chen, K. S., Lou, J. C., Hwang, S. W., Wang, W. C., Lai, C. H., and Tsai, M. Y.: Measurements and Mesoscale Modeling of Autumnal Vertical Ozone Profiles in Southern Taiwan, *Terrestrial Atmospheric and Oceanic Sciences*, 19, 505-514, 10.3319/tao.2008.19.5.505(a), 2008.
- 655 Sillman, S.: The relation between ozone, NO<sub>x</sub> and hydrocarbons in urban and polluted rural environments, *Atmospheric Environment*, 33, 1821-1845, 10.1016/s1352-2310(98)00345-8, 1999.
- Sills, D. M. L., Brook, J. R., Levy, I., Makar, P. A., Zhang, J., and Taylor, P. A.: Lake breezes in the southern Great Lakes region and their influence during BAQS-Met 2007, *Atmospheric Chemistry and Physics*, 11, 7955-7973, 10.5194/acp-11-7955-2011, 2011.
- 660 Stanier, C. O., Pierce, R. B., Abdi-Oskouei, M., Adelman, Z. E., Al-Saadi, J., Alwe, H. D., Bertram, T. H., Carmichael, G. R., Christiansen, M. B., Cleary, P. A., Czarnetzki, A. C., Dickens, A. F., Fuoco, M. A., Hughes, D. D., Hupy, J. P., Janz, S. J., Judd, L. M., Kenski, D., Kowalewski, M. G., Long, R. W., Millet, D. B., Novak, G., Roozitalab, B., Shaw, S. L., Stone, E. A., Szykman, J., Valin, L., Vermeuel, M., Wagner, T. J., Whitehill, A. R., and Williams, D. J.: Overview of the Lake Michigan Ozone Study
- 665 2017, *Bulletin of the American Meteorological Society*, 102, E2207-E2225, 10.1175/bams-d-20-0061.1, 2021.
- Sullivan, J. T., Berkoff, T., Gronoff, G., Knepp, T., Pippin, M., Allen, D., Twigg, L., Swap, R., Tzortziou, M., Thompson, A. M., Stauffer, R. M., Wolfe, G. M., Flynn, J., Pusede, S. E., Judd, L. M., Moore, W., Baker, B. D., Al-Saadi, J., and McGee, T. J.: The Ozone Water-Land Environmental
- 670 Transition Study: An Innovative Strategy for Understanding Chesapeake Bay Pollution Events, *Bulletin of the American Meteorological Society*, 100, 291-306, 10.1175/bams-d-18-0025.1, 2019.
- Sun, J. L., Desjardins, R., Mahrt, L., and MacPherson, I.: Transport of carbon dioxide, water vapor, and ozone by turbulence and local circulations, *Journal of Geophysical Research-Atmospheres*, 103, 25873-25885, 10.1029/98jd02439, 1998.
- 675 Tang, G. Q., Liu, Y. T., Huang, X., Wang, Y. H., Hu, B., Zhang, Y. C., Song, T., Li, X. L., Wu, S., Li, Q. H., Kang, Y. Y., Zhu, Z. Y., Wang, M., Wang, Y. M., Li, T. T., Li, X., and Wang, Y. S.: Aggravated ozone pollution in the strong free convection boundary layer, *Science of the Total Environment*, 788, 10.1016/j.scitotenv.2021.147740, 2021.
- Tang, Y. H., Lee, P., Tsidulko, M., Huang, H. C., McQueen, J. T., DiMego, G. J., Emmons, L. K.,
- 680 Pierce, R. B., Thompson, A. M., Lin, H. M., Kang, D. W., Tong, D., Yu, S. C., Mathur, R., Pleim, J. E., Otte, T. L., Pouliot, G., Young, J. O., Schere, K. L., Davidson, P. M., and Stajner, I.: The impact of chemical lateral boundary conditions on CMAQ predictions of tropospheric ozone over the continental United States, *Environmental Fluid Mechanics*, 9, 43-58, 10.1007/s10652-008-9092-5, 2009.
- Tanimoto, H., Zbinden, R. M., Thouret, V., and Nedelec, P.: Consistency of tropospheric ozone
- 685 observations made by different platforms and techniques in the global databases, *Tellus Series B-Chemical and Physical Meteorology*, 67, 10.3402/tellusb.v67.27073, 2015.
- Tarasick, D., Galbally, I. E., Cooper, O. R., Schultz, M. G., Ancellet, G., Leblanc, T., Wallington, T. J., Ziemke, J., Liu, X., Steinbacher, M., Staehelin, J., Vigouroux, C., Hannigan, J. W., Garcia, O., Foret, G., Zanis, P., Weatherhead, E., Petropavlovskikh, I., Worden, H., Osman, M., Liu, J., Chang, K. L.,
- 690 Gaudel, A., Lin, M. Y., Granados-Munoz, M., Thompson, A. M., Oltmans, S. J., Cuesta, J., Dufour, G., Thouret, V., Hassler, B., Trickl, T., and Neu, J. L.: Tropospheric Ozone Assessment Report: Tropospheric ozone from 1877 to 2016, observed levels, trends and uncertainties, *Elementa-Science of the Anthropocene*, 7, 10.1525/elementa.376, 2019.

- 695 Telg, H., Murphy, D. M., Bates, T. S., Johnson, J. E., Quinn, P. K., Giardi, F., and Gao, R. S.: A practical set of miniaturized instruments for vertical profiling of aerosol physical properties, *Aerosol Sci. Technol.*, 51, 715-723, 10.1080/02786826.2017.1296103, 2017.
- Tirado, J., Torti, A. O., Butterworth, B. J., Wangen, K., Voon, A., Kies, B., Hupy, J. P., de Boer, G., Pierce, R. B., Wagner, T. J., and Cleary, P. A.: Observations of coastal dynamics during lake breeze at a shoreline impacted by high ozone, DOI: 10.1039/D2EA00101B 2023.
- 700 Verhoelst, T., Compernelle, S., Pinardi, G., Lambert, J. C., Eskes, H. J., Eichmann, K. U., Fjaeraa, A. M., Granville, J., Niemeijer, S., Cede, A., Tiefengraber, M., Hendrick, F., Pazmino, A., Bais, A., Bazureau, A., Boersma, K. F., Bogner, K., Dehn, A., Donner, S., Elokhov, A., Gebetsberger, M., Goutail, F., de la Mora, M. G., Gruzdev, A., Gratsea, M., Hansen, G. H., Irie, H., Jepsen, N., Kanaya, Y., Karagkiozidis, D., Kivi, R., Kreher, K., Levelt, P. F., Liu, C., Muller, M., Comas, M. N., Piters, A.
- 705 J. M., Pommereau, J. P., Portafaix, T., Prados-Roman, C., Puentedura, O., Querel, R., Remmers, J., Richter, A., Rimmer, J., Cardenas, C. R., de Miguel, L. S., Sinyakov, V. P., Stremme, W., Strong, K., Van Roozendaal, M., Veeffkind, J. P., Wagner, T., Wittrock, F., Gonzalez, M. Y., and Zehner, C.: Ground-based validation of the Copernicus Sentinel-5P TROPOMI NO<sub>2</sub> measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandonia global networks, *Atmospheric Measurement*
- 710 *Techniques*, 14, 481-510, 10.5194/amt-14-481-2021, 2021.
- Vermeuel, M. P., Cleary, P. A., Desai, A. R., and Bertram, T. H.: Simultaneous Measurements of O-3 and HCOOH Vertical Fluxes Indicate Rapid In-Canopy Terpene Chemistry Enhances O-3 Removal Over Mixed Temperate Forests, *Geophysical Research Letters*, 48, 10.1029/2020gl090996, 2021.
- Vermeuel, M. P., Novak, G. A., Kilgour, D. B., Clafin, M. S., Lerner, B. M., Trowbridge, A. M., Thom,
- 715 J., Cleary, P. A., Desai, A. R., and Bertram, T. H.: Observations of biogenic volatile organic compounds over a mixed temperate forest during the summer to autumn transition, *Atmospheric Chemistry and Physics*, 23, 4123–4148, <https://doi.org/10.5194/acp-23-4123-2023>, 2023.
- Vermeuel, M. P., Novak, G. A., Alwe, H. D., Hughes, D. D., Kaleel, R., Dickens, A. F., Kenski, D., Czarnetzki, A. C., Stone, E. A., Stanier, C. O., Pierce, R. B., Millet, D. B., and Bertram, T. H.:
- 720 Sensitivity of Ozone Production to NO<sub>x</sub> and VOC Along the Lake Michigan Coastline, *Journal of Geophysical Research-Atmospheres*, 124, 10989-11006, 10.1029/2019jd030842, 2019.
- Wagner, T. J., Czarnetzki, A. C., Christiansen, M., Pierce, R. B., Stanier, C. O., Dickens, A. F., and Eloranta, E. W.: Observations of the Development and Vertical Structure of the Lake Breeze Circulation During the 2017 Lake Michigan Ozone Study, *Journal of the Atmospheric Sciences*, 79,
- 725 1005-1020, <https://doi.org/10.1175/JAS-D-20-0297.1>, 2022.
- Wainwright, C. E., Bonin, T. A., Chilson, P. B., Gibbs, J. A., Fedorovich, E., and Palmer, R. D.: Methods for Evaluating the Temperature Structure-Function Parameter Using Unmanned Aerial Systems and Large-Eddy Simulation, *Boundary-Layer Meteorology*, 155, 189-208, 10.1007/s10546-014-0001-9, 2015.
- 730 Wilson, K. L. and Birks, J. W.: Mechanism and elimination of a water vapor interference in the measurement of ozone by UV absorbance, *Environmental Science & Technology*, 40, 6361-6367, 10.1021/es052590c, 2006.
- Witte, J. C., Thompson, A. M., Smit, H. G. J., Vomel, H., Posny, F., and Stubi, R.: First Reprocessing of Southern Hemisphere ADDitional OZonesondes Profile Records: 3. Uncertainty in Ozone Profile and

- 735 Total Column, *Journal of Geophysical Research-Atmospheres*, 123, 3243-3268, 10.1002/2017jd027791, 2018.
- Wu, C., Liu, B., Wu, D., Yang, H. L., Mao, X., Tan, J., Liang, Y., Sun, J. Y., Xia, R., Sun, J. R., He, G. W., Li, M., Deng, T., Zhou, Z., and Li, Y. J.: Vertical profiling of black carbon and ozone using a multicopter unmanned aerial vehicle (UAV) in urban Shenzhen of South China, *Science of the Total Environment*, 801, 10.1016/j.scitotenv.2021.149689, 2021.
- 740 Wu, Y. L., Lin, C. H., Lai, C. H., Lai, H. C., and Young, C. Y.: Effects of Local Circulations, Turbulent Internal Boundary Layers, and Elevated Industrial Plumes on Coastal Ozone Pollution in the Downwind Kaohsiung Urban-Industrial Complex, *Terrestrial Atmospheric and Oceanic Sciences*, 21, 343-357, 10.3319/tao.2009.04.14.01(a), 2010.
- 745 Xu, Z. N., Huang, X., Nie, W., Shen, Y. C., Zheng, L. F., Xie, Y. N., Wang, T. Y., Ding, K., Liu, L. X., Zhou, D. R., Qi, X. M., and Ding, A. J.: Impact of Biomass Burning and Vertical Mixing of Residual-Layer Aged Plumes on Ozone in the Yangtze River Delta, China: A Tethered-Balloon Measurement and Modeling Study of a Multiday Ozone Episode, *Journal of Geophysical Research-Atmospheres*, 123, 11786-11803, 10.1029/2018jd028994, 2018.
- 750 Ye, J., Batista, C. E., Zhao, T., Campos, J., Ma, Y., Guimarães, P., Ribeiro, I. O., Medeiros, A. S. S., Stewart, M. P., Vilà-Guerau de Arellano, J., Guenther, A. B., Souza, R. A. F. d., and Martin, S. T.: River Winds and Transport of Forest Volatiles in the Amazonian Riparian Ecoregion, *Environmental Science & Technology*, 56, 12667-12677, 10.1021/acs.est.1c08460, 2022.
- Zhang, J., Ninneman, M., Joseph, E., Schwab, M. J., Shrestha, B., and Schwab, J. J.: Mobile Laboratory 755 Measurements of High Surface Ozone Levels and Spatial Heterogeneity During LISTOS 2018: Evidence for Sea Breeze Influence, *Journal of Geophysical Research-Atmospheres*, 125, 10.1029/2019jd031961, 2020.
- Zhang, K., Zhou, L., Fu, Q. Y., Yan, L., Bian, Q. G., Wang, D. F., and Xiu, G. L.: Vertical distribution of ozone over Shanghai during late spring: A balloon-borne observation, *Atmospheric Environment*, 760 208, 48-60, 10.1016/j.atmosenv.2019.03.011, 2019.