



Development and Evaluation of an Improved Off-Line Aerosol Mass Spectrometry Technique

Christina N. Vasilakopoulou^{1,2}, Kalliopi Florou², Christos Kaltsonoudis², Iasonas Stavroulas³, Nikolaos Mihalopoulos³ and Spyros N. Pandis^{1,2}

5 ¹Department of Chemical Engineering, University of Patras, Patras, Greece

²Institute of Chemical Engineering Sciences, ICE-HT, Patras, Greece

³Institute for Environmental Research and Sustainable Development, National Observatory of Athens, Athens, Greece

Correspondence to: Spyros N. Pandis (spyros@chemeng.upatras.gr)

Abstract. The off-line Aerosol Mass Spectrometry (AMS) technique is a useful tool for the source apportionment of organic aerosol (OA) in areas and periods during which an AMS is not available. However, the technique is based on the extraction of aerosol samples in water, while several atmospheric OA components are partially or fully insoluble in water. In this work an improved off-line technique was developed and evaluated in an effort to capture most of the partially soluble and insoluble organic aerosol material, reducing significantly the uncertainty of the corresponding source apportionment. A major advantage of the proposed approach is that no corrections are needed for the off-line analysis to account for the limited water solubility of some OA components. The improved off-line AMS analysis was tested in three campaigns: two during winter and one during summer. Collocated on-line AMS measurements were performed for the evaluation of the off-line method. Source apportionment analysis was performed separately for the on-line and the off-line measurements using Positive Matrix Factorization (PMF). The PMF results showed that the fractional contribution of each factor to the total OA differed between the on-line and the off-line PMF results by less than 15%. The differences in the AMS spectra of the factors of the two approaches could be significant suggesting that the use of factor profiles from the literature in the off-line analysis may lead to complications. Part of the good agreement between the on-line and the off-line PMF results is due to the ability of the improved off-line AMS technique to capture a bigger part of the OA, including insoluble organic material. This was evident by the significant fraction of submicrometer suspended insoluble particles present in the water extract, and by the reduced insoluble material on the filters after the extraction process. More than half of the elemental carbon (EC) was on average missing from the filters after the water extraction. Significant EC concentrations were measured in the produced aerosol that was used as input to the AMS during the off-line analysis.

10
15
20
25

1 Introduction

The High-Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS) has been used during the last fifteen years for the source apportionment of organic aerosol (OA) in field studies (DeCarlo et al., 2008; Sun et al., 2011). The corresponding results have dramatically improved our understanding of particulate matter and especially of organic aerosol. AMS is one of

30



the few instruments which can provide information for the size distribution of aerosols, their concentration, and their chemical composition automatically and in high temporal resolution (Drewnick et al., 2005; Jayne et al., 2000).

The AMS measurements can also be combined with source apportionment algorithms, such as Positive Matrix Factorization (PMF) (Paatero and Tapper, 1994) to estimate the contribution of the various sources of organic aerosol (OA).

35 The advantage of PMF compared to other receptor models is that the solutions are constrained to be non-negative, which makes it suitable for environmental applications. The multilinear engine algorithm (ME-2) can be also used in cases that PMF results in non-meaningful solutions or to quantify smaller contributions of sources (Paatero, 1999). The difference between PMF and ME-2, is that in the latter, the user can use a-priori information about the factor profiles as input to the algorithm forcing it to account for the specific source.

40 Even though the AMS has been used in many field campaigns around the world (Aiken et al., 2009; Saarikoski et al., 2012; Setyan et al., 2012) there are circumstances in which its use is impractical. Its weight and size make sometimes its move to the field challenging or even impossible for some sites. Also, its high cost makes its use in multiple locations in the same city or region impossible. The off-line AMS technique, developed by Daellenbach et al. (2016) is a possible solution to both these problems. In the off-line analysis the ambient particulate matter is collected in filters. Then the filter samples are extracted
45 in ultrapure water, and the water extract is filtered, aerosolized, dried, and finally measured with the AMS in the laboratory. Even though this technique provides valuable information about the OA, it has been designed to capture the water-soluble part of the measured OA. The ME-2 was proposed by Daellenbach et al. (2016) for the source apportionment of the corresponding off-line measurements. Comparisons between the on-line and the off-line measurements by Daellenbach et al. (2016) suggested the need for significant corrections for hydrocarbon-like OA (HOA), cooking OA (COA), and even biomass burning OA
50 (BBOA) to account for the partial recovery of the components. Many studies have used the corrections proposed by Daellenbach et al. (2016) for OA to correct the off-line results for this partial recovery (Bozzetti et al., 2017; Vlachou et al., 2018; Srivastava et al., 2021). However, these corrections may introduce significant uncertainties in the approach (Ge et al., 2017). They may be location specific (Xu et al., 2017) or they may be sensitive to small variations in the experimental method. Some studies do not use these correction factors (Sun et al., 2011; Mihara and Mohinda, 2011; Xu et al., 2015; Ye et al., 2017;
55 Li et al., 2021) and assume that the off-line results correspond only to the water-soluble part and not the total OA.

The application of the off-line technique may be limited by the OA mass loading of the filter. Niedeck et al. (2022) developed a technique in order to reduce the detection limits of the off-line AMS measurements. Micronebulization was performed with only 10 μ L of liquid used for the extraction. Organics and inorganics were quantitated by an isotopically labelled internal standard.

60 A significant difference between the on-line and the off-line measurements is the reduced temporal resolution of the latter. Typically, the on-line measurements are conducted every few minutes, in contrast to the filter collection which is in most cases performed daily. The effect of the reduced temporal resolution to the PMF results was addressed by Vasilakopoulou et al. (2022), who analyzed with PMF a five-month period in high (30 min) and low (daily) temporal resolution. The results



65 showed that the average contribution of each source to the total OA differed by less than 15% between the high and the low-resolution analysis. However, significant differences were observed in the factor spectra.

In this study we aim to improve the original off-line experimental procedure of Daellenbach et al. (2016) in order to capture a higher fraction of the OA and reduce significantly the uncertainty of the PMF results. The proposed improved off-line AMS technique is evaluated on three field campaigns in Greece, two during winter and one in summer. Comparisons with the on-line source apportionment results are used to evaluate the accuracy of the off-line source apportionment solution.

70 2 Methodology

2.1 Experimental

The proposed off-line method is based on the work of Daellenbach et al. (2016) but with changes in the extraction and atomization steps in an effort to characterise most of the OA. Particulate matter is collected daily in quartz filters by a high-volume sampler (Tisch Scientific). Filter punches (1.5 cm²) are extracted in 20 mL of ultrapure water, with the help of an ultrasonic sonicator (Elmasonic S80). The extracts are placed in a syringe pump working at a flow rate of 15 mL h⁻¹. The water sample is not filtered in an effort to analyse a higher fraction of the OA and to limit the losses of organic aerosol material. This is one of the major features of our proposed approach. The extract is then atomized using an atomizer (TSI, 3075) and the produced droplets are dried using a silica gel dryer. The resulting aerosol is characterized by a HR-ToF-AMS (Aerodyne Inc.). The AMS measures particles smaller than 1 µm. If there are larger particles (e.g. fragments of the quartz filter) in the produced aerosol these do not make it through the AMS aerodynamic lenses.

80 On average around 40 min of off-line measurements are performed with the AMS for each collected sample. The temporal resolution of the off-line AMS measurements was 3 min. Before and after each measurement, a blank measurement is conducted. Ultrapure water is atomized, and the produced droplets are dried and measured by the AMS, similarly with the actual sample, for around 30 minutes. The average of the two blank measurements is subtracted from the sample value.

85 The proposed technique was evaluated in three different campaigns in Greece. The number of samples used in each period was above 30. In two of the campaigns (winter 2020 and early summer 2019 in Patras) an HR-AMS was used for the on-line measurements and in the third (winter 2019 in Athens) an Aerosol Mass Speciation Monitor (ACSM) (Aerodyne Inc., USA) was used.

2.2 On-line source apportionment

90 The PMF analysis of the on-line measurements (on-line PMF from now on) was performed using SoFi Pro in resolution of 3 min for the AMS and 30 min for the ACSM. No a-priori information about the factor profiles was used. OA high resolution (HR) mass spectra were analyzed (m/z 12-300) (Canonaco et al., 2013) for the case of the AMS datasets. The “weak” signals were downweighted by a factor of 2 and the “bad” by a factor of 10. Also, the variables related to CO₂ (16, 17, 18, 28 and 44)



were downweighted by a factor of 2. The rotational ambiguity of the solution was explored using the F_{peak} . The minimum F_{peak}
95 value was -1, the maximum 1 and the step was 0.1.

For the Thissio campaign in Athens the unit mass resolution (UMR) spectra (m/z 12-125) were used, and they were pretreated in accordance to the procedure described above. Because of the low temporal resolution of the measurements (30 min) high concentration events were not removed.

2.3 Off-line source apportionment

100 The off-line PMF was also performed using SoFi Pro, without utilization of any a-priori information about the factor profiles. The high resolution MS data were used for the off-line PMF in all three datasets. Each sampling day was represented by an off-line spectrum that was the arithmetic average of the 40 min off-line AMS data. The error for each sample was represented by the arithmetic average of the error of the off-line measurements. The measured AMS spectrum was corrected based on the blank measurements. The blank concentration at each m/z was subtracted from the sample. The error matrix of the AMS
105 spectrum was also blank corrected using the same procedure.

The same downweighting of weak and bad signals, and the same F_{peak} approach as in the on-line measurements was used.

3 Field Campaigns

The improved off-line AMS technique was applied to more than 100 ambient daily filter samples from two urban areas in
110 Greece and during different periods of the year. Continuous online AMS measurements were also performed, and they were used for the evaluation of the approach. In this work we will focus on the results from two winter periods (Athens and Patra) and one summer period in Greece (Patra). The measurements from each campaign are analysed separately in the following sections. For the two campaigns performed in Patras (winter 2020 and summer 2019) an HR-AMS, with 3 min temporal resolution was used, while in the Athens campaign an ACSM was used. The temporal resolution of the ACSM was 30 min.

115 3.1 Winter campaign in Patras, 2020

Patras is the third biggest city of Greece, with almost 200 thousand inhabitants. The winter campaign was conducted at the University of the Peloponnese campus during January-February 2020. This urban background site is located around 4 km from the center of Patras. The main goal of the campaign was to study the biomass burning emissions and especially residential wood burning.

120 Both ambient measurements and mobile smog chamber experiments were conducted (Jorga et al., 2021) but in this study we focus on the ambient measurements. Together with several other instruments the HR-ToF-AMS was used to characterize the non-refractory PM_{10} concentration and composition continuously for one month. The V-mode was used, and the temporal resolution of the measurements was 3 min. The vaporizer temperature was 600°C and no drying of the sample



125 was performed. Collocated filter samples were also collected during the same period. The sampling started at 18:00 LT each day and lasted for 24 h. The black carbon (BC) mass concentration was measured using a Multi Angle Absorption Photometer (MAAP).

High PM_{10} concentrations were measured during the late afternoon and night hours when biomass burning for heating purposes was taking place. In some periods the PM_{10} mass concentration exceeded $100 \mu\text{g m}^{-3}$ (Fig. 1) with the organics accounting for around 70% of the total. Black carbon levels were also high during these periods exceeding $10 \mu\text{g m}^{-3}$.

130 3.2 Summer campaign in Patras, 2019

The summer campaign in Patras was conducted in the Institute of Chemical Engineering Sciences (ICE-HT), which is a suburb of Patras, 8 km away from the city centre. The area is surrounded by olive tree fields and there is limited anthropogenic activity within a radius of 1 km.

135 The campaign lasted from March until June 2019. On-line AMS measurements took place during a few days every week, and mostly during weekends, due to the use of the AMS in other laboratory experiments during that period. Measurements were performed during 10 days per month on average. Filter samples were collected during the same days. Low PM_{10} concentrations were observed during this measurement period. The aerosol levels were on average $4 \mu\text{g m}^{-3}$. Among the four months, March was the one with the highest average OA concentration ($6.9 \mu\text{g m}^{-3}$), and May had the lowest ($2 \mu\text{g m}^{-3}$) (Fig. 2).

140 3.3 Winter campaign in Athens, 2019

Athens is the biggest city of Greece, with a population of 3.2 million inhabitants. The winter campaign was performed at the National Observatory of Athens at Thessio, in the centre of the city. The measurements started in January and lasted until March 2019. The OA in this case was measured using an Aerosol Mass Speciation Monitor (ACSM) (Aerodyne Inc., USA) with a temporal resolution of 30 min.

145 The OA was on average $8.2 \mu\text{g m}^{-3}$ during the examined period. The highest average concentration ($10.8 \mu\text{g m}^{-3}$) was observed on February, and the lowest ($4.4 \mu\text{g m}^{-3}$) on January (Fig. 3). High concentration events were observed during the whole period late at night. The maximum half-hour OA concentration ($98 \mu\text{g m}^{-3}$) was observed on February 19 at midnight. The OA concentration started to increase from 22:00 LT on February 18 and remained high until 4:00 LT that night.



4 Source apportionment results

150 4.1 Source apportionment of the Patras 2020 winter campaign

4.1.1 On-line

The one- up to six-factor solutions were explored, and the five-factor solution was chosen as the one that could explain satisfactorily the OA sources. Biomass burning (BBOA) was the dominant source of the OA (Fig. 1). Two biomass burning factors were identified (responsible for 53% of the total OA), a cooking OA (COA) factor (12%), a hydrocarbon-like OA
155 (HOA) (10%) and an oxidized OA (OOA) factor (25%) (Fig. 1).

BBOA I and BBOA II had similar time series ($R^2=0.91$), but different spectra. BBOA I had strong peaks at m/z values 43, 60 and 73 (Ng et al., 2011). On the other hand, BBOA II was characterized by m/z 's 44 and 60 (Figs. S1-S2). The mass concentration of both BBOA I and BBOA II increased from 17:00 LT and remained at high levels until 1:00 LT (Fig. S3) which is consistent with the hours that fireplaces and wood stoves are used in the area. The highest BBOA concentration of
160 the day was observed around 20:00 LT, when BBOA was almost $40 \mu\text{g m}^{-3}$ on average.

The COA concentration peaked at the late evening hours (17:00- 23:00 LT) and had a weaker peak around midday (10:00-15:00 LT) (Fig. S3). The highest COA concentration was observed at 21:00 LT, which is the common wintertime dinner time in Greece, and was about $5 \mu\text{g m}^{-3}$ on average. The COA spectrum was characterized by peaks at m/z values 41, 43, 55, 57 consistent with the results of Kaltsonoudis et al. (2017). The m/z 55 signal was almost twice that at m/z 57, a feature
165 related to charbroiling in previous studies (Lanz et al., 2007; Allan et al., 2010).

The HOA concentration increased at 9:00 LT which is the local morning rush hour and during the evening at 18:00 LT. The HOA spectrum was characterized by the m/z values 41, 43, 55, 57, 67, 69 and 81 which are hydrocarbon fragments of typical traffic-related emissions (Aiken et al., 2009).

The OOA was characterized by a strong signal at m/z 44 (mostly due to CO_2^+) and another one at m/z 28, which is
170 also reported in previous studies (Florou et al., 2017). The average diurnal pattern of OOA showed less variation compared to the primary factors. A slight increase of OOA was present during 10:00-14:00 LT, possibly due to the higher photochemical activity during that period.

4.1.2 Off-line

During the 2020 winter campaign in Patras, 31 ambient aerosol samples were collected on quartz filters using a medium-
175 volume sampler (Tisch Scientific, 220 L min^{-1} flowrate). The sampling started at 18:00 (LT) each day and lasted 24 h. The experimental procedure described in Section 2.1 was used for the off-line AMS analysis, and the off-line PMF was conducted according to the procedure described in Section 2.3.

The results of the PMF analysis for the off-line AMS method showed similar fractional contributions with the on-line (differences below 15%) (Fig. 4). The off-line BBOA was responsible for 48% of the total OA (Fig. 4). The second highest
180 contributor to the total OA was COA (19%) and then HOA and OOA were responsible for 16% and 17% of the total OA



respectively. These results are quite encouraging showing that our new off-line approach can reproduce the on-line results without the significant corrections (e.g. a factor of 10 for the HOA) needed in the original method.

The only minor weakness of the above off-line analysis results is the different split of the two BBOA factors. The on-line PMF analysis indicated that BBOA I was 22% and BBOA II 31% of the total OA, while the off-line PMF analysis resulted in 32% for BBOA I and 16% for BBOA II (Fig. 4). However, the difference between the on-line and the off-line contribution to the total BBOA is below 5%, which shows that the off-line analysis can still provide an estimate of the importance of BBOA on the total OA that is consistent with the on-line results. This different split in the BBOA factors is probably of secondary importance given our limited understanding of the differences of BBOA I and BBOA II.

The uncertainty of the off-line PMF results was estimated using bootstrap analysis. The results showed that BBOA I varied the most among the five factors. The estimated average contribution of BBOA I to the total OA was 32%, and ranged between 18% and 47%. All the other factor contributions to the total OA varied less than 20%. The BBOA II contribution to the total OA was 16% and varied between 11% and 21%. The total BBOA estimated contribution to the total OA was 48%, and ranged from 34% to 67%. COA was on average 19% of the total OA and ranged between 11% and 28%. The HOA contribution to the total OA was 16%, ranging from 12% to 21%, and OOA was on average 17%, ranging from 11% to 23% (Fig. S4).

Even though the average contribution of each factor to the total OA showed good agreement between the on-line and the off-line PMF results, higher discrepancies were observed on a daily basis (Fig. S5). For example, the off-line COA showed two different behaviors during the period, one of overestimation and one of underestimation compared to the on-line COA. This is evidence that the off-line AMS technique proposed in this work can give relatively accurate (within 15%) contributions of each factor for big periods, but on a daily basis the uncertainty can be really high. This is due to a large extent to the temporal resolution of the off-line measurements (Vasilakopoulou et al., 2022).

Even though the factor contributions of the off-line and the on-line PMF results differed by less than 15%, the factor profiles were significantly different. The off-line PMF spectra were compared with the on-line, using the theta angle approach (Kostenidou et al., 2009). Theta is the angle that the two factors have if they are treated as vectors. The highest difference was observed for HOA ($\theta=40^\circ$) (Fig. S6), showing that the off-line HOA spectrum is quite different from the on-line. This is not surprising given the limited water solubility of the HOA components. However, the HOA tracers (m/z values 41, 43, 55, 57 and 69) were present in the off-line HOA spectrum making its identification possible. These differences are probably due to the partial inclusion of the HOA components in the aerosol that makes it to the AMS. Therefore, the extraction process appears to modify the AMS spectrum, but still the PMF reaches a reasonably accurate estimate, at least in this case, with the “processed” AMS spectrum. Also, high theta angles were observed for the other primary factors ($\theta>25^\circ$). The off-line COA spectrum differed by 34° from the on-line one, the off-line BBOA I by 27° , and the off-line BBOA II by 36° from the on-line BBOA II. The OOA had the most similar spectrum profile between the off-line and the on-line results ($\theta=17^\circ$).

These differences in the spectra profiles are a consequence of two facts. First, the lower temporal resolution that the off-line results have (24 h) compared to the on-line (3 min) can affect the profiles of each factor leading to high theta angles



215 that can reach up to 30° (Vasilakopoulou et al., 2022). Secondly, the extraction efficiency of the different OA components can also result in increased angles.

In order to understand better the differences between the off-line and the on-line PMF analyses, we compared the total organic mass spectra from the two techniques. The off-line organic mass spectra were blank corrected. The theta angle between the average off-line and the average on-line organic mass spectrum was 21° (Fig. 5). Differences were observed at
220 m/z values below 20, between the on-line and the off-line spectra. If these values are excluded, the theta angle drops to 17°, and the two spectra appear more similar to each other. Comparing the signals at four major HOA fragments (m/z 55, 57, 67, 69) we estimated that on average the off-line technique captured 64% of the $C_xH_y^+$ and 82% of the $C_xH_yO^+$ (Fig. 6). The average off-line to on-line ratio for these HOA markers was 0.73 (Fig. S7).

A day-by-day analysis has also been performed between the on-line (24 h averages) and the off-line samples. The
225 organic mass spectra comparison between the off-line and the on-line results showed that the best agreement was observed on January 20 and their angle was 16°. The worst agreement was observed on January 30 and the theta angle was 26° (Fig. S8).

The atomic oxygen to carbon ratio (O:C) of the off-line spectra was on average 0.47 and the on-line was 0.50, showing good agreement between the two for the whole period (Fig. S9). The fractional error of the off-line compared to the on-line O:C was 12%.

230 The organics to sulfate ratio (OA/Sulf) was also compared between the on-line and the off-line results. The off-line OA/Sulf ratio followed the same trend over time of the on-line ratio (Fig. 7). The two ratios agreed within 30%. The off-line results showed a surprisingly small overestimation of the OA/Sulf ratio compared to the on-line with a bias of 1.05 and a fractional bias of 0.18. The average off-line OA/Sulf ratio was 8.0, while the on-line was 7.0.

4.2 Source apportionment of the Patras 2019 summer campaign

235 4.2.1 On-line

Solutions with one up to four factors were examined and the three-factor solution was chosen as the one which explains better the ambient OA spectral variation. One primary (HOA) and two secondary (MO-OOA, LO-OOA) factors were identified. The OA was highly oxidized during the examined period, as the secondary factors accounted for 80% of the total OA (LO-OOA 48% and MO-OOA 32%). The estimated average HOA contribution to the total OA was 20% (Fig. 2).

240 The HOA spectrum was characterized by m/z 's 41, 43, 55, 57, 67, 69 and 81 (Fig. S10). Its diurnal profile had one peak at 10:00 LT and one during the evening at 22:00 LT (Fig. S11). The HOA O:C was 0.22, which is on the high end of the ranges reported in the literature. Since HOA and COA have relatively similar AMS spectra, distinguishing between the two is not always possible. The HOA of this study agrees well with the HOA-2 reported in Kostenidou et al. (2015), which also included meat cooking emissions.

245 MO-OOA was the most oxidized of the two secondary factors. Its diurnal profile had small variation, showing that it was not affected by local sources. On the other hand, the LO-OOA diurnal profile was more variable and was characterized



250 by a first peak at 10:00 LT and a second one during the evening (22:00 LT). Both the MO-OOA and LO-OOA mass spectra were characterized by a strong peak at m/z 44. However, the LO-OOA spectrum had also peaks at m/z values, 39, 41 and 55. These peaks were weak in the MO-OOA spectrum. Also, the MO-OOA O:C was a little higher (1.0) than the O:C of the LO-OOA (0.94).

4.2.2 Off-line

We focus here on the analysis of the 31 daily samples collected during the early summer period (March to June 2020) in Patras. A low-volume sampler (6.7 L min^{-1}) was used for the sample collection on quartz filters and the sampling period was 24 hours. Because of the low mass loading of the filter half of the initial filter (47 mm) was used in this off-line AMS analysis.

255 The off-line PMF analysis resulted in a three-factor solution with one primary (HOA) and two secondary (MO-OOA and LO-OOA) factors. The off-line results showed that the OA was highly oxidized, which is in agreement with the on-line results. OOA was responsible for 68% of the total OA, while HOA represented 32% of the total OA, which is 12% more than in the on-line solution. The contribution of each factor to the total OA differed by less than 17% with the on-line estimation (Fig. 8). The off-line HOA contribution to the total OA was 32% and the bootstrap analysis showed that it ranged from 23%
260 to 42% (5th and 95th percentiles). The MO-OOA was 37%, ranging from 24% to 57%, and the LO-OOA was 31% (ranging from 26% to 35%) (Fig. S12). These uncertainties suggest that the estimates of the two approaches were practically in agreement with each other considering their uncertainty.

The off-line PMF spectra were characterized by the specific markers of each factor. The off-line HOA spectrum was characterized by peaks related to aliphatic hydrocarbons i.e., 41, 43, 55, 57, 67 and 69 (Fig. S13). The MO-OOA spectrum
265 was characterized by the m/z 44 peak, and the LO-OOA by the m/z peaks at 43 and 44. The different m/z 43/44 ratio among the two secondary factors was used for their identification. The off-line spectrum of each factor was different compared with the on-line spectrum. The off-line HOA spectrum differed by 34° from the on-line HOA (Fig. S14). A high theta angle was also observed for the LO-OOA, as the off-line LO-OOA differed by 34° with the on-line LO-OOA. The MO-OOA spectra were more similar, as the off-line one differed by 12° with the on-line MO-OOA spectrum.

270 The average total off-line to on-line spectra comparisons in the summer period, showed better agreement compared with the winter campaign. The total OA spectra for the off-line and the on-line results had an average theta angle of 12° (Fig. S15). The best agreement was observed on May 8, when the two spectra differed by 7° . The worst was observed on May 18 and the angle between the off-line and the on-line OA spectra was 16° .

275 The off-line technique captured a big part of the HOA on-line signal as indicated by the comparison of five HOA markers (Fig. 9). For these HOA markers the average $\text{C}_x\text{H}_y\text{O}^+$ off-line to on-line ratio was 0.77, and the average C_xH_y^+ off-line to on-line ratio was 1.1 (Fig. 9). This good agreement between the off-line and the on-line HOA markers could be responsible for the lower theta angle between the off-line and the on-line HOA spectrum in the PMF analysis, compared to the winter campaign.



4.3 Source apportionment of the Thissio 2019 winter campaign

280 4.3.1 On-line

Five factors were identified, three primary (HOA, COA, BBOA) and two secondary (MO-OOA, LO-OOA). The primary factors represented on average 53% of the total OA, and the secondary 47%. The contribution of the secondary factors to the total OA was relatively high for a winter period in the center of a big city. The bootstrap analysis suggested low uncertainty for the contribution of each factor to the total OA (below 15%) (Fig. S16).

285 The BBOA (11% of the total OA) had high concentrations during January and February and decreased in March (Figs. S17-S18), a behavior that is consistent with the temperatures in Greece during the winter period. The highest hourly BBOA mass concentration ($26 \mu\text{g m}^{-3}$) was observed on February 19 at midnight, which was also the night with the highest OA mass concentration. The same night the HOA mass concentration was the highest of the three-month period ($49 \mu\text{g m}^{-3}$). On average the HOA mass concentration was $2.1 \mu\text{g m}^{-3}$ and it represented 26% of the total OA. On the other hand, COA
290 (16% of the total OA) remained at the same levels during the three-month period examined. The LO-OOA (21% of the total OA) and the MO-OOA (26% of the total OA) were less variant during the three-month period compared to the primary factors.

4.3.2 Off-line

In this campaign 33 daily samples were collected in Thissio, at the National Observatory of Athens. A high-volume sampler (DH-77, Digital) was used for the sample collection. One of the additional objectives of this test was to study the difference
295 that may occur between HR-AMS off-line measurements and unit mass resolution ACSM online results.

The average off-line mass spectrum was similar to the on-line ($\theta=16^\circ$) (Fig. S20) for the examined days. The off-line PMF solution resulted in five factors, just like the on-line solution. Three primary (HOA, COA and BBOA) and two secondary (MO-OOA, LO-OOA) factors were identified. Each factor was identified using the specific tracer m/z 's that have been already discussed in the previous sections. The off-line spectrum of each factor was different compared to the one in the on-line PMF
300 solution (Fig. S21). The difference in the estimated spectra are partially due to the different sampling temporal resolution (30 min in the on-line, daily in the off-line) (Vasilakopoulou et al., 2022) and could also be partially due to the different instrumentation used for the on-line and the off-line analysis. In the off-line PMF results of this study, the two secondary factors start to mix at low temporal resolution PMF results, as the O:C of the off-line MO-OOA is lower compared to the on-line, while the O:C of the LO-OOA is higher. This is also characteristic of the reduced temporal resolution analysis
305 (Vasilakopoulou et al., 2022).

The difference between the on-line and the off-line contributions to the total OA were below 15% (Fig. 10), which is evidence of consistency between the off-line HR-AMS analysis with the on-line ACSM results. The highest difference was observed for BBOA, which was 13% higher (24% vs 11%) than the on-line BBOA, and the lowest was observed for COA, which was 1% lower than the on-line COA (15% vs 16%). The off-line HOA was 8% lower (18% vs 26%). The off-line LO-



310 OOA contribution to the total OA was only 4% higher (25% vs 21%) than the on-line LO-OA. The off-line MO-OOA contribution to the total OA differed by 9% (17% vs 26%) from the on-line MO-OOA.

In order to estimate the uncertainty of the off-line PMF results a bootstrap analysis of 1000 runs has been performed. The mean BBOA contribution to the total OA was 24% and ranged from 14% to 36% (Fig. S22). The HOA contribution to the total OA ranged from 14% to 23% (the mean value was 18%). The estimated average COA contribution to the total OA
315 was 15%, and it ranged from 9% to 21%. The MO-OOA ranged from 15% to 20% (mean value 18%) and LO-OOA ranged from 16% to 37% (mean value 25%).

A sensitivity analysis has also been performed, in which the off-line high-resolution AMS spectra were averaged to unit resolution and were used as inputs to the PMF. The UMR PMF analysis resulted in more mixed profiles than the HR analysis and the factors were not identified as easily as in the HR analysis. The contribution of each factor to the total OA
320 between the HR and the UMR off-line PMF results differed by less than 20% (Suppl. Mat. Section S4). The highest difference was observed for BBOA and was 17% (24% for the HR analysis and 7% for the UMR analysis). Comparing the UMR off-line solution with the on-line solution the highest difference observed was 14% and it referred to the COA. The UMR off-line COA was 21% of the total OA, while the on-line COA was 17%.

5 Transfer of OA from the filters to the AMS

325 Our results so far have indicated good agreement between the off-line and collocated on-line AMS measurements. This is a rather surprising result, because a significant fraction of the organics analysed (components of cooking aerosol or components of the HOA coming from transportation) are practically insoluble in water. Our hypothesis is that during the extraction process, the water-soluble material of the particles dissolves, the remaining small water-insoluble particle cores leave the filter and get suspended in the water during the sonication phase and then are transferred with the solution to the atomizer. After they enter
330 the atomizer, they are included in the formed water droplets together with the dissolved material and become part of the produced aerosol after the drying. In this way they are measured by the AMS. The different aspects of this hypothesis are tested in the following sections.

5.1 Measurements of suspended particles in the liquid phase

In order to test our hypothesis, we first quantified the concentration of the particles in the aqueous phase after the extraction and the filtration phase. If our hypothesis is not valid, there should be no particles suspended in the solution, just the background. We have used a Zeta-Sizer (Malvern Panalytical) that can measure the size distribution of nanoparticles in liquids using dynamic light scattering. The particle size distribution was measured in the ultrapure water used for the extraction and in the solution produced by the extraction of ambient aerosol samples. The Zeta-Sizer was calibrated using monodisperse polystyrene spheres (PSL) of 100 nm and 200 nm (Suppl. Mat. Section S5).
335



340 The method was applied to 20 ambient samples, which were collected during winter at the University of Peloponnese campus, in Patra. The results support our hypothesis as significant concentrations of particles of sizes in the 100-200 nm size range were detected in the extract of the ambient aerosol samples (Fig. 11). These particles were not present in the blanks.

5.2 Insoluble material removed from the filter after the extraction process

345 Since the Zeta-Sizer results showed that particles were detected in the water extracts, the concentration of insoluble material on the filter after the extraction process, should be reduced compared to the initial concentration.

To test this, we used a Sunset OC/EC analyzer for 20 ambient filter samples. In these additional experiments the EC was measured before and after the extraction. The filters after the extraction were dried at 100°C for 5 min before the EC measurement, because the humidity of the sample increases significantly the uncertainty of the EC measurement. The ambient EC concentration for the examined period was on average 0.7 $\mu\text{g m}^{-3}$ and ranged from 0.2 to 1.4 $\mu\text{g m}^{-3}$. A significant fraction
350 of EC was missing from the filter after the extraction process. The average EC on the filter after the extraction was 44% of the initial (Fig. 12).

5.3 BC measurements using the off-line technique

To test if the insoluble material was transferred to the AMS, we measured the BC in the resulting particles with a Single Particle Soot Photometer-Extended Range (SP2-XR) (Droplet Measurement Technologies). Two filter punches (3 cm² in total)
355 from ambient samples were extracted in 10 mL of ultrapure water, following the off-line procedure. The samples were placed in an ultrasonic sonicator for 30 min and no filtration phase was performed. The water extracts were then atomized and dried, as in the off-line AMS procedure, and then the resulting aerosol was detected with the SP2-XR. Significant BC concentrations were detected in the samples (Fig. 13), while in the blank (atomized water) the BC was practically zero. These tests strongly support our hypothesis that a significant fraction of even the most insoluble material (elemental carbon) in the ambient
360 particles, leaves the filter, gets suspended in the aqueous solution as particles in the 50-300 nm size range and makes it to the AMS.

6 Conclusions

The off-line AMS technique can be a powerful tool for characterizing OA in areas and periods when an AMS is not available. However, so far it could capture only the water-soluble part of OA if no corrections were used. This could lead to significant
365 differences of the source apportionment results compared to on-line measurements. In this work an improved off-line aerosol mass spectrometer analysis technique has been developed, which can capture a significant part of the insoluble and the partially soluble fraction of OA.

The improved off-line AMS technique has been evaluated in three different campaigns in Greece, two during winter (Athens and Patras) and one in the summer (Patras). PMF analysis was performed for each campaign, separately for the on-



370 line and for the off-line results. The fractional contribution of each source to the total OA differed by less than 15% between
the on-line and the off-line PMF results, which shows that the two methods are in good agreement (considering their
uncertainty) without any corrections required for the off-line results.

The AMS spectra, however, showed significant differences between the on-line and the off-line PMF results. This is
due to both the reduced temporal resolution in the off-line results compared to the on-line, and to the uncertainty introduced
375 by aspects of the experimental procedure, such as the extraction efficiency etc. However, the PMF appears to be able to adjust
the factors and still give relatively accurate fractional contributions of each factor to the total OA. Comparing summer with
winter results, better agreement was observed in summer between the on-line and off-line OA spectra. This could be due to
the prevalence of the secondary material that coats most of the primary particles and potentially facilitates their suspension as
it dissolves in water.

380 The proposed improved off-line AMS technique can mobilize a significant fraction of the water insoluble material
and allow the AMS to measure it. This reduces significantly the uncertainty compared to the original method. The suspension
of the insoluble material in the improved off-line technique, was evident by the considerable number of particles detected in
the water extract at the range of 50-300 nm, and by the lack of more than half of the EC from the filter after our water extraction
process. In addition, a significant BC concentration was detected in the aerosol samples before the AMS, which is evidence
385 that insoluble material can be the extracted from the filter, can be atomized and can get analysed by the AMS with the proposed
approach.

Data availability. Measurement data are available by request by ch.vasilakopoulou@chemeng.upatras.gr.

390 *Author contributions.* CNV and SP designed the study and wrote the paper. CNV did the off-line measurements and analysis.
CNV, KF, CK, IS and NM obtained and provided the on-line measurements. All authors edited the manuscript.

Acknowledgments. We would like to thank Dr. Aikaterini Bougiatioti and Dr. Despina Paraskevopoulou for providing the
filters from the Thissio 2019 winter campaign.

395

Financial support. This research has been supported by the EU H2020 RI-URBANS project (grant 101036245) and by the
Hellenic Foundation for Research & Innovation (HFRI) under project CHEVOPIN, grant agreement no. 1819.

Competing interests. None.



400 References

- Aiken, A. C., Salcedo, D., Cubison, M. J., Huffman, J. A., DeCarlo, P. F., Ulbrich, I. M., Docherty, K. S., Sueper, D., Kimmel, J. R., Worsnop, D. R., Trimborn, A., Northway, M., Stone, E. A., Schauer, J. J., Volkamer, R. M., Fortner, E., De Foy, B., Wang, J., Laskin, A., Shutthanandan, V., Zheng, J., Zhang, R., Gaffney, J., Marley, N. A., Paredes-Miranda, G., Arnott, W. P., Molina, L. T., Sosa, G. and Jimenez, J. L.: Mexico City aerosol analysis during MILAGRO using high resolution
405 aerosol mass spectrometry at the urban supersite (T0) - Part 1: Fine particle composition and organic source apportionment, *Atmos. Chem. Phys.*, 9, 6633–6653, doi:10.5194/acp-9-6633-2009, 2009.
- Allan, D., Williams, P. I., Morgan, W. T., Martin, C. L., Flynn, M. J., Lee, J., Nemitz, E., Phillips, G. J., Gallagher, M. W. and Coe, H.: Contributions from transport, solid fuel burning and cooking to primary organic aerosols in two UK cities, *Atmos. Chem. Phys.*, 10, 647–668, doi:10.5194/acp-10-647-2010, 2010.
- 410 Bozzetti, C., El Haddad, I., Salameh, D., Daellenbach, K. R., Fermo, P., Gonzalez, R., Minguillón, M. C., Iinuma, Y., Poulain, L., Elser, M., Müller, E., Slowik, J. G., Jaffrezo, J. L., Baltensperger, U., Marchand, N. and Prévôt, A. S. H.: Organic aerosol source apportionment by offline-AMS over a full year in Marseille, *Atmos. Chem. Phys.*, 17, 8247–8268, doi:10.5194/acp-17-8247-2017, 2017.
- Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U. and Prévôt, A. S. H.: SoFi, an IGOR-based interface for the efficient
415 use of the generalized multilinear engine (ME-2) for the source apportionment: ME-2 application to aerosol mass spectrometer data, *Atmos. Meas. Tech.*, 6, 3649–3661, doi:10.5194/amt-6-3649-2013, 2013.
- Daellenbach, K. R., Bozzetti, C., Křepelová, A., Canonaco, F., Wolf, R., Zotter, P., Fermo, P., Crippa, M., Slowik, J. G., Sosedova, Y., Zhang, Y., Huang, R. J., Poulain, L., Szidat, S., Baltensperger, U., El Haddad, I. and Prévôt, A. S. H.: Characterization and source apportionment of organic aerosol using offline aerosol mass spectrometry, *Atmos. Meas.*
420 *Tech.*, 9, 23–39, doi:10.5194/amt-9-23-2016, 2016.
- DeCarlo, P. F., Dunlea, E. J., Kimmel, J. R., Aiken, A. C., Sueper, D., Crounse, J., Wennberg, P. O., Emmons, L., Shinozuka, Y., Clarke, A., Zhou, J., Tomlinson, J., Collins, D. R., Knapp, D., Weinheimer, A. J., Montzka, D. D., Campos, T. and Jimenez, J. L.: Fast airborne aerosol size and chemistry measurements above Mexico City and Central Mexico during the MILAGRO campaign, *Atmos. Chem. Phys.*, 8, 4027–4048, doi:10.5194/acp-8-4027-2008, 2008.
- 425 Drewnick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian, K. L., Borrmann, S. and Worsnop, D. R.: A new time-of-flight aerosol mass spectrometer (TOF-AMS) - Instrument description and first field deployment, *Aerosol Sci. Technol.*, 39, 637–658, doi:10.1080/02786820500182040, 2005.
- Florou, K., Papanastasiou, D. K., Pikridas, M., Kaltsonoudis, C., Louvaris, E., Gkatzelis, G. I., Patoulias, D., Mihalopoulos, N. and Pandis, S. N.: The contribution of wood burning and other pollution sources to wintertime organic aerosol levels in
430 two Greek cities, *Atmos. Chem. Phys.*, 17, 3145–3163, doi:10.5194/acp-17-3145-2017, 2017.



- Ge, X., Li, L., Chen, Y., Chen, H., Wu, D., Wang, J., Xie, X., Ge, S., Ye, Z., Xu, J. and Chen, M.: Aerosol characteristics and sources in Yangzhou, China resolved by offline aerosol mass spectrometry and other techniques, *Environ. Pollut.*, 225, 74–85, doi:10.1016/j.envpol.2017.03.044, 2017.
- Jayne, J. T., Leard, D. C., Zhang, X., Davidovits, P., Kenneth, A., Kolb, C. E., Worsnop, D. R., Jayne, J. T., Leard, D. C.,
435 Zhang, X., Davidovits, P., Smith, A., Kolb, C. E., Worsnop.: Development of an aerosol mass spectrometer for size and composition analysis of submicron particles, *Aerosol Science & Technology*, 6826, doi:10.1080/027868200410840, 2000.
- Jorga, S. D., Florou, K., Kaltsonoudis, C., Kodros, J. K., Vasilakopoulou, C., Cirtog, M., Fouqueau, A., Picquet-Varrault, B., Nenes, A. and Pandis, S. N.: Nighttime chemistry of biomass burning emissions in urban areas: A dual mobile chamber study, *Atmos. Chem. Phys.*, 21, 15337–15349, doi:10.5194/acp-21-15337-2021, 2021.
- 440 Kaltsonoudis, C., Kostenidou, E., Louvaris, E., Psichoudaki, M., Tsiligiannis, E., Florou, K., Liangou, A., and Pandis, S. N.: Characterization of fresh and aged organic aerosol emissions from meat charbroiling, *Atmos. Chem. Phys.*, 17, 7143–7155, <https://doi.org/10.5194/acp-17-7143-2017>, 2017
- Kostenidou, E., Lee, B. H., Engelhart, G. J., Pierce, J. R. and Pandis, S. N.: Mass spectra deconvolution of low, medium, and high volatility biogenic secondary organic aerosol, *Environ. Sci. Technol.*, 43, 4884–4889, doi:10.1021/es803676g, 2009.
- 445 Kostenidou, E., Florou, K., Kaltsonoudis, C., Tsiflikiotou, M., Vratolis, S., Eleftheriadis, K. and Pandis, S. N.: Sources and chemical characterization of organic aerosol during the summer in the eastern Mediterranean, *Atmos. Chem. Phys.*, 15, 11355–11371, doi:10.5194/acp-15-11355-2015, 2015.
- Lanz, V. A., Alfara, M. R., Baltensperger, U., Buchmann, B., Hueglin, C. and Prévôt, A. S. H.: Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra, *Atmos. Chem. Phys.*,
450 7, 1503–1522, doi:10.5194/acp-7-1503-2007, 2007.
- Li, H., Zhang, Q., Jiang, W., Collier, S., Sun, Y., Zhang, Q. and He, K.: Characteristics and sources of water-soluble organic aerosol in a heavily polluted environment in Northern China, *Sci. Total Environ.*, 758, 143970, doi:10.1016/j.scitotenv.2020.143970, 2021.
- Mihara, T. and Mochida, M.: Characterization of solvent-extractable organics in urban aerosols based on mass spectrum
455 analysis and hygroscopic growth measurement, *Environ. Sci. Technol.*, 45, 9168–9174, doi:10.1021/es201271w, 2011.
- Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Zhang, Q., Ulbrich, I. M. and Worsnop, D. R.: Real-time methods for estimating organic component mass concentrations from aerosol mass spectrometer data, *Environ. Sci. Technol.*, 45, 910–916, doi:10.1021/es102951k, 2011.
- Niedek, C. R., Mei, F., Zawadowicz, M. A., Zhu, Z., Schmid, B. and Zhang, Q.: Quantitative Chemical Assay of Nanogram-Level PM Using Aerosol Mass Spectrometry: Characterization of Particles Collected from Uncrewed Atmospheric
460 Measurement Platforms, *Atmos. Meas. Tech. Discuss.*, 1–23, doi: 10.5194/amt-2022-246, 2022.
- Paatero, P.: The Multilinear Engine—A table-driven, least squares program for solving multilinear problems, including the n-way parallel factor analysis model, *J. Comput. Graph. Stat.*, 8, 854–888, doi:10.1080/10618600.1999.10474853, 1999.



- Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error
465 estimates of data values, *Environmetrics*, 5, 111–126, doi:10.1002/env.3170050203, 1994.
- Saarikoski, S., Carbone, S., Decesari, S., Giulianelli, L., Angelini, F., Canagaratna, M., Ng, N. L., Trimborn, A., Facchini, M.
C., Fuzzi, S., Hillamo, R. and Worsnop, D.: Chemical characterization of springtime submicrometer aerosol in Po Valley,
Italy, *Atmos. Chem. Phys.*, 12, 8401–8421, doi:10.5194/acp-12-8401-2012, 2012.
- Setyan, A., Zhang, Q., Merkel, M., Knighton, W. B., Sun, Y., Song, C., Shilling, J. E., Onasch, T. B., Herndon, S. C., Worsnop,
470 D. R., Fast, J. D., Zaveri, R. A., Berg, L. K., Wiedensohler, A., Flowers, B. A., Dubey, M. K. and Subramanian, R.:
Characterization of submicron particles influenced by mixed biogenic and anthropogenic emissions using high-resolution
aerosol mass spectrometry: Results from CARES, *Atmos. Chem. Phys.*, 12, 8131–8156, doi:10.5194/acp-12-8131-2012,
2012.
- Srivastava, D., Daellenbach, K. R., Zhang, Y., Bonnaire, N., Chazeau, B., Perraudin, E., Gros, V., Lucarelli, F., Villenave, E.,
475 Prévôt, A. S. H., El Haddad, I., Favez, O. and Albinet, A.: Comparison of five methodologies to apportion organic aerosol
sources during a PM pollution event, *Sci. Total Environ.*, 757, 143168, doi:10.1016/j.scitotenv.2020.143168, 2021.
- Sun, Y. L., Zhang, Q., Schwab, J. J., Demerjian, K. L., Chen, W. N., Bae, M. S., Hung, H. M., Hogrefe, O., Frank, B., Rattigan,
O. V. and Lin, Y. C.: Characterization of the sources and processes of organic and inorganic aerosols in New York city
with a high-resolution time-of-flight aerosol mass spectrometer, *Atmos. Chem. Phys.*, 11, 1581–1602, doi:10.5194/acp-
480 11-1581-2011, 2011.
- Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R. and Jimenez, J. L.: Interpretation of organic components from
Positive Matrix Factorization of aerosol mass spectrometric data, *Atmos. Chem. Phys.*, 9, 2891–2918, doi:10.5194/acp-9-
2891-2009, 2009.
- Vasilakopoulou, C., Stavroulas, I., Mihalopoulos, N. and Pandis, S. N.: The effect of the averaging period for PMF analysis
485 of aerosol mass spectrometer measurements during off-line applications, *Atmos. Meas. Tech.*, 15, 6419–6431,
doi:10.5194/amt-15-6419-2022, 2022.
- Vlachou, A., Daellenbach, K. R., Bozzetti, C., Chazeau, B., Salazar, G. A., Szidat, S., Jaffrezo, J. L., Hueglin, C.,
Baltensperger, U., El Haddad, I. and Prévôt, A. S. H.: Advanced source apportionment of carbonaceous aerosols by
coupling offline AMS and radiocarbon size-segregated measurements over a nearly 2-year period, *Atmos. Chem. Phys.*,
490 18, 6187–6206, doi:10.5194/acp-18-6187-2018, 2018.
- Xu, L., Guo, H., Weber, R. J. and Ng, N. L.: Chemical characterization of water-soluble organic aerosol in contrasting rural
and urban environments in the southeastern United States, *Environ. Sci. Technol.*, 51, 78–88, doi:10.1021/acs.est.6b05002,
2017.
- Ye, Z., Liu, J., Gu, A., Feng, F., Liu, Y., Bi, C., Xu, J., Li, L., Chen, H., Chen, Y., Dai, L., Zhou, Q. and Ge, X.: Chemical
495 characterization of fine particulate matter in Changzhou, China, and source apportionment with offline aerosol mass
spectrometry, *Atmos. Chem. Phys.*, 17, 2573–2592, doi:10.5194/acp-17-2573-2017, 2017.




Figure 1: Time series of a) PM_{10} organics and b) BC during the winter 2020 campaign in Patra. The fractional contribution of each factor to the total OA is also shown.

500

505




Figure 2: Organic aerosol time series and fractional composition of the sources derived from the on-line PMF analysis for the summer 2019 Patras campaign. Measurements were performed only during specific days in this three-month period.

510

515

520




Figure 3: Organic aerosol concentration for the Athens 2019 winter campaign.

525

530

535

540





Figure 4: Contribution of each factor to the total OA for the on-line and the off-line PMF results in Patras during the winter of 2020. The contribution of each one of the two BBOA factors to the total OA is also shown. A different scale is used for the
545 BBOA.

550




555

Figure 5: Comparison of the average on-line and off-line organic mass spectrum for the Patras 2020 winter campaign.


560

565



570

Figure 6: Off-line to on-line ratio of various HOA markers ($C_xH_y^+$ on top and $C_xH_yO^+$ at the bottom) for the winter 2020 Patras campaign.



575 **Figure 7:** Organics to sulfate ratio comparison between the off-line and the on-line results for the TEI winter 2020 campaign.

580




Figure 8: Comparison of the contribution of each factor to the total OA between the on-line and the off-line PMF results for the summer 2019 Patras campaign.

585

590

595



600





Figure 9: Off-line to on-line fraction various $C_xH_y^+$ and $C_xH_yO^+$ HOA markers for the summer 2019 Patras campaign.

605

610



615

Figure 10: Comparison of the contribution of each factor to the total OA between the on-line and the off-line PMF analysis for the Thissio 2019 winter campaign.

620

625

630





Figure 11: Number concentration of suspended particles in the water extract for the sample (in red) and the blank (in blue).

635

640



645

Figure 12: Elemental carbon before and after the extraction process.

650

655

660

665




Figure 13: BC concentration time series of the off-line SP2 test. The blanks are shown in blue background and the samples in white.