

1 **Testing ground based observations of wave activity in the (lower and upper) atmosphere**
2 **as possible (complementary) indicators of streamer events**

3 **Michal Kozubek¹, Lisa Kuchelbacher², Jaroslav Chum¹, Tereza Sindelarova¹, Franziska**
4 **Trinkl^{2, a}, Katerina Podolska¹**

5 ¹Institute of Atmospheric Physics CAS, Bocni II 1401, Prague, 14100, Czech Republic

6 ² Earth Observation Center, Deutsches Zentrum für Luft- und Raumfahrt, 82234 Weßling,
7 Germany

8 ^a now at Karlsruhe Institute of Technology (KIT), Institute of Meteorology and Climate
9 Research, Karlsruhe, Germany

10 **Correspondence: Michal Kozubek, kom@ufa.cas.cz**

11 **Keywords:** gravity waves, streamer events, infrasound, Doppler measurements

12

13 **Abstract:** For a better understanding of atmospheric dynamics, it is very important to know
14 the general condition (dynamics and chemistry) of the atmosphere. Planetary waves (PWs) are
15 global scale waves, which are well-known as main drivers of the large-scale weather patterns
16 in mid-latitudes on time scales from several days up to weeks in the troposphere. When PWs
17 break, they often cut pressure cells off the jet stream. A specific example are so-called
18 streamer events, which occur predominantly in the lower stratosphere at mid- and high-
19 latitudes. For streamer events we check, whether there are any changes of gravity wave (GW)
20 or infrasound characteristics related to these events in ionospheric and surface measurements
21 (continuous Doppler soundings, two arrays of microbarometers) in the Czech Republic.

22 **Phenomena in infrasound arrival parameters undoubtedly related with streamer events were**
23 **not identified in observations of two stations located in Central Europe. Simulations of**
24 **infrasound propagation show influences of the streamer events on the waveguide formed near**
25 **the tropopause. Microbarom propagation is influenced by the tropopause waveguide in a**
26 **limited azimuth sector and at limited distances. Due to the typical occurrence of the streamer**
27 **events over the North Atlantic, infrasound stations in Western Europe can be of particular**
28 **interest for future studies of streamer event signatures in infrasound arrivals. Arrivals to**
29 **Central Europe are through the waveguide formed between the ground and the upper**
30 **stratosphere. The upper stratosphere waveguide is not influenced by the streamer events.**

31 Supplementary ground-based measurements of GW using the WBCI array in the troposphere
32 showed that GW propagation azimuths were more random during streamer and streamer-like
33 events compared to those observed during calm conditions. GW propagation characteristics
34 observed in the ionosphere by continuous Doppler soundings during streamer events did not
35 differ from those expected for the given time period.

36

37 **1) Introduction**

38 For a better comprehension of climate change it is fundamentally important, how well we
39 understand the climate system in general, and the dynamics of the atmosphere in particular.
40 The dynamical processes relevant in this context in the atmosphere take place over a
41 comparatively wide range of scales in space and time. They include in particular both,
42 planetary and gravity waves. Planetary waves are the main drivers of the extratropical
43 circulation. When they break, they lead to an irreversible exchange of air masses between the
44 equatorial and polar region due to an amplification of their amplitudes (e.g. McIntyre &
45 Palmer, 1983; Polvani & Plumb, 1992). In the lower stratosphere ozone can be used as a tracer
46 for these large-scale motions, as it has a comparatively long life-time. When planetary waves
47 break tropical air masses of low ozone concentration are mixed poleward into the
48 surrounding atmosphere of the mid and higher latitudes (e.g. Leovy et al., 1985).

49 The term "streamer" lacks a precise definition, as noted by Krüger et al. (2005). They
50 discuss various aspects of streamers, including their impact on mixing and the divergent
51 definitions associated with them. Offermann et al. (1999) describe streamers as large-scale
52 tongue-like structures formed by the meridional deflection of air masses. Streamers are
53 characterized by irreversible mixing of air masses between equatorial and polar regions which
54 is why **they** might be linked to planetary wave breaking (Waugh, 1993). Eyring et al. (2003)
55 give a climatology of the seasonal and geographical distribution of streamer events. They
56 show, that streamers often occur over the Northern Atlantic and can be identified by either
57 high NO₂ or low ozone concentration, which is why we select streamers by total ozone
58 column measurements. They show that streamer events occur most often during winter and
59 least during July and August in the Northern Hemisphere. During a streamer event the wind
60 field changes rather strong over a comparatively small distance. Since a streamer event shows
61 a strong wind shear at its flanks, it is expected that it excites GW (e.g. Kramer et al., 2015 and
62 2016 or Peters et al., 2003).

63 It is well-known that enhanced wind gradients or anticyclones can lead to the
64 excitation of gravity waves (GW) in the atmosphere (e.g. Pramitha et al., 2015; Kai et al.,
65 2010; Kramer et al., 2015, 2016 and Gerlach et al., 2003). GW have typical vertical
66 wavelengths from a few 100 m to several kilometres (Wüst & Bittner, 2006), and horizontal
67 wavelengths over tens of km (Wüst et al., 2018), and longer (Rauthe et al., 2006); their
68 fluctuations in the upper troposphere / lower stratosphere typically show amplitudes of 5–10
69 m/s at maximum (e.g., Kramer et al., 2015). Those waves transport energy and momentum
70 horizontally and vertically through the atmosphere and deposit them especially in the
71 stratosphere and mesosphere but also above and below this height region. The propagation of
72 GWs is strongly dependent on the wind conditions in the stratosphere since the wind speed of
73 the middle atmosphere (10–100 km) reaches its maximum there. That is why monitoring
74 waves in upper parts of the atmosphere, e.g. based on Doppler observations in the ionosphere,
75 can provide additional information about stratospheric conditions (for details see Fritts and
76 Alexander, 2003).

77 Using pressure recordings at a microbarograph array, GWs and infrasound at the ground can
78 be observed. Ground based observations of GWs at a large aperture microbarograph array are
79 utilized in the present study as an independent data source for the analysis of GW activity
80 during streamer events. Infrasound propagation is influenced by wind and temperature fields
81 in the atmosphere. Three regions play an important role in long-distance infrasound
82 propagation: (1) the lower thermosphere; (2) the stratosphere; (3) the jet stream near the
83 tropopause and inversion layers in the troposphere (Evers and Haak, 2010). Infrasound
84 observed at the ground and emitted by distant sources mostly propagates in the stratospheric
85 waveguide (Ceranna et al., 2019). The thermospheric waveguide is not as efficient as the
86 stratospheric waveguide in the long-range infrasound propagation. Besides signal loss due to
87 geometrical spreading, infrasound absorption is important in the upper atmosphere (Bittner et
88 al., 2010). Infrasound absorption is proportional to the frequency; higher frequencies,
89 particularly those above 1 Hz undergo stronger absorption in the thermosphere (Sutherland
90 and Bass, 2004). Signal attenuation is low at frequencies of the order of 10^{-3} – 10^{-2} Hz (Blanc,
91 1985; Georges, 1968).

92 A number of case studies have proved that stratospheric dynamics can be deduced from
93 microbarograph measurements at the ground (Assink et al., 2014; Blixt et al., 2019; Evers and
94 Siegmund, 2009; Evers et al., 2012; Garcès et al., 2004; Le Pichon and Blanc, 2005; Le
95 Pichon et al., 2006 and 2009; Smets and Evers, 2014). Streamer events are significant

96 transient disturbances to circulation patterns in the tropopause/lower stratosphere region;
97 modifications of the stratospheric waveguide can therefore be expected. A feasibility study on
98 utilisation of ground infrasound measurements in research of streamer events is performed. Its
99 aim is to identify phenomena in infrasound detections related to the streamers; we focus on
100 deviations of the azimuth of signal arrivals, trace velocity, signal amplitude, and frequency.
101 The dedicated studies demonstrated that from the observed signal trace velocity, information
102 about the signal refraction height can be derived (Lonzaga, 2015). If the source of received
103 signals is well defined in time and space, mean atmospheric cross-winds along the signal
104 propagation path can be estimated from back-azimuth deviations and time of signal
105 propagation (Blixt et al., 2019). Fluctuations of signal frequency and amplitude are, besides
106 variability of the signal source influenced by atmospheric filtering (Sutherland and Bass,
107 2004).

108 Our study will focus on possible utilization Doppler sounding and microbarographs for
109 description and analysis of GW behaviour and propagation in the stratosphere.

110 The structure of the paper is as follows: After introduction the description of the used dataset
111 and method can be found in the second section. Then we describe our results and in the last
112 section we discuss the possible connection to previous studies.

113

114 **2) Data and methods**

115 The selection of streamer events is based on the visual inspection of global maps of total
116 ozone column (TO3), accessible through a service provided by DLR
117 (<https://atmos.eoc.dlr.de/>) measured by the Tropospheric Monitoring Instrument (TROPOMI)
118 aboard the Sentinel 5 Precursor (S5P) mission. See Veefkind et al., 2012 for details about
119 TROPOMI/S5P. In cases where TROPOMI/S5P data is unavailable, measurements from the
120 Global Ozone Monitoring Experiment-2 (GOME-2) on the Metop series of satellites are
121 utilized. Both instruments operate in a nadir-viewing configuration on near-polar sun-
122 synchronous orbits. Further specifics regarding TO3 measurements by TROPOMI/S5P are
123 elaborated by Spurr et al. (2022). The TO3 retrieval process is built upon the predecessor
124 instrument's processor, with GOME-2 on Metop-AB, see Munro et al. (2006) and Munro et al.

125 (2016). For detailed information on the GOME-2 retrieval algorithm, refer to Loyola et al.
126 (2011).

127 We define a streamer as such when the ozone column concentration of the finger-like
128 structure above the Northern Atlantic/Western Europe is lower than 300 DU and persists for
129 at least 3 days. The longitudinal extension is of approx. 15 to 30 degrees in the mid-latitudes
130 (between 30 to 70°N). The northernmost point of a streamer exceeds 50°N. Fig. 1 shows a
131 streamer event above the Northern Atlantic, indicated by the blue color which represent the
132 low ozone concentrations. The streamer shown in Fig. 1 reaches latitudes beyond 70°N,
133 which indicates a large example. At the western and eastern flanks of the streamer, the ozone
134 concentration exceeds 350 DU, defining distinct boundaries. This is also visible in Fig. 1
135 represented by the green colors at the eastern coast of Northern America and western Europe.
136 So, there is a gradient of the ozone concentration of about 50 DU / 5°. Furthermore, the
137 streamer exhibits a discernible pattern of circulation, with air masses being meridionally
138 deflected, contributing to its formation and maintenance. These air masses, characterized by
139 their movement from south to north at the eastern flank and from north to south at the western
140 flank, play a significant role in the streamer's dynamics. This is the reason why equatorial low
141 ozone concentration is transported northward. In contrast, the calm periods, representing the
142 opposite dynamic situation to the streamer events, are characterized by only very few
143 meridionally deflected air masses. During these periods, the ozone concentration in the mid-
144 latitudes above the Northern Atlantic is consistently higher than 350 DU, indicating stable
145 atmospheric conditions and minimal perturbations in the ozone distribution. An example for a
146 calm period is shown in Fig. 2.

147 The streamer events are selected by eye for this study (results see Table 1) considering the
148 TO3 global maps from January 2020 and March 2021. As planetary waves are permanently
149 disturbing the atmospheric dynamic of the higher troposphere / lower stratosphere, especially
150 smaller scale streamers can be observed almost every day and the identification of streamer
151 events becomes subjective. We therefore focus on few events which are comparatively strong
152 in their evolution from our perspective. Moreover, we focus on streamer events above the
153 Northern Atlantic. Whenever another streamer event occurs somewhere other than over the
154 Northern Atlantic region with comparable spatiotemporal extent, we do not consider this date
155 as a streamer event. We assume that the effects of the streamer superimpose and a distinct
156 backtrack to the streamer over the Northern Atlantic will not be possible. This means, that the
157 analysis of the streamer events can be blurred to some extent.

158 We consider dates from January 2020 to April 2021. In general, planetary waves drive the
 159 Brewer Dobson Circulation in the stratosphere during winter and ozone-poor airmasses are
 160 transported northward. Streamer events are therefore detected between September and March.
 161 The streamer events are distinguished if they have a large spatial size, high intensity (low
 162 TO3 concentration) and if air masses are irreversibly mixed into the surrounding atmosphere.
 163 All the selected events persist for several days, but no longer than 10 days.

164 To evaluate whether streamer events effect the smaller-scale atmospheric dynamics, calm
 165 events are identified as well by subjective criteria. These events serve as a reference to
 166 streamer events, as large-scale spatial structures are hardly visible in the TO3. The events are
 167 selected when the ozone concentration shows a meridional gradient from the equator to polar
 168 region on the Northern Hemisphere with almost no longitudinal variation. The examples of
 169 calm atmospheric dynamics are listed in Table 1 (right).

170


Streamer events		Calm periods	
From	To	From	To
06.02.2020	10.02.2020	02.03.2020	08.03.2020
11.2.2020	13.2.2020	09.03.2020	14.03.2020
31.08.2020	03.09.2020	28.03.2020	10.04.2020
05.09.2020	11.09.2020	19.04.2020	27.05.2020
03.11.2020	07.11.2020	9.11.2020	15.11.2020
21.11.2020	25.11.2020	12.12.2020	22.12.2020
23.02.2021	27.02.2021	30.12.2020	06.01.2021
09.03.2021	12.03.2021	21.01.2021	20.02.2021
		28.02.2021	07.03.2021
		13.03.2021	24.03.2021
		29.03.2021	07.04.2021

--	--	--	--

171 **Table 1** Streamer events above Northern Atlantic from January 2020 until March 2021 and
 172 related start and end dates. The right part shows calm periods.

173

174 Figure 1 shows the TO3 by TOPOMI/S5P integrated from November 3rd to November 5th
 175 2020. Ozone-poor airmasses (blue) are located above the Northern Atlantic from 30°N to
 176 70°N next to smaller scale ozone-poor airmasses above western North America and Central
 177 Asia. The TO3 concentration is disturbed by planetary waves along the latitudes, which lead
 178 to wave structures visible especially at the transition of blue to green colors. A large streamer
 179 event of ozone-poor airmasses is detected over the Northern Atlantic. A small streamer can be
 180 detected over western North America. There are also ozone-poor air masses above eastern
 181 Europe. The temporal evolution shows, that the ozone-poor air masses above eastern Europe
 182 are due to a decaying streamer which evolved several days earlier. As planetary waves are
 183 more or less permanently disturbing the atmospheric dynamics, especially smaller scale
 184 streamers can be detected almost every day. In this example, the streamer event above the
 185 Northern Atlantic is largest. Therefore, we consider this event for the further analysis.




187 Fig. 1. TO3 by TROPOMI/S5P from November 3rd to November 5th 2020 shows ozone poor
 188 airmasses above the Northern Atlantic as an example of a streamer event for the further
 189 analysis. Colors (from violet to red) indicate the total ozone column concentrations (from low
 190 to high) in Dobson Units. Source: DLR, CC-BY 3.0

191

192 Figure 2 shows the TO3 by TOPOMI/S5P from February 11th to February 13th 2020. The
 193 event is characterized by a strong meridional gradient from the equatorial to polar region on
 194 the Northern Hemisphere with almost no longitudinal variation. Therefore, we consider this
 195 event for the further analysis.

196



198 Fig. 2. TO3 by TROPOMI/S5P from February 11th to February 13th 2020 as an example of
 199 calm atmospheric dynamics. A clear meridional gradient of ozone can be observed on the
 200 Northern Hemisphere without large-scale wave structures. Colors (from violet to red) indicate
 201 the total ozone column concentrations (from low to high) in Dobson Units. Source: DLR, CC-
 202 BY 3.0

203 Two stations of the Czech microbarograph network (Bondar et al., 2022) are involved in the
 204 study – the large aperture array WBCI (50.25°N 12.44°E) and the small aperture array PVICI
 205 (50.52°N 14.57°E). To study propagation of GW and long-period infrasound (from acoustic
 206 cut-off up to about 2.5 s) pressure recordings at WBCI are utilized. Four sensors of the WBCI
 207 array are arranged in a tetragon. The inter-element distances of 4 – 10 km define an optimum
 208 performance of the array in the infrasound frequency range from the acoustic cut-off

209 frequency of 0.0033 to 0.0068 Hz (Garcès, 2013). The WBCI array with its large inter-
 210 element distances has a unique configuration compared to the arrays of the International
 211 Monitoring System of the Comprehensive Nuclear Test Ban Treaty Organisation intended
 212 for infrasound monitoring in the frequency band of 0.02 – 4 Hz (Marty, 2019). Each array
 213 element at WBCI is equipped with an absolute microbarometer of the type Paroscientific
 214 6000-16B-IS with parts-per-billion resolution. A GPS receiver is used for time stamping. Data
 215 are stored with a sampling rate of 50 Hz. For infrasound monitoring, WBCI data are
 216 resampled at 10 Hz sampling rate. To detect and analyze GW, 1-min mean values of the
 217 absolute pressure data are used.

218 The small aperture array PPCI provides optimal precision of detections in the frequency
 219 range of 0.14 – 3.4 Hz (Garcès, 2013). Three sensors are arranged in an equilateral triangle;
 220 the array aperture is 200 m. The differential sensors of the type Infrasound Gage ISGM03
 221 manufactured by the Scientific and Technical Centre give a flat response in the frequency
 222 range of 0.02 – 4 Hz. A GPS receiver is used for time stamping. The data are stored with a
 223 sampling frequency of 25 Hz. This sampling rate is also used in regular processing of
 224 infrasound detections at PPCI.

225 Infrasound detections are processed using the **DTK-GPMCC** software the core of which is the
 226 Progressive Multi-Channel Correlation (PMCC) detection algorithm (Cansi, 1995; Le Pichon
 227 and Cansi, 2003). PMCC analyses pressure recordings from an infrasound array and looks for
 228 coherent signals in overlapping time windows in several frequency bands (Le Pichon and
 229 Cansi, 2003). An elementary detection with the PMCC, or the detection pixel is declared in
 230 the time-frequency window, when signal correlation and consistency criteria are met.
 231 Detection pixels are grouped into the detection families based on similar time, frequency,
 232 azimuth of signal arrival, and signal trace velocity (Brachet et al., 2010). The arrival
 233 parameters of the detected infrasound are stored in the detection bulletins. The parameters of
 234 interest for the present study include time of arrival, azimuth of arrival, trace velocity,
 235 frequency, and amplitude. The PMCC configuration is set on an individual basis and is
 236 optimized for the given array (Brachet et al., 2010; Garcès, 2013; Szuberla et al., 2004); main
 237 parameters of the **DTK-GPMCC** settings for the arrays PPCI and WBCI are given in Table 2.

Station	PPCI	WBCI
Detection range	0.09-7 Hz	0.0033-0.4 Hz
Number of detection bands	19	11

Length of the detection window; frequency dependent	412.84-6.44 s	2555-118 s
Adjacent windows overlap	95 %	90 %
Consistency	0.1 s	3 s
Azimuth tolerance for families forming	10°	3°
Family size	10-50 pixels	15-50 pixels
Frequency range analysed in the study of streamer events	0.09-0.4 Hz	0.0033-0.4 Hz

238

239 **Table 2.** Main parameters of the DTK-GPMCC configurations for the arrays PPCI and
240 WBCI.

241

242 InfraGA/GeoAc raytracing tools are employed to study infrasound ducting in the
243 atmosphere (Blom and Waxler, 2012; Blom, 2019). Infrasound raytracing can reveal
244 possible modifications of atmospheric waveguides above the Eastern Atlantic and Western
245 Europe during streamer events. Streamer events influence the tropopause and lower
246 stratosphere. Hence, modifications of the stratospheric waveguide are expected rather than
247 its entire reversals or collapse. Raytracing can identify azimuths and distances from the
248 source that are influenced by a streamer event. And so, it can reveal whether these
249 influences reach Central Europe or the signals are ducted to the region through the
250 waveguide in the upper stratosphere or thermosphere like in quiet periods. InfraGA/GeoAc
251 provides simulations of signal propagation from a point source; propagation through the
252 range dependent atmosphere is modelled in the present study. Atmospheric characteristics
253 are obtained from the G2S model (Drob et al. 2003). Vertical profiles of temperature, zonal
254 and meridional winds, density and pressure are an input for the InfraGA/GeoAc. The grid
255 of profiles covers the area from 45° to 65°N and from 30°W to 22.5°E; latitudinal step is 5°
256 and longitudinal step is 7.5°.

257 Propagation of GW in the thermosphere/ionosphere is studied using the multi-point and multi-
258 frequency continuous Doppler sounding system located in Czechia. Its advantage is a high
259 time resolution (around 10 s) compared with ionospheric sounders (ionosondes) that measure
260 the profile of electron densities in the ionosphere. The frequency shift is due to the motion and
261 electron density changes in the ionospheric plasma, caused for example by interaction with

262 atmospheric waves propagating in the neutral atmosphere, with which the ionosphere (above
263 ~ 80 km) merges. The sounding radio signal is reflected at the height, where its frequency
264 matches the so called local plasma frequency, which is determined by the local electron
265 density. Therefore, the reflection height changes during the day and depends on the sounding
266 frequency. Significant Doppler shifts, usable for analysis, are obtained if the signal is
267 reflected from the so called F2 layer (approximately 200 – 300 km). Several sounding
268 frequencies are used in Czechia. The 3.59 MHz sounding was mostly effective at night, while
269 the 4.65 MHz sounding provided good daytime data during the period analyzed. The
270 propagation characteristics of GWs are calculated from the time delays between signals
271 observed at the respective sounding paths (reflection points for each transmitter-receiver
272 pairs) assuming that the reflection points are in the midpoints between each transmitter and
273 receiver. A 60 or 90 min long time interval is usually used to calculate the velocities and
274 azimuth of the observed waves. The methods are in detail described by Chum and Podolska
275 (2018). The two-dimensional (2-D) version (propagation analysis in horizontal plane only) is
276 anticipated for most of the studies, since a 3-D analysis requires simultaneous observation and
277 signal correlation at different frequencies, which is often not the case, especially during solar
278 minimum. Results of statistical investigation have been recently published (Chum et al.,
279 2021). Identical methods of propagation analysis have been applied to investigate
280 propagation of GWs in the troposphere based on data from large-aperture array WBCI (here
281 the time delays are related to the locations of individual microbarometers). All analyses will
282 be done with respect to the streamer events and calm periods shown in Table 1.

283 3) Results

284 3.1 Infrasound observations at ground microbarograph arrays WBCI and PVICI in 285 November 2020 and in March 2021

286 Wave activity in the infrasound frequency range of 0.0033-0.4 Hz is investigated combining
287 observations at the stations WBCI and PVICI. Infrasound detections at WBCI are processed in
288 the frequency band of 0.0033 – 0.4 Hz. The operational range of the array is extended above
289 the upper limit of the optimum array range; the degraded performance of WBCI at
290 frequencies higher than 0.0068 Hz shall be considered. The upper limit of the analysed band
291 is intentionally set to 0.4 Hz to cover microbaroms. PVICI detections are analysed in the
292 frequency range of 0.09 – 0.4 Hz. The band partly overlaps with the detection range of the
293 WBCI array and at frequencies of 0.12 – 0.35 Hz it is dominated by microbaroms (e.g.,

294 Campus and Christie, 2010). Unlike WBCI, PPCI provides an optimal performance in the
295 microbarom band.

296 Microbaroms are infrasound signals generated by a non-linear interaction of ocean waves
297 travelling in opposite directions. Microbaroms form a wide peak around 0.2 Hz in the
298 infrasound spectrum; their frequency corresponds to twice the frequency of sea waves. A
299 powerful source of microbaroms is located in the North Atlantic and the signals are regularly
300 detected by European stations (Hupe et al., 2019). The detection capability of microbaroms
301 from the North Atlantic is particularly high from October to March when the source becomes
302 stronger due to stormy weather above the ocean and signal propagation to the East from the
303 source is supported by the stratospheric waveguide (Landès et al., 2012). From the global
304 point of view, microbaroms are permanently present in recordings of infrasound stations
305 worldwide.

306 Streamer events often occur above the North Atlantic. Thus, microbaroms propagating from
307 the North Atlantic to the continental Europe can travel through the region influenced by a
308 streamer event and the detections at infrasound stations in Europe can show signatures of
309 streamer events.

310 We analyse infrasound observations from 3rd to 25th November 2020 and from 28th
311 February to 25th March 2021 with focus on microbaroms. In these time intervals adjacent
312 streamers and calm periods occurred (Table 1). Streamers and the calm period in the
313 November 2020 time window are evaluated separately from those in the March 2021 time
314 window to avoid seasonal influences. While a well-developed eastward stratospheric
315 waveguide can be expected in November, its efficiency can decrease in March due to the
316 seasonal reversal of stratospheric winds.

317


318 3.1.1 Infrasound observations from 3rd to 25th November 2020

319 Two streamer events developed in November 2020. The first streamer occurred from 3rd
320 to 7th November and the second one from 21st to 25th November. The streamers were
321 separated by a calm period from 9th to 15th November.

322 The most important phenomena found in the infrasound arrival parameters are fluctuating
323 signal frequency and fluctuating signal amplitude.

324 WBCI provides rather sparse detections during both streamer events and only two
325 detection families are obtained during the seven-day calm period (Figure 3). The signal
326 frequencies near 0.2 Hz and back-azimuths of 290° – 350° indicate that the observed
327 signals are likely microbaroms from the North Atlantic. A decrease of the signal frequency

328 is observed during the first streamer event. On 5th – 6th November from 20 to 05 UTC, the
 329 mean frequency of the north-west arrivals drops down to 0.04 Hz, below the microbarom
 330 frequency range. During the second streamer event from 21st to 25th November, the signal
 331 frequency is stable around 0.22 Hz. An increase of the amplitude from the mean value of
 332 0.019 Pa to 0.035 Pa is observed from 23rd November, 18 UTC until the end of the analysed
 333 time period on 25th November at 24 UTC.
 334




335
 336 Fig. 3. Infrasound observations at WBCI on 3rd - 25th November 2020. Azimuth of signal
 337 arrivals is shown; the colorbar refers to the mean frequency of the detection family. One circle
 338 in the plot represents one detection family. Green background marks the streamer events, grey
 339 background marks the calm period.

340
 341 Similar to the back-azimuths at WBCI, PPCI detects arrivals from the north-west in the
 342 analysed frequency range of 0.09 – 0.4 Hz (Figure 4). Fluctuating signal amplitudes are
 343 observed. Values around 0.020 Pa occur on 3rd November. From 4th November, 18 UTC to
 344 7th November, 22:30 UTC, the signals are of amplitudes around 0.089 Pa. The amplitudes
 345 decrease to the values around 0.046 Pa during the following quiet period on 9th – 15th
 346 November. Microbarom amplitudes fluctuate between 0.013 and 0.036 Pa (1st decile and 9th

347 decile, respectively) during the streamer event on 21st – 25th November. Publicly available
 348 data – meteorological charts provided by Deutscher Wetterdienst and the
 349 WAVEWATCHIII[®] wave-action model (The WAVEWATCHIII[®] Development Group,
 350 2016) indicate that there are maritime storms in the North Atlantic within the analysed time
 351 window from 3rd to 25th November 2020. The storms can cause fluctuating intensity of the
 352 microbarom source and as a consequence, fluctuating microbarom amplitudes are observed
 353 at the infrasound stations.

354
 355
 356




357
 358 Fig.4. Infrasound observations at PVC.I on 3rd - 25th November 2020. Azimuth of signal
 359 arrival is shown; the colorbar refers to the signal amplitude. Green background marks the
 360 streamer events, grey background marks the calm period.

361

362 To study propagation of signals from sources located at the surface of the North Atlantic
 363 the InfraGA/GeoAc tools are employed. The fictitious point sources are located (1) at 55°N
 364 and 15°W, (2) at 55°N and 5°W, and (3) at 60°N and 0°longitude. The coordinates of the
 365 sources are estimated based on the position of the tropopause jet stream disturbance. Point

366 (1) is located under the northward jet-stream, point (3) under the southward jet-stream, and
367 point (2) is located between those two opposing branches of the jet stream disturbance see
368 Figure 5.



369
370

371 Fig.5. Wind field at the pressure level of 250 hPa on 06 November 2020 at 00 UTC. A
372 disturbance of the jet-stream above the eastern North Atlantic and the British Isles is caused
373 by the streamer event.

374

375 A multi azimuth simulation is run on 6th November at 00 UTC, during the streamer event,
376 Taking into account the mutual locations of the sources and the receiving arrays, eastward
377 signal propagation is modelled. The azimuth limits are set to 0° and 180°, the azimuth step
378 is 3°. Signal inclinations 2° – 45° are considered in 2° resolution. Information is obtained
379 through which waveguides the signal can possibly arrive to the infrasound stations and their
380 surroundings. The reason why arrivals to the larger areas are considered is that signal
381 propagation from three fictitious point sources stands in for an areal source, the surface of
382 the North Atlantic where microbaroms are generated. Therefore, the model outputs must be
383 taken as an approximation of the real situation. The turning height and ground reflections of
384 the 0.2 Hz signal are obtained in the multi azimuth simulation. The results are visualised in
385 Figure 6 and in supplementary materials. The red asterisk represents the point source. The
386 concentric sectors of circles show regions of ensonification, regions where the signal
387 emitted by the source can be recorded at an infrasound station. The dots, signal ground
388 reflections are organized in a radial pattern. Each of the lines of this pattern represents one
389 azimuth of signal propagation for which the multi azimuth simulation is run; the azimuth
390 step is 3°. The colours of the dots inform about the turning height of the ray and thus
391 provide information about signal ducting in the waveguides. Depending on the turning
392 height, infrasound is subject to attenuation of variable strength when it propagates through

393 the atmosphere. Infrasound attenuation is low in the stratospheric waveguide. Strong
394 absorption occurs in the thermospheric waveguide; the absorption is higher at higher signal
395 frequencies (Sutherland and Bass, 2004). To obtain the view of signal attenuation along the
396 raypath in the vertical plain a single azimuth simulation is employed. The single azimuth
397 simulation is run along the azimuths from the fictitious sources (1) – (3) to the stations
398 WBCI and PPCI; it is obtained for the frequencies of 0.04 Hz and 0.2 Hz. As a reference, a
399 multi azimuth propagation of the 0.2 Hz signal is modelled from a source at 55°N and
400 15°W on the calm day 12th November at 00 UTC.


401 First, we focus on infrasound propagation from the North Atlantic to Central Europe.
402 Stratospheric and thermospheric ducting are possible from the sources at 55°N 15°W and
403 55°N 5°W to Central Europe during the streamer event as well as on the calm day
404 (supplementary materials). Signal propagation only through the thermospheric waveguide
405 is enabled from the source at 60°N and 0°longitude (Figure 6). The distances between the
406 fictitious sources and the stations are 1300 – 2000 km. The amplitude loss of the 0.2 Hz
407 signal in the thermospheric waveguide at these distances is 100 dB relative to the amplitude
408 at a distance of 1 km from the source. According to the simulations, observations of the
409 thermospheric arrivals of microbaroms are unlikely at PPCI and WBCI due to strong signal
410 attenuation. Microbaroms apparently arrive to Central Europe through the waveguide
411 formed in the upper stratosphere during the streamer events as well as on the calm day.
412 Indeed, arrivals from the back-azimuths of 285° - 315° are dominant at PPCI on 3rd to 7th
413 November. Those back-azimuths correspond to the sources at 55°N 15°W and at 55°N
414 5°W, while the back-azimuth to the source at 60°N and 0°latitude is 325°. The amplitude
415 loss of the 0.04 Hz signal at the distances of 1300 – 2000 km from the source is 60 – 80 dB.
416 In general, thermospheric arrivals of this low frequency signal are not strictly rejected.
417 However, in our case the 0.04 Hz signal arrives with trace velocity around 0.330 km/s at
418 WBCI. The low trace velocity indicates signal ducting in the troposphere/lower
419 stratosphere waveguide (Lonzaga, 2015).

420 Next, we study the influences of the streamer event related disturbance anywhere in the
421 modelled region. The disturbance of the jet stream can modify signal propagation up to
422 distances of several hundreds to a thousand km from the source; the influenced azimuth
423 range is limited. Signals from the source at 55°N and 15°W can propagate in the tropopause
424 waveguide in azimuths between 10° and 60° up to the distance of ~1000 km. The amplitude
425 loss of the 0.2 Hz signal at a distance of 1000 km is 60-70 dB relative to the amplitude at 1
426 km from the source. The southward branch of the jet-stream disturbance enables infrasound

427 propagation in the tropospheric waveguide in azimuths of 100 - 160° from the source at
428 60°N 0°longitude. Maximum distance which the signal can travel in the south-east direction
429 is ~600 km. The amplitude loss of the 0.2 Hz signal at a distance of 600 km is 60 dB
430 relative to the amplitude at 1 km from the source.

431 The observations and the model outputs during the November 2020 event can be
432 summarized as follows: infrasound arrives from sources in the North Atlantic to Central
433 Europe mainly through the stratospheric waveguide formed between the ground and upper
434 stratosphere. The jet-stream disturbance above the eastern North Atlantic does not have an
435 impact on infrasound arrivals in Central Europe on 6th November 2020 at 00 UTC.
436 Fluctuating signal amplitudes are likely a consequence of fluctuating intensity of the
437 microbarom source during maritime storms. The decrease of signal frequency at WBCI is
438 not caused by a transient change in signal ducting and by the related signal filtering in the
439 thermospheric waveguide.

440



441

442 Fig.6. Modelled infrasound propagation from a point source located at 60°N and 0°longitude
443 (red asterisk) during the streamer event on 6th November 2020 at 00 UTC. Colobar refers to
444 the turning height (maximum height) of the signal. Red indicates signal propagation in the

445 waveguide formed near the tropopause (altitudes around 10 km), arrivals through the
446 thermospheric waveguide are in blue (altitudes above 100 km). Black triangles represent
447 infrasound arrays WBCI (the left triangle) and PVICI (the right triangle).

448

449 3.1.2 Infrasound observations from 28th February to 24th March 2021


450 A streamer event occurred from 9th to 12th March 2021 preceded and followed by calm
451 periods from 28th February to 7th March and from 13th to 24th March, respectively.

452 The most important phenomenon identified in the infrasound arrival parameters is a
453 fluctuating trace velocity.

454 Both WBCI and PVICI detect signals arriving from the north-west, from back-azimuths of
455 285° – 310°. An increase of signal trace velocities is observed in some of the detections at
456 WBCI during the streamer event compared to calm periods (Figure 7). On 10th March at 00 –
457 06 UTC, trace velocities of 0.460 km/s and 0.380 km/s are observed from back-azimuths of
458 270° and 310° respectively. It is by 0.05 – 0.13 km/s higher than on the calm days. On the
459 other hand, signals from the back-azimuth of 288° arrive with the trace velocity of 0.330 km/s
460 within the same time window, the velocity corresponds to that on the calm days. Effects of
461 spatial aliasing shall be taken into account when evaluating the detections. The signal
462 frequencies are around 0.2 Hz, well above the range of array optimum performance. The
463 observed different trace velocities at WBCI can therefore be a processing bias rather than a
464 consequence of variations in signal ducting.

465 In contrast to the WBCI observations, PVICI records a decrease in trace velocities on 10th
466 March at 00 – 06 UTC (Figure 8). Trace velocities of 0.377 km/s are observed compared to
467 0.413 km/s and 0.395 km/s during the calm periods before and after the streamer,
468 respectively.


469



470

471 **Fig.7.** Infrasound observations at WBCI on 28th February – 24th March 2021. Azimuth of
 472 signal arrival is shown; the colorbar refers to the signal trace velocity. Green background
 473 marks the streamer event, grey background marks the calm periods.

474



475

476 **Fig.8.** Infrasound observations at PVC I on 28th February – 24th March 2021. Azimuth of
 477 signal arrival is shown; the colorbar refers to the signal trace velocity. Green background
 478 marks the streamer event, grey background marks the calm periods.

479

480 **Changes of the trace velocity can indicate changes of the refraction altitude of the signal**
 481 **(Lonzaga, 2015). The exact limits of the trace velocity for the given atmospheric waveguide**
 482 **depend on the current state of the atmosphere. We use the thresholds determined for a**
 483 **model atmosphere in Lonzaga (2015) as helpful hints for our further consideration: Trace**
 484 **velocities below 0.340 km/s indicate signal refraction in the troposphere and lower**
 485 **stratosphere. Trace velocities between 0.340 and 0.380 km/s are typical for signals ducted**
 486 **in the waveguide between the ground and the upper stratosphere. Signals traveling in the**
 487 **thermospheric waveguide arrive with trace velocities larger than 0.380 km/s.**

488 **The high trace velocities recorded at PVC I disprove signal refraction in the lower**
 489 **stratosphere. Hence, it is unlikely that the signals arrive through a waveguide that can form**
 490 **at the tropopause – lower stratosphere by the effect of the streamer event.**

491 **Like in the November 2020 case, signal propagation above the eastern North Atlantic and**
 492 **Western and Central Europe is investigated using the InfraGA/GeoAc tools.** Propagation of
 493 the 0.2 Hz signal is modelled **on 10th March at 03 UTC.** A source is located at 55°N 15°W


494 at a distance of ~2000 km from the stations. This scenario represents signal propagation
495 from the central North Atlantic. The other source is located at 55°N 0°latitude representing
496 propagation of microbaroms from the North Sea. The distance from the stations is ~1000
497 km.

498 Eastward signal ducting is enabled in the stratospheric and thermospheric waveguides
499 from both sources to the stations. Strong signal absorption in the thermospheric waveguide
500 likely disables thermospheric arrivals to the PPCI and WPCI. We assume that signals
501 ducted in the upper stratosphere are detected. The other eastward waveguide occurs near
502 the tropopause, formed by the eastward to south-eastward jet-stream above the eastern
503 North Atlantic and Western Europe at latitudes 50 – 60°N Figure 9. Signals from the source
504 in the North Atlantic are predicted to travel in the tropopause waveguide to distances of
505 1000-1100 km. The signal attenuation is low in the tropopause waveguide; the relatively
506 short distance under the waveguide influence is determined by the location and extent of
507 the jet-stream disturbance. The tropopause/lower stratosphere arrivals can be detected
508 mainly on the British Isles. The waveguide does not reach to PPCI and WPCI stations (see
509 supplementary materials).


510 Signals emitted by a source in the North Sea can propagate through the tropopause
511 waveguide. The signals propagate to the south-east and are predicted to reach Central
512 Europe. The tropopause/lower stratosphere arrivals are represented by red dots in Figure
513 10. The influenced regions are to the south-west from PPCI and WPCI, several hundreds
514 of kilometres distant from the stations. The approximation of infrasound propagation
515 obtained from the raytracing is in accord with observations. The trace velocities at PPCI of
516 0.377 km/s indicate infrasound propagation in the waveguide formed between the ground
517 and the upper stratosphere rather than in the waveguide near the tropopause.

518 Like in the November 2020 case, infrasound arrivals from the North Atlantic to the
519 stations PPCI and WPCI in Central Europe are not influenced by the waveguide at the
520 tropopause – lower stratosphere. Observed trace velocities fluctuate within or close above
521 the limits that indicate infrasound propagation in the upper stratosphere during the streamer
522 event and both adjacent quiet periods.

523



524
 525 Fig.9. Wind field at the pressure level of 250 hPa on 10 March 2021 at 03 UTC. A
 526 disturbance of the jet-stream above the eastern North Atlantic and the British Isles is caused
 527 by the streamer event.
 528



529
 530 **Fig.10 Modelled** infrasound propagation from a point source located at 55°N and 0°longitude
 531 (red asterisk) on 10th March 2021 at 03 UTC. **Colobar refers to the turning height (maximum**
 532 **height) of the signal. Red indicates signal propagation in the waveguide formed near the**
 533 **tropopause (altitudes around 10 km), arrivals through the stratospheric waveguide are in**

534 yellow (altitudes around 40-50 km) and arrivals through the thermospheric waveguide are in
535 blue (altitudes above 100 km). Black triangles represent infrasound arrays WBCI (the left
536 triangle) and PSCI (the right triangle).

537

538


539 3.2 Results and discussion of gravity waves in the troposphere and ionosphere

540

541 3.2.1 Investigation of GWs measured on the ground by WBCI array of micro- 542 barometers.


543 Figure 12 shows the RMS amplitudes of pressure fluctuations in the period range 5-60 min
544 recorded from November 1 to November 9, 2020. This interval covers a distinct streamer
545 event that occurred from November 3 to November 7. The results of propagation analysis are
546 shown in Figure 13, which displays the phase velocities and azimuths of GWs. Only results
547 that satisfied the criterion ($dv/v < 0.5$) and ($dAZ < 10^\circ$) and ($p_{RMS} > 0.02$ Pa) are presented,
548 where dv/v , dAZ , p_{RMS} are the relative uncertainty of GW phase velocity, uncertainty of
549 azimuth and root mean square value of pressure fluctuations in the analysed time interval.
550 Figure 13 demonstrates that there is a tendency for higher phase velocities and occurrence of
551 different azimuths during the streamer event. Therefore, it is useful to compare the GW
552 characteristics during streamer events and calm conditions.

553 Figure 14 shows histograms obtained by a statistical analysis. The RMS amplitudes of
554 pressure fluctuations in the period range 5 – 60 min, phase velocities and azimuths were
555 investigated separately for calm conditions (upper plots) and for streamer events listed in
556 Table 1 (bottom plots) with a 1-hour time resolution. The solid vertical lines mark lower (Q1)
557 and upper (Q3) quartiles. The dashed vertical lines depict boundaries for large ($Q3 + 1.5 \cdot (Q3 -$
558 $Q1)$) and extreme ($Q3 + 3 \cdot (Q3 - Q1)$) values. A difference between histograms for RMS
559 pressure fluctuations and azimuths obtained for calm and disturbed conditions is obvious.
560 During the streamer events the azimuths are distributed more randomly and more extreme
561 pressure amplitudes can be observed. A minor difference is also observed for phase
562 velocities.




563

564 **Figure 12** Amplitude of GWs recorded by WBCI from 2020-11-01 to 2020-11-09



565

566 **Figure 13** Propagation velocity and azimuth of GWs recorded by WBCI from 2020-11-01 to
 567 2020-11-09



568


569 **Figure 14** GW characteristics (RMS of pressure fluctuations, phase velocity and azimuth)
 570 for calm periods (upper plots) and streamer and streamer like events (bottom plots) for 2020
 571 and winter 2021. The red vertical lines mark lower (Q1) and upper (Q3) quartiles. The dashed
 572 magenta vertical lines depict boundaries for large ($Q3 + 1.5 \cdot (Q3 - Q1)$) and extreme ($Q3 + 3 \cdot (Q3 -$
 573 $Q1)$) values.

574

575 **3.2.2 Investigation of GWs measured in the ionosphere by continuous Doppler**
 576 **sounding system (CDS)**

577 The 2D propagation analysis of GWs was performed using the 2D versions of methods
 578 mentioned in Section 2 and in detail described by Chum and Podolská (2018). As discussed in
 579 Section 2 and by (Chum et al., 2021), the 2D propagation analysis makes it possible to
 580 analyze much larger number of time intervals than the 3D analysis. The propagation analysis
 581 obtained for the interval from 1st November to 9th November 2020, which covers the
 582 significant streamer event that occurred from 3rd November 2020 to 7th November 2020, is
 583 presented in Figure 15. Only results that satisfied the criteria ($dv/v < 0.2$) and ($dAZ < 20^\circ$) and

584 ($f_{\text{DRMS}} > 0.05$ Hz) and ($C_{\text{max}} < 0.5$) are presented, where dv/v is the relative uncertainty of GW
585 phase velocity, dAZ is the azimuth uncertainty, f_{DRMS} is the root mean square of the Doppler
586 shift in the analysed time interval and C_{max} is the maximum in the normalized energy map for
587 the best beam (slowness) search; C_{max} is 1 for identical signals (Chum and Podolská, 2018). It
588 is considered that signals are not sufficiently correlated (coherent) for reliable propagation
589 analysis if $C_{\text{max}} < 0.5$ (Chum et al., 2021). The velocities and azimuth obtained by observation
590 at 3.59 MHz are in red, whereas the values based on measurements at 4.65 MHz are in blue.
591 Obviously, the observations at 3.59 MHz mostly correspond to the nighttime, whereas
592 observations at 4.65 MHz were mostly made during the daytime. The 4.65 MHz signal did not
593 reflect from the ionosphere (escaped to the outer space) at night due to the low critical
594 frequency of the ionosphere. On the other hand, the 3.59 MHz signal mostly reflected during
595 the day from the ionospheric E layer and the Doppler shift was negligible, difficult to
596 analyse. The GWs usually propagated roughly poleward at night and roughly equatorward
597 during the daytime. This is fully consistent with the statistical investigation (Chum et al.,
598 2021) which showed that propagation directions of GWs in the ionosphere exhibit diurnal and
599 seasonal behaviour and are mainly controlled by the neutral winds in the thermosphere.



600

601 **Figure 15** Propagation velocity and azimuth of GWs in the ionosphere obtained using CDS
602 measurements from 2020-11-01 to 2020-11-09. The velocities and azimuth obtained by
603 observation at 3.59 MHz are by red, whereas the values based on measurements at 4.65 MHz
604 are by blue.

605 Based on the analysis of the GW observed in the ionosphere during the streamer event and
606 on the previous statistical analysis, we conclude that no obvious signature related to streamer
607 event was observed for the propagation of GW the ionosphere.

608 It should be also mentioned that the phase velocities of GW measured on the ground (Figure
609 8) and at heights around 200 km in the ionosphere differ. There are several reasons for that.
610 First, the observed horizontal phase velocities depend on the elevation angle of GW
611 propagation and on the ambient temperature as follows from the dispersion relation (the
612 temperature enters the dispersion relation via the buoyancy frequency and the scale height).
613 The temperature in the ionosphere/thermosphere is several times higher than in the
614 troposphere. The elevation angles might change during the upward propagation of GWs,
615 depending on the wind and temperature profile. Second, GWs propagate with a tilt, not
616 vertically upward. It is therefore highly probable that the sources of the GWs observed in the
617 troposphere and ionosphere are different. Moreover, GW can break during their propagation
618 upward and secondary gravity waves might be observed in the ionosphere.

619 **4) Conclusion and discussion**

620 The focus of this study was to test independent types of observations like Doppler sounding
621 and microbarograph measurements for an analysis of GW behavior during streamer events,
622 which are strongly connected with PW or GW and the large scale mass transport of ozone and
623 that is why it can be very interesting for studies of atmospheric dynamics.

624 We also investigate effects of the streamer events on infrasound propagation. Streamer
625 events are significant disturbances to circulation in the tropopause/lower stratosphere region.
626 Modifications of infrasound ducting in the atmosphere can therefore be expected. Indeed, the
627 InfraGA/GeoAc raytracing tools predict that a waveguide develops at the tropopause during
628 the analyzed streamer events in November 2020 and in March 2021 the direction of which is
629 determined by the disturbed jet-stream and varies from event to event. The tropopause
630 waveguide ducts infrasound up to distances of several hundreds to a thousand of km from the
631 source in a limited azimuth range. The azimuth sector of the extent of 50 – 60° is influenced
632 in the analysed cases. In accord with the model predictions, streamer event related phenomena

633 were not found in infrasound detection at the infrasound stations PPCI and WBCI in Central
634 Europe during the studied cases. The observability of streamer event signatures in infrasound
635 arrival parameters therefore strongly depends on the mutual position of the source, the
636 streamer event disturbance of the tropopause jet-stream and the infrasound station. It can be
637 recommended for future studies to use a dense network of infrasound arrays that covers
638 various directions and distances from the streamer event. Due to the typical occurrence of the
639 streamer events over the North Atlantic, infrasound stations in Western Europe can be of
640 particular interest.

641 Supplementary ground-based measurements of GW using the WBCI array in the
642 troposphere showed that GW propagation azimuths were more random during streamer and
643 streamer-like events compared to those observed during calm conditions **as can be seen from**
644 **the plots in Figure 14.** On the other hand, the GW propagation characteristics observed in the
645 ionosphere by CDS during streamer events did not differ from those expected for the given
646 time period, based on previous statistical studies (Chum et al., 2021).

647 The results therefore indicate that streamers in the stratosphere might lead to changes in wave
648 propagation in the troposphere. The impact on the ionosphere was not confirmed, but cannot
649 be excluded due to sparse and localized observations of GW activity. In general, to validate the
650 preliminary results obtained in this study, a denser measurement network and more streamer
651 events need to be analyzed.

652

653 **Data availability:**

654 ozone column measurements (TO3) which are available as a service by DLR at
655 <https://atmos.eoc.dlr.de/>

656 Ground to space model vertical atmospheric profiles were obtained at
657 <https://g2s.ncpa.olemiss.edu/>; accessed on 27 January – 4 February 2024

658

659 The WAVEWATCHIII[®] wave-action model data were accessed via ftp at
660 polar.ncep.noaa.gov/waves/JCOMM/2020 on 13-14 March 2023.

661

662 The Deutscher Wetterdienst synoptic charts were accessed at
663 https://www2.wetter3.de/archiv_dwd_dt.html on 3 February 2024.

664

665 **Author contributions**

666 MK and LK create the idea of manuscript; JCh, MK, TS, LK, and KP suggest the datasets and
667 methods; TS, JCh, LK, KP and FT analyzed the data; MK wrote the manuscript draft; JCh,
668 TS, LK and KP reviewed and edited the manuscript.

669 **Competing interests**

670 The authors declare that they have no conflict of interest.

671

672 **Acknowledgement**

673 The DTK-GPMCC software was kindly provided by Commissariat à l'énergie atomique et
674 aux énergies alternatives, Centre DAM-Île-de-France, Département Analyse, Surveillance,
675 Environnement, Bruyères-le-Châtel, F91297 Arpajon, France.

676 The authors are grateful to Dr. Phil Blom and Los Alamos National Laboratory for opening
677 the InfraGA/GeoAc tools to the public.

678 **Financial support:** This study is supported by LISA project- Lidar measurements to
679 Identify Streamers and analyze Atmospheric waves, AEOLUS-INNOVATION, Contract No.
680 4000133567/20/I-BG

681

682 **References**

683 Assink, J.D., Waxler, R., Smets, P., Evers, L.G. (2014). Bidirectional infrasonic ducts
684 associated with sudden stratospheric warm-ing events. *J. Geophys. Res. Atmos.* 119,1140-
685 1153.

686 Bittner, M., Höppner, K., Pilger, C., Schmidt, C. (2010). Mesopause temperature
687 perturbations caused by infrasonic waves as a potential indicator for the detection of
688 tsunamis and other geo-hazards. *Nat. Hazards Earth Syst. Sci.*, 10, 1431-1442. [www.nat-](http://www.nat-hazards-earth-syst-sci.net/10/1431/2010/doi:10.5194/nhess-10-1431-2010)
689 [hazards-earth-syst-sci.net/10/1431/2010/doi:10.5194/nhess-10-1431-2010](http://www.nat-hazards-earth-syst-sci.net/10/1431/2010/doi:10.5194/nhess-10-1431-2010)

690 Blanc, E. (1985). Observations in the upper atmosphere of infrasonic waves from natural or
691 artificial sources: A summary. *Ann. Geophys.*, 3, 673-688.

692 Blixt, E.M., Nasholm, S.P., Gibbons, S.J., Evers, L.G., Charlton-Perez, A.J., Orsolini, Y.J.,
693 Kvaerna, T. (2019). Estimating tropo-spheric and stratospheric winds using infrasound from
694 explosions. *J. Acoust. Soc. Am.* 146:2.

695 Blom, P., Waxler, R. (2012). “Impulse propagation in the nocturnal boundary layer: Analysis
696 of the geometric component”. *J. Acoust. Soc. Am.*, **131**, 3680 – 3690. doi:
697 [10.1121/1.3699174](https://doi.org/10.1121/1.3699174).

698 Blom, P. (2019). “Modeling infrasonic propagation through a spherical atmospheric layer:
699 Analysis of the stratospheric pair.” *J. Acoust. Soc. Am.*, **145**, 2198–2208. doi:
700 [10.1121/1.5096855](https://doi.org/10.1121/1.5096855).

701 Bondár I., T. Šindelářová, D. Ghica, U. Mitterbauer, A. Liashchuk, J. Baše, J. Chum, C.
702 Czanik, C. Ionescu, C. Neagoe, M. Pásztor, A. Le Pichon (2022), Central and Eastern
703 European Infrasound Network: Contribution to Infrasound Monitoring, *Geophys. J. Int.*,
704 ggac066, <https://doi.org/10.1093/gji/ggac066>

705 Brachet, N., Brown, D., Le Bras R., Cansi, Y., Mialle, P., Coyne, J. (2010). Monitoring the
706 Earth’s Atmosphere with the Global IMS Infrasound Network. In: Le Pichon, A., Blanc, E.,
707 Hauchecorne A. (Eds.), *Infrasound Monitoring for Atmospheric Studies*. Springer
708 Science+Business Media B.V., 77-118. Doi: 10.1007/978-1-4020-9508-5_3

709 Campus, P., Christie, D.R. (2010). Worldwide Observations of Infrasonic Waves. In: Le
710 Pichon, A., Blanc, E., Hauchecorne A. (Eds.), *Infrasound Monitoring for Atmospheric*
711 *Studies*. Springer Science+Business Media B.V., 185234-118. Doi: 10.1007/978-1-4020-
712 9508-5_6

713 Cansi, Y., 1995. An automatic seismic event processing for detection and location: The
714 P.M.C.C. method. *Geophys. Res. Lett.* 22, 1021-1024. doi: 10.1029/95GL00468

715 Ceranna, L., Matoza, R., Hupe, P., Le Pichon, A., Landès, M., (2019). Systematic Array
716 Processing of a Decade of Global IMS Infrasound Data. In: Le Pichon, A., Blanc, E.,
717 Hauchecorne, A. (eds) *Infrasound Monitoring for Atmospheric Studies*. Challenges in
718 *Middle Atmospheric Dynamics and Societal Benefits*. Springer Nature Switzerland AG.

719 Chum J, Podolská K (2018) 3D analysis of GW propagation in the ionosphere. *Geophysical*
720 *Research Letters*, 45, 11,562–11,571, <https://doi.org/10.1029/2018GL07969>

721 Chum, J., Podolská, K., Rusz, J., Baše, J., Tedoradze, N. (2021), Statistical investigation of
722 gravity wave characteristics in the ionosphere. *Earth Planets Space* 73, 60,
723 <https://doi.org/10.1186/s40623-021-01379-3>

724 Czech microbarograph network, <https://doi.org/10.7914/SN/C9>

725 Drob, D. P., Picone, J. M., Garcés, M. (2003). Global morphology of infrasound propagation.
726 *J. Geophys. Res. Atmospheres*, **108** (D21). doi: [10.1029/2002JD003307](https://doi.org/10.1029/2002JD003307).

727 Evers, L. G., Siegmund, P. (2009). Infrasonic signature of the 2009 major sudden
728 stratosphericwarming, *Geophys. Res. Lett.*, 36, L23808, doi:10.1029/2009GL041323

729 Evers, L.G., Haak, H.W. (2010). The Characteristics of Infrasound, its Propagation and Some
730 Early History. In: Le Pichon, A., Blanc, E., Hauchecorne, A. (eds) *Infrasound Monitoring for*
731 *Atmospheric Studies*. Springer, Dordrecht.

732 Evers, L. G., van Geyt, A. R. J. , Smets, P., Fricke, J.T. (2012). Anomalous infrasound
733 propagation in a hot stratosphere and the existence of extremely small shadow zones, *J.*
734 *Geophys. Res.*, 117, D06120, doi:10.1029/2011JD017014.

735

736 Eyring, V., Dameris, M., Grewe, V., Langbein, I., & Kouker, W. (2002). Climatologies of
737 streamer events derived from a transport model and a coupled chemistry-climate model.

738 Fritts, D.C. & Alexander, M.J., (2003). Gravity wave dynamics and effects in the middle
739 atmosphere. *Rev. Geophys.*, 41 (1), 1003.

740 Garcès, M., Willis, M., Hetzer, C., Le Pichon , A., Drob, D., (2004). On using ocean swells
741 for continuous infrasonic measurements of winds and temperature in the lower, middle, and
742 upper atmosphere. *Geophys. Res. Lett.* 31, L19304. doi: 10.1029/2004GL020696

743 Garcès, M.A., (2013). On infrasound standards, part 1: Time, frequency, and energy scaling.
744 *InfraMatics* 2, 13-35. doi: 10.4236/inframatics.2013.22002

745 Georges, T.M. (1968). H. F. Doppler studies of travelling ionospheric disturbances. *J.*
746 *Atmos.Terr. Phys.*, 30, 735-746.

747 Gerlach, C., Földvary, L., Švehla, D., Gruber, T., Wermuth, M., Sneeuw, N., ... &
748 Steigenberger, P. (2003). A CHAMP-only gravity field model from kinematic orbits using the
749 energy integral. *Geophysical Research Letters*, 30(20).

750 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., ... &
751 Thépaut, J. N. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal*
752 *Meteorological Society*, 146(730), 1999-2049.

753 Hupe, P., Ceranna, L., Pilger, C., de Carlo, M., Le Pichon, A., Kaifler, B., Rapp, M. (2019).
754 Assessing middle atmosphere weather models using infrasound detections from microbaroms.
755 *Geophys. J. Int.*, 216, 1761–1767 doi: 10.1093/gji/ggy520

756 James, P. M. (1998): A climatology of ozone mini-holes over the Northern Hemisphere.
757 *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 18, 12:
758 12871303

759 Kramer, R, S. Wüst, and M. Bittner (2016). Investigation of gravity wave activity based on
760 operational radiosonde data from 13 years (1997-2009): Climatology and possible induced
761 variability, *Journal of Atmospheric and Solar-Terrestrial Physics* 140, 23–33;
762 <http://dx.doi.org/10.1016/j.jastp.2016.01.014>

763 Kramer, R., S. Wüst, C. Schmidt, and M. Bittner (2015). Gravity wave characteristics in the
764 middle atmosphere during the CESAR campaign at Palma de Mallorca in 2011/2012: Impact
765 of extratropical cyclones and cold fronts, *Journal of Atmospheric and Solar-Terrestrial*
766 *Physics* 128 (2015) 8–23, <http://dx.doi.org/10.1016/j.jastp.2015.03.001>

767 Kai Ming Huang, Shao Dong Zhang, Fan Yi, (2010). Reflection and transmission of
768 atmospheric gravity waves in a stably sheared horizontal wind field, *Journal of Geophysical*
769 *Research: Atmospheres*, 10.1029/2009JD012687, **115**, D16,

770 Landès, M., Ceranna, L., Le Pichon, A., & Matoza, R. S. (2012). Localization of microbarom
771 sources using the IMS infrasound network. *Journal of Geophysical Research:*
772 *Atmospheres*, 117(D6).

773 Le Pichon, A., Cansi, Y. (2003). PMCC for infrasound data processing. *InfraMatics* 02, 1-9.

774 Le Pichon, A., Blanc, E., (2005). Probing high-altitude winds using infrasound. *J. Geophys.*
775 *Res.*, 110, D20104. doi: 10.1029/2005JD006020

776 Le Pichon, A., Ceranna, L., Garcès, M., Drob, D., Millet, C., (2006). On using infrasound
777 from interacting ocean swells for global continuous measurements of winds and temperature
778 in the stratosphere. *J. Geophys. Res.*, 111, D11106. doi: 10.1029/2005JD006690

779 Le Pichon, A., Vergoz, J., Blanc, E., Guilbert, J., Ceranna, L., Evers, L., Brachet, N., (2009).
780 Assessing the performance of the International Monitoring System's infrasound network:
781 Geographical coverage and temporal variabilities. *J. Geophys. Res.* 114, D08112. doi:
782 10.1029/2008JD010907

783 Leovy, C. B., Sun, C. R., Hitchman, M. H., Remsberg, E. E., Russell III, J. M., Gordley, L.
784 L., ... & Lyjak, L. V. (1985). Transport of ozone in the middle stratosphere: Evidence for
785 planetary wave breaking. *Journal of Atmospheric Sciences*, 42(3), 230-244.

786 Lonzaga, J.B., (2015). A theoretical relation between the celerity and trace velocity of
787 infrasonic phases, *J. Acoust. Soc. Am.*, 138, EL242-EL247.
788 <http://dx.doi.org/10.1121/1.4929628>

789 Loyola D.G., Koukouli M.E., Valks P., Balis D.S., Hao N., van Roozendaal M., Spurr R.J.D.,
790 Zimmer W., Kiemle S., Lerot C., Lambert J.-C. (2011) The GOME-2 total column ozone
791 product: Retrieval algorithm and ground-based validation, *Journal of Geophysical Research*,
792 vol. 116, D07302, Wiley-Blackwell

793 Marty, J., (2019). The IMS Infrasound Network: Current Status and Technological
794 Developments, in: Le Pichon, A., Blanc, E., Hauchecorn, A. (Eds.), *Infrasound Monitoring*
795 *for Atmospheric Studies. Challenges in Middle Atmosphere Dynamics and Societal Benefits.*
796 Springer Nature Switzerland AG, pp. 3–62. doi:10.1007/978-3-319-75140-5_1

797 McIntyre, M. E., & Palmer, T. N. (1983). Breaking planetary waves in the stratosphere.
798 *Nature*, 305(5935), 593-600.

799 Munro, R., Eisinger, M., Anderson, C., Callies, J., Corpaccioli, E., Lang, R., ... & Albinana,
800 A. P. (2006, June). GOME-2 on MetOp. In *Proc. of The 2006 EUMETSAT Meteorological*
801 *Satellite Conference, Helsinki, Finland (Vol. 1216, p. 48).*

802 Munro, R., et al. (2016): The GOME-2 instrument on the Metop series of satellites:
803 instrument design, calibration, and level 1 data processing – an overview, *Atmos. Meas.*
804 *Tech.*, 9, 1279–1301, <https://doi.org/10.5194/amt-9-1279-2016>.

805 Peters, D., Hoffmann, P., & Alpers, M. (2003). On the appearance of inertia-gravity waves on
806 the north-easterly side of an anticyclone. *Meteorologische Zeitschrift*, 12(1), 25-35

807 Polvani, L. M., & Plumb, R. A. (1992). Rossby wave breaking, microbreaking, filamentation,
808 and secondary vortex formation: The dynamics of a perturbed vortex. *Journal of Atmospheric*
809 *Sciences*, 49(6), 462-476.

810 Pramitha, M., Venkat Ratnam, M., Taori, A., Krishna Murthy, B. V., Pallamraju, D., and
811 Vijaya Bhaskar Rao, S. (2015). Evidence for tropospheric wind shear excitation of high-
812 phase-speed gravity waves reaching the mesosphere using the ray-tracing technique, *Atmos.*
813 *Chem. Phys.*, 15, 2709–2721, <https://doi.org/10.5194/acp-15-2709-2015>.

814 Rauthe, M., Gerding, M., Höffner, J., & Lübken, F. J. (2006). Lidar temperature
815 measurements of gravity waves over Kühlungsborn (54° N) from 1 to 105 km: A winter-
816 summer comparison. *Journal of Geophysical Research: Atmospheres*, 111(D24).

817

818 Wüst, S., & Bittner, M. (2006). Non-linear resonant wave–wave interaction (triad): Case
819 studies based on rocket data and first application to satellite data. *Journal of atmospheric and*
820 *solar-terrestrial physics*, 68(9), 959-976.

821

822 Wüst, S., Offenwanger, T., Schmidt, C., Bittner, M., Jacobi, C., Stober, G., Yee, J.H.,
823 Mlynczak, M. G. & Russell III, J. M. (2018). Derivation of gravity wave intrinsic parameters
824 and vertical wavelength using a single scanning OH (3-1) airglow spectrometer. *Atmospheric*
825 *Measurement Techniques*, 11(5), 2937-2947.

826

827 Smets, P.S.M., Evers, L.G. (2014). The life cycle of a sudden stratospheric warming from
828 infrasonic ambient noise observations, *J. Geophys. Res. Atmos.*, 119, 12,084-12,099

829 Spurr, R., Loyola, D., Heue, K. P., Van Roozendaal, M., & Lerot, C. (2022). S5P/TROPOMI
830 Total Ozone ATBD. Deutsches Zentrum für Luft- und Raumfahrt (German Aerospace
831 Center), Weßling, Germany, Tech. Rep. S5P-L2-DLR-ATBD-400A.

832 Sutherland, L.C., Bass, H.E., (2004). Atmospheric absorption in the atmosphere up to 160
833 km. *J. Acoust. Soc. Am.*, 115, 1012–1032. <https://doi.org/10.1121/1.1631937>

834 Szuberla, C.A.L., Olson, J.V., (2004). Uncertainties associated with parameter estimation in
835 atmospheric infrasound rays. *J. Acoust. Soc. Am.* 115, 253-258. doi: 10.1121/1.1635407

836 Veefkind, J. P., Aben, I., McMullan, K., Förster, H., De Vries, J., Otter, G., ... & Levelt, P. F.
837 (2012). TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global
838 observations of the atmospheric composition for climate, air quality and ozone layer
839 applications. *Remote sensing of environment*, 120, 70-83.

840

841