

Evaluation of the first year of Pandora NO₂ measurements over Beijing and application to satellite validation

Ouyang Liu^{1,2}, Zhengqiang Li^{1,*}, Yangyan Lin^{2,3}, Cheng Fan¹, Ying Zhang¹, Kaitao Li¹, Peng Zhang⁴, Yuanyuan Wei¹, Tianzeng Chen^{2,4}, Jiantao Dong^{1,5}, Gerrit de Leeuw^{1,6}

5 ¹ State Environmental Protection Key Laboratory of Satellite Remote Sensing & State Key Laboratory of Remote Sensing Science, Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing 100101, China

² University of Chinese Academy of Sciences, Beijing 100049, China

³ Key Laboratory of Watershed Geographic Sciences, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, Nanjing 210008, China

10 ⁴ State Key Joint Laboratory of Environment Simulation and Pollution Control, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

⁵ Satellite Application Center for Ecology and Environment, Ministry of Ecology and Environment of People's Republic of China, Beijing 100094, China

⁶ Royal Netherlands Meteorological Institute (KNMI), R & D Satellite Observations, 3730 AE De Bilt, The Netherlands

15 *Correspondence to:* Zhengqiang Li (lizq@radi.ac.cn)

Abstract. Nitrogen dioxide (NO₂) is a highly photochemically reactive gas, has a lifetime of only a few hours, and at high concentrations it is harmful to human beings. Therefore, it is important to monitor NO₂ with high-precision, time-resolved instruments. To this end, a Pandora spectrometer has been installed on the roof of the laboratory building of the Aerospace Information Research Institute of the Chinese Academy of Sciences in the Olympic Park, Beijing, China. The concentrations of trace gases (including NO₂, HCHO, O₃) measured with Pandora are made available through the open-access Pandora data base (<https://data.pandonia-global-network.org/Beijing-RADI/Pandora171s1/>). In this paper, an overview is presented of the Pandora total and tropospheric NO₂ vertical column densities (VCDs) and surface concentrations collected during the first year of operation, i.e., from August, 2021, to July, 2022. The data show that NO₂ concentrations were high in the winter and low in the summer, with diurnal cycle where the concentrations reach a minimum during day time. The concentrations were significantly lower during the 2022 Winter Olympics in Beijing, showing the effectiveness of the emission control measures during that period. The Pandora observations show that during northerly winds clean air is transported to Beijing with low NO₂ concentrations, whereas during southerly winds pollution from surrounding areas is transported to Beijing and NO₂ concentrations are high. The contribution of tropospheric NO₂ to the total NO₂ VCD varies significantly on daily to seasonal time scales, i.e., monthly averages vary between 50% and 60% in the winter and between 60% and 70% in the spring and autumn. The comparison of Pandora-measured surface concentrations with collocated in situ measurements using a Thermo Scientific 42i-TL Analyzer shows that the Pandora data are low and that the relationship between Pandora-derived surface concentrations and in situ measurements are different for low and high NO₂ concentrations. Explanations for these differences are offered in terms of measurement techniques and physical (transport) phenomena. The use of Pandora total and tropospheric NO₂ VCDs for validation of collocated TROPOMI data, resampled to 100×100 m², shows that although on

35 average the TROPOMI VCDs are slightly lower, they are well within the expected error for TROPOMI of $0.5 \text{ Pmolec} \cdot \text{cm}^{-2} + (0.2 \text{ to } 0.5) \cdot \text{VCD}_{\text{trop}}$ ($1 \text{ Pmolec} \cdot \text{cm}^{-2} = 1 \times 10^{15} \text{ molec} \cdot \text{cm}^{-2}$). The location of the Pandora instrument within a sub-orbital TROPOMI pixel of $3.5 \times 5.5 \text{ km}^2$ may result in an error in the TROPOMI-derived tropospheric NO_2 VCD between 0.223 and 0.282 $\text{Pmolec} \cdot \text{cm}^{-2}$, i.e., between 1.7% and 2%. In addition, the data also show that the Pandora observations at the Beijing-RADI site are representative for an area with a radius of 10 km.

40 **1 Introduction**

NO_2 is a trace gas that plays an important role in atmospheric chemistry (Seinfeld and Pandis, 1998), such as the O_3 - NO_x -VOC sensitivity (e.g., Wang et al., 2021; Liu and Shi, 2021), the formation of aerosols (Behera and Sharma, 2011) and thus air quality (de Leeuw et al., 2021). NO_2 is also a precursor for the production of aerosol and ozone and therefore indicated as an essential climate variable (ECV) (<https://gcos.wmo.int/en/essential-climate-variables/precursors>; last visited 11th July
45 2023). Sources of tropospheric NO_2 include anthropogenic sources such as coal-fired power plants, motor vehicle emissions and industrial chemical production (Carslaw and Beevers, 2005; Felix and Elliott, 2014), and natural sources such as stratospheric NO_x intrusion, bacterial and volcanic action (Mather et al., 2004), and lightning (Zhang et al., 2020). NO_2 is removed by chemical reactions and diluted by horizontal and vertical transport (Jorba et al., 2012; Dai et al., 2022). Thus, the NO_2 concentration decreases with distance to the emission sources. Most of the NO_2 is located at altitudes between 0 and 3
50 km with a small amount in the upper troposphere and stratosphere due to chemical reactions such as the oxidation of nitrous oxide (Brasseur and Nicolet, 1973; Grenfell et al., 2006; Dirksen et al., 2011; Herman et al., 2019). NO_2 is a short-lived trace gas with a chemical lifetime of $3.8 \pm 1.0 \text{ h}$ during the summer (Liu et al., 2016), increasing during colder conditions in winter and at higher altitudes (Herman et al., 2018). Together with the multitude of localized (power plants, factories) and diffuse (traffic) sources, this short lifetime results in a large spatial and temporal variation of the NO_2 concentrations (Weber and
55 Bylicki, 1987; Sivakumaran et al., 2001; Zhao et al., 2022a). To catch this variability, observations are needed with high temporal resolution together with good spatial resolution.

Information on NO_2 concentrations can be obtained by using different methods and techniques, i.e., ground-based in situ or remote sensing measurements or satellite remote sensing. Each of these techniques provides different types of data with their own advantages and disadvantages. Instruments used for ground-based observations have the advantage that they can easily
60 be serviced, have high and well-known accuracy, can measure continuously and thus provide good temporal coverage during day and night. In China, ground-based in situ measurements of NO_2 concentrations are available, together with other species, from the National Real-time Air Quality Publishing Platform public website for air quality (AQ) monitoring data, maintained by the China National Environmental Monitoring Center (CNEMC) of the Ministry of Ecology and Environment of China (<http://www.cnemc.cn/>; last access 11th July 2023) (Zhai et al., 2019; Li et al., 2020; Xie et al., 2005). This network includes
65 more than 2000 stations all over China, but they are mainly located in densely populated and urban centers. In situ data are representative for concentrations near the surface and within a certain distance from the observation site, especially for a

short-lived species like NO₂ with many local emission sources such as traffic using fossil fuel powered engines and households. This leaves large gaps in the data coverage, i.e., there are no data for large areas outside the urban and industrial agglomerations.

70 Ground-based remote sensing of atmospheric NO₂ concentrations can be made using instruments such as MAX-DOAS (Wagner et al, 2010) or Pandora (Herman et al., 2009). Both instruments are spectrometers with a mode viewing the sun directly, and another one viewing scattered radiation (sky) at different angles, providing information on the vertical distribution. In the current study, data from a Pandora instrument will be used (see Section 2.2 for more detail on the instrument).

75 The gaps in the spatial distributions can be filled with data from satellite observations, i.e., using dedicated satellite-based instruments (spectrometers) providing wide spatial coverage. Such instruments have been developed during the last three decades with enormous improvement in both spatial resolution and accuracy, i.e., from the Global Ozone Monitoring Experiment (GOME, launched on the European Remote Sensing satellite ERS-2 in 1995), the SCanning Imaging Absorption spectroMeter for Atmospheric Cartography (SCIAMACHY; launched on the Environmental Satellite ENVISAT in 2002),
80 the Ozone Monitoring Instrument (OMI; launched on the AURA satellite in 2004), GOME-2 (first launched on the Meteorological Operational satellite METOP-A in 2006), to the TROPOspheric Monitoring Instrument (TROPOMI; launched on the Sentinel-5 Precursor (S-5p) satellite in 2017). However, although these sensors provide data with daily near-global coverage, they have only one single overpass each day (more at high latitudes) which does not capture diurnal variation (Bovensmann et al., 1999; Burrows et al., 1999; Levelt et al., 2006; Levelt et al., 2018; Verhoelst et al., 2021). The
85 primary NO₂ product from such satellites is the total vertical column density (VCD), (i.e., column-integrated NO₂ concentrations) which can be separated into tropospheric and stratospheric VCDs by using a chemical transport model (e.g., van Geffen et al., 2022b). The accuracy of satellite data is usually less good than from ground-based data because corrections need to be made for contributions from the surface and from other atmospheric constituents contributing to the reflected solar radiation measured by the instrument at the top of the atmosphere (TOA). In addition, satellite data need regular
90 validation to account for post-launch degradation. In this study, TROPOMI data will be used as discussed in Section 2.2.

For the validation of satellite-retrieved NO₂ data, the PANDONIA Global Network (PGN) has been established in 2018 to provide “long-term quality observations of total column and vertically resolved concentrations of a range of trace gases” (<http://www.pandonia-global-network.org/>, last accessed: 11th July 2023). The information on NO₂, O₃ and HCHO provided by PGN is obtained using Pandora instruments and is publicly available. Most PGN sites have been established in North
95 America and Europe, with fewer data from Japan, South Korea and some other countries, thus leaving large areas uncovered. In particular, until recently there were no publicly available data in China.

As a first step to fill this gap, a Pandora instrument has been installed in Beijing on the roof of the Aerospace Information Research Institute of the Chinese Academy of Sciences (AirCAS) laboratory building in the Olympic Park, Beijing, China (116.3786E, 40.0048N). The instrument has been running continuously since the end of July 2021. The observations and
100 data processing follow the Pandora protocol (Cede et al., 2021) and provides high-precision trace gas VCD data in near real-

time. The Pandora instrument, referred to as Beijing-RADI, is the first instrument in China which joined the PGN network and all data are publicly available via the PGN website (<https://data.pandonia-global-network.org/Beijing-RADI/Pandora171s1/>, last accessed: 11th July 2023), within one day of the observations.

In this paper we provide an overview of the results from the first year of NO₂ observations from the Beijing-RADI Pandora instrument, i.e., total VCDs, tropospheric VCDs and surface concentrations. The site and experimental methods are discussed in Section 2. The results are presented in Section 3, where we discuss the variations of NO₂ VCDs and surface concentrations and the contribution of tropospheric NO₂ to the total VCD on hourly, daily, monthly and seasonal time scales. Influences of wind speed and direction on the NO₂ concentrations at the Beijing-RADI site are discussed. Pandora-derived surface concentrations are evaluated by comparison with collocated in situ observations and differences are explained in terms of local production and long-range transport. Pandora NO₂ tropospheric and total VCDs are used for validation of TROPOMI NO₂ data and the spatial representativeness of the Pandora is evaluated. A brief summary of the study and the main conclusions are presented in Section. 4.


2 Data and methodology

2.1 Site description

Beijing is a Megacity with a population of over 21.7 million in 2023 (<https://worldpopulationreview.com/world-cities/beijing-population>, last accessed: 26 October 2023). The vast majority of the people in Beijing live within the fifth ring road (see Figure 1). Beijing is located in the north of the North China Plain (NCP), with the Yanshan and Taihang Mountains to the north and west, respectively. The NCP includes a highly industrialized and urbanized area where the weather conditions are often conducive to the accumulation of air pollution resulting in the frequent occurrence of haze episodes (e.g., Sundström et al., 2012; Sun et al., 2014; Li et al., 2016). The effects of pollution in the NCP on the air quality in Beijing varies with weather conditions. Southern airflow transports pollution to the city, whereas northern airflow transports clean air from the north. Hence the air quality situation may vary strongly, depending on large scale meteorological patterns (e.g., Hou et al., 2020). Furthermore, as the capital city of China, Beijing hosts the Chinese government and many national and international events. During such events, emissions from industry and traffic in and around Beijing are regulated to improve air quality (e.g., Wang et al., 2010; Wang et al., 2016; Fan et al., 2021; Chu et al., 2022).

The AirCAS laboratory building is located at the north side of Beijing between the Fourth and Fifth Ring Roads (see Figure 1). The Beijing-RADI site is located at the roof of this building, at a height of 21m above the ground. The site includes instrumentation for atmospheric observations such as a sun photometer, a lidar and a suite of instruments for in situ measurements of aerosol properties, trace gas concentrations, and for solar radiation. The Pandora instrument has been added in 2021, to measure NO₂ VCDs and surface concentrations. Due to its location in a science park at the edge of the Olympic Park, the Beijing-RADI site is considered a combined urban and suburban background observatory with small local

emissions and large influences from nearby highways, urban activities and industrial pollution advected from the south of Beijing and the NCP.



135 **Figure 1: Digital elevation map of Beijing and surroundings.** The solid black line is the provincial boundary and the solid grey line is the administrative boundary of Beijing on a district basis. The Beijing-RADI site is indicated by the black triangle, at about 12 km north of the administrative center of China (red pentagram). The inset shows Beijing with the ring roads 2-6 (from the center to outside; there is no ring road 1) overlaid on the TROPOMI-derived NO₂ tropospheric VCD re-gridded to a resolution of 100×100 m² and averaged over the period from August 1, 2021, to July 31, 2022.

140 2.2 Instrumentation and auxiliary data

2.2.1 Pandora

Pandora is a sun-viewing instrument at solar zenith angles (SZA) smaller than 80°. The instrument consists of a UV/VIS spectrometer connected to an optical head through a 400 μm core diameter fiber-optic cable (Herman et al., 2009; Cede, 2021). The optical head is mounted on a sun tracker for accurate pointing at the sun (precision 0.013°; Herman et al., 2009).

145 Solar radiation is collected by the front-end optics with a field of view (FOV) of 2.6° for direct-sun observations using a diffuser and with a FOV of 1.5° for sky observations without a diffuser. The vast majority of the light sampled in direct observation mode comes from an angle of 0.5°. The received solar radiation is transmitted through a fiber-optic cable to the spectrometer which is a 2048×64 pixels back-thinned Hamamatsu charge-coupled device (CCD), with a 50 μm entrance slit, and a grating with 1200 lines / mm.

150 Pandora measures spectrally resolved solar radiances at wavelengths between 290 nm and 380 nm, using a UV bandpass filter, and between 280 nm and 525 nm, with a spectral resolution of 0.6 nm and 4.5 times oversampling. In clear-sky conditions, about 4000 spectra are measured in about 80s, including about 20s of dark current measurements between each spectral measurement (Herman et al., 2009; Cede, 2021). These 4000 spectra are averaged to achieve very high signal to

noise ratios (Herman et al., 2019). The spectra are used to determine trace gas amounts using the differential optical
155 absorption spectroscopy (DOAS) technique, i.e., spectral fitting, as described in detail in the ATBD (Cede, 2021). For the
retrieval of NO₂ VCDs, the part of the spectrum between 400 and 440 nm is used.

Using direct sun measurements, information is obtained on the total VCDs of the trace gases. Diffuse (scattered) radiation is
measured at 5 pointing zenith angles (PZAs) in sky mode which, together with the direct sun measurement, provides
information on the tropospheric VCD and on the surface concentrations. The PZAs are 0°, 60°, 75°, 88° and a maximum
160 angle taken as 89°. The measurements are taken in a V shape (all angles are measured twice around a central angle) as
described in Cede (2021).

Details of the Pandora spectrometer instrument can be found from the Pandora project website
<https://Pandora.gsfc.nasa.gov/Instrument/> (last accessed: 11th July 2023), as well as the NASA Pandora website
https://avdc.gsfc.nasa.gov/pub/DSCOVER/Pandora/Web_Pandora/index.html (last accessed: 11th July 2023). The Beijing-
165 RADI Pandora is operated following the PGN operational procedures and data processing, described in detail in Cede (2021),
which also includes the ATBD. In this study, L2 data products for NO₂ (L2H files L2Tot for total VCD and L2Tro for
tropospheric NO₂ and surface concentrations) are used which include all measurements available. The data quality (DQ) has
been checked for a number of criteria which have to be satisfied. Only DQ0 (assured high quality), DQ1 (assured medium
quality), DQ10 (not-assured high quality) and DQ 11 (not-assured medium quality) data are used. DQ2 and DQ12 data (low
170 quality) are available in the data files but their use is not recommended (Cede, 2021). The estimated nominal accuracy is
about $\pm 2.67 \text{ Pmolec} \cdot \text{cm}^{-2}$ (error in fraction for 5.33%) . ($1 \text{ Pmolec} \cdot \text{cm}^{-2} = 1 \times 10^{15} \text{ molec} \cdot \text{cm}^{-2} = 3.745 \times$
 $10^{-2} \text{ DU (Dobson Unit)} = 7.639 \times 10^{-7} \text{ kg} \cdot \text{m}^{-2}$) (Herman et al., 2009).

2.2.2 TROPOMI

TROPOMI on-board the Sentinel-5 Precursor (S-5p) satellite of the European Space Agency (ESA) was launched on 13
175 October 2017 (Veefkind et al., 2012) to fill the gap between OMI (Schoeberl et al., 2006; Levelt et al., 2018) and future
payloads. TROPOMI has been designed to retrieve the slant column densities (SCDs) of key atmospheric species such as
NO₂, O₃, SO₂, HCHO, CH₄ and CO (e.g., <http://www.tropomi.eu/>) with a spatial resolution of 3.5×7 km² which was
further reduced to 3.5 × 5.5 km² from 6 August 2019. This spatial resolution is suitable for air quality monitoring at city
level (Lama et al., 2020). TROPOMI has a swath width of 2600 km which allows for daily global coverage, with an overpass
180 time at about 13:00 local time (Van Geffen et al., 2022b). Trace gas parameters are retrieved using six spectral channels
covering wavelengths from the ultraviolet to the near-infrared. The NO₂ total SCD is retrieved by application of the DOAS
method to the spectral radiation and irradiance measured in the UV-VIS (320-500nm) spectral channels (Van Geffen et al.,
2015; Boersma et al., 2011). The stratospheric and tropospheric SCDs are derived from the total SCD by using the TM5-MP
chemical transport model (CTM) (Dentener et al., 2003). The meteorological information required to run the model is
185 obtained from the European Centre for Medium-Range Weather Forecast (ECMWF) (Dee et al., 2011; Uppala et al., 2005).
The stratospheric and tropospheric SCDs are converted to VCDs using a height-dependent air mass factor (AMF) look-up

table with a spatial resolution of $1^\circ \times 1^\circ$ built using the double adding KNMI (DAK) radiative transfer model (Palmer et al., 2001; Boersma et al., 2004). It is worth noting that the NO_2 total VCDs provided by TROPOMI are the sum of tropospheric and stratospheric VCDs calculated as described above. The margin of error given by the data provider is $0.5 \text{ Pmolec} \cdot \text{cm}^{-2} + (0.2 \text{ to } 0.5) \cdot \text{VCD}_{\text{trop}}$ (Van Geffen et al., 2022b).

TROPOMI provides near-real time data (NRTI) and off-line data (OFFL). The OFFL data used in the current study are produced using observational meteorological data for assimilation in the TM5-MP CTM (Van Geffen et al., 2022b), whereas the NRTI are produced using meteorological forecast data. The TROPOMI OFFL data used in this study were retrieved using retrieval processor version 2.2.0 from 1 August 2021 to 14 November 2022, version 2.3.1 from 15 November 2021 to 17 July 2022 and version 2.4.0 after 18 July 2022 (van Geffen et al., 2022a; van Geffen et al., 2022b). Only data were used with high quality, i.e., with $\text{QA} > 0.75$, which disqualifies scenes with a cloud radiance fraction > 0.5 , some scenes covered by snow or ice and scenes that have been determined to include errors or problematic retrievals. TROPOMI NO_2 VCD data were downloaded from the ESA website: <https://s5phub.copernicus.eu/> (last accessed: 11th July 2023).

2.2.3 In situ measurements of NO_2 concentrations

In addition to NO_2 surface concentrations obtained from Pandora observations, data from an in situ instrument are available for a short period of time, i.e., from a Thermo Scientific 42i-TL Analyzer (<https://www.thermofisher.com/order/catalog/product/42ITL>, last accessed 11th July 2023). This instrument was located at the roof of the AirCAS building at approximately 20 m from the Pandora instrument, during the period January 10-30, 2022 (20 days). The Thermo Scientific 42i-TL Analyzer measures the chemiluminescence caused by the reaction of NO with ozone, where NO is produced from the dissociation of NO_2 ($2\text{NO}_2 \leftrightarrow 2\text{NO} + \text{O}_2$) on a heated molybdenum surface. The chemiluminescence intensity is proportional to the NO concentration (Kley and Mcfarland, 1980). More detailed information is provided in the instrument's manual (<https://www.manualslib.com/manual/1251056/Thermo-Scientific-42i.html> (last accessed: 11th July 2023)). The Thermo Scientific NO_2 data used in this study were screened for high quality using the quality flags and anomalous data were discarded.

2.2.4 Re-analysis data

ERA5 hourly reanalysis wind data, available from the European Centre for Medium-Range Weather Forecasts (ECMWF, <https://www.ecmwf.int/>), were used in the data analysis. This dataset provides hourly atmospheric and oceanic information such as wind speed, temperature, and specific humidity at different pressure levels. ERA5 hourly wind speed data at four pressure levels: 925, 950, 975 and 1000 hPa) were downloaded from the ECMWF website (ERA5 web: <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels?tab=form>, last accessed: 11th July 2023) and averaged (following, e.g., Stein et al., 2015; Ialongo et al., 2020; Zhao et al., 2022a) for use in the analysis presented below.

2.3 Methodology

For the evaluation of the NO₂ observations different methods are used, such as time series to show the variability on different
 220 time scales or the effects of external parameters such as wind speed, averaging to reduce short-time variability, scatterplots
 for comparison with independent data sets. For the comparison between Pandora and TROPOMI NO₂ VCDs, the data need
 to be collocated. Furthermore, the difference in observation geometry needs to be taken into account. The Pandora bottom-up
 observations of direct and scattered (sky) solar radiation depend on the NO₂ vertical profile, the spatial distribution and the
 SZA (Boersma et al., 2011; Van Geffen et al., 2015). For example, for a situation when most of the total VCD is located
 225 below 2 km and the SZA is 45°, the total viewing area of Pandora with a FOV of $(2.6\pi / 180) \cdot (2 / \cos(SZA))$, is about
 $128 \times 128 \text{ m}^2$. The TROPOMI top-down observations from space provide NO₂ VCDs with a spatial resolution of 3.5×5.5
 km^2 . To match the observation volumes, the TROPOMI observations were resampled to a spatial resolution of $100 \times 100 \text{ m}^2$,
 using Google Earth Engine (GEE, (<https://developers.google.com/earth-engine/guides/scale>; last accessed: 11th July 2023)).
 For an accurate comparison between TROPOMI and Pandora, only high quality data were retained for the collocated
 230 TROPOMI/Pandora pixels. For time collocation, Pandora data obtained within 10 minutes before and after the TROPOMI
 overpass time were averaged. The resulting collocated TROPOMI and Pandora data sets were quantitatively evaluated using
 scatterplots and the statistical metrics mean difference (MD), mean absolute difference (MAD), mean relative difference
 (MRD), standard deviation (σ), correlation coefficient) and fitted slope.

MD is calculated by averaging the difference between two observations, as shown in Eq. (1):

$$235 \quad MD = \frac{1}{n} \sum_{i=1}^n (VCD_{\text{TROPOMI},i} - VCD_{\text{Pan},i}), \quad (1)$$

The MAD, given by Eq. (2), is defined as the mean of the absolute differences between TROPOMI and Pandora:

$$MAD = \frac{1}{n} \sum_{i=1}^n |VCD_{\text{TROPOMI},i} - VCD_{\text{Pan},i}|, \quad (2)$$

MRD, Eq. (3), is the mean of the differences between TROPOMI and Pandora when normalized with Pandora's VCDs.
 Positive and negative values of the MRD indicate the degree of overestimation or underestimation:

$$240 \quad MRD = \frac{1}{n} \sum_{i=1}^n \frac{VCD_{\text{TROPOMI},i} - VCD_{\text{Pan},i}}{VCD_{\text{Pan},i}} \times 100\%, \quad (3)$$

The Pearson correlation R (Pearson., 1895) is defined in Eq. (4).

$$R = \frac{\sum_{i=1}^n (VCD_{\text{TROPOMI},i} - \overline{VCD_{\text{TROPOMI}}})(VCD_{\text{Pan},i} - \overline{VCD_{\text{Pan}}})}{\sqrt{\sum_{i=1}^n (VCD_{\text{TROPOMI},i} - \overline{VCD_{\text{TROPOMI}}})^2} \sqrt{\sum_{i=1}^n (VCD_{\text{Pan},i} - \overline{VCD_{\text{Pan}}})^2}}, \quad (4)$$

The standard deviation σ is defined in Eq. (5).

$$\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n (VCD_{\text{TROPOMI},i} - VCD_{\text{Pan},i})^2}, \quad (5)$$

245 The dilution factor, D_f , is defined in Eq. (6), where $VCD_{TROPOMI,0.1}(\text{median})$ and $VCD_{TROPOMI,R}(\text{median})$ represent the median value of the TROPOMI VCD within a radius of 0.1 km and within a radius R around the location of Pandora, respectively (Chen et al., 2009; Griffin et al., 2019; Pinardi et al., 2020):

$$D_f = \frac{VCD_{TROPOMI,R}(\text{median})}{VCD_{TROPOMI,0.1}(\text{median})}, \quad (6)$$

3 Result and discussion

250 3.1 Pandora data overview

An overview of the Pandora data is presented in Figure S1 of the supplementary material, as time series for each month from August 2021 to July 2022, where the time scale is local time (i.e., Beijing Time or BJT, i.e., UTC+8) throughout this paper. As an example, the data for November and December are enlarged in Figure 2. Total NO₂ VCD, tropospheric NO₂ VCD and surface NO₂ concentrations are plotted in different colours and only high and medium quality data are included (DQ0, DQ1, 255 DQ10 and DQ11). Time series including also low quality data (DQ2 and DQ12) are presented in the supplementary material, Figure S2. Although the use of low quality data is not recommended, we have added this figure in the Supplementary material to show the high data density throughout the year. Overall, the NO₂ concentrations are low in the summer and high in the winter, as is commonly observed in satellite data (e.g., van der A et al, 2006; Wang et al., 2019; Fan et al., 2021). For the tropospheric VCDs, including low quality data adds 4681 data points to the total of 8620 data points during the one year 260 study period (i.e., 54.3% of all data are low quality). Among the total VCDs, 21767 data points out of a total of 80153 (27.2%) are low quality. The comparison of Figures S1 and S2 shows that low quality data occur in particular during the winter months when the NO₂ concentrations are high and aerosol (PM_{2.5}) concentrations are also high, the boundary layer is shallow, the sky is often cloudy, and air quality is bad. These are conditions when the data processing flags bad situations for some criteria, the reliability of the results decreases and hence the DQ is increased to DQ2.

265 3.1.1 Diurnal, day-to-day and episodic variations

The data in Figure 2 show the total VCDs are larger than the tropospheric VCDs. The data also show the diurnal variation of these parameters and the surface concentrations derived from Pandora. Although Pandora has the capability of using moonlight for the NO₂ observations (Cede, 2021), the Beijing-RADI data includes only daytime (solar) measurements. The NO₂ VCDs decrease in the morning to reach a daily minimum around local noon and then increase. The complete diurnal 270 cycle of surface concentrations measured with the Thermo Scientific 42i-TL Analyzer, discussed in section 3.4 and presented in Figure 7, confirms these observations and shows that the maximum is reached during the night.

The time series in Figure 2 show large variations of all three parameters over periods of several days, suggesting the occurrence of pollution episodes. An example is a long period from 11 to 19 November 2021, when the NO₂ parameters are increasing until 15 November, whereafter the total VCDs are not available and the tropospheric VCDs and surface

275 concentrations remain relatively high until 19 November. On other occasions the tropospheric VCDs and surface
concentrations are not available, but the total VCDs increase over a period of several days. For example, this is observed in
December 2021 during 5 different periods: 1-5 December, 7-9 December, 11-13 December, 16-22 December and 25-28
December. Comparison with data from the nearby national air quality monitoring site in the Olympic Forest Park in Beijing
(Figure S3) shows that these periods coincide with similar behaviour of the air quality (as indicated by the air quality index,
280 dominated by high aerosol (PM) concentrations) and the associated NO₂ time series (in situ). The variation of the NO₂
concentrations and the occurrence of haze episodes is strongly related to the wind direction and the wind speed, the influence
of which will be discussed in Section 3.2.

285





Figure 2: Time series of NO₂ data measured with Pandora at the Beijing-RADI site for November and December 2022: total VCD in green and tropospheric VCD in red (both on left vertical axes in Pmolec.cm⁻²) and surface concentrations in red (plotted on the right axis in μg.cm⁻³) Only data with DQ0, DQ1, DQ01 and DQ11 are plotted. Note that

290 the surface concentrations scale has been chosen such that they are plotted at the bottom of the VCDs.

3.1.2 Monthly mean NO₂ VCDs

As shown in Figure 2, apart from diurnal and short-term variations, there is also a clear seasonal variation of the NO₂ VCDs. This is further illustrated in Figure 3 where time series are plotted for the mean and median of monthly mean tropospheric VCDs (calculated from daily averages) derived from Pandora measurements during the period from August 2021 to July 2022. Only the data measured between 08:00 and 17:00 BJT, corresponding to daylight hours during the winter, have been included, with DQ0, 1, 10 and 11. This selection of high-quality data reduces the number of days for which data are available, but with ensures reliability. A similar figure including also DQ2 and DQ12 data is included in the Supplementary material (Figure S4) which shows that in the autumn and winter the NO₂ VCDs are high and that they are low in the spring and summer, in good agreement with the seasonal variation which is commonly observed in satellite data (e.g., van der A et al, 2006; Wang et al., 2019; Fan et al., 2021). The number of days for which data are available for each month is included in Table S1 and Figure S2 (including DQ2 and 12). Comparison shows that mainly the monthly mean VCD for December 2021 is affected by the data selection: the number of days is reduced from 31 to 9 and apparently mainly data with high VCDs, i.e., during days with bad AQ (Section 3.1.1), were removed by the DQ2 criteria. As a result, the annual cycle of the mean tropospheric NO₂ VCDs in not as obvious in the high-quality data as in Figure S4 which clearly shows the maximum occurring in the winter and the minimum in the summer. Such an annual variation pattern is in good agreement with near surface observations of in situ NO₂ measurements in Beijing. The mean, median, minimum and maximum values, and the standard deviation (σ), of the tropospheric NO₂ VCDs over Beijing for each month are summarized in Table 1. The difference between the median and mean values is significantly larger in the winter months than in the summer. The variability of the NO₂ VCDs over Beijing is much higher during the winter than in the summer. The minimum in February 2022, as compared to January and March, illustrates the effectiveness of the emission control policy in and around Beijing during the 2022 Winter Olympics (Chu et al., 2022). Time series of the monthly mean total VCDs, are presented in Figure S5 and shows similar variation as for the tropospheric VCDs in Figure 3.



315 **Figure 3: Time series of the monthly mean and median tropospheric NO₂ VCDs over Beijing from August 2021 to July 2022. Only high quality data (DQ 0, 1, 10 and 11) are included, which limits the number of days for which data are available as indicated in Table 1. The blue shaded area indicates the standard deviation for the monthly mean data. Monthly mean data were calculated from daily means.**

Table 1: Statistics of the Pandora-derived monthly mean NO₂ tropospheric VCDs. Only data with DQ0, 1, 10, 11 included.

	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul
Mean (Pmolec.cm ⁻²)	10.45	12.89	8.72	22.14	6.12	13.53	8.12	13.22	9.75	7.97	8.66	10.99
Median (Pmolec.cm ⁻²)	9.26	11.36	6.22	14.78	5.42	4.83	4.03	10.42	8.83	6.06	8.87	9.14
Minimum (Pmolec.cm ⁻²)	0.60	1.68	2.78	1.98	2.91	1.47	0.82	0.60	1.31	1.41	2.10	1.44
Maximum (Pmolec.cm ⁻²)	33.59	50.06	38.46	60.13	10.83	59.31	46.50	45.76	35.66	27.47	16.81	27.28
1σ (Pmolec.cm ⁻²)	6.09	7.24	7.72	18.99	2.28	15.26	9.14	11.11	6.50	5.54	3.33	6.95
Number of days with high quality data	29	17	14	23	9	19	26	29	25	21	10	12

320

3.2 Impact of wind direction and wind speed on NO₂ VCDs in Beijing

The wind influences the concentrations of atmospheric constituents, and thus air quality, in several different ways. The wind transports aerosols and trace gases away from their sources and thus disperses them, leading to lower concentrations. This includes both vertical mixing due to wind-generated turbulence and advection (horizontal transport). This also implies that at very low wind speed the constituents accumulate which leads to enhanced concentrations (e.g., Feng et al., 2014) and, in particular due to (photochemical) reactions, the formation of haze (e.g., An et al., 2019). The wind direction and its history indicate the transport pathways, determined by large scale weather phenomena (e.g., You et al., 2018; He et al., 2018; Hou et al., 2020; Li et al., 2021; Zhao et al., 2022b). When the transport pathways cross source regions, the constituents are transported to the receptor point. Hence, depending on wind direction, the wind transports polluted or clean air over long distances (e.g., Sundström et al., 2012).

To address the effect of advection on the NO₂ tropospheric VCDs at the Beijing RADI site, a polar map has been created of NO₂ tropospheric VCDs versus wind speed and wind direction (Figure 4). To separate effects of wind direction and wind speed on NO₂ VCDs from the large diurnal variations (Figure 2), only data have been plotted at 13:00 BJT when the concentrations are close to their daily minimum. The data in Figure 4 show that, overall, the tropospheric VCDs are smaller during north-westerly winds, in particular when wind speeds exceed about 4 ms⁻¹. This confirms that wind from these directions transport cleaner air to Beijing. As discussed in Section 2.1, the Yanshan and Taihang mountains are situated to the north and west of Beijing and the area between Beijing and the mountains is mainly agricultural with sparse population

and little industry. Thus anthropogenic and industrial emissions over this area are small. Hence, during low wind speeds from this direction, relatively clean air is transported to Beijing. When wind speeds are higher, northwesterly winds from the Siberian plains bring clean air, thus greatly improving Beijing's air quality. As a result, the NO_2 concentrations in the northwesterly wind sector are generally low, as shown in Figure 4. However, the data in Figure 4 show some exceptions when NO_2 concentrations are high. Further analysis shows that these observations were all made during the winter and are likely due to NO_x emissions from natural gas companies located in the Changping district in the northwest of Beijing. Natural gas is provided for, e.g., heating in the winter, and NO_x is produced during the combustion process (Pan et al., 2023). Thus, in the winter, during northwesterly winds, NO_2 is transported to the Pandora site. This explains the observations of high NO_2 concentrations, due to local emissions. (More details of high concentration number and time during north-west wind please see Table S2.)

During southerly winds, the NO_2 VCDs are substantially higher. The high industrial activity and associated traffic and high degree of urbanization in the south of Beijing and the NCP leads to high emissions and thus, during southerly winds, transport of large amounts of NO_2 to Beijing. In addition, during low wind speed, the pollutants are accumulated resulting in the observed very high pollution levels at low wind speed, in particular during southerly winds.




Figure 4: Polar diagram of NO_2 tropospheric VCDs, wind direction and wind speed during the TROPOMI overpass time. Wind speed is indicated by the radius of the circles, ranging from 0 to 12 $\text{m}\cdot\text{s}^{-1}$; wind direction corresponds to the angle in polar coordinates, clockwise from north. Note that the south wind direction indicates that the wind comes from the south.

3.3 Tropospheric contribution to total NO_2 VCDs

The total NO_2 VCDs are determined from the Pandora direct sun measurements and the tropospheric NO_2 VCDs are determined from the Pandora sky measurements. Hence the total and tropospheric NO_2 VCDs are independently determined,

and can be used to obtain information on atmospheric processes. The data in Figure 2 show the strong diurnal variation of both the total and tropospheric NO_2 columns, but the time series of the ratio of the tropospheric to the total NO_2 VCD (see e.g., Figure 5, where the ratios for February are shown as an example, and Figure S6 for all months) show that they do not vary in the same way. Because direct sun measurements are made more frequent than sky measurements, the ratios have been calculated using the first tropospheric VCD sky measurement available after a direct sun measurement and then averaged to obtain daily and monthly means (Figure 6). The results show that the fraction of tropospheric NO_2 to total NO_2 VCDs varies between about 0.2 and 1.05 (tropospheric VCDs should be smaller than or equal to the total VCDs, a value larger than 1 indicates the uncertainties in the data). The large diurnal and day-to-day variations of these ratios indicate that different processes in the troposphere and above affect NO_2 concentrations. The data show that the tropospheric / total ratio decreases to a minimum around the middle of the day, i.e., early in the day the tropospheric NO_2 VCDs decrease faster than the total, before it increases in the afternoon. This is due to a combination of processes including sources and sinks, of (photo)chemical nature (Herman et al. 2009), transport influenced by meteorological phenomena such as variations in wind speed and wind direction (discussed above in Section 3.2) and variations in boundary layer height, while also the temperature profile changes throughout the day, influencing reaction rates and chemical balance (cf. Kang et al., 2022, for a brief overview). Likely all of these are different between the troposphere and above and therefore influence the ratio tropospheric / total NO_2 VCD and its daily evolution.

375




Figure 5: Time series of the ratios of tropospheric VCDs to total VCDs during February 2022.

The monthly mean ratios of the NO_2 tropospheric to total VCDs derived from the Pandora measurements at the Beijing-RADI site are plotted as time series in Figure 6, together with the number of days for which data are available in each month. The results in Figure 6 show that the monthly mean ratios vary between about 0.38 and about 0.75, and there is some inter-monthly variation. Considering only months with 10 or more days for which data are available, the monthly averages show

380

that, on average, the tropospheric NO₂ contribution is 50% to 60% in the winter and 60% to 70% in the spring and autumn. The smaller ratio in the winter may be related to the frequent occurrence of haze days when tropospheric NO₂ is converted to fine particulate matter (e.g., Zheng et al., 2015; Xie et al., 2015; Wang et al., 2020), whereas the larger ratio in spring can be derived from reduced stratospheric concentrations due to enhanced solar shortwave radiation (Cheng et al., 2016; Müller, 2021). Similar to ozone, stratospheric intrusion could be a possible reason for the springtime increase in tropospheric NO₂ concentrations (Lin et al., 2015), because the higher values in the stratosphere have been observed in many studies (Sioris et al., 2003; Hendrick et al., 2004; Preston et al., 1998). Also, the larger standard deviations in spring (especially in March when it was larger than 0.2) indicate a larger day-to-day variability than in other seasons, which may be due to more active photochemical reactions in response to enhanced radiation intensity.





Figure 6: Box and whisker plot of the monthly mean ratios of NO₂ tropospheric VCDs to total VCDs from August 2021 to July 2022, with the number of valid data in each month plotted as a blue dot (right hand axis). For each data point, the dot represents the mean value, the horizontal line represents the median value, the top and bottom edges of the boxes are the 25% and 75% quartiles and the whiskers are the values corresponding to 10% and 90% of the data volume.

3.4 Comparison of Pandora-derived surface NO₂ concentrations with ground-based in-situ measurements

During the study period, the Thermo Scientific 42i-TL Analyzer was operated side by side with the Pandora instrument during a period of 20 days in January 2022. The collocation of these two instruments provides an excellent opportunity for the evaluation of the ground-based NO₂ concentrations derived from Pandora sky measurements, using the in situ data from the Thermo Scientific 42i-TL Analyzer as reference. Figure 7 shows the time series of these two data sets. The in-situ observations were made continuously, 24 hours per day, but the Pandora measurements were only made during daylight hours. In addition, there are many gaps in the Pandora time series because only the best quality data (QA = 0, 1, 10, 11) were retained. To fill part of these gaps, Pandora-derived total VCDs, for which many more high quality data are available as shown in Figure S2 and discussed in Section 3.1.1, have been included in Figure 7.




410 **Figure 7: Time series of NO₂ surface concentrations measured using the Thermo Scientific 42i-TL Analyzer (gray) and derived from the Pandora sky data (blue) and total VCDs (red) during 20 days: 10 - 30 January 2022. Yellow vertical dashed lines indicate Beijing Time 0:00. Note that the NO₂ total VCDs are plotted on the secondary vertical axis and scaled to match the Thermo 42i-TL Analyzer surface concentration data.**

415 The continuous time series of the ground-based in situ measurements shows that the NO₂ concentrations are usually high during the night, decrease after sunrise to a minimum in the afternoon and increase thereafter. The observed diurnal variation of the NO₂ concentrations is due to complex photochemical reactions between atmospheric gases such as nitrogen oxides (NO_x), O₃, and volatile organic compounds (VOCs) (Law et al., 2002; Xue et al., 2016). On days with large variations of the NO₂ surface concentration, such as on 11, 14, 16 and 17 January 2022 (Figure 7), the total VCDs measured with Pandora traced the surface concentrations well (note that in Figure 7 the total NO₂ VCDs data were scaled to match the Thermo 42i-TL Analyzer surface concentrations). In addition, the data show that on 11 and 16 January, the total VCDs increased earlier than the surface concentrations, suggesting that the increase of the NO₂ VCDs on these two days resulted from atmospheric transport. On 14 January, the surface concentrations changed before the total VCDs increased, suggesting the occurrence of an NO₂ source near the surface from where NO₂ was vertically mixed resulting in the subsequent increase of the tropospheric VCDs.

425 The data in Figure 7 further show that the Pandora-derived surface NO₂ concentrations are overall relatively low as compared with the in situ concentrations measured with the Thermo Scientific 42i-TL Analyzer. This is further illustrated in the scatterplot of the Pandora-derived NO₂ surface concentrations versus the Thermo Scientific 42i-TL Analyzer in situ data (Figure 8). Figure 8 shows that there are two data regimes: for low in situ NO₂ concentrations, up to about 22 μg·cm⁻³, the Pandora-derived surface concentrations trace the in situ concentrations but vary over a much smaller range (between about 5 and 10 μg·cm⁻³). For larger NO₂ concentrations, the Pandora-derived surface concentrations are closer to the in situ data but

430

with a substantial underestimation. Similar differences have been reported between NO₂ concentrations retrieved from MAX-DOAS measurements, which use a similar technique as used for Pandora to derive surface concentrations, and NO₂ point measurements using instruments mounted in a mast at three heights above the surface (ASL), which, like the Thermo Scientific 42i-TL Analyzer, derive the NO₂ concentrations from chemiluminescence measurements (Kang et al., 2021).
 435 These authors report that MAX-DOAS substantially underestimates the in situ concentrations, by about 65% at 60m and 160 m ASL and by 33% at 280 m ASL, but do not offer an explanation. Below we discuss the Pandora and chemiluminescence data and influencing factors which may contribute to the observed differences.



440 **Figure 8: Scatterplot of NO₂ surface concentrations: Pandora data versus Thermo Scientific 42i-TL Analyzer data.**

The Pandora surface concentration is derived from the ratio of the difference between the SCDs of NO₂ at the largest and smallest SZA and the same difference for air, i.e., $(SCD_{\text{Pandora,max}} - SCD_{\text{Pandora,0}}) / (SCD_{\text{AIR,max}} - SCD_{\text{AIR,0}})$ (Cede, 2021). In view of the exponential decrease of NO₂ concentrations with height as reported by Kang et al. (2021), the slant
 445 optical path at Pandora's maximum observation angle, which is slightly less than 90°, implies that the NO₂ concentration is integrated over a certain height with a negative NO₂ concentration gradient. As a result, the measured concentration is lower than that at the surface which is measured by the Thermo Scientific 42i-TL Analyzer.

Furthermore, Pandora measures the average NO₂ concentration along the (slant) optical path, whereas the Thermo Scientific 42i-TL Analyzer provides a local value at the location of the instrument (Kang et al., 2021) which may be more sensitive to
 450 local variations, such as short-term emissions by passing cars or turbulent perturbations of plumes.

Another factor influencing the comparison of Pandora and in situ measurements of NO₂ concentrations using a Thermo Scientific 42i-TL Analyzer, is that this instrument may overestimate atmospheric NO₂ concentrations due to interference from other atmospheric constituents which react with ozone to produce chemiluminescence, such as peroxyacetyl nitrate and nitric acid (Dunlea et al., 2007; Steinbacher et al., 2007) or alkenes (Alam et al., 2020). In particular, during the Beijing winter the nitrate concentrations are high (Luo et al., 2019; Wang et al., 2021) and may contribute to overestimation of the NO₂ concentrations measured by the thermo-chemiluminescence method.

3.4.1 Difference between pandora and in situ surface NO₂ concentrations for different concentration regimes

As mentioned above, Figure 8 shows two groups of data which are separated according to low and high NO₂ concentrations. Comparison with the polar diagram in Figure 4 shows that, for the TROPOMI overpass time, the low NO₂ concentrations mainly occur during northerly winds, whereas high concentrations are mainly observed during southerly winds, and more specific for wind directions between about 135° and 315°. The scatterplot in Figure 8 includes all observations for a short time period. To further investigate the different NO₂ regimes revealed in Figure 8, wind direction has been added to the time series plot of Figure 7, as well as AOD, and the results are presented in the Supplementary material, Figure S7. Figure S7 confirms that the conclusions from Figure 4, for TROPOMI overpass times, also apply to the observations at other times, i.e., that the low surface NO₂ concentrations (<10 µg.m⁻³) derived from Pandora almost all occur during northerly winds, are smaller than the in situ concentrations measured at the same time with the Thermo Scientific 42i-TL Analyzer, and trace the latter well. This is well-illustrated on 26 January when the surface concentrations initially decrease (from about 15 and 24 µg.m⁻³ for Pandora and Thermo, respectively) to a minimum, then increase. This variation may be due to a combination of the diurnal variation described in Section 3.1 and the change in wind direction and thus transport pathway. During southerly winds the Pandora-derived surface NO₂ concentrations are substantially higher (>15-20 µg.m⁻³) and increase with increasing in situ data. In particular for in situ concentrations between about 30 and 40 µg.m⁻³ there is a good correlation although Pandora data are still underestimated as discussed above. The data in Figure S7 also show the large changes occurring during changes in wind direction, in combination with diurnal variations.

The difference between the relations between Pandora-derived and in situ measured surface NO₂ concentrations for the low and high NO₂ concentration regimes may be explained as follows. During northerly winds, the contribution to the NO₂ concentrations from transport over long distances is small because the air masses are transported over relatively clean areas. Thus, the main contribution to the concentration is due to local production of NO₂ from anthropogenic activities near the surface, which is first observed by the in-situ sensor. Pandora surface NO₂ concentration is an integrated value over the height range covered by the largest SZA. Mixing of the surface-produced NO₂ over this height range takes some time, and thus there is a concentration gradient resulting in a lower value of the Pandora-derived surface concentration. In contrast, during southerly winds, the air mass transported to the measurement location has passed through an area with high NO₂ concentrations and the contribution from local production of NO₂ near the surface is relatively small. Thus, the NO₂ concentration measured by the in-situ sensor is relatively little enhanced due to local production as compared to that in

northerly winds. Therefore, the Pandora-derived surface NO₂ concentration compares better to the in-situ data when the
485 concentrations are high than when they are small.

3.4.2 Comparison with AOD

Figure S7 shows that the AOD and NO₂ total VCD (both are column integrated quantities measured using collocated ground-
based instruments), trace each other very well. Hence the different relationships between MAX-DOAS - and Tower -
averaged NO₂ concentrations for low and high AOD reported by Kang et al. (2021) are due to the same reason as the
490 different relationship for low and high NO₂ concentrations shown in Figure 8 and discussed above. It is noted that the
measurements reported by Kang et al. (2021) were made on a tower located in Beijing at ca. 5 km from the Beijing-RADI
site and the conditions at both sites are similar.

3.5 Validation of TROPOMI NO₂ VCDs over Beijing using Pandora

The Pandora instrument and the Pandora Global Network were designed for satellite validation, i.e., providing VCDs as
495 independent reference for the validation and evaluation of satellite data. Here Pandora data are used for the validation of
TROPOMI NO₂ total and tropospheric VCDs, re-sampled to a spatial resolution of 100×100 m². Scatterplots of NO₂ total
and tropospheric VCDs from TROPOMI versus Pandora VCDs and histograms of the TROPOMI-Pandora differences are
presented in Figure 9. For TROPOMI, OFFL data were used and for Pandora only high quality data with DQ0, 1, 10 and 11.
The Pandora quality control limits the amount of suitable data, especially in June, July, August, and September (Section 3.1).
500 The scatterplots in Figures 9a and c show the good correlation of both the TROPOMI-derived NO₂ total and tropospheric
VCDs with the Pandora data. For the total VCDs, R= 0.95, the slope of the LSQ fit is 0.96 and the RD is -8.44%, well below
the expected deviation of $0.5 \text{ Pmolec} \cdot \text{cm}^{-2} + (0.2 \text{ to } 0.5) \cdot \text{VCD}_{\text{trop}}$. (Van Geffen et al., 2022b). Figure 9a shows that for
small values of the total VCDs the data points are very close to the identity line, whereas for VCDs > 10 Pmolec.cm⁻² the
differences are larger and the data points are rather evenly scattered around the identity line. The histogram of the differences
505 between the TROPOMI and Pandora NO₂ total VCDs in Figure 9b shows a uniform distribution with the characteristics of a
Gaussian with the mean or median as the symmetry axis. On average, the mean TROPOMI total VCDs are slightly smaller
than those from Pandora, with an MD of -1.16 Pmolec.cm⁻², a MAD of 3.14 Pmolec.cm⁻² and a standard deviation of 4.44
Pmolec.cm⁻². This uncertainty is in part due to the large SZA, which leads to uncertainty in the air mass factor (AMF)
(Herman et al., 2019; Griffin et al., 2019; Ialongo et al., 2020). In addition, a 10% error in the AMF estimate due to cloud
510 effects was reported in the ATBD document (Van Geffen et al., 2022b).

The validation of the TROPOMI-derived NO₂ tropospheric VCDs is presented as scatterplots and histograms in Figures 9c
and 9d. As for the total VCDs, the TROPOMI tropospheric VCDs trace the Pandora data well and the scatterplot shows the
good correlation between both data sets, with a correlation coefficient of 0.92, slightly smaller than for the total VCDs, and a
somewhat smaller slope of 0.94, with a RD of 16.15%. The histogram in Figure 9d shows that the tropospheric VCDs are

515 roughly symmetrically distributed around 0, and the number of large negative deviations (deviation < -10 Pmolec.cm⁻²) is slightly larger than the number of positive deviations (deviation > 10 Pmolec.cm⁻²).

The above analysis confirms the good performance of the TROPOMI NO₂ total and tropospheric VCDs and provides confidence for the use of the TROPOMI data. For this analysis, the TROPOMI data were re-sampled to a spatial resolution of 100×100 m², similar to the Pandora observation area.

520





Figure 9: Validation of TROPOMI total and tropospheric NO₂ VCDs, re-sampled to a spatial resolution of 100×100 m², using Pandora observation as reference data: (a, c) scatter plots of total and tropospheric TROPOMI vs Pandora data together with statistical metrics; (b, d) histograms of the differences between TROPOMI and Pandora NO₂ total/tropospheric VCDs.

525

3.6 Spatial representativeness of Pandora vertical column density

Considering the good performance of the TROPOMI-retrieved NO₂ VCDs discussed in section 3.5, these data have been used to evaluate the spatial representativeness of the Pandora observations at the Beijing-RADI site. Figure 10a shows the

spatial distribution of the TROPOMI-derived NO₂ tropospheric VCD over a wide area centered at the Beijing-RADI site averaged over the first year of Pandora operations (Figure 1). Large differences are observed with high tropospheric NO₂ VCDs over Beijing, Tianjin and its highly industrialized surroundings and several other locations. Over the mountains to the north and west of the study area the tropospheric NO₂ VCDs are much lower, by a factor of 10 or more. The Beijing-RADI Pandora site is located in the high NO₂ tropospheric VCD area (as also shown in the inset of Figure 1), while strong NO₂ gradients are observed toward the north and west. Obviously, because of the variation in the NO₂ VCD, the correlation between the TROPOMI and Pandora data is influenced by the choice of the area over which the TROPOMI data are averaged. In particular, this may happen when a separation between clean and polluted air occurs over Beijing, as reported, for instance, by Sun et al. (2016) (their Figure 7). The effect of the size of the area over which TROPOMI data are averaged is illustrated in Figure 10b, showing the annual mean (black) and median (red) tropospheric NO₂ VCDs averaged over areas centered at the Beijing RADI Pandora site with increasing radius. The data show that, for this specific case when the TROPOMI NO₂ VCDs are averaged over annual mean values, the area-averaged value remains constant within a radius of 10 km from the Pandora site, and decreases when the radius is further increased. This is further illustrated by the data in Table 2, showing that for a radius larger than 10 km, the standard deviation (σ) increases as the radius increases. For less homogeneous NO₂ distributions the area for which the situation at the site is representative will be much smaller than 10 km. This also implies that the location of the Pandora site influences the validation results. The data in Table 2 show that for a circle of 3 km the σ is 0.223 Pmolec.cm⁻² and for a circle of 5 km it is 0.282 Pmolec.cm⁻². This means that when the Pandora site is at a different location in an overpass pixel, the 3.5×5.5 km² FOV produces an error in the NO₂ tropospheric VCD between 0.223 and 0.282 Pmolec.cm⁻², i.e., between 1.7% and 2%.



550

Figure 10: Illustration of the spatial variation of the NO₂ tropospheric VCD over a wide area around the Beijing-RADI (Pandora-171) site and the evaluation of the representativeness of the Pandora data for a larger area. (a) spatial distribution of the

555 TROPOMI-derived NO₂ tropospheric VCD re-gridded to a resolution of 0.1×0.1 km² over the study area indicated in Figure 1, averaged over the period from August 1, 2021, to July 31, 2022. The three purple circles indicate a radius of 10, 30 and 50 km around the site. (b) annual mean (black) and median (red) tropospheric NO₂ VCD averaged over circles centered at the Beijing RADI Pandora site with increasing radius, as a function of that radius (expressed in km), with variability (1 standard deviation) given as a light blue area around the mean.

Table 2: Statistical metrics for the tropospheric VCDs averaged over circles with increasing radius around the Pandora site.

Radius (Km)	0.1	1	2	3	5	10	20	30	40	50	80
Mean (Pmolec.cm ⁻²)	12.83	12.90	12.98	12.97	13.98	12.97	12.58	11.84	10.88	10.05	8.58
Median (Pmolec.cm ⁻²)	12.85	12.89	12.98	13.01	13.01	12.99	12.68	12.15	11.40	10.77	9.10
Standard deviation σ (Pmolec.cm ⁻²)	0.037	0.110	0.164	0.223	0.282	0.407	0.901	1.444	2.087	2.563	3.210
Dilution factor (D_f)	1	1.003	1.010	1.012	1.012	1.011	0.987	0.945	0.887	0.838	0.710

560

4 Summary and conclusions

The first operational Pandora site in China has been established on the roof of the laboratory building of the Aerospace Information Research Institute of the Chinese Academy of Sciences (AirCAS) in Beijing, at the end of July 2021. The Pandora instrument provides continuous observations of NO₂ total and tropospheric VCDs together with surface concentrations from direct sun and sky measurements. The Beijing-RADI site is part of the PANDONIA Global Network (PGN), where the Beijing-RADI data are publicly accessible from the PGN website. In this paper, an overview has been presented of the first year of data from the Beijing-RADI Pandora, i.e., their variations on time scales from hours to seasonal, and the influences of wind speed and direction as well as chemical reactions on these variations. The NO₂ surface concentrations have been compared with independent measurements from a collocated Thermo Scientific 42i-TL Analyzer, using a different physical principle for NO₂ concentration measurements. Explanations were offered to explain the differences between Pandora and in situ NO₂ surface concentrations. The Pandora NO₂ VCD data have been used as an independent reference for the evaluation of TROPOMI retrieved NO₂ VCDs over Beijing. This study leads to the following conclusions:

- 575 1. The Pandora observations show that NO₂ concentrations in Beijing are high during the winter and low during the summer, with a diurnal cycle where the concentrations reach a minimum during day time; the reduced concentrations during the 2022 Winter Olympics show that the emission control during that period was highly effective.

2. The fraction of tropospheric NO₂ contributing to the total NO₂ VCDs varies between 0.2 to 1, with large diurnal to seasonal variations, is high in the morning and afternoon with a minimum around noon. Monthly averages show that the tropospheric NO₂ contribution is 50% to 60% in the winter and 60% to 70% in the spring and autumn. The relatively small amount of valid data does not allow us to assess the tropospheric contribution in summer.
3. During northerly winds, the tropospheric NO₂ VCD is small, especially when the wind speed is larger than 4 m.s⁻¹. When the wind speed is low, clean air from areas between the Yanshan and Taihang mountains is transported to Beijing. When the wind speed is higher, northwesterly winds transport clean air from the Siberian plains, greatly improving Beijing's air quality. During southerly winds, polluted air is transported from adjacent areas with high NO₂ emissions resulting in high NO₂ pollution in Beijing. In addition, at low wind speeds, pollutants accumulate which results in increased NO₂ concentrations which are most obvious during southerly winds.
4. Comparison of NO₂ surface concentrations derived from Pandora measurements with in-situ data from a Thermo Scientific 42i-TL Analyzer show that the Pandora-derived concentrations are substantially smaller than the in-situ data. Reasons for this discrepancy have been identified: 1) The Thermo Scientific 42i-TL Analyzer is sensitive to other sources of chemiluminescence contributing to the NO₂ signal, such as peroxyacetyl nitrate and nitric acid, or alkenes. 2) The Thermo Scientific 42i-TL Analyzer represents the local in-situ NO₂ concentration near the sampling port whereas Pandora represents the average concentration along a slant optical path, instead of a horizontal optical path for near-surface measurements, because the Pandora observation zenith angle is slightly smaller than 90 degrees. The concentrations are therefore an average over a small vertical range with, considering that the NO₂ source is at the surface, a vertical concentration gradient. This results in a an average NO₂ concentration which is smaller than the actual concentration at the surface.
5. The comparison of the Pandora and in-situ surface NO₂ concentrations further shows that the data are separated in two clusters with different relations between the Pandora and in situ data: one for low NO₂ concentrations during northerly winds where the Pandora concentrations show very little variation with in-situ data and another regime for high NO₂ concentrations where the Pandora data are still smaller than the in-situ concentrations but with a definite positive correlation. These differences are explained in terms of transport and local production.
6. Using the Pandora data for validation of the TROPOMI NO₂ total and tropospheric VCDs shows the good performance of the TROPOMI retrieval over Beijing with R= 0.95 and 0.92, a LSQ fit with a slope of 0.96 and 0.94 and a standard deviation well below the expected $0.5 \text{ Pmolec} \cdot \text{cm}^{-2} + (0.2 \text{ to } 0.5) \cdot \text{VCD}_{\text{trop}}$.
7. The location of the Pandora instrument within a sub-orbital TROPOMI pixel of 3.5×5.5 km² may result in an error in the NO₂ tropospheric VCD between 0.223 and 0.282 Pmolec.cm⁻², i.e., between 1.7% and 2%. The analysis shows that the Pandora observations at the Beijing-RADI site are representative for an area with a radius of 10 km.

610 **Data availability.**

The TROPOMI data are available via

https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S5P_OFFL_L3_NO2.

The Pandonia data are available via <http://data.pandonia-global-network.org/>.

The ERA5 reanalysis data are available via <https://cds.climate.copernicus.eu/>.

615

Auth contributions.

OL, YL, CF and GL were involved in the research design. OL, GL, YZ and CF analysed the data. OL, GL, YZ, CF and YL prepared the manuscript. JD provides support to the subsequent data analysis. KL operated and managed the Beijing-RADI Pandonia measurements. PZ, TZ and YW provide data of Thermo Scientific 42i-TL instrument. ZL and YZ provided critical
620 comments which substantially improved the paper. All authors have read and agreed to the published version of the manuscript.

Competing interests.

The authors declare that they have no conflict of interest.

625

Financial support.

This work was supported by the National Distinguished Youth Foundation of China (Grant No. 41925019) and National Natural Science Foundation of China (Grant No. 42101365).

630 **References**

Alam, M. S., Crilley, L. R., Lee, J. D., Kramer, L. J., Pfrang, C., Vázquez-Moreno, M., Ródenas, M., Muñoz, A., and Bloss, W. J.: Interference from alkenes in chemiluminescent NO_x measurements, *Atmos. Meas. Tech.*, 13, 5977–5991, <https://doi.org/10.5194/amt-13-5977-2020>, 2020.

- An, Z., Huang, R.-J., Zhang, R., Tie, X., Li, G., Cao, J., Zhou, W., Shi, Z., Han, Y., Gu, Z., and Ji, Y.: Severe haze in northern China: A synergy of anthropogenic emissions and atmospheric processes. *PNAS*, 116, 8657-8666, doi.org/10.1073/pnas.1900125116, 2019.
- Behera, S. N. and Sharma, M.: Degradation of SO₂, NO₂ and NH₃ leading to formation of secondary inorganic aerosols: An environmental chamber study, *Atmospheric Environment*, 45, 4015-4024, https://doi.org/10.1016/j.atmosenv.2011.04.056, 2011.
- Boersma, K. F., Eskes, H. J., and Brinksma, E. J.: Error analysis for tropospheric NO₂ retrieval from space, *Journal of Geophysical Research-Atmospheres*, 109, 10.1029/2003jd003962, 2004.
- Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitao, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO₂ column retrieval algorithm for the Ozone Monitoring Instrument, *Atmospheric Measurement Techniques*, 4, 1905-1928, 10.5194/amt-4-1905-2011, 2011.
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noel, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission objectives and measurement modes, *Journal of the Atmospheric Sciences*, 56, 127-150, 10.1175/1520-0469(1999)056<0127:smoamm>2.0.co;2, 1999.
- Brasseur, G. and Nicolet, M.: Chemospheric processes of nitric oxide in the mesosphere and stratosphere, *Planetary and Space Science*, 21, 939-961, https://doi.org/10.1016/0032-0633(73)90141-4, 1973.
- Burrows, J. P., Weber, M., Buchwitz, M., Rozanov, V., Ladstätter-Weißenmayer, A., Richter, A., DeBeek, R., Hoogen, R., Bramstedt, K., Eichmann, K.-U., Eisinger, M., and Perner, D.: The Global Ozone Monitoring Experiment (GOME): Mission Concept and First Scientific Results, *Journal of the Atmospheric Sciences*, 56, 151-175, 10.1175/1520-0469(1999)056<0151:Tgomeg>2.0.Co;2, 1999.
- Carslaw, D. C. and Beevers, S. D.: Estimations of road vehicle primary NO₂ exhaust emission fractions using monitoring data in London, *Atmospheric Environment*, 39, 167-177, https://doi.org/10.1016/j.atmosenv.2004.08.053, 2005.
- Cede, A.: Manual for Blicik Software Suite 1.8, available at: https://www.pandonia-global-network.org/wp-content/uploads/2021/09/BlickSoftwareSuite_Manual_v1-8-4.pdf (last accessed: 11th July 2023), 2021.
- Chen, D., Zhou, B., Beirle, S., Chen, L. M., and Wagner, T.: Tropospheric NO₂ column densities deduced from zenith-sky DOAS measurements in Shanghai, China, and their application to satellite validation, *Atmos. Chem. Phys.*, 9, 3641-3662, 10.5194/acp-9-3641-2009, 2009.
- Cheng, Y., Zheng, G., Wei, C., Mu, Q., Zheng, B., Wang, Z., Gao, M., Zhang, Q., He, K., Carmichael, G., Pöschl, U., and Su, H.: Reactive nitrogen chemistry in aerosol water as a source of sulfate during haze events in China, *Science advances*, 2, e1601530, 10.1126/sciadv.1601530, 2016.
- Chu, F., Gong, C., Sun, S., Li, L., Yang, X., and Zhao, W.: Air Pollution Characteristics during the 2022 Beijing Winter Olympics, *International journal of environmental research and public health*, 19, 10.3390/ijerph191811616, 2022.

- Dai, Y., Cai, X., Zhong, J., Mazzeo, A., and MacKenzie, A. R.: Chemistry, transport, emission, and shading effects on NO₂ and Ox distributions within urban canyons, *Environmental Pollution*, 315, 120347, <https://doi.org/10.1016/j.envpol.2022.120347>, 2022.
- 670 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly*
675 *Journal of the Royal Meteorological Society*, 137, 553-597, 10.1002/qj.828, 2011.
- Dentener, F., van Weele, M., Krol, M., Houweling, S., and van Velthoven, P.: Trends and inter-annual variability of methane emissions derived from 1979-1993 global CTM simulations, *Atmospheric Chemistry and Physics*, 3, 73-88, 10.5194/acp-3-73-2003, 2003.
- de Leeuw, G., van der A, R., Bai, J., Xue, Y., Varotsos, C., Li, Z., Fan, C., Chen, X., Christodoulakis, I., Ding, J., Hou, X.,
680 Kouremadas, G., Li, D., Wang, J., Zara, M., Zhang, K., and Zhang, Y.: Air Quality over China, *Remote Sensing*, 13, 3542, <https://doi.org/10.3390/rs13173542>, 2021.
- Dirksen, R. J., Boersma, K. F., Eskes, H. J., Ionov, D. V., Bucseba, E. J., Levelt, P. F., and Kelder, H. M.: Evaluation of stratospheric NO₂ retrieved from the Ozone Monitoring Instrument: Intercomparison, diurnal cycle, and trending, 116, <https://doi.org/10.1029/2010JD014943>, 2011.
- 685 Dunlea, E. J., Herndon, S. C., Nelson, D. D., Volkamer, R. M., San Martini, F., Sheehy, P. M., Zahniser, M. S., Shorter, J. H., Wormhoudt, J. C., Lamb, B. K., Allwine, E. J., Gaffney, J. S., Marley, N. A., Grutter, M., Marquez, C., Blanco, S., Cardenas, B., Retama, A., Ramos Villegas, C. R., Kolb, C. E., Molina, L. T., and Molina, M. J.: Evaluation of nitrogen dioxide chemiluminescence monitors in a polluted urban environment, *Atmos. Chem. Phys.*, 7, 2691-2704, 10.5194/acp-7-2691-2007, 2007.
- 690 Fan, C., Li, Z., Li, Y., Dong, J., van der A, R., and de Leeuw, G.: Variability of NO₂ concentrations over China and effect on air quality derived from satellite and ground-based observations, *Atmos. Chem. Phys.*, 21, 7723–7748, <https://doi.org/10.5194/acp-21-7723-2021>, 2021.
- Felix, J. D. and Elliott, E. M.: Isotopic composition of passively collected nitrogen dioxide emissions: Vehicle, soil and livestock source signatures, *Atmospheric Environment*, 92, 359-366, <https://doi.org/10.1016/j.atmosenv.2014.04.005>, 2014.
- 695 Feng, X., Li, Q., Zhu, Y., Wang, J., Liang, H., Xu, R: Formation and dominant factors of haze pollution over Beijing and its peripheral areas in winter: *Air Pollution Research*, 4, 528-538. doi:10.5094/APR.2014.062, 2014.
- Grenfell, J. L., Lehmann, R., Mieth, P., Langematz, U., and Steil, B.: Chemical reaction pathways affecting stratospheric and mesospheric ozone, *J. Geophys. Res.*, 111, D17311, <https://doi.org/10.1029/2004JD005713>, 2006.
- Griffin, D., Zhao, X. Y., McLinden, C. A., Boersma, F., Bourassa, A., Dammers, E., Degenstein, D., Eskes, H., Fehr, L.,
700 Fioletov, V., Hayden, K., Kharol, S. K., Li, S. M., Makar, P., Martin, R. V., Mihele, C., Mittermeier, R. L., Krotkov, N.,

- Sneep, M., Lamsal, L. N., ter Linden, M., van Geffen, J., Veefkind, P., and Wolde, M.: High-Resolution Mapping of Nitrogen Dioxide With TROPOMI: First Results and Validation Over the Canadian Oil Sands, *Geophysical Research Letters*, 46, 1049-1060, 10.1029/2018gl081095, 2019.
- 705 He, J., Gong, S., Zhou, C., Lu, S., Wu, L., Chen, Y., Yu, Y., Zhao, S., Yu, L., Yin, C.: Analyses of winter circulation types and their impacts on haze pollution in Beijing. *Atmospheric Environment* 192, 94–103, doi.org/10.1016/j.atmosenv.2018.08.060, 2018.
- Hendrick, F., Barret, B., Van Roozendaal, M., Boesch, H., Butz, A., De Mazière, M., Goutail, F., Hermans, C., Lambert, J. C., Pfeilsticker, K., and Pommereau, J. P.: Retrieval of nitrogen dioxide stratospheric profiles from ground-based zenith-sky UV-visible observations: validation of the technique through correlative comparisons, *Atmos. Chem. Phys.*, 4, 2091-2106, 710 10.5194/acp-4-2091-2004, 2004.
- Herman, J., Cede, A., Spinei, E., Mount, G., Tzortziou, M., and Abuhassan, N.: NO₂ column amounts from ground-based Pandora and MFDOAS spectrometers using the direct-sun DOAS technique: Intercomparisons and application to OMI validation, *Journal of Geophysical Research-Atmospheres*, 114, 10.1029/2009jd011848, 2009.
- 715 Herman, J., Abuhassan, N., Kim, J., Kim, J., Dubey, M., Raponi, M., and Tzortziou, M.: Underestimation of column NO₂ amounts from the OMI satellite compared to diurnally varying ground-based retrievals from multiple PANDORA spectrometer instruments, *Atmos. Meas. Tech.*, 12, 5593-5612, 10.5194/amt-12-5593-2019, 2019.
- Herman, J., Spinei, E., Fried, A., Kim, J., Kim, J., Kim, W., Cede, A., Abuhassan, N., and Segal-Rozenhaimer, M.: NO₂ and HCHO measurements in Korea from 2012 to 2016 from Pandora spectrometer instruments compared with OMI retrievals and with aircraft measurements during the KORUS-AQ campaign, *Atmos. Meas. Tech.*, 11, 4583-4603, 10.5194/amt-11-720 4583-2018, 2018.
- Hou, X., Zhu, B., Kumar, K. R., de Leeuw, G., Lu, W., Huang, Q., and Zhu, X.: Establishment of Conceptual Schemas of Surface Synoptic Meteorological Situations Affecting Fine Particulate Pollution Across Eastern China in the Winter, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD033153, <https://doi.org/10.1029/2020JD033153>, 2020.
- 725 Ialongo, I., Virta, H., Eskes, H., Hovila, J., and Douros, J.: Comparison of TROPOMI/Sentinel-5 Precursor NO₂ observations with ground-based measurements in Helsinki, *Atmos. Meas. Tech.*, 13, 205-218, 10.5194/amt-13-205-2020, 2020.
- Jorba, O., Dabdub, D., Blaszcak-Boxe, C., Pérez, C., Janjic, Z., Baldasano, J., Spada, M., Badia, A., and Gonçalves, M. J. J. o. G. R. A.: Potential significance of photoexcited NO₂ on global air quality with the NMMB/BSC chemical transport model, *J. Geophys. Res.*, 117, D13301, doi:10.1029/2012JD017730, 2012.
- 730 Kang, Y., Tang, G., Li, Q., Liu, B., Cao, J., Hu, Q., and Wang, Y.: Evaluation and Evolution of MAX-DOAS-observed Vertical NO₂ Profiles in Urban Beijing, *Advances in Atmospheric Sciences*, 38, 1188-1196, 10.1007/s00376-021-0370-1, 2021.
- Kang, H., Zhu, B., de Leeuw, G., Yu, B., van der A, R. J., and Lu, W.: Impact of urban heat island on inorganic aerosol in the lower free troposphere: a case study in Hangzhou, China, *Atmos. Chem. Phys.*, 22, 10623–10634, <https://doi.org/10.5194/acp-22-10623-2022>, 2022.

- 735 Kley, D. and McFarland, M.: Chemiluminescence detector for NO and NO₂, *Atmos. Technol.*, 12, <https://www.osti.gov/biblio/6457230>, 1980.
- Lama, S., Houweling, S., Boersma, K. F., Eskes, H., Aben, I., Denier van der Gon, H. A. C., Krol, M. C., Dolman, H., Borsdorff, T., and Lorente, A.: Quantifying burning efficiency in megacities using the NO₂/CO ratio from the Tropospheric Monitoring Instrument (TROPOMI), *Atmos. Chem. Phys.*, 20, 10295-10310, 10.5194/acp-20-10295-2020, 2020.
- 740 Law, M., Kind, H., Messer, B., Kim, F., and Yang, P.: Photochemical Sensing of NO₂ with SnO₂ Nanoribbon Nanosensors at Room Temperature, *Angewandte Chemie International Edition*, 41, 2405-2408, [https://doi.org/10.1002/1521-3773\(20020703\)41:13<2405::AID-ANIE2405>3.0.CO;2-3](https://doi.org/10.1002/1521-3773(20020703)41:13<2405::AID-ANIE2405>3.0.CO;2-3), 2002.
- Levelt, P. F., Oord, G. H. J. v. d., Dobber, M. R., Malkki, A., Huib, V., Johan de, V., Stammes, P., Lundell, J. O. V., and Saari, H.: The ozone monitoring instrument, *IEEE Transactions on Geoscience and Remote Sensing*, 44, 1093-1101, 10.1109/TGRS.2006.872333, 2006.
- 745 Levelt, P. F., Joiner, J., Tamminen, J., Veefkind, J. P., Bhartia, P. K., Stein Zweers, D. C., Duncan, B. N., Streets, D. G., Eskes, H., van der A, R., McLinden, C., Fioletov, V., Carn, S., de Laat, J., DeLand, M., Marchenko, S., McPeters, R., Ziemke, J., Fu, D., Liu, X., Pickering, K., Apituley, A., González Abad, G., Arola, A., Boersma, F., Chan Miller, C., Chance, K., de Graaf, M., Hakkarainen, J., Hassinen, S., Ialongo, I., Kleipool, Q., Krotkov, N., Li, C., Lamsal, L., Newman, P.,
- 750 Nowlan, C., Suleiman, R., Tilstra, L. G., Torres, O., Wang, H., and Wargan, K.: The Ozone Monitoring Instrument: overview of 14 years in space, *Atmos. Chem. Phys.*, 18, 5699-5745, 10.5194/acp-18-5699-2018, 2018.
- Lin, M., Fiore, A. M., Horowitz, L. W., Langford, A. O., Oltmans, S. J., Tarasick, D., and Rieder, H. E.: Climate variability modulates western US ozone air quality in spring via deep stratospheric intrusions, *Nature Communications*, 6, 7105, 10.1038/ncomms8105, 2015.
- 755 Liu, F., Beirle, S., Zhang, Q., Dörner, S., He, K., and Wagner, T.: NO_x lifetimes and emissions of cities and power plants in polluted background estimated by satellite observations, *Atmos. Chem. Phys.*, 16, 5283–5298, <https://doi.org/10.5194/acp-16-5283-2016>, 2016.
- Liu, C. and Shi, K.: A review on methodology in O₃-NO_x-OC sensitivity study: *Environmental Pollution* 291, 118249, <https://doi.org/10.1016/j.envpol.2021.118249>, 2021.
- 760 Li, K., Jacob, D. J., Shen, L., Lu, X., De Smedt, I., and Liao, H.: Increases in surface ozone pollution in China from 2013 to 2019: anthropogenic and meteorological influences, *Atmos. Chem. Phys.*, 20, 11423-11433, 10.5194/acp-20-11423-2020, 2020.
- Li, M., Yao, Y., Simmonds, I., Luo, D., Zhong, L., and Pei, L.: Linkages between the atmospheric transmission originating from the North Atlantic Oscillation and persistent winter haze over Beijing, *Atmos. Chem. Phys.*, 21, 18573–18588, <https://doi.org/10.5194/acp-21-18573-2021>, 2021.
- 765 Li, Z., Zhang, Y., Shao, J., Li, B., Hong, J., Liu, D., Li, D., Wei, P., Li, W., Li, L., Zhang, F., Guo, J., Deng, Q., Wang, B., Cui, C., Zhang, W., Wang, Z., Lv, Y., Xu, H., Chen, X., Li, L., and Qie, L.: Remote sensing of atmospheric particulate mass

- of dry PM_{2.5} near the ground: Method validation using ground-based measurements, *Remote Sensing of Environment*, 173, 59-68, <https://doi.org/10.1016/j.rse.2015.11.019>, 2016.
- 770 Luo, L., Wu, Y., Xiao, H., Zhang, R., Lin, H., Zhang, X., and Kao, S.-j.: Origins of aerosol nitrate in Beijing during late winter through spring, *Science of The Total Environment*, 653, 776-782, <https://doi.org/10.1016/j.scitotenv.2018.10.306>, 2019.
- Mather, T. A., Pyle, D. M., and Allen, A. G.: Volcanic source for fixed nitrogen in the early Earth's atmosphere, *Geology*, 32(10), 905-908, <https://doi.org/10.1130/G20679.1>, 2004.
- 775 Palmer, P. I., Jacob, D. J., Chance, K., Martin, R. V., Spurr, R. J. D., Kurosu, T. P., Bey, I., Yantosca, R., Fiore, A., and Li, Q. B.: Air mass factor formulation for spectroscopic measurements from satellites: Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *Journal of Geophysical Research-Atmospheres*, 106, 14539-14550, 10.1029/2000jd900772, 2001.
- Pan, H., Geng, S., Yang, H., Zhang, G., Bian, H., and Liu, Y.: Influence of H₂ blending on NO_x production in natural gas
780 combustion: Mechanism comparison and reaction routes, *International Journal of Hydrogen Energy*, 48, 784-797, <https://doi.org/10.1016/j.ijhydene.2022.09.251>, 2023.
- Pinardi, G., Van Roozendael, M., Hendrick, F., Theys, N., Abuhassan, N., Bais, A., Boersma, F., Cede, A., Chong, J., Donner, S., Drosoglou, T., Dzhola, A., Eskes, H., Frieß, U., Granville, J., Herman, J. R., Holla, R., Hovila, J., Irie, H., Kanaya, Y., Karagkiozidis, D., Kouremeti, N., Lambert, J. C., Ma, J., Peters, E., Piders, A., Postlyakov, O., Richter, A.,
785 Remmers, J., Takashima, H., Tiefengraber, M., Valks, P., Vlemmix, T., Wagner, T., and Wittrock, F.: Validation of tropospheric NO₂ column measurements of GOME-2A and OMI using MAX-DOAS and direct sun network observations, *Atmos. Meas. Tech.*, 13, 6141-6174, 10.5194/amt-13-6141-2020, 2020.
- Preston, K. E., Fish, D. J., Roscoe, H. K., and Jones, R. L.: Accurate Derivation of Total and Stratospheric Vertical Columns of NO₂ from Ground-Based Zenith-Sky Measurements, *Journal of Atmospheric Chemistry*, 30, 163-172,
790 10.1023/A:1006019628406, 1998.
- Schoeberl, M., Douglass, A., Hilsenrath, E., Bhartia, P., Beer, R., Waters, J., Gunson, M., Froidevaux, L., Gille, J., Barnett, J., Levelt, P., and DeCola, P.: Overview of the EOS Aura mission, *Geoscience and Remote Sensing, IEEE Transactions on*, 44, 1066-1074, 10.1109/TGRS.2005.861950, 2006.
- Seinfeld, J.H.; Pandis, S.N.: *From Air Pollution to Climate Change*, *Atmos. Chem. Phys.*, ISBN 0-471-17815-2,
795 <https://doi.org/10.1063/1.882420>, 1998.
- Sioris, C. E., Haley, C. S., McLinden, C. A., von Savigny, C., McDade, I. C., McConnell, J. C., Evans, W. F. J., Lloyd, N. D., Llewellyn, E. J., Chance, K. V., Kurosu, T. P., Murtagh, D., Frisk, U., Pfeilsticker, K., Bösch, H., Weidner, F., Strong, K., Stegman, J., and Mégie, G.: Stratospheric profiles of nitrogen dioxide observed by Optical Spectrograph and Infrared Imager System on the Odin satellite, 108, <https://doi.org/10.1029/2002JD002672>, 2003.
- 800 Sivakumaran, V., Subramanian, K. P., and Kumar, V.: Lifetime measurement of NO₂ at 423–462 nm, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 69, 513-518, [https://doi.org/10.1016/S0022-4073\(00\)00092-3](https://doi.org/10.1016/S0022-4073(00)00092-3), 2001.

- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Journal Bulletin of the American Meteorological Society*, 96, 2059-2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- 805 Steinbacher, M., Zellweger, C., Schwarzenbach, B., Bugmann, S., Buchmann, B., Ordóñez, C., Prevot, A. S. H., and Hueglin, C.: Nitrogen oxide measurements at rural sites in Switzerland: Bias of conventional measurement techniques, *J. Geophys. Res.*, 112, D11307, <https://doi.org/10.1029/2006JD007971>, 2007.
- Sun, L., Li, R.-B., Tian, X.-P., and Wei, J.: Analysis of the Temporal and Spatial Variation of Aerosols in the Beijing-Tianjin-Hebei Region with a 1 km AOD Product, *Aerosol and Air Quality Research*, 17, 923-935, [10.4209/aaqr.2016.05.0185](https://doi.org/10.4209/aaqr.2016.05.0185), 2017.
- 810 Sun, Y., Jiang, Q., Wang, Z., Fu, P., Li, J., Yang, T., and Yin, Y.: Investigation of the sources and evolution processes of severe haze pollution in Beijing in January 2013, 119, 4380-4398, <https://doi.org/10.1002/2014JD021641>, 2014.
- Sundström, A. M., Kolmonen, P., Sogacheva, L., and de Leeuw, G.: Aerosol retrievals over China with the AATSR dual view algorithm, *Remote Sensing of Environment*, 116, 189-198, <https://doi.org/10.1016/j.rse.2011.04.041>, 2012.
- 815 Uppala, S. M., Kallberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Van De Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Holm, E., Hoskins, B. J., Isaksen, I., Janssen, P., Jenne, R., McNally, A. P., Mahfouf, J. F., Morcrette, J. J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Quarterly Journal of the Royal Meteorological Society*, 131, 2961-3012, [10.1256/qj.04.176](https://doi.org/10.1256/qj.04.176), 2005.
- van der A, R. J., D. H. M. U. Peters, H. Eskes, K. F. Boersma, M. Van Roozendael, I. De Smedt, and H. M. Kelder.: Detection of the trend and seasonal variation in tropospheric NO₂ over China, *J. Geophys. Res.*, 111, D12317, [doi:10.1029/2005JD006594](https://doi.org/10.1029/2005JD006594), 2006.
- 825 van Geffen, J., Boersma, K. F., Van Roozendael, M., Hendrick, F., Mahieu, E., De Smedt, I., Sneep, M., and Veefkind, J. P.: Improved spectral fitting of nitrogen dioxide from OMI in the 405-465 nm window, *Atmospheric Measurement Techniques*, 8, 1685-1699, [10.5194/amt-8-1685-2015](https://doi.org/10.5194/amt-8-1685-2015), 2015.
- van Geffen, J., Eskes, H., Compernelle, S., Pinardi, G., Verhoelst, T., Lambert, J. C., Sneep, M., ter Linden, M., Ludewig, A., Boersma, K. F., and Veefkind, J. P.: Sentinel-5P TROPOMI NO₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data, *Atmos. Meas. Tech.*, 15, 2037-2060, [10.5194/amt-15-2037-2022](https://doi.org/10.5194/amt-15-2037-2022), 2022a.
- 830 van Geffen, J. H. G. M., Eskes, H. J., Boersma, K. F. and Veefkind, J. P. TROPOMI ATBD of the total and tropospheric NO₂ data products, [https://sentinel.esa.int/documents/247904/2476257/Sentinel-5P-TROPOMI-ATBD-NO₂-data-products](https://sentinel.esa.int/documents/247904/2476257/Sentinel-5P-TROPOMI-ATBD-NO2-data-products) (last accessed: 9th March 2023), 2022b.
- Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B.,
- 835

- Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens Environ*, 120, 70-83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
- Verhoelst, T., Compernelle, S., Pinardi, G., Lambert, J. C., Eskes, H. J., Eichmann, K. U., Fjæraa, A. M., Granville, J.,
840 Niemeijer, S., Cede, A., Tiefengraber, M., Hendrick, F., Pazmiño, A., Bais, A., Bazureau, A., Boersma, K. F., Bogner, K.,
Dehn, A., Donner, S., Elokho, A., Gebetsberger, M., Goutail, F., Grutter de la Mora, M., Gruzdev, A., Gratsea, M., Hansen,
G. H., Irie, H., Jepsen, N., Kanaya, Y., Karagkiozidis, D., Kivi, R., Kreher, K., Levelt, P. F., Liu, C., Müller, M., Navarro
Comas, M., Piters, A. J. M., Pommereau, J. P., Portafaix, T., Prados-Roman, C., Puentedura, O., Querel, R., Remmers, J.,
Richter, A., Rimmer, J., Rivera Cárdenas, C., Saavedra de Miguel, L., Sinyakov, V. P., Stremme, W., Strong, K., Van
845 Roozendaal, M., Veefkind, J. P., Wagner, T., Wittrock, F., Yela González, M., and Zehner, C.: Ground-based validation of
the Copernicus Sentinel-5P TROPOMI NO₂ measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandonia
global networks, *Atmos. Meas. Tech.*, 14, 481-510, [10.5194/amt-14-481-2021](https://doi.org/10.5194/amt-14-481-2021), 2021.
- Wang, H., Lu, K., Chen, S., Li, X., Zeng, L., Hu, M., and Zhang, Y.: Characterizing nitrate radical budget trends in Beijing
during 2013–2019, *Science of The Total Environment*, 795, 148869, <https://doi.org/10.1016/j.scitotenv.2021.148869>, 2021.
- 850 Wang, T., Huang, X., Wang, Z., Liu, Y., Zhou, D., Ding, K., Wang, H., Qi, X., and Ding, A.: Secondary aerosol formation
and its linkage with synoptic conditions during winter haze pollution over eastern China, *Science of The Total Environment*,
730, 138888, <https://doi.org/10.1016/j.scitotenv.2020.138888>, 2020.
- Wang, T., Nie, W., Gao, J., Xue, L. K., Gao, X. M., Wang, X. F., Qiu, J., Poon, C. N., Meinardi, S., Blake, D., Wang, S. L.,
Ding, A. J., Chai, F. H., Zhang, Q. Z., and Wang, W. X.: Air quality during the 2008 Beijing Olympics: secondary pollutants
855 and regional impact, *Atmos. Chem. Phys.*, 10, 7603-7615, [10.5194/acp-10-7603-2010](https://doi.org/10.5194/acp-10-7603-2010), 2010.
- Wang, Y., Zhang, Y., Schauer, J. J., de Foy, B., Guo, B., and Zhang, Y.: Relative impact of emissions controls and
meteorology on air pollution mitigation associated with the Asia-Pacific Economic Cooperation (APEC) conference in
Beijing, China, *Science of The Total Environment*, 571, 1467-1476, <https://doi.org/10.1016/j.scitotenv.2016.06.215>, 2016.
- Wagner, T., Ibrahim, O., Shaiganfar, R., and Platt, U.: Mobile MAX-DOAS observations of tropospheric trace gases, *Atmos.*
860 *Meas. Tech.*, 3, 129–140, <https://doi.org/10.5194/amt-3-129-2010>, 2010.
- Wang, C., Wang, T. and Wang, P.: The Spatial–Temporal Variation of Tropospheric NO₂ over China during 2005 to 2018,
Atmosphere, 10(8), 444, <https://doi.org/10.3390/atmos10080444>, 2019.
- Wang, Y.-L., Song, W., Yang, W., Sun, X.-C., Tong, Y.-D., Wang, X.-M., Liu, C.-Q., Bai, Z.-P., and Liu, X.-Y.: Influences
of Atmospheric Pollution on the Contributions of Major Oxidation Pathways to PM_{2.5} Nitrate Formation in Beijing, *J.*
865 *Geophys. Res.*, 124, 4174-4185, <https://doi.org/10.1029/2019JD030284>, 2019.
- Wang, W., van der A, R., Ding, J., van Weele, M., and Cheng, T.: Spatial and temporal changes of the ozone sensitivity in
China based on satellite and ground-based observations, *Atmos. Chem. Phys.*, 21, 7253–7269, <https://doi.org/10.5194/acp-21-7253-2021>, 2021.

- Weber, H. G. and Bylicki, F.: NO₂ lifetimes by hanle effect measurements, *Chemical Physics*, 116, 133-140, 870 [https://doi.org/10.1016/0301-0104\(87\)80075-7](https://doi.org/10.1016/0301-0104(87)80075-7), 1987.
- Xie, S., Yu, T., Zhang, Y., Zeng, L., Qi, L., and Tang, X.: Characteristics of PM₁₀, SO₂, NO_x and O₃ in ambient air during the dust storm period in Beijing, *Science of The Total Environment*, 345, 153-164, <https://doi.org/10.1016/j.scitotenv.2004.10.013>, 2005.
- Xie, Y., Ding, A., Nie, W., Mao, H., Qi, X., Huang, X., Xu, Z., Kerminen, V.-M., Petäjä, T., Chi, X., Virkkula, A., Boy, M., 875 Xue, L., Guo, J., Sun, J., Yang, X., Kulmala, M., and Fu, C.: Enhanced sulfate formation by nitrogen dioxide: Implications from in situ observations at the SORPES station, *Journal of Geophysical Research: Atmospheres*, 120, 12679-12694, <https://doi.org/10.1002/2015JD023607>, 2015.
- Xue, L., Gu, R., Wang, T., Wang, X., Saunders, S., Blake, D., Louie, P. K. K., Luk, C. W. Y., Simpson, I., Xu, Z., Wang, Z., Gao, Y., Lee, S., Mellouki, A., and Wang, W.: Oxidative capacity and radical chemistry in the polluted atmosphere of Hong 880 Kong and Pearl River Delta region: analysis of a severe photochemical smog episode, *Atmos. Chem. Phys.*, 16, 9891-9903, [10.5194/acp-16-9891-2016](https://doi.org/10.5194/acp-16-9891-2016), 2016.
- You, T., R. G. Wu, and G. Huang: Differences in meteorological conditions between days with persistent and non-persistent pollution in Beijing, China. *J. Meteor. Res.*, 32(1), 81–98, [10.1007/s13351-018-7086-x](https://doi.org/10.1007/s13351-018-7086-x), 2018.
- Zhai, S., Jacob, D., Wang, X., Shen, L., Ke, L., Zhang, Y., Gui, K., Zhao, T., and Liao, H.: Fine particulate matter (PM_{2.5}) 885 trends in China, 2013–2018: separating contributions from anthropogenic emissions and meteorology, *Atmospheric Chemistry and Physics*, 19, 11031-11041, [10.5194/acp-19-11031-2019](https://doi.org/10.5194/acp-19-11031-2019), 2019.
- Zhang, X., Yin, Y., van der A, R., Lapierre, J. L., Chen, Q., Kuang, X., Yan, S., Chen, J., He, C., and Shi, R.: Estimates of lightning NO_x production based on high-resolution OMI NO₂ retrievals over the continental US, *Atmos. Meas. Tech.*, 13, 1709-1734, [10.5194/amt-13-1709-2020](https://doi.org/10.5194/amt-13-1709-2020), 2020.
- 890 Zhao, X., Fioletov, V., Alwarda, R., Su, Y., Griffin, D., Weaver, D., Strong, K., Cede, A., Hanisco, T., and Tiefengraber, M. J. R. S.: Tropospheric and Surface Nitrogen Dioxide Changes in the Greater Toronto Area during the First Two Years of the COVID-19 Pandemic, *Remote Sensing*, 14, 1625, <https://doi.org/10.3390/rs14071625>, 2022a.
- Zhao, W., Chen, S., Zhang, H., Wang, J., Chen, W., Wu, R., Xing, W., Wang, Z., Hu, P., Paio, J., and Ma, T.: Distinct Impacts of ENSO on Haze Pollution in the Beijing–Tianjin–Hebei Region between Early and Late Winters. *J. Climate*, 35, 895 687-704, doi.org/10.1175/JCLI-D-21-0459.1, 2022b.
- Zheng, G. J., Duan, F. K., Su, H., Ma, Y. L., Cheng, Y., Zheng, B., Zhang, Q., Huang, T., Kimoto, T., Chang, D., Pöschl, U., Cheng, Y. F., and He, K. B.: Exploring the severe winter haze in Beijing: the impact of synoptic weather, regional transport and heterogeneous reactions, *Atmos. Chem. Phys.*, 15, 2969-2983, [10.5194/acp-15-2969-2015](https://doi.org/10.5194/acp-15-2969-2015), 2015.