

Fast retrieval of XCO₂ over East Asia based on the OCO-2 spectral measurements

Fengxin Xie^{1,a}, Tao Ren¹, Changying Zhao¹, Yuan Wen², Yilei Gu², Minqiang Zhou³, Pucai Wang³, Kei Shiomi⁴, and Isamu Morino⁵

¹China-UK Low Carbon College, Shanghai Jiao Tong University, Shanghai, China

²Shanghai Institute of Satellite Engineering, Shanghai, China

³Institute of Atmospheric Physics, Chinese Academy of Science, Beijing, China

⁴Earth Observation Research Center, Japan Aerospace Exploration Agency, 2-1-1 Sengen, Tsukuba, Ibaraki 305-8505, Japan

⁵Satellite Remote Sensing Section and Satellite Observation Center, Earth system Division, National Institute for Environmental Studies, Onogawa 16-2, Tsukuba, Ibaraki 305-8506, Japan

^aCurrently at: Atmosphere and Ocean Research Institute, the University of Tokyo, Chiba, Japan

Correspondence: Tao Ren (tao.ren@sjtu.edu.cn)

Abstract. The increase in greenhouse gas concentrations, particularly CO₂, has significant implications for global climate patterns and various aspects of human life. Spaceborne remote sensing satellites play a crucial role in high-resolution monitoring of atmospheric CO₂. However, the next generation of greenhouse gas monitoring satellites is expected to face challenges, particularly in terms of computational efficiency in atmospheric CO₂ retrieving and analysis. To address these challenges, this study focuses on improving the speed of retrieving the column-averaged dry air mole fraction of carbon dioxide (XCO₂) using spectral data from the OCO-2 satellite, while still maintaining retrieval accuracy. A novel approach based on neural network (NN) models is proposed to tackle the nonlinear inversion problems associated with XCO₂ retrievals. The study employs a data-driven supervised learning method and explores two distinct training strategies. Firstly, training is conducted using experimental data obtained from the inversion of the operational optimization model, which are released as the OCO-2 satellite products. Secondly, training is performed using a simulated dataset generated by an accurate forward calculation model. The inversion and prediction performance of the machine learning model for XCO₂ are compared, analyzed, and discussed for the observed region over East Asia. The results demonstrate that the model trained on simulated data accurately predicts XCO₂ in the target area. Furthermore, when compared to OCO-2 satellite product data, the developed XCO₂ retrieval model not only achieves rapid predictions (<1 ms) with good accuracy (2 ppm or approximately 0.5%), but also effectively captures sudden increases of XCO₂ plumes near industrial emission sources. The accuracy of the machine learning model's retrieval results is validated against reliable data from TCCON sites, demonstrating its capability to capture CO₂ seasonal variations and annual growth trends effectively.

1 Introduction

Since the industrial revolution, human activities have released large amounts of greenhouse gases, primarily carbon dioxide, into the atmosphere. This continual increase in emissions has led to global warming and disrupted human societies and ecosys-

tems (Zehr, 2015). Accurately estimating atmospheric carbon fluxes is critical for implementing effective emission reduction strategies at national and regional levels. However, precise carbon flux estimates require assimilating carbon dioxide concentration data across regions, using measurements of atmospheric column-averaged dry air mole fraction of carbon dioxide (XCO_2) (Jin et al., 2021). Direct measurement methods like balloons or aircraft have challenges obtaining global-scale data.

25 Currently, the main monitoring approach uses spectrometers to record spectra in CO_2 absorption bands, followed by inversion algorithms to derive XCO_2 . The two primary monitoring methods are ground-based monitoring stations and satellite remote sensing.

The Total Carbon Column Observing Network (TCCON) provides ground-based monitoring of atmospheric carbon dioxide through a global network of high-precision Fourier transform spectrometers (Wunch et al., 2011, 2015). However, TCCON sites

30 are sparsely distributed and cannot be deployed in regions with unfavorable geography or harsh climates. Consequently, the network lacks the extensive spatial coverage required for comprehensive global carbon monitoring and carbon cycle analysis. Nevertheless, the ultra-high spectral resolution of TCCON spectrometers enables highly accurate retrievals of XCO_2 . Under clear sky conditions, TCCON precision can reach 0.1% (<0.4 ppm). Under relatively clear conditions with minimal clouds and aerosols, precision remains within 0.25% (<1 ppm) (Messerschmidt et al., 2011). Due to such high precision and accuracy,

35 TCCON data are invaluable for validating satellite-based XCO_2 products (Cogan et al., 2012; Wunch et al., 2017; Liang et al., 2017) and comparing them to carbon cycle models. However, the spatial limitations of the network underscore the need for satellite remote sensing to provide regular global measurements of atmospheric carbon dioxide.

High-spectral-resolution greenhouse gas monitoring satellites employ spectrometers on orbit to measure solar radiation spectra after interaction with the Earth's atmosphere and ground surface (Meng et al., 2022). Unlike ground monitoring, satellite remote sensing offers broader spatial coverage and more flexible temporal observation globally. Consequently, satellite remote sensing has become vital for greenhouse gas monitoring worldwide. Notable ongoing passive CO_2 observation missions include China's TanSat (Liu et al., 2018), Japan's GOSAT (2009) and GOSAT-2 (2018) (Hamazaki et al., 2005; Kuze et al., 2009; Imasu et al., 2023), and the United States' OCO-2 (2014) and OCO-3 (2018) (Crisp et al., 2017; Eldering et al., 2019). Upcoming missions are France's MicroCarb by CNES (Cansot et al., 2023), ESA's CO_2M (Sierk et al., 2021) and

45 GOSAT-GW (Matsunaga and Tanimoto, 2022). The next-generation greenhouse gases monitoring satellites mainly address the challenge of improving the spatial and temporal resolutions of observations. However, single satellites still have resolution, coverage, and meteorological limitations for regional emission monitoring. Enhancing satellite sensor performance alone cannot produce datasets sufficient for monitoring carbon sources and sinks. Improving the accuracy and efficiency of satellite data inversion is also crucial. Integrating data from multiple satellites into a coordinated system is necessary to fully capture

50 dynamic changes in regional carbon sources and sinks. Developing new high-precision, high-throughput inversion methods to efficiently derive accurate greenhouse gas concentration distributions from satellite data is a key challenge needing attention.

The mainstream inversion algorithms (O'Dell et al., 2012; Crisp et al., 2012; Yoshida et al., 2013) for retrieving greenhouse gas concentrations from high-spectral-resolution satellite measurements are based on nonlinear Bayesian optimization theory (Rodgers, 2000) and full physics models. In essence, these algorithms operate by iteratively adjusting estimated gas concentration profiles and other atmospheric and surface parameters in a radiative forward model to minimize the mismatch

55

between simulated and observed spectra. More specifically, the inversion process starts with an initial atmospheric state guess, including trace gas concentration profiles as functions of pressure/altitude. Radiative transfer equations are then solved to simulate the top-of-atmosphere radiance spectrum observed by the satellite for this atmospheric state. The simulated spectrum is compared to the actual observed spectrum, calculating the difference, covariance and cost function. The input gas profiles and model parameters are iteratively adjusted to reduce the cost function over multiple rounds of radiative transfer simulations. Once simulated spectra closely match observations, the model state is output as the retrieved concentration profile. However, executing these complex optimizations requires computationally expensive interpolation of high-spectral-resolution gas absorption reference data and solving the radiative transfer equations in each iteration. Running the radiative forward model repeatedly for every adjusted atmospheric state also leads to slow overall inversion. Consequently, optimization-based retrievals struggle to match increasing satellite observation volumes and throughput needs. This inherent inefficiency has become a major obstacle to operational greenhouse gas monitoring using current and planned high-resolution spectrometers. While rigorous, standard nonlinear optimization retrievals lack the speed and scalability required for high-precision real-time or near-real-time greenhouse gas mapping satellite-based greenhouse gas mapping. Overcoming this bottleneck necessitates new inversion approaches that can ingest high-resolution spectral data and retrieve concentrations with both accuracy and computational efficiency.

In recent years, machine learning has demonstrated exceptional performance across various research fields, with the discovery of potential nonlinear relationships between data being one of its fundamental and crucial applications. Regarding the important applications of carbon dioxide (CO_2) retrieving, Carvalho et al. (2010) attempted to retrieve the vertical CO_2 profiles using spectral data from SCIAMACHY's 6 channels (1000-1700 nm). The overall precision and bias of the retrieved results were estimated to be approximately 1.0% and less than 3.0%, respectively. Gribanov et al. (2010) developed a two-hidden-layer multilayer perceptron (MLP) model to retrieve CO_2 vertical concentrations by reflected solar radiation measured by the GOSAT TANSO-FTS sensor, achieving an inversion accuracy better than 1 ppm for CO_2 column-averaged values and better than 4 ppm for surface CO_2 concentrations for the test samples. In the study conducted by Zhao et al. (2022), a two-step machine learning approach was developed for retrieving atmospheric XCO_2 using spectral data from the GOSAT weak- CO_2 band. They established a direct one-dimensional line-by-line forward model to simulate GOSAT's observed spectra within the 6180-6280 cm^{-1} spectral interval, forming the foundation for training their machine learning model. The retrieval model operates by initially obtaining the atmospheric spectral optical thickness, followed by extracting XCO_2 from this optical thickness spectra. As a proof-of-concept, the method was tested in Australia under clear sky conditions using GOSAT's spectra, demonstrating an accuracy of approximately 3 ppm for XCO_2 retrieval. The study also discussed potential enhancements to further refine the accuracy of this retrieval method. Keely et al. (2023) employed the machine learning method of Extreme Gradient Boosting (Chen and Guestrin, 2016) to develop a nonlinear bias correction approach for OCO-2 version 10 product, significantly reducing systematic errors in CO_2 measurements and improving data quality, with an increase in sounding throughput by 14%. David et al. (2021) and Bréon et al. (2022) attempted to establish correlations between XCO_2 in the European Centre for Medium-Range Weather Forecasts' CAMS (Copernicus Atmosphere Monitoring Service) database and OCO-2 satellite monitoring spectra using multilayer perceptron artificial neural network models. However, their recent research (Bacour et al.,

2023) indicates that when the test dataset extends beyond the time range covered by the training dataset, the predicted results show a slight bias, approximately 2.5 ppm per year. Practical deployment of machine learning techniques for remote sensing demands additional research into the generalization performance of models on new observational data distributions beyond those encountered during training.

95 In the present paper, a proof-of-concept study demonstrates a novel machine learning strategy to accurately and efficiently retrieve atmospheric XCO₂ value from OCO-2 satellite spectral measurements. The model rapidly retrieves XCO₂ directly from OCO-2 spectral data, eliminating the need for repetitive radiative transfer simulations required by traditional nonlinear optimization retrieval algorithms. Additionally, the model enables the prediction of future XCO₂ values. The method was validated by comparing the retrieved XCO₂ against OCO-2 satellite version-10r products and ground-based TCCON mea-
100 surements, confirming the accuracy of our efficient spectral inversion approach. The model also successfully demonstrated its capability to detect local plume features, indicating its potential utility in monitoring and analyzing specific emission sources. A major innovation in the present study is using accurate radiative transfer simulations to generate the training data, rather than relying solely on experimental data products. This simulation-based training approach could help overcome limitations in existing experimental data. Additionally, our neural network model achieves XCO₂ retrieval speeds orders of magnitude
105 faster than traditional methods, reducing computation time from multiple seconds to less than 1 millisecond. This dramatic improvement in retrieval efficiency could enable real-time processing of the massive data volumes expected from next-generation greenhouse gas monitoring satellites. Importantly, our model achieves a precision of around 2 ppm, competitive with the current state-of-the-art in retrieval accuracy. We also demonstrate the ability to accurately capture temporal variations and trends in XCO₂ by validating against reliable TCCON ground-based data. This level of verifiable performance is an important capa-
110 bility. This provides an effective solution for rapid inversion of large-scale, high-spectral-resolution remote sensing data in the future.

2 The machine learning based XCO₂ retrieval model

2.1 Targeted area and data screening

This proof-of-concept study aims to develop and validate an accurate and efficient machine learning-based XCO₂ retrieval
115 model applied to the long OCO-2 time series for the East Asian region. Currently, similar global XCO₂ retrieval models rely on computationally intensive physical models. Our goal is to demonstrate a more efficient data-driven approach using MLP neural networks.

Before developing the machine learning based fast retrieval model, we implemented several preprocessing steps on the OCO-2 observational dataset (OCO-2 Science Team et al., 2020a) for the target East Asian area spanning between 20°N-
120 45°N and 110°E-145°E, as shown in Fig. 1. Specifically, we filtered the data both spatially and temporally to focus only on observations within this geographic region and time period of interest (2016-2021). Additionally, we filtered the data to only include Nadir mode observations marked as “Good” based on the quality flag indicator (“xco2_quality_flag” = 0 in OCO-2 Lite v10r files (OCO-2 Science Team et al., 2020b)), as these represent the highest quality OCO-2 measurements.

Several TCCON ground stations located in this region (e.g. Hefei, Saga, Tsukuba, Xianghe, Anmyeondo and Rikubetsu),
125 as shown in Fig. 1, provide valuable ground-truth XCO₂ data for validating the MLP model predictions. If the model can
accurately reproduce the TCCON observations from corresponding OCO-2 measurements, it suggests the model has learned
meaningful relationships between the satellite data and underlying CO₂ concentrations.

Furthermore, the successful demonstration of accurate XCO₂ retrieval over East Asia is a first step toward expanding this
approach globally. The model could be retrained or supplemented with additional regional data to extend coverage. By combin-
130 ing reliable regional MLP models, global XCO₂ maps could be retrieved. This “jigsaw puzzle” strategy would further validate
the feasibility of global-scale machine learning-based XCO₂ retrievals from satellite observations.

2.2 The artificial neural network architecture

This study introduces a multilayer perceptron (MLP) neural network model for estimating XCO₂ from OCO-2 satellite obser-
vations. Inspired by David et al. (2021) and Bréon et al. (2022), the “MLP-XCO₂” model input layer is designed based on the
135 measurement principles of OCO-2 and atmospheric radiative transfer effects on the observed spectra, the artificial neural net-
works architect is shown in Fig. 2. Specifically, the MLP model input layer consists of spectral information, surface pressure,
the corresponding year, and geographical observation information as summarized in Table 1 and explained below.

Spectral Information: The OCO-2 satellite instrument measures high-resolution spectra in three spectral bands centered
around 0.76, 1.6, and 2.0 μm, referred to as the O₂-A, weak CO₂ (WCO₂), and strong CO₂ (SCO₂) bands, respectively (OCO-2
140 Science Team et al., 2019a). However, only the WCO₂ and SCO₂ bands are used as inputs for current XCO₂ retrievals. The O₂-
A band is excluded as it lacks significant information needed to directly estimate XCO₂, based on radiative transfer principles.
Instead, the O₂-A band is primarily used in OCO-2’s operational full-physics algorithm for rapid cloud and aerosol screening
prior to CO₂ retrieval (O’Dell et al., 2012), saving substantial computational costs. Each OCO-2 spectral band is sampled
by 1024 detector pixels. However, over time some detectors have degraded or become unstable in the space environment,




Figure 1. The target area for the East Asia region, distribution of observation points (from OCO-2 L2std v10r files) of OCO-2 Nadir mode
in January 2016, and the distribution of TCCON sites in this area.




Figure 2. Schematic diagram of the MLP model. The input layer includes two interpolated radiance of WCO₂ and SCO₂ band, the geographical observation information, surface pressure, and the corresponding year.

Table 1. Detailed lists of the input layer of the MLP-XCO₂ model

| Input elements | Variables | Number |
|--------------------------|------------------|--------|
| Spectral information | WCO ₂ | 525 |
| | SCO ₂ | 755 |
| Geographical information | Solar zenith | 1 |
| | Relative azimuth | 1 |
| Others | Year | 1 |
| | Pressure | 1 |
| Total | | 1284 |

145 resulting in pixels being flagged as “bad samples” in quality filters (Marchetti et al., 2019). To maximize high-quality training data, additional preprocessing is performed on the WCO₂ and SCO₂ bands. Specifically, the beginning and ending spectral ranges corresponding to the most degraded detectors are removed. The remaining good quality spectra are re-sampled into 525 and 755 wavelength points for the WCO₂ and SCO₂ bands, respectively (spectral points in wavelength are detailed in Table 2). To enhance the CO₂ absorption line information, each input spectrum is normalized by dividing the mean radiance
150 within a nearby spectrally transparent window lacking absorption features (1.6056-1.6059 μm using 10 points for WCO₂; 2.0602-2.0607 μm using 15 points for SCO₂).

Geographical Information: The model is designed to accept two key observation geometry angles that are determined by the relative positions of the Sun, satellite, and ground observation point. These include solar zenith angle and relative azimuth angle. The solar zenith angle (SZA) features prominently as a cosine term in the radiative transfer equation that defines the
155 atmospheric radiative processes. Thus, SZA is pre-converted to its cosine form for model input. The relative azimuth angle is a comprehensive angle that jointly combines the solar azimuth angle and the satellite azimuth angle. It is important to emphasize

Table 2. Wavelength spacing of the input spectra

| Band | Spectral range [μm] | Spectral points [μm] |
|------------------|----------------------------------|---|
| WCO ₂ | 1.5990-1.6151 | $\lambda_1 = 1.5990, \lambda_{i+1} = \lambda_i + 10^{-4} (6.10 - 3.60\lambda_i), i = 1-524$ |
| SCO ₂ | 2.0478-2.0779 | $\lambda_1 = 2.0478, \lambda_{i+1} = \lambda_i + 10^{-4} (7.58 - 3.48\lambda_i), i = 1-754$ |

that the satellite zenith angle is not utilized in this study. Our current research is based on the Nadir mode of the OCO-2 satellite observation. In the Nadir observing mode, the satellite zenith angle is assumed to be nearly perpendicular to the Earth’s surface, theoretically approaching zero degrees.

160 **Other parameters:** In addition to the primary inputs, two other parameters play critical roles in the MLP-XCO₂ model: the surface pressure, and the corresponding year (e.g., 2016 or 2017, etc.). The surface pressure as a critical parameter, is not directly available and is typically retrieved based on the O₂-A band. To accurately provide surface pressure information to the neural network model, we first retrieve the surface pressure from the L2MET file (OCO-2 Science Team et al., 2019b) using another compact machine learning model, referred to as MLP-P, before inputting it into the MLP-XCO₂ model. The
165 MLP-P model, trained on historical OCO-2 product data, provides stable future predictions of surface pressure. The inclusion of the corresponding year is due to the gradual degradation and necessary adjustments over time to the OCO-2 satellite’s instrument response parameters. These parameters include the instrument line shape function, bad pixel list, and spectral dispersion coefficients. While these parameters may remain consistent over short periods, they exhibit significant variations over the years, which can affect the long-term reliability of the model.

170 3 Satellite product data based machine learning model

In this section, we first developed the MLP-XCO₂ model using the OCO-2 v10r product dataset. The primary goal was to optimize the hyperparameters of the MLP-XCO₂ network. On one hand, we aimed to confirm whether the “slow bias”, as shown in Bacour et al. (2023) is a universal issue across machine learning models with similar architectures. On the other hand, by fixing the hyperparameters of the MLP-XCO₂ network structure, we sought to develop a comparable model using
175 simulated data in later sections. In theory, MLP models using identical hyperparameters should possess the same fitting and generalization abilities. By first presenting results from a model trained solely on satellite product data, we can demonstrate the limitations of these satellite data-based models. This then motivates the development of new machine learning strategies to overcome these limitations, as discussed in later sections.

Following the target areas and data screening methods discussed previously, we collected observational data from the OCO-2
180 v10r L1B database and XCO₂ results from the L2std database. Specifically, we obtained data from March, June, September and December spanning the years 2016 to 2020. This timeframe was chosen to provide a comprehensive training and testing set for our analysis. In total, the dataset encompassed 194150 samples collected over this five-year period. The year-wise distribution

of the samples is as follows: 38626 samples from 2016, 39850 from 2017, 35945 from 2018, 36452 from 2019, and 43277 from 2020.

185 Before training the MLP-XCO₂ model, we should firstly focus on the development and training of the MLP-P model, which is crucial for accurate surface pressure retrieval and subsequent input into the MLP-XCO₂ model. The MLP-P model, with only {500, 200, 50} hidden layer nodes and the same input layer composition as the MLP-XCO₂. All hidden layers use ReLU activation functions. Trained on historical OCO-2 product data from 2016 to 2018, the MLP-P model has demonstrated the ability to provide stable predictions of surface pressure for the years 2019 to 2020, as illustrated in Fig. 3.

190 Following the successful implementation and validation of the MLP-P model, we proceeded to construct the MLP-XCO₂ model. To balance model complexity and performance, the MLP-XCO₂ architecture (Fig. 2) comprises five hidden layers, with 1000, 500, 300, 100 and 20 nodes, respectively. All hidden layers also use ReLU activation functions. The output layer contains a single node to predict XCO₂ values, with a linear activation function. Upon developing the MLP-XCO₂ model architecture, in this section, we independently trained two versions of the MLP-XCO₂ model, each based on the aforementioned model
195 structure but with different training and testing datasets.




Figure 3. Comparison of pressure results predicted by the MLP-P model, trained with OCO-2 v10r product from 2016-2018, versus results retrieved by OCO-2 v10r product from 2019-2020.

Historical Data Training: The first MLP-XCO₂ model based on OCO-2 product data was trained using historical XCO₂ collected from 2016 to 2018. The test set for this model comprised product data from the years 2019 and 2020. This setup allows us to assess the model’s predictive performance using a straightforward historical data approach.

200 **Skipped-Year Training:** The second version of the model was trained using data from the years 2016, 2018, and 2020. The test set for this MLP-XCO₂ model included the skipped years, 2017 and 2019. This unique approach enables a clearer and direct comparison of the potential limitations of relying solely on historical data for future predictions.

Figure 4 presents the results for the two trained MLP-XCO₂ models on their respective 10% out-of-sample testing datasets. The first subplot illustrates the predictions of the historical data training model from the 2016-2018 data, and the second subplot shows similar predictions for the skipped-year training model. Both models achieve high accuracy on these testing datasets, with a Root Mean Square Error (RMSE) close to 1 ppm and R-squared score (R²) larger than 0.91. These results demonstrate the robust interpolation capabilities of both models within their respective training periods, indicating their effectiveness in handling known observed scenarios.




Figure 4. Comparison of 10% out-of-sample XCO₂ testing cases predicted by the MLP-XCO₂ model versus results retrieved by OCO-2 v10r product. Panel (a) is for the historical data training model, while (b) is for the skipped-year training model. The solid red lines in the figure correspond to perfect agreement, where shadow areas around the solid red lines represent ±1% of XCO₂ deviations.

Figure 5 evaluates the generalization capabilities of each MLP-XCO₂ model on testing sets comprising years not included in their respective training datasets. These test sets represent periods outside the range of years used for training. Here, we solely observed a noticeable positive bias in the predictions from the historical data training model. In contrast, the skipped-year training model did not exhibit this bias. Performance remains highly accurate on these out-of-range points, further validating the model's robustness for XCO₂ prediction within skipped years.

Globally, the average XCO₂ in the atmosphere shows a stable annual increase, with an observed rise of approximately 2.5-3 ppm per year. However, despite the inclusion of the corresponding year in the input layer as a high-correlation parameter, there remains a limitation in capturing the atmospheric CO₂'s potential rising trend. This highlights the limitations of models trained solely on historical satellite data, motivating the development of new techniques to incorporate external information about temporal CO₂ dynamics.




Figure 5. Comparison of XCO₂ results predicted by the MLP-XCO₂ model versus results retrieved by OCO-2 v10r product on test sets consisting of years not included in the training periods. Panels (a-1) and (a-2) are for the historical data training model on the 2019 and 2020 test sets, respectively. Panels (b-1) and (b-2) are for the skipped-year training model on the 2019 and 2020 test sets, respectively. The solid red lines in the figure correspond to the perfect agreement, where shadow areas around the solid red lines represent $\pm 1\%$ of XCO₂ deviations.

4 Simulation data based machine learning model

In the previous section, the MLP-XCO₂ model showed excellent interpolation within the training data range but exhibited bias when predicting outside this period. To eliminate this bias, we propose using an accurate forward model to simulate representative training data that covers future atmospheric conditions. If we can pre-generate atmospheric profiles that capture possible future states, and simulate their corresponding spectral radiance using an accurate forward model, the MLP-XCO₂ model can pre-learn future satellite observations. This could prevent the incremental annual bias and enable accurate XCO₂

prediction. The effectiveness of this approach depends on the forward model accuracy and representativeness of the simulated
 225 atmospheres (Zhao et al., 2022).

It is therefore critical to select an appropriate radiative transfer forward model with proven reliability in simulating spectral
 radiance under varying atmospheric conditions. The model must precisely capture the relationship between trace gas concen-
 230 trations, meteorological states, and resulting spectral signatures. With accurate simulations, the machine learning model can
 generalize robustly to future atmospheric scenarios. The representative training data should span the expected range of atmo-
 spheric variability in XCO₂ and interfering species like water vapor. A broad sampling of the state space is key for the model
 to learn a robust mapping to XCO₂ across multiple atmospheric regimes. The following sections describe our approach for
 accurate spectral radiative transfer simulations and possible (realistic) atmospheric profile generations.

4.1 Forward model

In this study, we developed a forward radiative transfer calculation model using the ReFRACtor (Reusable Framework for
 235 Retrieval of Atmospheric Composition) software (McDuffie et al., 2018). ReFRACtor is an extensible framework for multi-
 instrument atmospheric radiative transfer and retrieval, originally derived from the operational OCO-2 retrieval program. Al-
 though ReFRACtor contains both radiative transfer and retrieval capabilities, we only utilized the radiative transfer component.
 Specifically, we configured ReFRACtor to simulate top-of-atmosphere radiance spectra that would be observed by OCO-2.
 These simulated observations were then used to generate a large training dataset for our machine learning model, MLP-XCO₂.

240 The OCO-2 satellite primarily observes the radiative spectra in the short-wave infrared (SWIR) band. Over the range of
 SWIR, the impact of thermal emission can be ignored when simulating the spectra (Crisp et al., 2021). To simulate OCO-2's
 observed spectra in the WCO₂ band around 1.6 μm and the SCO₂ band around 2.06 μm, the ReFRACtor model numerically
 solves the Eq. (1) of the radiative transfer equation (RTE) (Modest and Mazumder, 2021):

$$\mu \frac{dI(\tau, \mu, \phi)}{d\tau} = -I(\tau, \mu, \phi) + J(\tau, \mu, \phi) \quad (1)$$

245 where I_η is the observed spectra, μ is the cosine of the observation zenith angle (e.g., $\mu = \cos\theta$), τ is the vertical optical depth
 which can be column-integrated from the molecular absorption coefficients and optical path, ϕ is the azimuthal angle relative
 to the observation point for the satellite and the sun, and J represents the scattering components and inhomogeneous source
 term, describing both single scattering and multiple scattering contributions. The term J in RTE can be expressed as Eq. (2):

$$J(\tau, \mu, \phi) = \frac{\omega}{4\pi} \int_{-1}^1 \int_0^{2\pi} P(\tau, \mu, \phi; \mu', \phi') I(\tau, \mu', \phi') d\mu' d\phi' + \frac{\omega}{4\pi} P(\tau, \mu, \phi; \mu', \phi') I_0 \exp(-\tau/\mu_0) \quad (2)$$

250 where ω is the single scattering albedo, P is the scattering phase function, μ' and ϕ' are the cosine and azimuth angle of the
 incident direction angle in each direction, μ_0 is the cosine of the solar zenith, and I_0 is the solar intensity in the top of the
 atmosphere.

The ReFRACtor model uses a hybrid model to solve RTE. Specifically, the radiative transfer software LIDORT (Spurr, 2008)
 is applied for the scalar and Jacobian computation. Concurrently, the two-order scattering model (Natraj and Spurr, 2007)

255 is utilized for the additional radiance correction. Within this framework, the ReFRACtor model comprehensively considers five types of scatter particles for each sounding: two types of clouds, two types of tropospheric aerosols, and one type of stratospheric aerosol. The single scattering optical properties for each cloud and aerosol particle, including cross-section, single scattering albedo, and scattering phase matrix, have been pre-computed and tabulated for the forward calculations. Furthermore, the model determines surface reflectance as a quadratic spectral albedo for each band which is derived from the
260 bidirectional reflectance distribution function (BRDF).

An essential step for developing the forward calculation model is referencing the pre-computed look-up table of H₂O and CO₂ to obtain the required spectral absorption coefficients. In this study, the ABSCO v5.1 database (Absorption Coefficient Table (Payne et al., 2020)) was applied for this purpose. Additionally, we identified and corrected an overestimation of the solar continuum in ReFRACtor compared to the OCO-2 Level 2 algorithm (Crisp et al., 2021). Without this correction, there
265 would have been approximately 3 % overestimation in the 1.6 μm band and 6.5 % in the 2.06 μm band. By reducing the solar continuum, our forward model aligned better with the OCO-2 spectral measurements. These configurations of the absorption coefficients and solar continuum were essential to accurately simulate OCO-2 spectra for generating training data across a variety of observing conditions.

To assess the performance of the forward model, we selected four distinct global locations in the year of 2017. The goal was
270 to replicate the OCO-2 observed spectra for both the WCO₂ 1.6 μm absorption band and the SCO₂ 2.06 μm absorption band at the four locations. By accessing the OCO-2 L2std database, we acquired atmospheric conditions and pertinent geographical data (including spectral albedo, surface pressure, and observation angles) specific to these chosen locations. The outcomes of our simulations for these four locations are visually depicted in Fig. 6 and Fig. 7, respectively for the two bands, with accompanying residual plots displayed in the lower panels. It is worth noting that the simulated results exhibit a high level of
275 agreement with the observed OCO-2 spectra; the relative error remains under 1%, underlining the robustness of the established forward model. The remarkable agreements between the observed and simulated spectra indicates the excellent performance of the forward radiative transfer model. This performance is particularly evident in accurately replicating the satellite observations from OCO-2. As a result, this forward model serves as a reliable tool for the development of machine learning models trained using simulated spectral data.

280 4.2 Training data generation

To optimize the training of the MLP-XCO₂ model, it is essential that the input training vectors cover a wide range of realistic variations. Although the idea of randomizing all input parameters to enhance diversity might appear attractive, simulating satellite spectra involves managing a multitude of interdependent variables. In addition to the CO₂ vertical profile, factors such as surface pressure, temperature profile, water vapor, aerosols, and observation geometry must be accurately represented.
285 Randomizing all of these parameters would require an impractical amount of data and could result in combinations that have no real-world relevance. For example, the four viewing angles determined by the sun, observation point, and the OCO-2 satellite have fixed combinations during the satellite's regular operation. Therefore, randomly selecting angle combinations lacks practical significance. To ensure that the training data covers valid variations, we conducted an analysis of historical




Figure 6. Comparisons of the OCO-2 observed spectra with the simulated ones from the modified ReFRACtor forward calculation model in WCO₂ band. The lower panel shows the relative error between the spectrum observed by the OCO-2 satellite and that simulated by the forward calculation model. Subplots (a)-(d) correspond to test samples from four different regions. The input vectors for the ReFRACtor model were derived from OCO-2 L2std retrieved results.

OCO-2 retrievals. This analysis revealed consistent seasonal patterns and year-to-year trends in most parameters. This supports
 290 the idea of selecting representative samples from statistical distributions rather than relying on complete randomization. By
 carefully considering the relationships between parameters and the realities of satellite observations, we can create a reasonably
 sized training dataset that effectively captures the range of expected predictions.

The generation of the vertical CO₂ profile is especially critical among all input parameters. This dataset theoretically deter-
 mines the generalization domain of the MLP-XCO₂ model. In the forward model based on the ReFRACtor model, the
 295 atmospheric CO₂ profile is segmented into 20 sub-layers by pressure. By statistically analyzing the OCO-2 retrieved CO₂
 profiles in the target East Asia area from 2016-2018, the box plots for atmospheric CO₂ concentration in each sub-layer are
 shown in Fig. 8(a), and the historic XCO₂ results from the OCO-2 product data are showing in Fig. 8(b). From the upper atmo-




Figure 7. Comparisons of the OCO-2 observed spectra with the simulated ones from the corrected ReFRACtor forward calculation model in SCO₂ band. The lower panel shows the relative error between the spectrum observed by the OCO-2 satellite and that simulated by the forward calculation model. Subplots (a)-(d) correspond to test samples from four different regions. The input vectors for the ReFRACtor model were derived from OCO-2 L2std retrieved results.

sphere down to the ground surface, the variability of CO₂ concentrations gradually increases. This challenges the ability for the standardization of atmospheric CO₂ profiles, particularly closer to the Earth's surface. Fortunately, a consistent year-on-year rise in CO₂ concentrations in each sub-layer has been observed over time. Consequently, in our research, we have proposed a method for generating subsequent CO₂ atmospheric profiles. We incrementally increase the CO₂ concentration by 2.5 ppm annually, starting from the 2016 OCO-2 retrieved CO₂ vertical profile. This approach ensures that we encompass a range of plausible atmospheric CO₂ distributions with realistic shapes, enabling the generation of simulated spectra for the designated training years.

In addition to the CO₂ profile, Fig. 9 illustrates the year-to-year trends of various observed parameters essential for the forward calculation model in the East Asian region. These parameters, although they display seasonal variations, consistently




Figure 8. Panel (a) is the boxplot of the vertical distribution of CO₂ profiles (from OCO-2 L2std files) retrieved by the OCO-2 satellite over East Asia in Nadir mode from 2016 to 2018. The horizontal-axis represents the atmospheric layers from layer 1 (top of atmosphere) to layer 20 (near-surface). The upper and lower bounds of each box show the maximum and minimum CO₂ concentrations recorded within that layer for each year. Panel (b) is the scatter plot of historic XCO₂ results retrieved by the OCO-2 inversion program (from L2std files)

exhibit annually cyclic patterns. Given that the OCO-2 satellite conducts global observations in cycles of approximately half a month (15-16 days), this study employed observation parameters and priori data for atmospheric profiles, except for CO₂, from the year 2016 as a reference. These reference data were repetitively utilized for generating simulations in subsequent years.

310 Regarding the quadratic spectral albedo, the constant term in the training data samples is uniformly set to 1 (to be normalized before being processed by the neural network). The slope and the quadratic coefficient are stochastically sampled within the range of values corresponding to the retrieval results based on the OCO-2 L2 products.

Based on 60,000 uniformly sampled observation data points exclusively from the OCO-2 satellite throughout 2016, we randomly separated it into six sets of 10,000 data points each. Each set represents CO₂ profiles from 2016 to the end of 2021,

315 with a yearly increase of 2.5 ppm added to the original data, reflecting projected future profiles. The forward model was used to generate the corresponding simulated spectra for each set. These simulated samples serve as the foundational dataset for training the new MLP-XCO₂ machine learning model. It's important to note that this new model relies solely on the data recorded by the OCO-2 satellite in 2016 as its reference. However, it is essential to acknowledge that real-world observations by the OCO-2 satellite involve parameters that are not predetermined in future simulations, such as the Empirical Orthogonal

320 Functions (EOF) parameters, signal-to-noise ratio (SNR), bad sample lists, and the degradation of grating pixels. Therefore, our new model is trained not only on the 60,000 simulated datasets but also on the 2016 historical data. According to the data selection criteria outlined in Section 2.1, we identified a total of 38,626 sets of historical data in 2016, comprising spectral measurements from OCO-2 and the corresponding XCO₂ products. These historical experimental datasets are integrated with the simulated data, enriching the training datasets. This dual combination and data augmentation techniques ensure that the

325 model is well-equipped to handle both potential future atmospheric conditions and the current realities of instrument and




Figure 9. Scatter plots of atmospheric parameters required for forward calculation models (excluding CO₂ profiles) from 2016 to 2020, sourced from the OCO-2 L2 product. Panel (a) is the surface pressure, (b) is the surface temperature, (c) is the near-surface water vapor concentration, (d) is the solar zenith, (e) is the Sun-Earth distance, and (f) is the Earth-satellite relative velocity.

spectral measurement capabilities. By doing so, we provide a more comprehensive training strategy that captures both the

anticipated future scenarios to accurately and efficiently perform XCO₂ retrieval for the “future” years from 2017 to the end of 2020.

5 Results and discussions

330 5.1 Comparison with the OCO-2 satellite product data

To evaluate the retrieval capability of the MLP-XCO₂ model trained on a combined dataset of simulated data and historical 2016 OCO-2 satellite data, the neural network architecture and hyperparameters were intentionally kept identical to the previous model trained solely on actual OCO-2 satellite product data. Keeping these factors constant isolates the training data source as the only major difference between the models. This enables a direct, apples-to-apples comparison of how the training
335 data affects model performance.

Figure 10 (a) shows the retrieval results on 10% out-of-sample testing data that was excluded from model training. Setting aside this test subset is a standard technique for evaluating model performance on new examples. The accurate predictions of the MLP-XCO₂ model on the test data suggest the model has learned generalizable patterns not overfit to the training data. Figure 10 (b) shows the comparison of the retrieval results of the MLP-XCO₂ model on real OCO-2 satellite spectral
340 observations in 2016. Figure 11 displays XCO₂ predictions from 2017 to 2020 using test data consistent with Fig. 4 and Fig. 5. As the simulated training data was generated based on 2016 OCO-2 measurements, testing on 2017-2020 data evaluates the model’s ability to make predictions beyond the time frame of the training data. The scatter plots demonstrate the MLP-XCO₂ model trained on simulated data can accurately and stably predict the annual XCO₂ growth trend, maintaining RMSE around 2 ppm (0.5%). Compared to models trained relying solely on historical satellite product data, the key advantage is the ability
345 to make reasonable forecasts of future atmospheric XCO₂ levels.

Table 3 offers a detailed spatio-temporal comparison of the results presented in Figure 11, enhancing our understanding of the MLP-XCO₂ model’s performance across distinct subregions within East Asia. This table specifically focuses on a finer spatial segmentation within the broad East Asian longitude and latitude range, dividing it into four subregions. These are defined based on the geographical demarcation of 35°N and 130°E, categorized as Northeast (NE), Northwest (NW), Southeast (SE),
350 and Southwest (SW) regions, respectively. The results demonstrate that, regardless of the distribution of sample sizes across these subregions and their varied topographical characteristics (land or ocean), the model maintains a consistent and stable performance in each subregion. Furthermore, the error metrics for these individual subregions align closely with the overall regional errors, indicating a uniformity in the model’s predictive accuracy and reliability across different spatial scales within East Asia.

355 Considering these results, by generating possible realistic future prior information for the atmospheric conditions and using an accurate forward model to simulate the corresponding spectra, the approach avoids inherent biases when extrapolating beyond the distribution of the training data. Rather than simply extending trends, the model is constrained by fundamental physical relationships to interpolate within realistic bounds. This transforms the prediction task into a well-posed interpolation problem versus an unconstrained extrapolation. The simulated data provides a physical regularization that makes the model’s




Figure 10. The comparison of XCO₂ results predicted by the MLP-XCO₂ model from 10% test data (not involved in training). Panel (a) shows the predicted XCO₂ values for the test data that are derived from simulated dataset, and panel (b) shows the test data that are derived from OCO-2 2016 L2 XCO₂ data.

Table 3. Spatio-temporal comparison of XCO₂ predicted by MLP-XCO₂ model results versus results retrieved by OCO-2 across four subregions. These subregions are delineated based on the geographical demarcation of 35°N and 130°E as: Northeast (NE), Northwest (NW), Southeast (SE), and Southwest (SW) regions, respectively.

| Year | Full [Number / ME / RMSE] | NE | NW | SE | SW |
|------|---------------------------|-----------------------|------------------------|----------------------|-----------------------|
| 2016 | 38626 / -0.05 / 1.024 | 2714 / -0.065 / 1.033 | 26111 / -0.049 / 1.017 | 451 / -0.005 / 1.033 | 9350 / -0.052 / 1.033 |
| 2017 | 39850 / 0.548 / 1.743 | 1235 / 0.488 / 1.786 | 30774 / 0.540 / 1.744 | 244 / 0.594 / 1.742 | 7597 / 0.589 / 1.734 |
| 2018 | 35945 / 0.197 / 1.710 | 1854 / 0.218 / 1.731 | 27288 / 0.195 / 1.717 | 745 / 0.167 / 1.719 | 6058 / 0.203 / 1.668 |
| 2019 | 36452 / 0.016 / 2.090 | 1777 / 0.066 / 2.074 | 26082 / 0.013 / 2.095 | 304 / 0.062 / 2.082 | 8289 / 0.013 / 2.080 |
| 2020 | 43277 / 0.081 / 2.010 | 1841 / 0.070 / 2.010 | 34971 / 0.079 / 2.007 | 270 / 0.147 / 1.981 | 6195 / 0.090 / 2.028 |

360 outputs to be scientifically sound. By training on synthetic data spanning potential future scenarios, the model learns robust representations not tightly coupled to specifics of the training data time period. This enables high-fidelity inversion and prediction of XCO₂ even for future time periods beyond available measurements.

5.2 Detecting plume features from the OCO-2 observation

365 In a further effort to deeply analyze the ability of our MLP-XCO₂ model to capture key XCO₂ information from spectral data, we focused on plume detection at sites of potential high emissions, such as thermal power plants, in our target regions from 2017 to 2020. Utilizing the data in the work of Li et al. (2023), we sourced test samples from multiple instances of XCO₂




Figure 11. Comparison of XCO₂ results predicted by the MLP-XCO₂ model versus results retrieved by OCO-2 v10r product from 2017-2020. Panel (a), (b), (c), and (d) display the predictions of the MLP-XCO₂ model from 2017 to 2020, respectively.

enhancements detected by the OCO-2 satellite in Nadir mode observations. These samples were located in close proximity to known large power plants, providing an ideal scenario for assessing retrieval accuracy in detecting localized emission sources.

Figure 12 presents a geographical map that highlights XCO₂ predictions in the test samples from the MLP-XCO₂ model and compares them with results retrieved by the OCO-2 v10r product. This map clearly marks power plants with red triangles, establishing a visual link between industrial emission sources and observed points where elevated XCO₂ levels are detected. Figure 13 further explores this relationship by presenting a longitude-based comparison of XCO₂ results. This figure plots the same data points from Figure 12 against their corresponding longitude coordinates. This arrangement facilitates a direct and intuitive comparison of the trends in XCO₂ enhancements as captured by our model and as reported by the OCO-2 product.

In both figures above, it is visually evident that observation points near power plants show sudden increases in XCO₂ values, aligning with the trend from the OCO-2 v10r product. This trend is particularly pronounced when compared to points

farther away from these emission sources. Considering that these samples are nearly identical in terms of observation angles and times, such consistency is a powerful confirmation to our model's capability to retrieve genuine atmospheric XCO₂ from OCO-2 spectral data.




Figure 12. Geographical map of XCO₂ predictions by the MLP-XCO₂ model compared with OCO-2 v10r product results. The potential plume enhancements and the large power plants (marked by red triangles) were screened in Nadir mode OCO-2 observations as reported in the work of Li et al. (2023).




Figure 13. Longitude-based scatter comparison of XCO₂ predicted by the MLP-XCO₂ model versus results retrieved by OCO-2 v10r product. The potential plume enhancements were screened in Nadir mode OCO-2 observations as reported in the work of Li et al. (2023).

380 5.3 Comparison with the TCCON data

A comparison of the retrieved results from the OCO-2 satellite showed that the RMSE of our developed MLP-XCO₂ model was around 2 ppm. In other words, our results could be worse or better than OCO-2 satellite, requiring further comparison with ground-based measurements. To further validate the accuracy of the MLP-XCO₂ model, we compared the XCO₂ retrievals from the OCO-2 v10r Nadir mode products, the MLP-XCO₂ model outputs, and ground-based measurements from five TCCON sites within the study region (Fig. 1). As summarized in Table 4, spatiotemporal screening was applied to the TCCON and OCO-2 data to obtain comparable observations. The five TCCON sites included were: Tsukuba (Morino et al., 2022b), Saga (Shiomi et al., 2022), Hefei (Liu et al., 2022), Xianghe (Zhou et al., 2022) and Rikubetsu (Morino et al., 2022a). The Anmyeondo site was excluded from this analysis as the XCO₂ data was not updated in the TCCON GGG2020 database, and was only available until early 2018 in the GGG2014 database.

Table 4. Spatio-temporal screening conditions for TCCON sites and OCO-2 satellite Nadir mode observations

| TCCON site | Local time | Observed location | Sample number | Reference |
|------------|-------------------|---------------------------------|---------------|-----------------------|
| Tsukuba | 12 : 48 – 12 : 58 | 36.05°N ± 0.5°, 140.12°E ± 0.5° | 2078 | Morino et al. (2022b) |
| Saga | 13 : 30 – 13 : 40 | 33.24°N ± 0.5°, 130.29°E ± 0.5° | 87 | Shiomi et al. (2022) |
| Hefei | 13 : 20 – 13 : 30 | 31.90°N ± 0.5°, 117.17°E ± 0.5° | 984 | Liu et al. (2022) |
| Xianghe | 13 : 15 – 13 : 25 | 39.80°N ± 0.2°, 116.96°E ± 0.2° | 2770 | Zhou et al. (2022) |
| Rikubetsu | 13 : 20 – 13 : 30 | 43.46°N ± 0.2°, 143.77°E ± 0.2° | 723 | Morino et al. (2022a) |

390 Figure 14-1 presents time series comparisons of XCO₂ retrievals from the different TCCON sites, MLP-XCO₂ model, and OCO-2 Nadir observations. Figure 14-2 displays the box plots of the differences between the MLP-XCO₂ model results, OCO-2 products, and TCCON site data. The plots at each of the five TCCON sites demonstrate the simulated data-trained MLP-XCO₂ model accurately predicts XCO₂ from the OCO-2 spectra. The model successfully captures seasonal variations and the long-term XCO₂ growth trend over the 4-year study period. The reliable performance over time and across multiple TCCON sites further validates the model has learned generalizable representations of carbon cycle processes rather than overfitting to specifics of the simulated training data. By using realistic future simulations for training, the model provides robust and unbiased XCO₂ retrievals across a range of atmospheric conditions.

5.4 Retrieval efficiency

In this study, the ReFRACtor forward model required 12.16 seconds per simulation case (two absorption bands) using an AMD Ryzen-7 5800X computer. The OCO-2 retrieval based on Bayesian optimization typically needs over three iterations to converge, indicating at least 36.48 seconds per retrieval. In contrast, the MLP-XCO₂ model demonstrated remarkable efficiency on the same hardware. It required just 1.14 seconds total to retrieve XCO₂ from 6642 OCO-2 test spectra across all five TCCON sites, averaging 0.17 milliseconds per sample with GPU RTX 3080Ti. This rapid inversion drastically reduces processing times




Figure 14. Comparisons of XCO_2 results from 2017 to 2020 across five TCCON sites. Panel (a-1)-(e-1) show the time series comparisons of XCO_2 retrievals from the different TCCON sites, MLP- XCO_2 model, and OCO-2 L2Lite Nadir observations for the Tsukuba, Saga, Hefei, Xianghe and Rikubetsu site, respectively, with data screening conditions as defined in Table 4. Panel (a-2)-(e-2) present the boxplots depicting the differences (ΔXCO_2) between the MLP- XCO_2 model and OCO-2 products in comparison to the TCCON results for each year. The boxes showing the middle half of the data, from the 25% to the 75% percentiles. The median (50%) is represented by the line within each box. The whiskers encompass the central 90% of the data, extending from the 5% to the 95% percentiles.

compared to traditional methods. While machine learning models need significant upfront time for training data generation and hyperparameter tuning, the prediction is extremely fast once deployed. This enables near real-time processing ideal for operational satellite data streams. Furthermore, the precision and efficiency of neural networks makes them well-suited to meet future demands of high-resolution global greenhouse gas monitoring, enabling millisecond-scale XCO₂ retrievals suitable for large-scale satellite analysis.

6 Conclusions

This proof-of-concept study aims to use the efficient regression inversion capability of the machine learning method to develop machine learning models based on simulated atmospheric radiative transfer data for efficient inversion of satellite observed spectra to retrieve XCO₂. This helps overcome the low efficiency in traditional optimization-based iterative algorithms for XCO₂ retrievals. In the presented study, XCO₂ inversion models using both satellite product based and simulation based data were developed, trained and tested. Long time series inversion and prediction of OCO-2 observations over East Asia were also performed using the developed models. The results were compared with OCO-2 and TCCON retrievals, showing the simulation data based machine learning models can effectively eliminate lagging biases while achieving millisecond-level (<1 ms) inversion efficiency, good accuracy (around 2 ppm), local emission source capture, and long-term prediction stability. It needs to be noted that our current MLP-XCO₂ model does not provide direct uncertainty estimates, estimating prediction intervals is an important next step for future improvement.

Code availability. The ReFRACtor model and its OCO retrieval implementation can be accessed from the Github ReFRACtor repository (<https://github.com/ReFRACtor>, last accessed in August 2023). The codes and models used in this study have been uploaded to GitHub and can be accessed at: https://github.com/TaoRen-Rad/XCO2_retrieval. For access to the dataset, please send requests to Tao Ren (tao.ren@sjtu.edu.cn).

Data availability. The OCO-2 products (including OCO-2 L1B, Met, L2std and L2Lite files) are available from Goddard Earth Sciences Data and Information Services Center (<https://disc.gsfc.nasa.gov/datasets/>, last access: March 2023). The TCCON site products are available from TCCON DATA ACHIEVE (<https://tcocondata.org/>, last access: March 2023).

Financial support. This research has been supported by the National Natural Science Foundation of China (Grants Nos. 52276077 and 52120105009)

Author contributions. FX and TR designed the study. FX made updates and modifications for the Refractor forward model and developed

430 the machine learning code, carried out the tests and result analysis under the supervision of TR. FX and TR prepared the manuscript. All authors reviewed the manuscript.

Competing interests. The corresponding author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

435 *Acknowledgements.* We gratefully acknowledge the TCCON Data Archive hosted by CaltechDATA at <https://tccodata.org> for providing the TCCON data for our study. Our thanks also go to the OCO-2 Science Team for the OCO-2 project. We appreciate our colleagues for their feedback and our affiliated institutions for their unwavering support.

References

- Bacour, C., Bréon, F.-M., and Chevallier, F.: On the challenge posed by the estimation of XCO₂ from OCO-2 observations in near-real time based on artificial neural network, IWGGMS-19, 2023.
- Bréon, F.-M., David, L., Chatelana, P., and Chevallier, F.: On the potential of a neural-network-based approach for estimating XCO₂ from OCO-2 measurements, *Atmospheric Measurement Techniques*, 15, 5219–5234, <https://doi.org/10.5194/amt-15-5219-2022>, 2022.
- Cansot, E., Pistre, L., Castelnau, M., Landiech, P., Georges, L., Gaeremynck, Y., and Bernard, P.: MicroCarb instrument, overview and first results, in: *International Conference on Space Optics — ICSO 2022*, edited by Minoglou, K., Karafolas, N., and Cugny, B., vol. 12777, p. 1277734, *International Society for Optics and Photonics, SPIE*, <https://doi.org/10.1117/12.2690330>, 2023.
- Carvalho, A. R., Ramos, F. M., and Carvalho, J. C.: Retrieval of carbon dioxide vertical concentration profiles from satellite data using artificial neural networks, *Trends in Computational and Applied Mathematics*, 11, 205–216, <https://doi.org/10.5540/tema.2010.011.03.0205>, 2010.
- Chen, T. and Guestrin, C.: Xgboost: A scalable tree boosting system, in: *Proceedings of the 22nd acm sigkdd international conference on knowledge discovery and data mining*, pp. 785–794, 2016.
- Cogan, A., Boesch, H., Parker, R., Feng, L., Palmer, P., Blavier, J.-F., Deutscher, N. M., Macatangay, R., Notholt, J., Roehl, C., et al.: Atmospheric carbon dioxide retrieved from the Greenhouse gases Observing SATellite (GOSAT): comparison with ground-based TCCON observations and GEOS-Chem model calculations, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2012JD018087>, 2012.
- Crisp, D., Fisher, B., O'Dell, C., Frankenberg, C., Basilio, R., Bösch, H., Brown, L., Castano, R., Connor, B., Deutscher, N., et al.: The ACOS CO₂ retrieval algorithm—part II: global XCO₂ data characterization, *Atmospheric Measurement Techniques*, 5, 687–707, <https://doi.org/10.5194/amt-5-687-2012>, 2012.
- Crisp, D., Pollock, H. R., Rosenberg, R., Chapsky, L., Lee, R. A., Oyafuso, F. A., Frankenberg, C., O'Dell, C. W., Bruegge, C. J., Doran, G. B., et al.: The on-orbit performance of the Orbiting Carbon Observatory-2 (OCO-2) instrument and its radiometrically calibrated products, *Atmospheric Measurement Techniques*, 10, 59–81, <https://doi.org/10.5194/amt-10-59-2017>, 2017.
- Crisp, D., O'Dell, C., Eldering, A., Fisher, B., et al.: Orbiting carbon observatory (OCO) - 2 level 2 full physics algorithm theoretical basis document Version 3.0 – Rev 1, https://docserver.gesdisc.eosdis.nasa.gov/public/project/OCO/OCO_L2_ATBD.pdf, 2021.
- David, L., Bréon, F.-M., and Chevallier, F.: XCO₂ estimates from the OCO-2 measurements using a neural network approach, *Atmospheric Measurement Techniques*, 14, 117–132, <https://doi.org/10.5194/amt-14-117-2021>, 2021.
- Eldering, A., Taylor, T. E., O'Dell, C. W., and Pavlick, R.: The OCO-3 mission: measurement objectives and expected performance based on 1 year of simulated data, *Atmospheric Measurement Techniques*, 12, 2341–2370, <https://doi.org/10.5194/amt-12-2341-2019>, 2019.
- Gribanov, K., Imasu, R., and Zakharov, V.: Neural networks for CO₂ profile retrieval from the data of GOSAT/TANSO-FTS, *Atmospheric and Oceanic Optics*, 23, 42–47, <https://doi.org/10.1134/S1024856010010094>, 2010.
- Hamazaki, T., Kaneko, Y., Kuze, A., and Kondo, K.: Fourier transform spectrometer for greenhouse gases observing satellite (GOSAT), in: *Enabling sensor and platform technologies for spaceborne remote sensing*, vol. 5659, pp. 73–80, *SPIE*, <https://doi.org/10.1117/12.581198>, 2005.
- Imasu, R., Matsunaga, T., Nakajima, M., Yoshida, Y., Shiomi, K., Morino, I., Saitoh, N., Niwa, Y., Someya, Y., Oishi, Y., et al.: Greenhouse gases Observing SATellite 2 (GOSAT-2): mission overview, *Progress in Earth and Planetary Science*, 10, 33, <https://doi.org/10.1186/s40645-023-00562-2>, 2023.

- 475 Jin, Z., Tian, X., Han, R., Fu, Y., Li, X., Mao, H., Chen, C., and GAO, J.: Tan-Tracker global daily NEE and ocean carbon fluxes for 2015-2019 (TT2021 dataset), <https://doi.org/10.11888/Meteoro.tpd.271317>, 2021.
- Keely, W. R., Mauceri, S., Crowell, S., and O'Dell, C. W.: A nonlinear data-driven approach to bias correction of XCO₂ for NASA's OCO-2 ACOS version 10, *Atmospheric Measurement Techniques*, 16, 5725–5748, 2023.
- Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the Greenhouse Gases Observing Satellite for greenhouse gases monitoring, *Applied optics*, 48, 6716–6733, <https://doi.org/10.1364/AO.48.006716>, 2009.
- 480 Li, Y., Jiang, F., Jia, M., Feng, S., Lai, Y., Ding, J., He, W., Wang, H., Wu, M., Wang, J., et al.: Improved estimation of CO₂ emissions from thermal power plants based on OCO-2 XCO₂ retrieval using inline plume simulation, *Science of The Total Environment*, p. 169586, 2023.
- Liang, A., Gong, W., Han, G., and Xiang, C.: Comparison of satellite-observed XCO₂ from GOSAT, OCO-2, and ground-based TCCON, *Remote Sensing*, 9, 1033, <https://doi.org/10.3390/rs9101033>, 2017.
- 485 Liu, C., Wang, W., Sun, Y., and Shan, C.: TCCON data from Hefei, China, Release GGG2020R0. TCCON data archive, hosted by Caltech-DATA, California Institute of Technology, Pasadena, CA, U.S.A., <https://doi.org/10.14291/tcon.ggg2020.hefei01.R0>, 2022.
- Liu, Y., Wang, J., Yao, L., Chen, X., Cai, Z., Yang, D., Yin, Z., Gu, S., Tian, L., Lu, N., et al.: The TanSat mission: preliminary global observations, *Science Bulletin*, 63, 1200–1207, <https://doi.org/10.1016/j.scib.2018.08.004>, 2018.
- 490 Marchetti, Y., Rosenberg, R., and Crisp, D.: Classification of anomalous pixels in the focal plane arrays of Orbiting Carbon Observatory-2 and-3 via machine learning, *Remote Sensing*, 11, 2901, <https://doi.org/10.3390/rs11242901>, 2019.
- Matsunaga, T. and Tanimoto, H.: Greenhouse gas observation by TANSO-3 onboard GOSAT-GW, in: *Sensors, Systems, and Next-Generation Satellites XXVI*, vol. 12264, pp. 86–90, SPIE, <https://doi.org/10.1117/12.2639221>, 2022.
- McDuffie, J., Bowman, K. W., Hobbs, J., Natraj, V., Val, S., Sarkissian, E., and Thill, M. D.: ReFRACtor: Reusable Software Framework for Retrieval of Satellite Atmospheric Composition, in: *AGU Fall Meeting Abstracts*, vol. 2018, pp. A11F–2282, 2018.
- 495 Meng, G., Wen, Y., Zhang, M., Gu, Y., Xiong, W., Wang, Z., and Niu, S.: The status and development proposal of carbon sources and sinks monitoring satellite system, *Carbon Neutrality*, 1, 32, 2022.
- Messerschmidt, J., Geibel, M., Blumenstock, T., Chen, H., Deutscher, N., Engel, A., Feist, D. G., Gerbig, C., Gisi, M., Hase, F., et al.: Calibration of TCCON column-averaged CO₂: the first aircraft campaign over European TCCON sites, *Atmospheric Chemistry and Physics*, 11, 10 765–10 777, <https://doi.org/10.5194/acp-11-10765-2011>, 2011.
- 500 Modest, M. F. and Mazumder, S.: *Radiative heat transfer*, Academic press, 2021.
- Morino, I., Ohyama, H., Hori, A., and Ikegami, H.: TCCON data from Rikubetsu, Hokkaido, Japan, Release GGG2020R0. TCCON data archive, hosted by CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A., <https://doi.org/10.14291/tcon.ggg2020.rikubetsu01.R0>, 2022a.
- 505 Morino, I., Ohyama, H., Hori, A., and Ikegami, H.: TCCON data from Tsukuba, Ibaraki, Japan, 125HR, Release GGG2020R0. TCCON data archive, hosted by CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A., <https://doi.org/10.14291/tcon.ggg2020.tsukuba02.R0>, 2022b.
- Natraj, V. and Spurr, R. J.: A fast linearized pseudo-spherical two orders of scattering model to account for polarization in vertically inhomogeneous scattering-absorbing media, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 107, 263–293, <https://doi.org/10.1016/j.jqsrt.2007.02.011>, 2007.
- 510

- OCO-2 Science Team, Gunson, M., and Eldering, A.: OCO-2 Level 1B calibrated, geolocated science spectra, Retrospective Processing V10r, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), <https://doi.org/10.5067/6O3GEUK7U2JG>, accessed: [August 2023], 2019a.
- OCO-2 Science Team, Gunson, M., and Eldering, A.: OCO-2 Level 2 meteorological parameters interpolated from global assimilation model
515 for each sounding, Retrospective Processing V10r, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), <https://doi.org/10.5067/OJZZW0LIGSDH>, accessed: [August 2023], 2019b.
- OCO-2 Science Team, Gunson, M., and Eldering, A.: OCO-2 Level 2 bias-corrected XCO₂ and other select fields from the full-physics retrieval aggregated as daily files, Retrospective processing V10r, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), <https://doi.org/10.5067/6SBROTA57TFH>, accessed: [August 2023], 2020a.
- OCO-2 Science Team, Gunson, M., and Eldering, A.: OCO-2 Level 2 geolocated XCO₂ retrievals results, physical model, Retrospective Processing V10r, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC),
520 <https://doi.org/10.5067/E4E140XDMPO2>, accessed: [August 2023], 2020b.
- O'Dell, C., Connor, B., Bösch, H., O'Brien, D., Frankenberg, C., Castano, R., Christi, M., Eldering, D., Fisher, B., Gunson, M., et al.: The ACOS CO₂ retrieval algorithm—Part 1: Description and validation against synthetic observations, *Atmospheric Measurement Techniques*,
525 5, 99–121, <https://doi.org/10.5194/amt-5-99-2012>, 2012.
- Payne, V. H., Drouin, B. J., Oyafuso, F., Kuai, L., Fisher, B. M., Sung, K., Nemchick, D., Crawford, T. J., Smyth, M., Crisp, D., et al.: Absorption coefficient (ABSCO) tables for the Orbiting Carbon Observatories: version 5.1, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 255, 107–217, <https://doi.org/10.1016/j.jqsrt.2020.107217>, 2020.
- Rodgers, C. D.: *Inverse methods for atmospheric sounding: theory and practice*, vol. 2, World scientific, 2000.
- Shiomi, K., Kawakami, S., Ohyama, H., Arai, K., Okumura, H., Ikegami, H., and Usami, M.: TCCON data from Saga, Japan, Release GGG2020R0. TCCON data archive, hosted by CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A.,
530 <https://doi.org/10.14291/tcon.ggg2020.saga01.R0>, 2022.
- Sierk, B., Fernandez, V., Bézy, J.-L., Meijer, Y., Durand, Y., Courrèges-Lacoste, G. B., Pachot, C., Löscher, A., Nett, H., Minoglou, K., et al.: The Copernicus CO₂M mission for monitoring anthropogenic carbon dioxide emissions from space, in: *International Conference on Space Optics—ICSO 2020*, vol. 11852, pp. 1563–1580, SPIE, <https://doi.org/10.1117/12.2599613>, 2021.
- Spurr, R.: LIDORT and VLIDORT: Linearized pseudo-spherical scalar and vector discrete ordinate radiative transfer models for use in remote sensing retrieval problems, *Light scattering reviews 3: Light scattering and reflection*, pp. 229–275, https://doi.org/10.1007/978-3-540-48546-9_7, 2008.
- Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network, *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 369(1943), 2087–2112,
540 <https://doi.org/10.1098/rsta.2010.0240>, 2011.
- Wunch, D., Toon, G. C., Sherlock, V., Deutscher, N. M., Liu, C., Feist, D. G., and Wennberg, P. O.: The Total Carbon Column Observing Network's GGG2014 Data Version, Pasadena, California, <https://doi.org/10.14291/TCCON.GGG2014.DOCUMENTATION.R0/1221662>, 2015.
- Wunch, D., Wennberg, P. O., Osterman, G., Fisher, B., Naylor, B., Roehl, C. M., O'Dell, C., Mandrake, L., Viatte, C., Kiel, M., et al.: Comparisons of the orbiting carbon observatory-2 (OCO-2) XCO₂ measurements with TCCON, *Atmospheric Measurement Techniques*,
545 10, 2209–2238, <https://doi.org/10.5194/amt-10-2209-2017>, 2017.

- Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G., Wunch, D., et al.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data, *Atmospheric Measurement Techniques*, 6, 1533–1547, <https://doi.org/10.5194/amt-6-1533-2013>, 2013.
- 550 Zehr, S.: The sociology of global climate change, *Wiley Interdisciplinary Reviews: Climate Change*, 6, 129–150, <https://doi.org/10.1002/wcc.328>, 2015.
- Zhao, Z., Xie, F., Ren, T., and Zhao, C.: Atmospheric CO₂ retrieval from satellite spectral measurements by a two-step machine learning approach, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 278, 108 006, <https://doi.org/10.1016/j.jqsrt.2021.108006>, 2022.
- 555 Zhou, M., Wang, P., Nan, W., Yang, Y., Kumps, N., Hermans, C., and De Mazière, M.: TCCON data from Xianghe, <https://doi.org/10.14291/tcon.ggg2020.xianghe01.R0>, 2022.