

Supplement of

Geostationary Environment Monitoring Spectrometer (GEMS) polarization characteristics and correction algorithm

5 Haklim Choi ^{1,a}, Xiong Liu ², Ukkyo Jeong ^{3,*}, Heesung Chong ^{4,b}, Jhoon Kim ⁴, Myung Hwan Ahn ⁵,
Dai Ho Ko ⁶, Dong-won Lee ⁷, Kyung-Jung Moon ⁷, and Kwang-Mog Lee ^{1,*}

¹Department of Astronomy and Atmospheric Sciences, Kyungpook National University, Daegu, Republic of Korea

²Center for Astrophysics | Harvard & Smithsonian, Cambridge, MA, USA

10 ³Division of Earth Environmental System Science, Major of Spatial Information Engineering, Pukyong National University,
Busan, Republic of Korea

⁴Department of Atmospheric Sciences, Yonsei University, Seoul, Republic of Korea

⁵Department of Climate and Energy Systems Engineering, Ewha Womans University, Seoul, Korea`

⁶Korea Aerospace Research Institute, Daejeon, Korea

15 ⁷National Institute of Environmental Research, Environmental Satellite Center, Incheon, Republic of Korea

^anow at: Kyungpook Institute of Oceanography, Kyungpook National University

^bnow at: Center for Astrophysics | Harvard & Smithsonian

Correspondence to: Kwang-Mog Lee (kmlee@knu.ac.kr) and Ukkyo Jeong (ukkyo.jeong@pknu.ac.kr)





Figure S1: The satellite-sun geometry for (a) solar zenith angle (SZA), viewing zenith angle (VZA), and relative azimuth angle (RAA) to simulate the synthetic data. The synthetic data was simulated for the January 15, 2016 at 03:00 UTC.



25

Figure S2: Comparison of spatial distributions of Q and U at 432 nm (a, b) simulated by a radiative transfer model (RTM) for synthetic data with those (c, d) estimated based on look-up tables (LUTs) as described in the main manuscript.