



The Pyrenean Platform for Observation of the Atmosphere: Site, long-term dataset and science

Marie Lothon¹, François Gheusi¹, Fabienne Lohou¹, Véronique Pont¹, Serge Soula¹, Solène Derrien¹, Yannick Bezombes¹, Emmanuel Leclerc¹, Gilles Athier¹, Antoine Vial¹, Alban Philibert^{4,1}, Bernard Campistron¹, Frédérique Saïd¹, Corinne Jambert¹, Jeroen Sonke², Julien Amestoy⁵, Erwan Bargain¹, Pierre Bosser⁶, Damien Boulanger³, Guillaume Bret^{1,†}, Renaud Bodichon⁹, Laurent Cabanas¹, Guylaine Canut⁷, Jean-Bernard Estrampes³, Eric Gardrat¹, Zaida Gomez Kuri¹, Jérémy Gueffier¹, Fabienne Guesdon³, Morgan Lopez¹⁰, Olivier Masson¹¹, Pierre-Yves Meslin⁴, Yves Meyerfeld¹, Nicolas Pascal⁸, Eric Pique¹, Michel Ramonet¹⁰, Felix Starck³, and Romain Vidal¹¹

¹Laboratoire d'Aérodynamique, University of Toulouse, CNRS, UPS, Toulouse

²Geosciences Environnement Toulouse, University of Toulouse, CNRS, UPS, Toulouse

³Observatoire Midi-Pyrénées, University of Toulouse, UPS, Toulouse

⁴Institut de Recherche en Astrophysique et Planétologie, Université de Toulouse, CNRS, UPS, France

⁵CEA, DAM, DIF, F-91297, Arpajon-Cédex, France

⁶Lab-STICC UMR 6285 CNRS/M3, ENSTA Bretagne/HOP, 29200 Brest, France

⁷CNRM-Université de Toulouse, Météo-France/CNRS, Toulouse, France

⁸AERIS/ICARE, CNRS/Université de Lille, Villeneuve d'Ascq, France

⁹ESPRI, Institut Pierre-Simon Laplace (IPSL), Paris, France

¹⁰Laboratoire des Sciences du Climat et de l'Environnement (LSCE), IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France

¹¹Institut de Radioprotection et de Sécurité Nucléaire IRSN, Saint-Paul-lez-Durance, France

[†]deceased, 22 June 2016

Correspondence: Marie Lothon (marie.lothon@aero.obs-mip.fr)

Abstract.

The Pyrenean Platform for Observation of the Atmosphere (P2OA) is a coupled plain-mountain instrumented platform in southwest France. It is composed of two physical sites: The "Pic du Midi" mountain top observatory (2877 m a.s.l.) and the "Centre de Recherches Atmosphériques" (600 m a.s.l.). Both sites are complementarily instrumented for the monitoring of climate variables and the study of meteorological processes in a mountainous region. The scientific topics covered by the P2OA include surface-atmosphere interactions in heterogeneous landscape and complex terrain, physics and chemistry of atmospheric trace species at large scale, influence of local and regional-scale emissions and transport on the atmospheric composition, and transient luminous events above thunderstorms.

With a large number of instruments and a high hosting capacity, the P2OA contributes to atmospheric sciences in (i) building long-term series of atmospheric observations, (ii) hosting experimental field campaigns and instrumental tests, (iii) educational training in atmospheric observation techniques.

Here we present the complete instrumentation of the P2OA and the associated datasets, give a meteorological characterization of the platform, and illustrate the potential of the P2OA and its dataset with past or ongoing studies and projects.



1 Introduction

15 The Pyrenean Platform for Observation of the Atmosphere (P2OA) is a ground-based facility devoted to research in atmospheric sciences. It is composed of two physical sites: The "Pic du Midi" mountain top observatory (2877 m a.s.l., hereafter PDM) and, 28 km apart in the plain, the "Centre de Recherches Atmosphériques" (600 m a.s.l, hereafter CRA).

It is one of the five national multi-instrumented sites of the National Institute of Universe Science (INSU) at the National Centre of Scientific Research (CNRS), for the observation of the atmosphere, and one of the foundation stones of the national research infrastructure ACTRIS-Fr (Aerosol, Clouds and Trace gases Research InfraStructure, Pappalardo et al., 2018, - French component). The P2OA belongs to a vast variety of ground-based observational platforms settled around the world, which gather on the same location a large number of complementary instruments for a comprehensive exploration of atmospheric processes. Ground-based observations are complementary to airborne or spaceborne remote sensing atmospheric measurements, since they offer the possibility to operate big, heavy, high-precision, or care-demanding instruments, more easily, or over a longer term, than on most mobile platforms. Sedentary ground-based atmospheric research stations can nowadays
25 be found in almost all geographical environments on Earth:

- polar stations ((e.g. Concordia in Antarctica **Argentini** et al., 2005))
- maritime stations on isolated islands – often on top of high volcanoes, e.g. Mauna Loa in Hawaii (Keeling et al., 1976), Izaña in the Canary Islands (Gomez-Pelaez et al., 2019), Maïdo in the Reunion Island (Baray et al., 2013) – or on coastal
30 places (Mace Head in Ireland (Milroy et al., 2012), Cape Grim in Tasmania (Chambers et al., 2016), Barbados Cloud Observatory (Stevens et al., 2016))
- continental stations, like the largely known and used sites of Atmospheric Radiation Measurements (ARM) Southern Great Plains site in the USA (Mather and Voyles, 2013), or Cabauw (Bosveld et al., 2020) in the Netherlands.

ACTRIS (Pappalardo et al., 2018), a European research infrastructure focusing on climate-relevant short-lived atmospheric variables, aggregates no less than 79 ground-based observational platforms¹. Continental ACTRIS stations can be roughly categorized in flat-terrain vs. mountain stations. Flat-terrain stations are found in either rural e.g. Cabauw (Bosveld et al., 2020) or Hohenpeissenberg (Leuchner et al., 2015), peri-urban (SIRTA, Haeffelin et al., 2005) or urban environments (Qualair, in **Paris Ammoura** et al., 2016), depending on scientific purposes or operational constraints. Only three sedentary mountain ACTRIS stations can be currently found in the Pyrenean area (Collaud Coen et al., 2018): the P2OA is the only one located
40 on the north side of the chain and under oceanic influence; Montsec is a mid-altitude station (1570 m a.s.l.) located about 100 km southsoutheast of the P2OA, on the Pyrenean southern flank (Pandolfi et al., 2014); the Montseny station (700 m a.s.l.) is settled on a Mediterranean coastal mountain in the area of Barcelona (Pandolfi et al., 2011). The latter two stations are mainly devoted to aerosol and trace gases observations. P2OA is one of the rare ACTRIS platforms which address a broader spectrum of atmospheric issues and observational techniques, with a large number of instruments and a significant hosting capacity of
45 its infrastructure.

¹<https://www.actris.eu/facilities/national-facilities>



Despite operational difficulty, mountain sites have long been attractive for scientific experiments in atmospheric sciences (e.g. the proof of atmospheric pressure vertical gradient by Blaise Pascal at Puy de Dôme in 1648 – Baray et al., 2020) or climate monitoring (centennial meteorological data series at e.g. Puy de Dôme and Pic du Midi: Baray et al., 2020; Marengo et al., 1994, respectively). There are several reasons for such attractiveness:

- 50 – mountain meteorology as research topic in itself,
- need of cold or icing conditions,
- far horizontal visibility (e.g. for the optical observation of transient luminous events in the high atmosphere, see Section 3.5)
- thinner atmosphere above (e.g. for the study of cosmic rays interacting with the atmosphere, sun/moon photometer calibration , etc.),
- 55 – lesser influence of the continental boundary layer, access to free tropospheric conditions,
- situation far from human activity.

A better knowledge in mountain meteorology, especially concerning the small-scale transport processes of atmospheric trace species, is an important challenge. Subgrid-scale vertical transport due to complex topography is not accounted for in global or regional-scale chemistry-transport models (Rotach et al., 2014; Bamberger et al., 2017). This is a major issue since complex topography cover more than 50% of Earth’s continental surface, and small-scale vertical transport may affect, e.g., the global carbon balance at global scale (Rotach et al., 2014), but certainly also other species.

Composed of both a mountain-top and a lowland stations close to each other, with a rich instrumentation at CRA to profile the tropospheric dynamics above the site, and atmospheric composition measurement on both sites, the topographic and instrumental configuration at the P2OA is especially suitable to address this question, as illustrated in Hulin et al. (2019).

The scientific topics covered by the P2OA activity are based on the potential of the two sites in term of instrumentation, expertise and geographical embedding:

- atmospheric dynamics, surface-atmosphere interactions, planetary boundary-layer in complex terrain and heterogeneous surface;
- 70 – physics and chemistry of atmospheric trace species at large scale, and their climate impact;
- influence of local and regional-scale emissions and transport processes on the atmospheric composition;
- atmospheric electricity, especially transient luminous events (TLEs);
- bio- and geochemical cycles in the environment.

The P2OA contributes to atmospheric sciences in three major ways:



- 75
- building long-term observation series of climate-relevant variables from a large panel of complementary instruments;
 - hosting experimental field campaigns dedicated to atmospheric process studies or tests of new observation techniques;
 - educational training in the domain of atmospheric observation and instrumental techniques.

The goal of this article is to describe the platform, its instruments, and the associated long-term dataset. It also gives a meteorological characterization of the P2OA, and reviews past or ongoing scientific studies based on the P2OA infrastructure
80 or data, in order to illustrate the platform potentials.

2 A plain/mountain double platform in the Pyrenees

The P2OA is located on the north side of the Pyrenees, at similar distance as the crow flies from the Atlantic Ocean to the west (~150 km) and the Mediterranean Sea to the east (~200 km). Figure 1 shows the topography of the region and the location of the two sites. The Pyrenees main axis is mostly aligned along the west-east direction (more precisely, along the 300° - 120°
85 axis). The highest peak is the Pico Aneto, at 3404 m a.s.l., and about 200 summits above 3000 m, most of them concentrated in the central part of the massif. On the French side, the main valleys are generally N-S orientated, transverse to the chain axis, while in Spain, the geometry is more complex with many E-W aligned sierras and valleys. The terrain lowers much more abruptly on the French side than on the Spanish side.

The CRA site (43.128°N, 0.367°E) is located on the Plateau of Lannemezan, at about 600 m a. s. l., close to the exit of
90 the Aure Valley. The first high ridge to the south, about 15 km away, reaches 1900 m a. s. l. (Bassia Mountain), that is 1300 m above the site. Several small hills and valleys start from the Plateau northward down to the Gers (district) cultivated plain. The Plateau of Lannemezan is covered by grasslands (at ~30%, some of them wetlands or moors) and forests (at ~30%, both deciduous trees and conifers), as well as crops to a lesser degree (mostly wheat and corn). CRA was built in the 1960's by Henri Dessens (1911-1971) for the study of convection, and has served atmospheric research since then.

95 PDM (42.936°N, 0.142°E – about 28 km southwest of CRA) at 2877 m a. s. l., is prominent for its astronomical observatory (Roudier et al., 2021), and its historical meteorological observations, which started in a heroic way in the early 1880's (Dessens and Bücher, 1995). PDM is nowadays easily accessible by cable car – a touristic resort being exploited at the summit. In the scope of the P2OA, we consider only the instrumentation for atmospheric observation, installed on a dedicated platform in the summit buildings. PDM is the only high summit situated 15 km north of the chain of the highest peaks, the latter being
100 concentrated along the French-Spanish border and water divide. This makes PDM a belvedere dominating the French plain, and probably justifies its equipment in the 1880's by Charles de Nansouty (1815-1895) and Célestin-Xavier Vaussenat (1831-1891). From a meteorological point of view, it is under the influence of air masses typically coming from the west, and mostly representative of the free troposphere composition (Section 6.1.2). The top of the peak is mostly made of rocks and pastures below.

105 The population in radii of 10 and 50 km around PDM amount to about 13,000 and 412,000 inhabitants, respectively, concentrated in two main cities: Pau (217,000 inhabitants) and Tarbes (110,000 inhabitants), situated 55 km and 30 km away to the



northwest, respectively. A smaller town, Bagnères-de-Bigorre (7,000 inhabitants), is located 14 km north of PDM. Lannemezan (6,000 inhabitants) is located about 1.5 km east of CRA.

110 CRA and PDM are both equipped with complementary atmospheric instrumentation, described in the next section. Their infrastructures also allow the hosting of field experiments, education training and workshops, with lodging and meeting capacity. About 8 practical educational training sessions are hosted every year at P2OA, often organized as dedicated “micro-field campaign”.

3 Instrumentation and data processing

115 Table 1 draws the list of the instruments in continuous operations at the CRA and PDM. The main variables deduced from the measurements are indicated, as well as the algorithm used to obtain them from the raw data, and the network to which the instrument is connected. The list of instruments can also be found at <https://p2oa.aeris-data.fr/instruments/>, as well as a description of each instrument. In this section, we present more precisely those instruments and the associated algorithm. The corresponding observational networks are addressed in Section 4.




Figure 1. (a) Topography of Southwest France and location of the two P2OA sites, the "Pic du Midi" (PDM), and the "Centre de Recherches Atmospheriques" (CRA). ©IGN. Main towns are indicated in blue. (b) Picture of the PDM observatory in winter. (c) Picture of the CRA site. On this picture, the PDM is visible with the snow on its top. In (a) and (b), the sight direction is indicated on the top left of the picture.



Table 1. List of permanent instruments at P2OA, with from left to right : the site where the instrument is located, type of instrument, main deduced variables, start time of the long term series, algorithm with which the raw data are processed (if appropriate), French network or French research infrastructure to which the instrument participates, international network or infrastructure. In this table, P, T and Hu design Pressure, Temperature and Humidity, respectively.

| Site | Instrument | Main variable(s) | Start time or period | Process algorithm | FR network or Res. Infrastr. | Int. network or Res. Infrastr. |
|----------------------|-----------------------------------|---|-------------------------|----------------------|---------------------------------|-----------------------------------|
| CRA | 60-m instrumented tower | P, T, Hu, Wind, Radiation | 2010- | | ACTRIS-Fr | |
| | | Surface energy balance | 2010- | Eddy Pro | | |
| | 2-m flux station | Surface energy balance | 2014- | Eddy Pro | | |
| | Meteorological Station | P, T, Hu, Wind, Rain, Radiation | 1989- | | Météo-France | |
| | Meteorological Station | P, T, Hu, Wind, Rain, Radiation | 2016- | StatIC | InfoClimat | |
| | Instrumented pit | Soil temperature and moisture | 2019- | | ACTRIS-Fr | |
| | UHF radar wind profiler | Wind, TKE dissipation rate, BL height | 2010- | DESMAN | ACTRIS-Fr | E-PROFILE |
| | VHF radar wind profiler | Wind and tropopause height | 2001- | DESMAN | ACTRIS-Fr | E-PROFILE |
| | Full-Sky Camera - RAPACE | Cloud Cover | 2006- | ELIFAN | ACTRIS-Fr | |
| | Ceilometer CL61 | CBH, Z_i , LPR | 2022- | STRAT-Finder | ACTRIS-Fr | TOPROF |
| | GNSS Antenna | Integrated water vapor | 2011- | | Renag | |
| | DOBSON Spectrometer | O ₃ total column | 2004- | O3-DOBSON | ACTRIS-Fr | NDACC |
| | Reactive Gaz Analyzers | O ₃ , CO, NO _x | 2013- | | ACTRIS-Fr | |
| | Greenhouse-gas analyzer | CH ₄ , CO, CO ₂ | 2019- | | ICOS-Fr | GAW |
| | TLE Camera | TLE high sensitivity images | 2007- | UFOCapture | | |
| | Lightning Detector | VLF radiation | 2008- | Time Of Arrival | | Linet |
| | High-volume sampler | Radionuclide activities | 2014- | | IRSN Opera | Ro5 |
| | Proportional counter | Gamma eq. dose. rate | 2020- | | IRSN Teleray | |
| PDM | Meteorological Station | P, T, Hu, Wind | 1882- | | Météo-France | |
| | Meteorological Station | P, T, Hu, Wind, Radiation | 2004- | | ACTRIS-Fr | |
| | Full-Sky Camera - EKO | Cloud Cover | 2017-2019 | ELIFAN | ACTRIS-Fr | |
| | GNSS Antenna | IWV | 2011- | | Renag | |
| | Aerosol filter sampling | EC, OC, inorganic ions | 2002- | | ACTRIS-Fr | |
| | Aethalometer | BC mass conc. | 2013- | | ACTRIS-Fr | ACTRIS-Eu |
| | Nephelometer | Part. scattering coeff. | 2013- | | ACTRIS-Fr | ACTRIS-Eu |
| | Optical Particle Sizer | Part. size distrib. (0.5-10 μm) | 2010-2021 | | ACTRIS-Fr | |
| | SMPS | Part. size distrib. (10 nm - 0.8 μm) | 2020- | | ACTRIS-Fr | ACTRIS-Eu |
| | Total suspended particles counter | Part. numb. conc. | 2013- | | ACTRIS-Fr | ACTRIS-Eu |
| | Reactive Gaz Analyzers | O ₃ , CO | 2001-, 2004- | | ACTRIS-Fr | GAW, ACTRIS-Eu |
| | Mercury Speciation | GEM, GOM, PBM | 2011-2014 | | | GMOS, iGOS4M |
| | Flask samples | CO ₂ , CH ₄ , CO, N ₂ O, SF ₆ | 2001- | | ICOS-Fr | GAW |
| | Greenhouse-gas analyzer | CH ₄ , CO, CO ₂ | 2014- | | ICOS-Fr | GAW |
| | Radon detector | ²²² Rn volumic activity | 2017- | | ICOS-Fr | |
| High-volume sampler | Radionuclide activities | 2018- | | IRSN Opera | Ro5 | |
| Proportional counter | Gamma eq. dose. rate | 2012- | | IRSN Teleray | | |
| TLE Camera | TLE high sensitivity images | 2009- | UFoCapture | | | |



3.1 Meteorological standard variables, surface energy balance, and atmospheric dynamics

120 Standard meteorological stations from Météo-France have been hosted at the two sites for many years. At PDM, it is actually
an exceptional historical long time series which has been constituted, starting in 1882, maintained alternatively by Météo-
France and Observatoire Midi-Pyrénées (the latter especially helping during World War II), and witness of the ongoing climate
change (Dessens and Bücher, 1995). CRA and PDM stations measure the standard meteorological variables: temperature (2
m), humidity (2 m), pressure, wind (10 m), downward shortwave radiation (2 m) and precipitation at a time interval of 6
125 min. Another meteorological station is hosted from InfoClimat French participative science network², with all basic variables
measured at 2 m³.

The two sites are equipped with a GNSS antenna as part of the RENAG scientific GNSS network and are operational since
2011 (RESIF, 2017). The use of GNSS measurements for atmospheric water vapour measurements consists in estimating the
propagation delay of GNSS signals in the atmosphere from the raw measurements. The integrated water vapour content can
130 then be extracted from this delay (Bossler and Bock, 2021). The technique is well established and widely used in meteorology
and climatology with an uncertainty of less than 1 kg/m² (Guerova et al., 2016). The raw GNSS data acquired by the two
antennas are routinely analysed with a latency of 14 days as part of the ACTRS-FR project. Propagation delays are estimated
at a rate of 5 min and converted into integrated water vapour (IWV) using the methodology detailed in Hadad et al. (2018). The
method uses hydrostatic delays and mean wet column temperature, which are calculated from 6-hourly ECMWF (European
135 Centre for Medium Range Weather Forecasting) analyses and provided by Technische Universität Wien (Boehm et al., 2006).

One of the major instrumented structure of P2OA is the 60 m tower at CRA site, with 5 levels of slow meteorological
measurements (temperature, humidity and wind, at 0.1 Hz at 2 and 15m, and 1 Hz at 30, 45 and 60 m), and 3 levels of fast (10
Hz) measurements (temperature, humidity, wind at 30, 45 and 60 m). The four radiative components are also measured at 60 m
with pyranometers (downward and upward shortwave flux) and pyrgeometers (downward and upward longwave flux). All the
140 sensors installed on the 60 m tower are detailed in Table A1 in the Appendix. The 60 m tower is mainly surrounded by prairies,
but with also small forests and crops in the vicinity, which more or less also contribute to the footprint, according to the wind
and stability. Another flux station is installed at CRA site, at 2 m height within a prairie (fast measurements of temperature,
humidity and wind at 2 m). Contrary and complementarily to the high tower which integrates a large heterogeneous landscape,
this smaller tower measures flux at the scale of a land parcel. Fast measurements with sonic anemometers and hygrometers
145 (Licor open path hygrometers) allow us to calculate the turbulent fluxes and moments, including sensible and latent heat,
momentum flux, surface layer stability and other key turbulence indexes or scales, based on Monin-Obukhov theory. All terms
of the surface energy balance (SEB) between the earth surface and the atmosphere are thus measured with this instrumentation.
The turbulent moments are calculated with the eddy-covariance method on 10-min and 30-min samples, with the EddyPro®
Software (Version 6.2.0) from LI-COR Environmental. The data process options have been discussed within AERIS and
150 ACTRIS-FR, and are homogeneously applied to all ACTRIS-FR eddycovariance stations.

²https://www.lameteorologie.fr/issues/2023/122/meteo_2023_122_37

³<https://www.infoclimat.fr/observations-meteo/temps-reel/campistrous-centre-de-recherches-atmospheriques/000CE.html#highlight=15>



Complementarily, soil temperature and moisture are measured at 6 levels into the ground (5, 10, 20, 30, 60 and 120 cm), as well as ground flux (5 cm).

Two radar wind profilers complementarily measure the wind vertical profiles from 150 m to 16 km above the ground: a Very High Frequency (VHF) radar and a Ultra High Frequency (UHF) radar. The UHF radar is a Degreane-Horizon PCL-1300, working at 1274 MHz, that is 23.5 cm wavelength, with 5 beams (one vertical beam and four oblique beams at 75° elevation). The sources of echoes are mainly the fluctuations of air refractive index, but also the hydrometeors when it rains, and the insects or birds in some conditions. The three components of the wind are deduced every 2 min with a 75 m vertical resolution in a low acquisition mode, or 150 m resolution in a high acquisition mode, based on the Doppler radial velocities of the 5 beams, with the velocity and turbulence volume processing technique (Campistron et al., 1991), based on the original Velocity Azimuth Display (VAD) technique (Browning and Wexler, 1968). This radar detects the top inversion of the convective boundary layer (Heo et al., 2003) and also allows us to estimate the turbulent kinetic energy dissipation rate (Jacoby-Koaly et al., 2002). An improved algorithm for the retrieval of the CBL structure has been developed by Philibert et al. (2024), which can detect thermal internal boundary layers and residual layers, in addition to the current CBL growth. Its maximum vertical coverage varies from 1000 m a. g. l. in dry winter days to 9 km in deep convection, but always includes the planetary boundary layer depth. Minimum height of measurements is close to 150 m a. g. l.

The VHF radar is a partly in-house-developed instrument, based on Degreane Horizon and TOMCO systems. It works at 45 MHz, that is 6.66 m wavelength, with 5 beams as well (one vertical beam and four oblique beams at 75° elevation). The sources of echoes come almost exclusively from the fluctuations of air refractive index, with parasite echoes from airplanes, which are filtered by the process algorithm. The three components of the wind are deduced with the same velocity volume processing technique as for the UHF, with a radial resolution of 375 m and a temporal resolution of 15 min. This radar can measure the wind profile from 1.5 km to 16 km a. g. l. It allows us to estimate the tropopause height, based on the local maximum of reflectivity generated by the specular echo that occurs at this strong inversion (Campistron et al., 1999; Kim et al., 2001)).

The algorithm which produces all the geophysical variables discussed before from the UHF and VHF radar is named DES-MAN, and is related to a GIT-lab CeCIL license at AERIS data center. It is used as a homogeneous processing for several radar wind profilers settled in ACTRIS-FR sites.

All the instruments described previously are permanent instruments, operating continuously. In addition to this, P2OA is also equipped with mobile meteorological instrumentation, which can be used for educational training and for specific field experiments hosted at the site, or elsewhere: a MODEM radiosounding station, a tethered balloon with 5 meteorological probes, a flux station.

3.2 Clouds

Both sites are equipped with a total sky imager, which enables us to visualize the whole sky (2π steradians) above each site. They routinely store the full sky images of the local cloud cover as seen from the ground.

At PDM, the camera is an EKO-SRF02 manufactured system operated between 2017 and 2019. At CRA, a home-made system (named RAPACE) is in operation (Lothon et al., 2019), which consists in the association of a digital camera and a large



185 angle lens (fish-eye). The latter is protected by a plexiglas dome and controlled by a thermostat which keeps the temperature
nearby the lens around 10°C, in order to avoid condensation. RAPACE has been continuously operated at CRA since February
2006, with visible daytime and nighttime images every 15 minutes (before February 2017) or 5 minutes (since then). During
the night, a longer 15 s exposure time is used, for astronomy application. The acquisition frequency can be increased anytime
(up to 1 Hz) in the context of field experiments or specific demands, with the possibility of making movies.

190 An algorithm named ELIFAN has been developed (Lothon et al., 2019), in order to estimate the cloud cover from each
image. It is based on red/blue ratio thresholding, both with an absolute or a differential (with a clear sky library) approach.
All the pixels are thus attributed to either cloud, clear sky or ‘uncertain’. Note that roughly 5% of the pixels are attributed
as uncertain in partly cloudy images, which gives an estimation of ELIFAN uncertainty. ELIFAN has been generalized and
adapted to several other sky cameras like the EKO camera of the Pic du Midi, and other total-sky camera of ACTRIS-Fr
195 infrastructure. It is associated with a Git-lab CECIL license.

Cloud base height is estimated from a ceilometer. From 2016 to 2019, a ceilometer Vaisala 25K was present at the CRA
site, giving estimates of cloud base height at three potential levels. Starting in April 2022, a new Vaisala CL61 ceilometer
was installed, with improved capabilities: it can detect cloud base up to 16 km above ground, give information on the vertical
distribution of aerosols derived from the backscatter profiles, and also measure the linear depolarization ratio. The latter enables
200 us to distinguish spherical particles from dissymmetrical particles, like liquid water from ice within clouds or precipitation, and
types of aerosols. This ceilometer also enables to estimate the boundary layer top. Through the participation to E-PROFILE,
the StratFinder (Kotthaus et al., 2020) algorithm will be applied in the future, which supplies different interfaces from the
vertical structure, including the convective boundary layer top.

The knowledge of cloud occurrence at the PDM summit, that is whether it is immersed within cloud or not, is relevant for
205 in situ measurements of soluble gas concentrations or aerosol properties. Since April 2022, a binary index indicating cloudy
conditions at PDM (called the "in-cloud index") has been derived from images taken every 5 minutes with a webcam showing
details of the summit platform (from a few meters to 150 m away from the camera), as well as the background mountain
landscape. (A similar approach had been developed by Baray et al., 2019, for the Puy de Dôme observation platform.) The
algorithm is based on edge detection using a Canny filter on several areas in the image, selected at various distances. Areas
210 with sharp contours are considered free of cloud, while blurred contours are considered as the signature of fog between the
scene and the camera. These various pieces of information are then merged using a fuzzy logic algorithm finally returning 0
(false) as value for clear air, or 1 (true) for the presence of cloud.

3.3 Atmospheric trace gases

3.3.1 Reactive gases

215 *O₃ and CO*

In the decades following Schönbein’s discovery of ozone (O₃) in 1839 and the characterization of the molecule (trioxygen)
by Soret in 1865, the question arose of the presence of this gas in the atmosphere. Among the very first historical tropospheric



ozone measurements, some were conducted at the Sencours station – settled at a saddle-point 500 m below the PDM summit – as early as 1874, then at the new summit station from 1881 to 1909, using the Schönbein’s paper method (Marenco et al., 1994). Modern measurement series by UV-absorption analyzer were conducted at PDM in 1982 (Marenco, 1986), in 1990-1993 (Marenco et al., 1994), and then continuously since 2001 to present. Compilation of all those data series revealed an increase of tropospheric ozone by a factor 5 from the 1900’s to the early 1990’s (Marenco et al., 1994) but a stabilization around 45 nmol mol⁻¹ since then (Chevalier et al., 2007). As a chemical precursor of tropospheric ozone, carbon monoxide (CO) was also measured during the 1982 campaign (Marenco, 1986), and continuously since 2004 by trace-level IR-absorption analyzer (Gheusi et al., 2011). Technical details on the O₃ and CO instruments, as well as uncertainty calculations are given in Gheusi et al. (2011). The WMO GAW standard operation procedures (GAW, 2013, 2010, for O₃ and CO, respectively) are followed at PDM.

The stratospheric ozone is also monitored at P2OA, with a Dobson UV spectrometer which measures the total ozone column.

Mercury

In May 2011, automated atmospheric mercury speciation sensors were installed at PDM for gaseous elemental mercury (GEM), gaseous oxidized mercury (GOM) and particulate-bound mercury (PBM). The instrumentation has been operated until the end of 2014 by the Geosciences Environment Toulouse (GET) research institute, and was composed of a Tekran ambient mercury vapor analyzer (model 2537A/B), a mercury speciation unit (model 1130) and a particulate mercury unit (model 1135). PDM joined the Global Mercury Observation System (GMOS⁴) project as an external site in 2012.

3.3.2 Greenhouse gases (GHG)

Continuous measurements of the two main GHGs, carbon dioxide (CO₂) and methane (CH₄), as well as carbon monoxide (ancillary of CH₄ measurements) are conducted in the framework of the national RAMCES/ICOS network since May 2014 at PDM⁵, and April 2019 at CRA⁶. These measurements are made on both sites by means of Picarro G24 analyzers based on cavity ring-down spectroscopy (CRDS; Yver-Kwok et al., 2015). The applied QA/QC protocol (tracability of data and calibration chain; regular human-eye data check; etc.) is defined by the European programme ICOS.

Prior to these continuous GHG measurements, air flask sampling has been made weekly since 2002 at PDM. Flasks were then sent to the RAMCES service at LSCE for laboratory analyses of CO₂ (with speciation of isotopes ¹³C et ¹⁸O), CH₄, CO, N₂O and SF₆.

3.3.3 Radon and radioactivity

Radon is an inert radioactive gas emitted from ice-free soils with a half-life of 3.8 days, making it a useful tracer of the atmospheric boundary layer dynamics, and thus a reliable tracer to discriminate between air masses recently influenced by the continental boundary-layer, and the free troposphere (Chambers et al., 2013). A highly-sensitive radon monitor manufactured

⁴<https://gmos.aeris-data.fr/>

⁵<https://icos-atc.lsce.ipsl.fr/panelboard/PDM>

⁶<https://icos-atc.lsce.ipsl.fr/panelboard/CRA>



by the Australian Nuclear Science and Technology Organisation (ANSTO, model 1500L; Whittlestone and Zahorowski, 1998) is in operation at PDM since October 2017.

250 A continuous monitoring of radon concentrations has also been performed at three different heights of the CRA 60 m tower (at 1 m, 30 m, 60 m), with AlphaGuards ionization chambers, since 2018 (Amestoy, 2021). Atmospheric radon is also monitored by gamma-ray spectroscopy, using a RSX-5 NaI spectrometer mounted on the same tower, and soil concentrations of radon are measured by means of three Barasol probes (Amestoy et al., 2021).

P2OA hosts 2 of the 50 national aerosol samplers for the monitoring of atmospheric radioactivity, as part of the Permanent
255 Observatory of Atmospheric Radioactivity (IRSN OPERA network). A very high flow rate sampler ($900 \text{ m}^3 \text{ h}^{-1}$, before temperature and pressure corrections) equips the PDM, making it the most sensitive altitude station in terms of trace-level radionuclide measurements. At both P2OA sites, samples are taken on a weekly basis. Both samplers make it possible to detect naturally occurring radionuclides such as cosmogenic ones (e.g., ^7Be , ^{22}Na) as well as any long-lived artificial radionuclides (e.g., ^{137}Cs) resulting from past nuclear tests or shorter-lived ones which could indicate a nuclear accident release. Results can
260 be downloaded on the <https://www.mesure-radioactive.fr/#/> website. These devices closely complement the high-frequency measurements of automatic alert probes (TELERAY network) which would react instantly in the event of high contamination.

3.4 Physical, chemical and optical properties of aerosols

At PDM, the chemical composition of aerosols has been measured since 2002. Weekly filter samples are taken from a pumped
265 volume of air of approximately 400 cubic meters. They are then analyzed to provide the concentrations of elemental carbon (EC) and organic carbon (OC) through a thermo-optical analytical technique (EUSAAR protocol; Cavalli et al., 2010) on the one hand, and the concentrations of major inorganic ions (calcium, magnesium, sulfate, nitrate, potassium, chlorine, ammonium) through an analysis by ion chromatography on the other hand.

Aerosol optical properties have been measured since 2013. Scattering measurements were implemented for one wavelength
270 from 2013 to 2018 (Nephelometer ECOTECH M9003) and then for three wavelengths from 2018 to present (Nephelometer ECOTECH AURORA 3000). Aerosol absorption properties were measured first (2013 -2017) for one wavelength and then for seven wavelengths (2017 - present) with a Magee AE16 and then an AE 33 Aethalometer, respectively.

The total aerosol number concentration has been measured since 2008 by a condensation particle counter (TSI CPC 3010) until 2020 and then by a CPC 3750 thereafter.

275 Finally, the aerosol size distribution measurement was initially performed only in the coarse mode ($0.5 - 10 \mu\text{m}$) from 2010 to 2021 thanks to the implementation of an Optical Particle Sizer (TSI OPS 3330), then from 2020 the aerosol size distribution from 10nm to $0.8 \mu\text{m}$ has been performed by the implementation of a Size Mobility Particle Spectrometer (SMPS model 4S from Paolo Vilani société 4S).

3.5 Transient Luminous Events optical observations

280 Two low-light Watec 902H camera (minimum illumination of 10^{-4} lux) are installed at the two P2OA sites. They have a field of view (FOV) of 31° and are oriented to the storm with a pan-tilt unit that can be remotely controlled via the Internet. Thus,



the observations of Transient Luminous Events (TLEs) can be made in night conditions, above thunderstorms in a range of about 800 km around each site, at the altitudes of these phenomena (between 30 km and 90 km), and on alert according to the meteorological conditions. The cameras operate in a triggered mode provided by the UFOCaptureV2 software to capture luminous events with brightness above a given threshold. They provide videos with a time resolution of 25 frames (or 50
285 interlaced fields) per second, which corresponds to a time resolution of 20 milliseconds. The azimuth and elevation of the sprite events are determined with the software “Cartes du Ciel” (SkyCharts). The methodology for the analysis of the video imagery, the time and space synchronization with other data about the storm activity, especially the lightning flashes associated with the TLEs, the event terminology, are explained in Soula et al. (2017).

4 Data dissemination

290 4.1 Infrastructures and Networks

Most of the permanent instruments at P2OA participate to national or international atmospheric monitoring networks. Those are indicated in Table 1 for each instrument. All networks pursue common objectives: sharing expertise on improving data quality and standardizing the observing systems, procedures, data bases, data processing, and data dissemination. They also favour the research and development, educational training, technical assistance and production of essential output information
295 to end-users.

4.1.1 ACTRIS

P2OA is intrinsically linked with the national ACTRIS-France infrastructure – (hereafter ACTRIS-Fr), the French component of the European ACTRIS infrastructure (Pappalardo et al., 2018). ACTRIS is a distributed infrastructure in support of research on climate and air quality, for a better understanding of the evolution of processes and atmospheric composition. It supplies
300 information on variability of climate-relevant reactive species from multiple observational and exploration platforms. ACTRIS-Fr has a larger scope than ACTRIS-ERIC (European Research Infrastructure Consortium, also ACTRIS-Eu in Table 1), by also including fundamental variables of climate and meteorology: atmospheric dynamics and surface/atmosphere heat fluxes. The plain involvement of P2OA within ACTRIS-Fr explains why so many instruments participate to this infrastructure (see Table 1). Only aerosols and trace gas continuous monitoring made at PDM participate so far to the European scale of the infrastructure,
305 with involvement in two of the six ACTRIS topical centers: the European Center for Aerosol Calibration and Characterization (ECAC), and the Center for Reactive Trace Gases In Situ Measurements (CiGas).

4.1.2 ICOS-ATC

Greenhouse gas measured at both PDM and CRA participate to the Integrated Carbon Observation System - Atmospheric Thematic Center (ICOS-ATC) at the European scale (Heiskanen et al., 2022). ICOS produces standardised long-term obser-
310 vations to understand the carbon cycle and to monitor the greenhouse gases, for a better understanding of climate change and



its impacts. Atmospheric measurements of CO₂ and CH₄ at both PDM and CRA contribute to the European Obspack compilation updated once a year. The data are available on the ICOS Carbon Portal (<https://doi.org/10.18160/PEKQ-M4T1> and <https://doi.org/10.18160/9CQ4-W69K> respectively for CO₂ and CH₄).

4.1.3 Météo-France

315 The two meteorological stations of P2OA managed by Météo-France – **the national** meteorological service – are part of its synoptic network, providing operational observation data for assimilation by the Numerical Weather Prediction (NWP) models ARPEGE and AROME, and for climate monitoring.

4.1.4 E-PROFILE

320 The two wind profilers and the ceilometer participate to E-PROFILE, a program of the European network of National Meteorological Services, EUMETNET. E-PROFILE is part of the EUMETNET Composite Observing System (EUCOS), managing the European networks of radar wind profilers, lidars and ceilometers for the monitoring of vertical profiles of wind and aerosols. Near real time data of the two P2OA wind profilers are sent every 30 min to the network, with constraints of timeliness for the Global Forecast System (GFS) to assimilate the data on time. Monthly statistics monitoring of the model-observation departure is supplied by the network and by Météo-France.

325 4.1.5 GAW

The Global Atmosphere Watch (GAW) programme⁷ of the World Meteorological Organization (WMO) aims at improving the global understanding of atmospheric composition, and coordinates the collection of high-quality atmospheric composition observations from stations all over the world. Owing to the remote character of the site and the panel of long-term observations conducted there, PDM has been accepted as a GAW Regional Station since 2018.

330 Beforehand, PDM had already contributed to GAW databases for many years:

- to the World Data Center for Greenhouse Gases (WDCGG⁸, hosted by the Japan Meteorological Agency) since 2001,
- to the World Ozone and Ultraviolet radiation Data Center (WOUDC⁹, hosted by the Canadian government) since 2004 (see the NDACC section below),
- to the World Data Center for Reactive Gases (WDCRG, hosted at the NILU on the EBAS system¹⁰) since 2007 (ozone),
- 335 – to the World Data Center for Aerosols (WDCA, also on EBAS) since 2013.

⁷<https://community.wmo.int/en/activity-areas/gaw>

⁸<https://gaw.kishou.go.jp/>

⁹<https://woudc.org/home.php>

¹⁰<https://ebas-data.nilu.no/>



4.1.6 NDACC

The Network for the Detection of Atmospheric Composition Change¹¹ (NDACC) is an international collaboration and a world-wide network of ground-based stations for remote-sensing observation of (mainly) water-vapor and ozone in the whole atmosphere (from the troposphere to the mesosphere). At the national scale, observations are coordinated by the NDACC-France
340 service. P2OA is contributing to the NDACC database (and in the same time to the GAW WOUDC database) with data from a Dobson UV spectrometer (historical instrument #49) operated at CRA since 2004.

4.2 Data and physical access policies

4.2.1 Data policy

The data are available from the P2OA web portal (<https://p2oa.aeris-data.fr/data/>), but also accessible from the ACTRIS-Fr web
345 portal (<https://www.actris.fr/actris-fr-data-centre/>) and AERIS data catalogue (<https://www.aeris-data.fr/en/catalogue-en/>) for the concerned variables, with no difference on the pointed dataset according to the portal used. Long-term observation data collected on any P2OA site, are ruled by the P2OA data policy which is available here: <https://p2oa.aeris-data.fr/p2oa-data-policy/>. The data policy follows ACTRIS-Fr data policy for any variable laying in the contour or ACTRIS-Fr.

The main spirit of the P2OA and ACTRIS-Fr data policies is to offer free and unlimited access to P2OA data.

350 The users agree to contact the concerned local principal investigators (PIs) to offer an appropriate level of acknowledgement or collaboration. In all cases, publications using P2OA long-term observation data should include the acknowledgement formula proposed in the data policy (or an appropriate adaptation of it).

4.2.2 Site physical access – Hosting policy

Observation data collected on P2OA sites in the context of temporary campaigns are not concerned by the data policy, but by
355 the P2OA hosting policy, available at this address: <https://p2oa.aeris-data.fr/physical-access-form/>.

Beyond the respect of this policy, the temporary users of the P2OA experimental sites are beforehand invited to fill a physical access form where the technical needs of their experiment should be described in sufficient detail. Required at least one month in advance, this form allows the P2OA staff to assess the feasibility of the project and prepare best technical conditions for the experiment, in close interaction with the hosted research team.

360 Again, the users are requested to agree with the local PIs on the appropriate level of collaboration. As minimum requirement, the acknowledgement formula mentioning the P2OA should appear in any publication.

4.3 The ReOBS project

For a facilitated access to the multi-instrumental data collected at observatories, the ReOBS project¹² (Chiriaco et al., 2018) aims at gathering and synchronizing multiple data sets from a given observatory in one single netcdf file at a 1 h time res-

¹¹<https://ndacc.larc.nasa.gov/>

¹²<https://reobs.aeris-data.fr/en/p2oa-long-term-analysis/>



365 olution. Chiriaco et al. (2018) more specifically deal with the data of the SIRTA observatory (Instrumented Site for Remote
Sensing Atmospheric Reserch), from which the project has started. Since then, and in the context of ACTRIS-Fr and its aim
at harmonizing the data and their access from the national AERIS data center, the ReOBS project has been extended to other
observatories, P2OA among them. This is an ongoing work at P2OA, to make a single file of a large set of relevant data at 1-h
time resolution, for an easier use by external users like modelists, air quality services, etc. ReOBS also merges native resolution
370 data, for principal investigators or more specific use. It includes an additional data quality control and statistics.

5 Meteorological characterization of the P2OA area

Here we present a meteorological characterization of the P2OA based on the meteorological measurements performed over
a 8-year period at P2OA, from 2015 to 2022. A flow regimes study has been previously carried out by Gueffier (2023), over
the period 2015-2019, based on hierarchical ascendant classification, which gives support to our discussions below. Starting
375 in 2015 enables us to work with the most homogeneous dataset (there are UHF data from 2010 to 2015 which still need to
be processed in an homogeneous way). We first address the flow regimes, based on the radar wind profiler data and the 60 m
tower data. Then we illustrate the seasonal variability of thermodynamic variables, radiation, cloud cover, precipitation and
convective boundary layer depth. We end this section with the seasonal and diurnal variability of the atmospheric composition
observed at PDM.

380 5.1 Flow regimes

The meteorological conditions observed at P2OA are governed by the presence of the Pyrenees' chain. Dynamically, and at the
regional scale:

- the typical westerly or north-westerly synoptic winds are channeled along the chain;
- northerlies, often associated with anticyclonic conditions, can be blocked by the chain on the French side;
- 385 – southerly or south-westerly winds, often associated with a low located over the Atlantic, lead to mountain waves, flow
splitting and foehn situations;
- easterlies, although rare, are also channeled along the chain. They are sometimes linked with a low situated over the
Mediterranean sea, and with enhanced precipitations over the eastern part of the Pyrenees.

Figure 2 presents the wind roses found over the period 2015-2022, at three different altitude levels in the lower troposphere,
390 based on three different instruments operated at CRA (the VHF and UHF wind profilers, and the 60-m tower). The highest level
shown, at 4000 m a. s. l. shows the predominance of synoptic winds in a large southwest to northwest quadrant. At 750 m a. g. l.,
the wind rose looks totally different, and shows the channelling of the flow along the chain. Close to the surface (15 m above
the ground), a superposition of different flow regimes is visible: in addition to the dominant westerlies, thermally-induced
winds are frequent, characterized by southerly drainage (katabatic) flow from the mountain during the night, and northerly



395 upmountain (anabatic) breeze during the day (Hulin et al., 2019). Note that the surface wind at CRA is generally weak, except
in case of rare strong westerly fronts or southerly downslope wind storms. The area, centrally close to the foothills of the
Pyrenees, and protected by them, is one of the areas in France with the weakest wind.

The plain-mountain thermal wind is connected to the valley winds of the Aure Valley mentioned before. It usually sets up
in calm synoptic conditions, but still can establish under significative synoptic forcing, especially when the northern Pyrenean
400 foreland is sheltered by the mountains from southerly to southwesterly wind blowing in altitude. Hulin et al. (2019), based on
the 2006-2015 dataset, found that almost 30% of days with surface breeze at CRA occur in conditions with a southerly wind
component of more than 5 m s^{-1} aloft. They also found an overall occurrence of 27.5% for days with surface breeze at CRA.
Román-Cascón et al. (2019) have shown, for the year of 2017, that the katabatic winds at CRA lay in the sector $110\text{-}200^\circ$ and
have a large occurrence of more than 30% of time. Anabatic winds are less frequent, about 15%, and lay in the sector $300\text{-}50^\circ$.

405 Strong synoptic forcing (windspeed at 4000 m larger than 30 m s^{-1}), mostly corresponds to flows in the west- to northwest-
erly sector in altitude. Below the altitude of the Pyrenean crest and down to the surface, they are usually channeled toward
westerlies.

The foehn phenomenon, typical of the lee of mountains (Elvidge and Renfrew, 2016), is very well observed from the CRA
site. It is generated by synoptic southwesterly flows, generally associated with the approach of a front and with a low situated
410 off the Bay of Biscay over the Atlantic. When the wind upstream of the mountain is strong enough, the foehn wind on the lee
side can penetrate down to the surface (Scorer, 1955; Smith, 1985; Ólafsson and Bougeault, 1997). Oscillation of the whole
troposphere is typically observed at CRA in southwesterly synoptic forcing (Gueffier et al., 2024), generated by mountain lee
waves (Bougeault et al., 1990; Ólafsson and Bougeault, 1997)), with an occurrence of about 10% (Gueffier et al., 2024). Note
that the large occurrence of southerly winds in the windrose of Fig. 2c includes the katabatic downslope winds from the Aure
415 Valley, which is predominant at that height.




Figure 2. Wind roses at (a) 3975 m a. s. l, (b) 750 m a. g. l and (c) 15 m a. g. l at CRA over the period 2015-2022.



5.2 Seasonal variability

Figure 3 presents the seasonal and inter-annual variability of the monthly mean diurnal cycles of air temperature (at 2 m a. g. l.), water vapour mixing ratio (2 m) and downward shortwave radiation (60 m) at CRA, all based on the 60-m tower measurements. The monthly composite days are shown separately for each year of the 2015-2022 period, which also highlights the inter-annual
420 variability for a given month.

The downward shortwave radiation seasonal variability is typical of the mid-latitude. June-July-August are the hottest and most humid months, associated with the highest shortwave downward radiation. The diurnal cycle of temperature is marked in any month, even if of lesser amplitude in January and February. It is noticeable to see how it can remain high in September at the end of summer, and be significant even in December on some specific warmer years. Those months often show anticyclonic
425 dry and calm conditions, with clear sky and marked diurnal cycle. The diurnal cycle of moisture is less marked, partly because the composite is disturbed by non-diurnal variability like fronts or mesoscale advections.

The inter-annual variability is usually large, often of the same order of magnitude as the diurnal variability (December and February are good examples of large inter-annual variability here), except in summer here (it is very small in August for this period, but might be larger for another period).

Figure 4 shows the seasonal variability of the cloud cover (monthly averages), based on the RAPACE total sky imager and the ELIFAN algorithm (Lothon et al., 2019). The monthly standard deviation is expectedly large (not shown), about 40%, larger than the seasonal variability (less than 10%). Still, April and May have on average larger cloud cover than September and October. Considering monthly averages calculated in the morning (between 3 h and 4 h after sunrise) or in the afternoon (between 4 h and 3 h before sunset) reveals a slight diurnal cycle, explained by a significant number of days with afternoon
435 convection. Interestingly, this diurnal cycle vanishes in summer. This could be explained by the occurrence in summer of nighttime stratus clouds which dissipate in the late morning or afternoon, compensating days with clear mornings but afternoon convection. Those stratus clouds can notably come from blocked northerly moist flows, or from moisture coming from the daytime moist convection over the mountain which is concentrated at the bottom of the valleys and the plain during the night. Summer has a large occurrence of moist low layers in the northern Pyrenees, associated with a rather vegetated and moist
440 surface, relatively to the southern Spanish side. The diurnal cycle also disappears in winter, with the decreased shortwave radiation, drier air and suppressed convection.

In Figure 5, the precipitations for the period 2015-2022 are shown, through the mean monthly cumulative rain, the monthly mean rain intensities for precipitations larger than 3 mm, and the monthly maximum rainrate encountered. The use of those three variables enables to distinguish between frequent low precipitation events and convective intense rain events. November,
445 January and February have the largest accumulated rain, associated with largest number of rainy days. May, June, July, August are the month with the highest maximum rainrates encountered, associated with convective storms. May and June have the largest mean rainrates, when considering only significant rain events with precipitation rates larger than 3 mm. These months combine the chance of storms and the occurrence of moist blocked northerly flows. The driest months are September and October.




Figure 3. Seasonal and inter-annual variability of monthly-averaged diurnal cycles of (a) 2-m temperature, (b) 2-m water vapour mixing ratio, (c) downward short wave radiation.

450 Figure 6 shows the convective boundary layer (CBL) depth estimated from the UHF wind profiler with the algorithm developed by Philibert et al. (2024), based on the detection of a local maximum of the air refractive index structure coefficient and a minimum of vertical velocity standard deviation. The algorithm also takes temporal continuity into account. Note that only convective boundary layers are considered here, which implies that only cases with no rain and no fog allow for a CBL depth estimation. As expected, the cold winter months of November, December and January have significantly smaller CBL
455 depth relatively to other months. Spring and summer interestingly reach similar CBL depth, despite the variation of incident shortwave radiation from one season to the other. This can be explained by several features. In summer in the mountain area,




Figure 4. Seasonal and inter-annual variability of monthly-averaged cloud cover, for the period 2012-2019. Each dot is a monthly average. The black line is the averaged for morning hours (3h after sunrise), and the grey line for afternoon hours (3h before sunset).

the heating of the valley atmospheres is stronger than above flat terrain, mainly due to reduced volume (concept of topographic amplification, see Steinacker, 1984; Whiteman, 1990), and generates enhanced convection due to increased instability. Cumulus and congestus clouds develop actively all along the mountain ridge during fair weather days, inducing a mesoscale subsidence near the mountain base (De Wekker, 2008). Therefore, the CBL at CRA is often capped by a significant subsidence (Petersen et al., 2015; Blay-Carreras et al., 2014b) in this season, which limits the CBL growth. In April and May convection is less active over the mountain, while the sensible heat flux can be as large as in summer or even larger, because solar heating is significant while air masses are still cold (typically during post-frontal situations). This favors CBL growth. The month of September experiences similar CBL depth (and sensible heat fluxes) than April and May, in clearer sky, less rain, compensated by a smaller temperature gradient close to surface. February and October consistently show intermediate CBL depths, between the two groups winter/summer. Standard deviation of Z_i within one month ranges from 100 to 500 m, with larger values in summer and at midday. It is around 300 m on average.




Figure 5. Monthly accumulated rain amounts (blue, left y-axis), monthly mean rainrates for precipitations > 3 mm (green, right y-axis), and monthly maximum encountered rainrates (brown, right y-axis) for the 2015-2022 period.




Figure 6. Composite diurnal variation of Z_i , averaged over 2015-2022, for each month of the year. Z_i estimates shown here were retrieved by use of CALOTRITON algorithm (Philibert et al., 2024).



5.3 Atmospheric composition

The gas and particles measurements performed at PDM enable to characterize the atmospheric composition at this altitude, including reactive and greenhouse gases, as well as physical and chemical properties of suspended particles. Figure 7 shows the diurnal composites of ozone, carbon dioxide, and methane mole fractions, and total suspended particle number concentrations, separated on a seasonal basis over the period 2015-2019.

Figure 7a shows that the ozone concentration is larger in spring and summer, in link with increased photochemical activity in the troposphere during those seasons (e.g. Chevalier et al., 2007). A diurnal cycle is well visible in summer, but of lesser amplitude in spring and fall, and absent in winter. This cycle is mainly due to the influence of anabatic transport from the valleys and the plain to the mountain summits. This had firstly been identified at PDM by Marengo (1986) based on a 1-year measurement campaign in 1982. He raised the idea that air sampled at PDM during the warm season was a mix of boundary-layer and free-tropospheric air, with proportions varying along the day time – free tropospheric conditions being experienced during the night. The anabatic transport thus results in a decrease of ozone concentration during the daytime, as the air from the lower levels is poorer in ozone relative to the free troposphere at the continental scale (as evidenced for western Europe by Chevalier et al., 2007). The anabatic ozone decrease is stronger during the hot seasons where anabatic transport most occurs, and is indeed typical of mountain summit observatories (Tsamalis et al., 2014).

The influence of anabatic transport was studied for PDM specifically by Gheusi et al. (2011); Tsamalis et al. (2014); Hulin et al. (2019). These studies further illustrate that it affects other atmospheric species, as soon as a vertical concentration gradient exists in the regional background atmospheric profile. This is the case for carbon dioxide (CO_2), which undergoes a similar influence (Fig. 7b): during the daytime, the anabatic upflows transport air depleted in CO_2 relative to free tropospheric air at the height of PDM, due to increased photosynthetic activity near the surface in the valleys and the plain. This again results in a marked diurnal cycle of CO_2 during summer, of weak amplitude in spring and fall, and almost no diurnal variability during winter. Such influence is also observed at other mountain observatories in the world (e.g. Necki et al., 2003). At the seasonal scale, the tropospheric background in CO_2 decreases as the photosynthetic activity increases during the vegetation season, so concentrations are maximum in winter and minimum in summer. Note that a multi-annual increase of about 2 ppm per year was estimated for CO_2 at PDM over the period 2015-2019 by Gueffier et al. (2024), consistent with the global anthropogenic trend observed worldwide (Friedlingstein et al., 2023).

In the P2OA rural and mountainous region, methane (CH_4) mainly comes from agricultural activity. However, oxydation of methane by OH radicals results in a seasonal decrease of CH_4 when oxydation is most important (Necki et al., 2003). This explains the maximum of methane concentration in winter, and the minimum in summer (Fig. 7c). Similarly to other mountain observatories (Necki et al., 2003), CH_4 at PDM displays a marked diurnal cycle in summer with a maximum in the daytime, here again related to transport by anabatic flows, conveying air richer in CH_4 from the low lands. This diurnal cycle is again less pronounced in spring and fall, and absent in winter, consistent with the seasonal variability of the occurrence and intensity of thermally-driven circulations at PDM (Hulin et al., 2019).




Figure 7. Composite diurnal cycle by season of mole fractions of (a) ozone, (b) carbon monoxide, (c) carbon dioxide, and (d) total suspended particle number concentration observed at PDM (2015-2019): DJF = December-January-February; MAM = March-April-May; JJA = June-July-August; SON = September-October-November. The uncertainty ranges correspond to plus or minus the standard error of the sample. Mean values are considered in (a-c) but median values were preferred in (d), due to the skewness of distributions.

As at high Alpine sites, anabatic transport at PDM (Fig. 7d) is one of the most decisive factors contributing to aerosol concentration variability (Collaud Coen et al., 2011; Herrmann et al., 2015). Still for comparison, Sun et al. (2021) showed also



that at Zugspitze-Schneefernerhaus (2671 m) and Jungfrauoch (3580 m) higher concentration and stronger diurnal variability were observed in the warm season, while lower concentration and less distinct diurnal variability in the cold season.

505 6 Illustrative studies based on the P2OA

In this section, we present examples of various applications of the P2OA dataset and experimental sites, concerning:

- atmospheric process studies based on the long-term series or on hosted field experiments (Section 6.1);
- instrumentation test campaigns (Section 6.2);
- evaluation of numerical weather prediction models (Section 6.3).

510 All field experiments hosted at P2OA are listed online (<https://p2oa.aeris-data.fr/field-campaigns/>), with indications of the hosting site (PDM, CRA, both), addressed scientific topic, and involved research laboratories. Here in the following subsections, we focus on a few chosen projects or studies illustrative of the scientific potentials of the P2OA.

6.1 Process studies

6.1.1 Surface/atmosphere interaction

515 The instrumentation of P2OA, and in particular at CRA, is especially appropriate for the study of the boundary layer dynamics and surface/atmosphere interaction. In 2011, an international field experiment was hosted at the P2OA for the study of the transition from daytime convective boundary layer to the stable nocturnal boundary layer: the Boundary Layer Late Afternoon and Sunset Turbulence (BLLAST: Lothon et al., 2014, <https://bllast.aeris-data.fr>). This transitional phase of the diurnal cycle was the unique focus of this project and field campaign. During 3 weeks, research groups from Europe and the USA have
520 gathered a large number of complementary devices in order to densely observe the boundary layer from surface to the top, and from midday to the night. Radiosoundings, tethered balloons, manned aircrafts, unmanned airplanes were operated during 12 favorable days, and continuous measurements were with instrumented towers deployed over various vegetation covers, and with remote sensing devices (lidar, sodar, radiometer, ceilometer, camera), which all were added to the existing permanent instrumentation. From the collected dataset, a fine description of the turbulence decay was done (e. g. Lothon et al., 2014;
525 Darbieu et al., 2015; Nilsson et al., 2016a, b), which revealed how the decay remains normalized and faithful to the Deardoff model quasi-stationarily until the surface flux gets really too small to maintain the mixing. Starting then, the turbulence decay gets faster, the turbulence structure changes, with larger turbulence scales in the boundary layer and smaller in the surface layer. The shape of the energy spectra also changes. It was also shown that a pre-residual layer can be defined before the residual layer overlying the stable nocturnal layer. This pre-residual layer corresponds to a layer between the top inversion and the
530 top of the turbulence layer coupled to the surface. At that time, again, the surface fluxes are not strong enough to maintain the mixing up to the top inversion and previous daytime convective boundary layer top. The scheme in Fig. 8 summarizes those findings. Several other features were finely studied, revealing for example the importance of the advection of small scale



heterogeneities (Cuxart et al., 2016), the mystery of the Lifting Temperature Minimum a few tens of centimeters above surface during the stabilization process (Blay-Carreras et al., 2015), the difficulty of scaling the turbulence decay (Nilsson et al., 2016a; El Guernaoui et al., 2016), the occurrence of counter-gradient heat fluxes (Blay-Carreras et al., 2014a), the uncertainty of Monin-Obukhov Similarity Theory (MOST) (Kooijmans and Hartogensis, 2016), or the interactions between the drainage flows, gravity waves and turbulence (Román-Cascón et al., 2015).




Figure 8. A scheme of the boundary layer diurnal cycle and turbulence structure evolution, based on the results of the BLLAST project. \hat{t} is normalized time. It is based on the period from maximum to null buoyancy flux; i.e. it is zero at maximum buoyancy, and 1 at zero buoyancy flux. EA: Early Afternoon, LA: Late Afternoon and ET: Evening Transition are defined from there. TKE: Turbulent Kinetic Energy, SBL: Stable Boundary Layer. Ellipses represent the structure, size and coupling of vertical velocity eddies.

BLLAST only slightly addressed the role of surface heterogeneity, while the latter raised several questions, not only during the afternoon transition, but more generally. In 2023, P2OA hosted a new field experiment focused on the impact of surface heterogeneity on the atmosphere: the MOSAI project (Model and Observation Surface-Atmosphere Interactions, <https://mosai.aeris-data.fr/>, Lohou et al. (2024)). MOSAI addresses three questions :

- What is the representativeness of the 60 m tower flux measurements or other permanent measurements with respect to the surrounding heterogeneous landscapes (prairies, forests, crops, and small villages)?
- How can we develop appropriate methodologies to evaluate the NWP and Climate models on surface/atmosphere interactions?
- Can we improve the representation of surface-atmosphere coupling in NWP and Climate models?



To address those questions, three one-year field experiments were planned, at three instrumented sites in France: Meteopole (Canut et al., 2019) in 2021, SIRTA (Haeffelin et al., 2005) in 2022, and P2OA in 2023. Each time, the 4 or 5 most representative vegetation covers of a 5 km x 5 km model mesh around the permanent tower are instrumented with surface flux towers. In 2023 at P2OA, three 15-day intensive observation periods (in April, August and December) were added for the study of a transition forest-to-culture, with a tethered balloon, remotely piloted airplane systems (RPAS) and several towers instrumented at different levels above and within the forest. This should enable us to finely study the vertical structure of the local transition, with different sublayers (internal boundary layer, equilibrium boundary layer, etc., Bou-Zeid et al., 2020).

6.1.2 Atmospheric composition

555 – Impact of meteorology on atmospheric composition at PDM

It has been long recognized (e.g. Keeling et al., 1976) that atmospheric composition measurements conducted on top of high mountains are better representative of the troposphere at the global scale than continental low altitude stations, which undergo local influences. Comparing ozone mole fractions measured at PDM and other mountain stations in Europe on the one hand, versus airborne measurements in the free troposphere on the other hand, Chevalier et al. (2007) indeed found agreement within 8% with the reference airborne profiles for stations above 2000 m a.s.l., provided multi-year averages were considered.

Summit observatories may nevertheless be influenced, at least part of the time, by local or regional emissions transported by boundary-layer processes. Two field campaigns – ‘Pic 2005 and ‘Pic 2010’ – were designed to address this question for PDM. Pic 2005 (Gheusi et al., 2011) revealed that during summer fair-weather days, ozone measurements in the daytime at PDM were representative of the mixing of layers present between 1000 and 2000 m a.s.l. above the Pyrenean foreland (CRA). By means of a Lagrangian transport model integrating ozone photochemistry, Tsamalis et al. (2014) showed, for two fair-weather days during Pic 2005, that best adjustments to observed ozone diurnal variations at PDM were obtained when 14 to 57% (depending on the day) of air from the boundary-layer was incorporated into free-tropospheric air before reaching PDM.

During Pic 2010, meteorological and ozone radiosoundings were launched simultaneously from CRA and from a valley bottom very close to PDM (Hulin et al., 2019). These profiles allowed the characterization of a humid venting layer formed during the daytime above the Pyrenees by anabatic pumping of low-level air, then exported above the plain by the altitude return branch of the plain-to-mountain breeze below 2000 m. This is here well illustrated by a numerical simulation of an anabatic day (Fig. 9). In such conditions, PDM appears to be influenced during the afternoon by local weak southerly wind (which is typically observed) conveying the Pyrenean venting layer towards the plain.

To put the above studies in their climatological context, Hulin et al. (2019) also explored the occurrence of thermally-driven circulations at the P2OA over a 10-year period (2006-2015), and their impact on air composition at PDM. Detection methods of thermal circulations in the P2OA area allowed to separate ensembles of days with or without anabatic




Figure 9. High-resolution (200 m) simulation with Meso-NH (Lac et al., 2018) of a typical summer sunny afternoon with a well-developed plain-mountain thermal circulation (10 July 2010, 1500 UTC; PDM being located at the center of a 40x40 km² domain). This south-north vertical cross-section (yz plane) shows model fields averaged over 40 km along the zonal (x) direction. The vectors show the projection of the (x -averaged) wind on the section plane, while the color code emphasizes the y component. The brown thin line shows the terrain profile at the longitude of PDM, while the bold line represents the x -averaged terrain elevation in the model domain. PDM (represented at its real altitude) peaks markedly above the x -averaged terrain altitude, but only 100 m above the model terrain at this place.

580

influence at PDM, revealing contrasted diurnal evolutions of the observed atmospheric species (consistent with the diurnal composites shown in Fig.7).

Gueffier et al. (2024) went beyond the specific influence of thermally-driven circulations, and explored more generally the influence of synoptic meteorology on air composition at PDM. Considering 5-year of meteorological data collected at the P2OA (2015-2019), weather regimes were distinguished by hierarchical clustering. The most characteristic ensembles that emerged were: fair calm weather days (favorable to thermally-driven circulations); disturbed weather with



585 westerly advection; and south foehn conditions. Marked differences were found between the meteorological clusters
when considering the air contents in Rn, O₃, CO, CH₄, CO₂ and particles. Among other results, it was shown that (i)
air driven to PDM by south foehn had mostly a free-tropospheric signature; (ii) despite enhanced anabatic influence,
the regional free tropospheric influence remains dominant during anticyclonic fair weather days; (iii) disturbed weather
caused intense mixing of the lower troposphere at the regional scale, and thus the influence of regional emission sources
590 is stronger.

– Aerosol properties

Guillaume et al. (2008) succeeded in simulating with a global model (ORISAM-TM4, Guillaume et al., 2007) the tem-
poral evolution of black carbon (BC) and total organic carbon (OC) in the aerosol measured at PDM from weekly filter
samples. They thus showed that large-scale BC pollution is most prominent at this site compared to possible local in-
595 fluences, especially during heat waves as the major one during the 2003 summer. In addition, formation of secondary
organic aerosols was found to be a major component of OC in such meteorological conditions.

As an indubitable evidence of hemispheric transport to PDM, radioactive fallout due to explosions of three reactors of
the nuclear power plant in Fukushima (Japan) on 12-15 March 2011, was detected at PDM in an aerosol filter sample
(Evrard et al., 2012). The presence of ¹³¹-iodine in aerosols ($200 \pm 6 \mu\text{Bq m}^{-3}$) indicated that the radioactive cloud
600 reached France between 22 and 29 March, i.e. less than two weeks after the initial emissions.

More recently, Tinorua et al. (2023) performed a study based on two-years measurements (2019-2020) of BC micro-
physical and optical properties at PDM. They showed that among the worldwide existing long-term monitoring sites, the
PDM experiences only occasional influence of the planetary boundary layer, making it an ideal site for characterizing
free tropospheric BC. Moreover, their classification of the dominant aerosol type using the spectral aerosol optical prop-
605 erties indicated that BC was the predominant absorption component of aerosols at PDM, and controlled the variation of
Single Scattering Albedo throughout the two years.

– Atmospheric Mercury

Mercury (Hg) is a heavy metal that is dispersed globally in the gas phase following its emission from volcanoes and
human activities. The atmospheric lifetime of Hg is not well constrained due to its complex redox chemistry, making
610 model predictions of Hg deposition and ecosystem loading difficult. At the P2OA, we have generated from 2010-2014
one of the longest high-altitude atmospheric Hg datasets, using automated instruments that quantified gaseous elemental
Hg(0), gaseous oxidized Hg(II) and aerosol Hg(II) dynamics at a 2 h resolution (Fu et al., 2016a; Maruszczak et al.,
2017). Together with experimental rainfall Hg photoreduction rates (Yang et al., 2019), these data have helped to better
constrain atmospheric Hg redox reactions in global Hg chemistry and transport models (Saiz-Lopez et al., 2018). We
615 have also developed unique methods to sample and measure the stable isotope composition of atmospheric Hg(0) and



Hg(II) in gas, aerosol and precipitation, which inform on deposition pathways and fluxes (Fu et al., 2016b, 2021; Enrico et al., 2016). The data are freely available through the AERIS/GMOS¹³ and iGOS4M¹⁴ datahubs.

– Microplastics and other trace species

The P2OA also hosted punctual campaigns on the atmospheric chemistry and transport of diverse environment-impacting species such as halogens and selenium (Suess et al., 2019), formaldehyde (Prados-Roman et al., 2020) and microplastics (Allen et al., 2021). Over a period of four months in summer 2017, polymer fragments and fibers ranging in size from 3.5 to 53 micrometers were observed at relatively low (0.25 microplastics m⁻³) but significant levels. The polymers identified, polyethylene, polystyrene, polyvinyl chloride (PVC), polyethylene terephthalate (PET) and polypropylene, are all known for their use in packaging. The origin of these microplastics has been studied by back-trajectory modelling of the air masses observed at PDM. Many of these trajectories have their origin in Africa, the Atlantic Ocean and North America, indicating that an intercontinental transport is at the origin of the microplastics detected at PDM.

6.1.3 Exploring Transient Luminous Events

In the late 1980s, a new field of research opened with the discovery of the Transient Luminous Events (TLEs), which includes now sprites, ‘elves’ (Emission of Light and Very low-frequency perturbations from Electromagnetic pulse Sources), jets and other electrical phenomena occurring above thunderstorms (Füllekrug et al., 2006). The first European TLE-dedicated observations were obtained at PDM, in 2000 (Neubert et al., 2008). In the following years, several European teams joined in Eurosprite during summer and fall campaigns leading to over 700 TLE images being captured in the period from 2000 to 2008 (Neubert et al., 2008). The two observation sites of P2OA, equipped with a remote-controlled sensitive camera system, contributed to a climatology of TLEs in Europe (Arnone et al., 2020) and a large number of process studies (e.g. Soula et al., 2015, 2017; Gomez Kuri et al., 2021; Tomicic et al., 2021).

During five successive nights from 16 to 21 January 2017, several storm systems over the Mediterranean Sea, highly productive in TLEs, were monitored with the camera at PDM. A total of 589 TLE events were recorded and analyzed in the thesis by Gomez Kuri (2021). This large number of TLEs and the diversity of the storm systems capable of producing TLEs allowed to support previous observations or reinforce some theories on mechanisms and conditions of their production. Figure 10 shows the continuity of the cloud-to-ground (CG) lightning flash activity during five days of the winter period within the region pointed by the camera. There was an increasing flash rate during several cycles corresponding with individual cells at the beginning of the period and an evolution to larger developments of storms such as Mesoscale Convective Systems (MCS) in the second part of the period. The TLEs were observed over the five nights, with specific time intervals more favorable to some TLE categories like elves (green histogram in Fig. 10) when the CG flash rate was large (around 20:00 UT on January 20th) and sprites (red histogram) a few hours later while the flash rate decreased substantially. The analysis of the location of each TLE in its storm context shows that sprites were often produced over stratiform regions of MCSs, during their dissipating

¹³<https://gmos.aeris-data.fr/>

¹⁴<http://www.gos4m.org/igos4m/>



stage, and associated with a positive stroke, the peak current of which was on average about 107 kA. elves mainly occurred in systems of strong convection, during the mature stage which produced negative strokes of high peak current. elves are more likely to occur with clouds that have large vertical but small horizontal extensions whereas sprites occur over clouds of smaller vertical but larger horizontal extensions (Gomez Kuri, 2021). In the night of 19 January, out of 106 TLEs observed above an MCS developing between south of Ebro delta and the Balearic Islands, 97% of 29 elves occurred over sea while 45% of sprites occurred over land. It is in accordance to other observations and to larger peak current values for strokes over sea.




Figure 10. TLE and CG lightning flash activity between 17 January 0000 UTC and 22 January 0000 UTC (from Gomez Kuri, 2021). (a) TLE rate over 15 minutes intervals (histogram) and mean positive and negative peak current of CG strokes in the same time intervals. (b) CG flash rates (negative and positive with black and grey histograms, respectively) and the minimum Cloud Top Temperature (CTT) over the area of study over 15 minutes (line).



6.2 Instrumental and methodological validation

The rich panel of instruments and infrastructures of both sites make P2OA a hosting platform suitable for research groups to
655 test new instruments or sensors, which can be compared to reference measurements of the permanent instrumentation. Here
we give two examples of such experimentations.

6.2.1 Instrumented RPAS and balloons experimentations

With its open spaces (70 ha) mainly composed of prairies, its instruments on boundary layer dynamics and its infrastructure
(hangar, storage buildings, mechanical and electrical workshops, meeting and working rooms, lodging), CRA is particularly
660 appropriate for balloons and RPAS operations, and airborne instrumentation tests. In total, about 30 test campaigns involving
light aircrafts and airborne instruments have been hosted at P2OA since 2015, either with a tethered balloon or with RPAS (free
radiosounding balloons are not counted here). In the context of aerial activities and regulation, a Temporary Regulated Area is
activated for those operations when needed. The goal of the field campaign is either to test the flying vector, the fly strategy
or to test new sensors. Thus, P2OA was the test-site of RPAS and sensors pre-campaigns before international or national field
665 experiments like BACCHUS (Calmer et al., 2019), NEPHELAE (Hattenberger et al., 2022), EUREC4A (Maury et al., 2023).
Strategy of fleet flying (Hattenberger et al., 2022), catapult take off and thread landing have been tested. Sensors for turbulence
measurements (Calmer et al., 2018; Alaoui-Sosse et al., 2019, 2022) have been validated based on the P2OA 60-m tower
turbulence measurements.

Earlier during BLLAST, several new observational devices or methodologies had been tested or enabled : a technique of
670 frequently-launched radiosoundings with line-cutting system and re-usable probes (Legain et al., 2013), the SUMO RPAS,
which is a light tool for frequent profiling of meteorological variables (Reuder et al., 2016) and turbulence (Båserud et al.,
2016), a tethered-balloon-borne turbulence probe (Canut et al., 2016), a method for estimating heat fluxes based on frequent
profiling of the atmosphere (Båserud et al., 2020).

6.2.2 Improvement of airborne gamma-ray technique for radiological surveys and environmental applications

675 Gamma-ray spectrometry allows the identification and quantification of natural (U- and Th-decay products, ^{40}K) and artificial
(e.g., ^{137}Cs) radionuclides in the environment. Monitoring their time and space variations offers the possibility to study the
environmental factors responsible for these variations, such as soil humidity content and vertical profile (which have an effect
on gamma-ray attenuation), surface-atmosphere gas exchange, atmospheric boundary layer dynamics and synoptic transport
(which modulate radon flux and atmospheric concentration), dry and wet aerosols' scavenging (which affects the vertical
680 distribution of atmospheric radon decay products) and migration of radionuclides in soils. A suite of instruments dedicated
to these studies has been deployed at CRA by CEA/DAM and IRAP since 2018-2019: a 20L NaI(Tl) RSX-5 spectrometer
mounted on top of CRA's 60 m tower, atmospheric and soil radon monitors (3 AlphaGuards at 1 m, 30 m and 60 m ; 3 Barasol
probes at depths from 15 cm to 90 cm in the ground) and (Time Domain Reflectometry) TDR soil humidity probes. A specific
application consisted in quantifying the influence of environmental parameters (atmospheric radon, soil moisture, cosmic



685 radiation, atmospheric pressure) on the measurement of natural and artificial radioactivity by airborne gamma-ray spectrometry
(Amestoy et al., 2021), in order to improve the survey of sites with nuclear activities or radioactive fallouts, and to characterize
the evolution of surface deposits by remote sensing systems such as the HelinucTM system developed by CEA/DAM (see
Fig. 11). A campaign was conducted to monitor the combined evolution of these parameters during fourteen months, combined
with several helicopter flights over CRA, and resulting in the correction and validation of the protocol for airborne gamma-ray
690 spectrometry (called PASTHEL) that leads to high measurement precision (Amestoy, 2021). The simulation of a hovering flight
by installing the NaI(Tl) spectrometer at 50 m on the 60 m tower made it possible to continuously measure natural radioactivity
at a fixed point, focusing on environmental temporal variations. The addition of atmospheric radon detectors made it possible to
understand the dynamics of this radioactive gas and its influence on the gamma-ray signal. The installation of TDR probes, the
study of satellite images and the use of a pluviometer allowed the quantification of the influence of soil moisture and rainfall
695 on the gamma-ray signal. Thus, the effects of atmospheric radon, soil moisture and rainfall could be characterized and taken
into account to restore the gamma signal representative of the site's natural radioactivity.




Figure 11. (A) Principle of airborne gamma-ray spectrometry. (B) Simultaneous acquisition between the onboard spectrometer (RSX-4) and the one installed on the 60 m tower (RSX-5) at CRA, 8 September 2020.

6.3 Evaluation of numerical weather prediction models

6.3.1 Atmospheric dynamics variability

As mentioned before, mountain lee waves are typical of the area, generated by southwesterly flow over the Pyrenees. The
700 marked oscillations of the vertical velocity within the whole troposphere that are associated with them, the complex cloud
cover system formed, and the impact on temperature, moisture and transport of chemical species, are all difficult to capture in
a numerical model, mostly due to the complexity of the terrain itself, and to the challenge of taking the subgrid topography
into account.



The measurements of the VHF wind profiler radar at P2OA enable to test the ability of NWP models to simulate the
705 dynamics of the atmosphere in such cases. In particular, the vertical velocity observed by the radar in the mid-troposphere
can reach absolute values larger than 2 m s^{-1} in such case, while it is usually smaller than 30 cm s^{-1} in other situations
(except convective storms). Large vertical velocity variance can thus be observed over a few hours in case of mountain waves
in southerly flows, relatively to other typical synoptic situations. The use of a threshold on vertical velocity variance is actually
one possible diagnostic of their occurrence (Gueffier et al., 2024). Note that the VHF measurements correspond to rather large
710 scales: a few km horizontally (depending on height, due to beam divergence) and 375 m vertically.

One way to statistically evaluate the NWP models on this aspect is to compare the observed and modeled density energy
spectra of the vertical wind component. The two Météo-France NWP models are considered here, from which profiles are
continuously extracted at P2OA location, for model evaluations: ARPEGE (Courtier and Geleyn, 1988, 7 km horizontal res-
olution) and AROME (Seity et al., 2011, 1.3 km horizontal resolution). For AROME, 16 atmospheric columns are averaged
715 around CRA site, and 3 for ARPEGE. Model outputs are extracted at 1 h time resolution. Figure 12 shows the normalized
energy density spectra of both the zonal component of the wind (Fig. 12a) and the vertical velocity (Fig. 12b), for a 4-month
period of 2018, from January to April. This period was chosen as long enough for this typical comparison example, and with-
out any gap in the observational data. The chosen altitude is close to 3000 m a. s. l., that of the highest peaks at the border.
It remains representative of the process, since the whole troposphere, from 1 km or below, up to 8 km or above, coherently
720 oscillates vertically during mountain wave events (Gueffier et al., 2024).




Figure 12. Normalized energy density spectra of (a) the zonal component of the wind and (b) the air vertical velocity, at the altitude of 3325 m a. s. l. as observed by the P2OA VHF wind profiler radar, modeled by AROME and modeled by ARPEGE, over the 4-month period of January to April 2018.



Figure 12a first shows that the 2 models correctly represent the variability of the horizontal wind at timescales ranging from 1 h to 1 month. At large synoptic scales, ARPEGE and AROME are very similar, consistently with the forcing of AROME with ARPEGE. They slightly depart from observations around the scale of 30–40 days, which is difficult to interpret. Consistently with their horizontal resolution, ARPEGE slightly departs from the observation at scales smaller than 1 day, while AROME is remarkably faithful to the observations at scales smaller than 1 day.

Based on the quality of horizontal wind spectra, one can now consider the vertical velocity variability. Figure 12b reveals how lower is the energy in ARPEGE: the coarser model is not able to represent the variability of the vertical velocity, at any timescale, with energy density about 10 times smaller than observations. Among other processes, it always strongly underestimates the oscillations of the troposphere during foehn or mountain wave events (not shown), which occur around 10 % of the time (Gueffier et al., 2024). The finer model AROME, however, much better represents the vertical velocity variability. It has globally a very satisfying energy density level, even if it slightly underestimates the variance at time scales smaller than 2 days.

This example shows the potential of the 20-year long VHF wind profiler dataset for evaluation of the NWP models and the studies of dynamical processes that are typical of mountain regions.

6.3.2 Surface fluxes

As overlined by the MOSAI project (see section 6.1.1), there is a clear need to dig into the representation of surface fluxes in the NWP and climate models. The long-term flux series like those of CRA can now be taken as a reference for their evaluation. Figure 13 shows the difference between the modeled fluxes and the observed fluxes, for both the sensible and the latent heat fluxes. It gives an example only for one year (2021), and with monthly averages, so that inter-annual and intraseasonal variability are worked off here. The overestimation of the sensible and latent heat fluxes by the two models from April to September is consistent with what Couvreux et al. (2016) found during BLLAST experiment in June 2011. One plausible explanation was related to the land-use (with more forest than real land-use in several grid points, especially in ARPEGE) and soil moisture. In specific conditions during BLLAST though, which correspond to a heat wave and very small or negative surface fluxes, the models then underestimated the fluxes. In fall and winter here, we find in average an underestimation of the sensible heat flux by the models, while the latent heat flux remains generally overestimated. The former could be related to the underestimation of the 2 m temperature almost all year long (not shown), with larger negative bias in winter and fall, reaching down to -2.5 °C in average in December 2021 in ARPEGE for example. This reveals the difficulty of such comparisons when the environment (temperature, relative humidity, wind, radiation and clouds, soil conditions...) may itself differ between model and observation, leading to different surface heat fluxes. In this case, it is not possible to conclude on the capability of the models to correctly represent the surface fluxes.




Figure 13. Difference between the monthly (a) sensible and (b) latent heat fluxes observed at 30 m at CRA and the simulated fluxes of ARPEGE and AROME NWP models.

750 This type of comparison is thus of clear interest as a first step for model evaluation, but is not sufficient to conclude on true
 model biases, and their sources. One first need to fully understand the representativity of the measured flux, and the meaning
 of the modeled flux on a given grid point. It is, as a second step, important to consider an appropriate way to compare both
 together for model evaluation, different such a direct point-to-point comparison. This is the core question of MOSAI project
 (Lohou et al., 2024). In its context, Zouzoua et al. (2024) for example propose a new method for model evaluation, based on
 755 supervised neural networks.



6.3.3 Validation of a regional climate model on aerosol composition

A one-year sensitivity study was performed at the European scale with the Regional Climate Model RegCM (Filippo et al., 2012) over the year 2010. Aerosol radiative forcings and feedbacks are highly dependent of the physical, optical and chemical properties of aerosols, and of their spatial and temporal distributions. Aerosol sources, transport and sinks performances of RegCM were assessed for the 2010 year over the complex terrain of Pyrenees thanks to the PDM aerosol dataset.

Simulated monthly mean aerosol concentrations were evaluated against in situ aerosol concentrations of elemental carbon (EC), organic carbon (OC), aerosol sulfate (SO_4^{2-}), calcium ion (Ca^{2+}), magnesium ion (Mg^{2+}), chlorine ion (Cl^-), sodium ion (Na^+) from PDM. Ca^{2+} and Mg^{2+} are considered as dust tracers, and Cl^- and Na^+ as sea salt (SSLT) tracers. Monthly means of PDM aerosol tracer concentrations were calculated from weekly filter samplings. The weekly sampling integrated volumes of air pumped continuously for 7 days. The chemical analyses performed on these filters were: ion chromatography analysis for the inorganic fraction of aerosols (World Meteorological Organization Quality Assurance/Science Activity Centre (WMO QA/SAC)¹⁵, for laboratory number 700106); and thermo-optical analysis according to the IMPROVE protocol (Chow et al., 2007) for the organic fraction of aerosols for the year studied. The analytical errors were estimated to 5%.

In Fig. 14, monthly averages for 2010 are plotted, as well as the monthly averages over the period 2002-2018. Comparing these two trends allows the 2010 variation to be compared to an average variation over a 17-year dataset. The 2010 seasonal variation lays within one standard deviation around the multi-year average. Only the month of July 2010 shows an average of each chemical compound outside the multi-annual average and associated standard deviation, which is related to the fact that the number of samples contributing to the average is less important and represents specific episodes. The high standard deviations around the multi-annual mean of calcium and magnesium ions (Fig. 14b) shows the occasional occurrence of intense dust episodes over PDM.

The BC and OC seasonal variability is well reproduced by the model as the modeled seasonal variation of BC and OC compounds is comparable to the seasonal variation of measured EC and OC. The model seems to well reproduce the temporal occurrence of regional scale biomass burnings and secondary organic aerosol during the summer period (Fig. 14a). Only the intensities of these annual trends are not comparable, the model largely underestimating the concentrations. This is related to the overestimated dry deposition for carbonaceous aerosol parameterized in the model.

Modeled sulfate concentrations show a temporal evolution and an intensity of comparable in order of magnitude with sulfate measurements (Fig. 14c). Figure 14b shows that the model overestimates the dust aerosol concentrations all along the year with a large magnitude. Tsikerdekis et al. (2017) showed through a sensitivity study, that RegCM overestimates dust emission fluxes. The seasonal variation seems to present little consistency with the measurement during the months of March and April. Concerning sea-salt concentrations (Fig. 14d), the modeled concentrations show an annual variation that does not agree with the measurement of marine influence tracers at PDM. This reveals again the problem of overestimation of the marine aerosol sources parameterized in the model according to the intensity of the wind fields in cyclonic season over the oceanic area.

¹⁵<http://www.qasac-americas.org/>




Figure 14. Annual trends of monthly mean mass concentrations ($\mu\text{g m}^{-3}$) of aerosol compounds modeled with RegCM in the Pyrenean region, over the year 2010 with horizontal spatial resolution of $35\text{km} \times 35\text{km}$, compared to monthly mean concentrations measured in situ at PDM. (a) Modeled BC and OC mixing ratios with measured EC and OC. (b) Modeled DUST mixing ratio with Ca^{2+} and Mg^{2+} measured as main tracers of dust influence. (c) Modeled SO_4^{2-} mixing ratio with measured sulfate concentrations. (d) Modeled SSLT mixing ratio with Na^+ and Cl^- measured concentration as main tracers of marine influence.

7 Conclusions

To conclude, we have shown that a broad spectrum of scientific questions and applications can be addressed based on the long-term data set collected at P2OA, due to the rich set of instruments for meteorological dynamics, atmospheric composition or electricity. Emphasis is made on specific expertises like **boundary layer** dynamics, trace gas transport, aerosol chemical and physical properties, or transient luminous events. Most continuously operated instruments are connected to a French or international network, for weather forecast service, climate and air quality monitoring, or atmospheric process studies.

From the point of view of air composition, the location of P2OA near the Atlantic Ocean makes it weakly influenced by human activity on continental Europe. Further, its implementation near the Pyrenees makes it specific for complex terrain studies. The coupling of the high altitude site at PDM and the plain site at CRA favours the analysis of orographically-forced regimes and their impact on exchange and transport of trace species.



800 With such a set of instruments and the capability of its infrastructure (lodging, workshops, meeting and conference room, etc.), P2OA is also a well-equipped hosting facility. CRA is especially suited to operate light RPAS and balloons, and PDM for experiments in high altitude. But beyond these specificities, a broad spectrum of instrumental tests or field campaigns took place at P2OA. P2OA is also a favourable and recognized place for educational training in atmospheric research: several undergraduate or master level trainings are yearly organized, based on micro field campaigns (radiosoundings, surface energy balance station, etc.), and the use of the permanent instruments.

805 In the context of the main science topics addressed at P2OA and illustrated before, perspectives of new instrumentation is foreseen:

- A new high resolution Doppler lidar should be installed on CRA site by the end of 2024, which will be complementary of all other devices for the description of the boundary layer dynamics. Its fine temporal and spatial resolution should give more turbulence statistics in clear air, and open new horizons for the turbulence retrieval observational techniques.
 - As coming contribution to the European ACTRIS infrastructure, two monitoring systems of atmospheric short-lived reactive gases will be installed at PDM in 2024: a proton-transfer-reaction – time-of-flight mass spectrometer (PTR-TOF-MS) for measuring a selection of volatile organic compounds (VOCs), and a NO/NO₂ monitoring system. These instruments will complement existing aerosol and gas measurements at PDM, as the targeted species are chemical precursors of ozone and secondary aerosols. They will also allow for a much better characterization of emission sources affecting the sampled air masses: biomass burning (from agriculture or forest fires), anthropogenic benzene sources, oceanic influence, etc.
- 815

Data availability. All the dataset is available at <https://p2oa.aeris-data.fr/catalogue/>. Part of the data is also available from <https://www.actris.fr/actris-fr-data-centre/>, and from <https://www.aeris-data.fr/catalogue/>.



Appendix A: Instrumentation of the 60 m tower

Table A1. Instruments installed on the 60 m tower of P2OA-CRA, listed by height, with corresponding measured variables and sampling frequency.

| Height | Sensor | Main variable(s) | Frequency |
|--------|----------------------|--|-----------|
| 2 m | Barometer | Pressure | 0.1 Hz |
| | Rain gauge | Rainfall | 0.1 Hz |
| | HMP45 | Temperature, Humidity | 0.1 Hz |
| | Flux plates (3) | Ground heat flux | 0.1 Hz |
| 15 m | HMP45 | Temperature, Humidity | 0.1 Hz |
| | wind vane | Wind direction | 0.1 Hz |
| | wind anemometer | Wind speed | 0.1 Hz |
| 30 m | HMP45 | Temperature, Humidity | 1 Hz |
| | Sonic anemoter CSAT3 | 3D wind, Virtual temperature | 10 Hz |
| | Licor LI7500 | Water vapour and CO ₂ concentration | 10 Hz |
| 45 m | HMP45 | Temperature, Humidity | 1 Hz |
| | Sonic anemoter GILL | 3D wind, Virtual temperature | 10 Hz |
| | wind vane | Wind direction | 0.1 Hz |
| | wind anemometer | Wind speed | 0.1 Hz |
| 60 m | Radiometer CNR1 | Up. & Down. SW & LW radiation | 1 Hz |
| | HMP45 | Temperature, Humidity | 1 Hz |
| | Sonic anemoter CSAT3 | 3D wind, Virtual temperature | 10 Hz |

Author contributions. Marie Lothon and François Gheusi, as scientific coordinators of the P2OA, are the main redactors of this article.

820 The P2OA scientific PIs are contributing authors, notably for sections dealing on their specific topics: Fabienne Lohou for the meteorological and flux ground stations; François Gheusi for atmospheric composition; Véronique Pont for aerosol instrumentation and studies; Marie Lothon, Bernard Campistron, Frédérique Saïd for the wind profiler radars; Marie Lothon for the sky imager and ceilometer; Jeroen Sonke for mercury and microplastics; Serge Soula for atmospheric electricity; Corinne Jambert for NO_x and VOCs measurements; Michel Ramonet for the Greenhouse Gas monitoring; Pierre Bosser for the GNSS antennas; Pierre-Yves Meslin and Julien Amestoy for gamma

825 spectroscopy; Olivier Masson and Romain Vidal for radionuclide activity. Solène Derrien is responsible for the CRA instrumentation and data management. Solène Derrien, Emmanuel Leclerc, Antoine Vial, Yannick Bezombes, Eric Gardrat, Gilles Athier operate and maintain the P2OA instrumentation and ensure data quality control. Damien Boulanger, Nicolas Pascal, Renaud Bodichon, Yves Meyerfeld contribute or have contributed to the data production and dissemination, in the context of AERIS data service, and within SEDOO, ESPRI or ICARE

830 CALOTRITON algorithm (Philibert et al., 2024) and other data process algorithms, he provided the convective boundary-layer height composite; Gilles Athier is the author of the in-cloud index algorithm. Eric Pique, Jean-Bernard Estrampes, Fabienne Guesdon, Felix Starck, Guillaume Bret, Laurent Cabanas and Erwan Bargain contribute or have contributed to the technical maintenance of the P2OA site and instrumentation. Jérémy Gueffier provided the monthly composite diurnal cycles of atmospheric composition at PDM. Guylaine Canut has



835 made the comparison between the AROME and ARPEGE operational numerical weather predicting models with the P2OA flux measure-
ments. Zaida Gomez Kuri has made the analysis of the Transient Luminescent Events shown in this article. All coauthors provided internal
reviews.

Competing interests. The authors declare no competing interests.

Acknowledgements. P2OA facilities and staff are funded and supported by the University Paul Sabatier Toulouse 3, France, and CNRS
(Centre National de la Recherche Scientifique). P2OA is part of the national research infrastructure ACTRIS-France.

840 Several P2OA dataset are related to ACTRIS-Fr research infrastructure (<https://www.actris.fr/>). Associated products and services are
maintained by the French national center for Atmospheric data and services AERIS. ACTRIS data policy can be found here:

https://www.actris.eu/sites/default/files/Documents/ACTRIS%20PPP/Deliverables/ACTRIS%20PPP_WP2_D2.3_ACTRIS%20Data%20policy.pdf.

We thank OMP/SEDOO, IPSL/ESPRI, and ICARE data centers for their contribution to AERIS and P2OA data services. We also specially
thank Andry Andriatiana for his involvement in the P2OA data base and web site.

845 Rénag is member of the Research Infrastructure (RI) Résif-Epos, managed by CNRS-Insu. Résif-Epos is inscribed on the roadmap of the
Ministry of Higher Education, Research and Innovation, the Résif-Epos IR is a consortium of eighteen French research organizations and
institutions. Résif-Epos benefits from the support of the Ministry of Ecological Transition.

We thank Eric Bazile and Yann Seity from Météo-France/CNRM for providing the AROME and ARPEGE operational model outputs,
and for their expertise.



850 References

- Alaoui-Sosse, S., Durand, P., Medina, P., Pastor, P., Lothon, M., and Cernov, I.: OVLI-TA: An Unmanned Aerial System for Measuring Profiles and Turbulence in the Atmospheric Boundary Layer, *Sensors*, 19, 581, 2019.
- Alaoui-Sosse, S., Durand, P., Medina, P., Pastor, P., Gavart, M., and Pizziol, S.: BOREAL-A Fixed-Wing Unmanned Aerial System for the Measurement of Wind and Turbulence in the Atmospheric Boundary Layer, *J. Atmos. Oc. Tech.*, 39, 387–402, 2022.
- 855 Allen, S., Allen, D., Baladima, F., Phoenix, V., Thomas, J., Le Roux, G., and Sonke, J.: Evidence of free tropospheric and long-range transport of microplastic at Pic du Midi Observatory, *Nature Communications*, 12, 7242, <https://doi.org/10.1038/s41467-021-27454-7>, 2021.
- Amestoy, J.: Etude de l'influence des facteurs environnementaux sur la spectrométrie gamma aéroportée et application à la surveillance environnementale de sites., PhD Thesis, Université Toulouse III, <https://theses.hal.science/tel-03515339>., 2021.
- Amestoy, J., Meslin, P. Y., Richon, P., Delpuech, A., Derrien, S., Raynal, H., Pique, E., Baratoux, D., Chotard, P., Van Beek, P., Souhaut, M.,
860 and Zambardi, T.: Effects of environmental factors on the monitoring of environmental radioactivity by airborne gamma-ray spectrometry, *J. Env. Radioactivity*, 237, 106 695, <https://doi.org/10.1016/j.jenvrad.2021.106695>, 2021.
- Ammoura, L., Xueref-Remy, I., Vogel, F., Gros, V., Baudic, A., Bonsang, B., Delmotte, M., Té, Y., and Chevallier, F.: Exploiting stagnant conditions to derive robust emission ratio estimates for CO₂, CO and volatile organic compounds in Paris, *Atmospheric Chemistry and Physics*, 16, 15 653–15 664, <https://doi.org/10.5194/acp-16-15653-2016>, <https://acp.copernicus.org/articles/16/15653/2016/>, 2016.
- 865 Argentini, S., Viola, A., Sempreviva, A., and Petenko, I.: Summer boundary-layer height at the plateau site of Dome'C, antarctica, *Boundary-Layer Meteorology*, 115, 409–422, <https://doi.org/10.1007/s10546-004-5643-6>, 2005.
- Arnone, E., Bór, J., Chanrion, O., Barta, V., Dietrich, S., Enell, C. F., Farges, T., Füllekrug, M., Kero, A., Labanti, R., Mäkelä, A., Mezuman, K., Odzimek, A., Popek, M., Prevedelli, M., Ridolfi, M., Soula, S., Valeri, D., van der Velde, O., Yair, Y., Zanotti, F., Zoladek, P., and Neubert, T.: Climatology of Transient Luminous Events and Lightning Observed Above Europe and the Mediterranean Sea, *Surveys in*
870 *Geophysics*, 41, 167–199, <https://doi.org/10.1007/s10712-019-09573-5>, 2020.
- Bamberger, I., Oney, B., Brunner, D., Henne, S., Leuenberger, M., Buchmann, N., and Eugster, W.: Observations of Atmospheric Methane and Carbon Dioxide Mixing Ratios: Tall-Tower or Mountain-Top Stations?, *Boundary-Layer Meteorology*, 164, 135–159, <https://doi.org/10.1007/s10546-017-0236-3>, 2017.
- Baray, J.-L., Courcoux, Y., Keckhut, P., Portafaix, T., Tulet, P., Cammas, J.-P., Hauchecorne, A., Godin Beekmann, S., De Mazière, M.,
875 Hermans, C., Desmet, F., Sellegri, K., Colomb, A., Ramonet, M., Sciare, J., Vuillemin, C., Hoareau, C., Dionisi, D., Dufлот, V., Vérèmes, H., Porteneuve, J., Gabarrot, F., Gaudo, T., Metzger, J.-M., Payen, G., Leclair de Bellevue, J., Barthe, C., Posny, F., Ricaud, P., Abchiche, A., and Delmas, R.: Maïdo observatory: a new high-altitude station facility at Reunion Island (21° S, 55° E) for long-term atmospheric remote sensing and in situ measurements, *Atmospheric Measurement Techniques*, 6, 2865–2877, <https://doi.org/10.5194/amt-6-2865-2013>, <https://amt.copernicus.org/articles/6/2865/2013/>, 2013.
- 880 Baray, J.-L., Bah, A., Cacault, P., Sellegri, K., Pichon, J.-M., Deguillaume, L., Montoux, N., Noel, V., Seze, G., Gabarrot, F., Payen, G., and Dufлот, V.: Cloud Occurrence Frequency at Puy de Dôme (France) Deduced from an Automatic Camera Image Analysis: Method, Validation, and Comparisons with Larger Scale Parameters, *Atmosphere*, 10, <https://doi.org/10.3390/atmos10120808>, <https://www.mdpi.com/2073-4433/10/12/808>, 2019.
- Baray, J.-L., Deguillaume, L., Colomb, A., Sellegri, K., Freney, E., Rose, C., Van Baelen, J., Pichon, J.-M., Picard, D., Fréville, P., Bouvier, L.,
885 Ribeiro, M., Amato, P., Banson, S., Bianco, A., Borbon, A., Bourcier, L., Bras, Y., Brigante, M., Cacault, P., Chauvigné, A., Charbouillot, T., Chaumerliac, N., Delort, A.-M., Delmotte, M., Dupuy, R., Farah, A., Febvre, G., Flossmann, A., Gourbeyre, C., Hervier, C., Hervo,



- 890 M., Huret, N., Joly, M., Kazan, V., Lopez, M., Mailhot, G., Marinoni, A., Masson, O., Montoux, N., Parazols, M., Peyrin, F., Pointin, Y., Ramonet, M., Rocco, M., Sancelme, M., Sauvage, S., Schmidt, M., Tison, E., Vaitilingom, M., Villani, P., Wang, M., Yver-Kwok, C., and Laj, P.: Cézeaux-Aulnat-Opme-Puy De Dôme: a multi-site for the long-term survey of the tropospheric composition and climate change, *Atmospheric Measurement Techniques*, 13, 3413–3445, <https://doi.org/10.5194/amt-13-3413-2020>, <https://amt.copernicus.org/articles/13/3413/2020/>, 2020.
- Blay-Carreras, E., Pardyjak, E. R., Pino, D., Alexander, D. C., Lohou, F., and Lothon, M.: Countergradient heat flux observations during the evening transition period, *ATMOSPHERIC CHEMISTRY AND PHYSICS*, 14, 9077–9085, 2014a.
- Blay-Carreras, E., Pino, D., de Arellano, J. V.-G., van de Boer, A., De Coster, O., Darbieu, C., Hartogensis, O., Lohou, F., Lothon, M., and 895 Pietersen, H.: Role of the residual layer and large-scale subsidence on the development and evolution of the convective boundary layer, *ATMOSPHERIC CHEMISTRY AND PHYSICS*, 14, 4515–4530, 2014b.
- Blay-Carreras, E., Pardyjak, E. R., Pino, D., Hoch, S. W., Cuxart, J., Martinez, D., and Reuder, J.: Lifted temperature minimum during the atmospheric evening transition, *ATMOSPHERIC CHEMISTRY AND PHYSICS*, 15, 6981–6991, 2015.
- Boehm, J., Werl, B., and Schuh, H.: Troposphere mapping functions for GPS and very long baseline interferometry from European Centre for Medium-Range Weather Forecasts operational analysis data, *Journal of Geophysical Research*, 11, 2406–, 900 <https://doi.org/10.1029/2005JB003629>, 2006.
- Bosser, P. and Bock, O.: IWV retrieval from ground GNSS receivers during NAWDEX, *Advances in Geosciences*, 55, 13–22, <https://doi.org/10.5194/adgeo-55-13-2021>, 2021.
- Bosveld, F., Baas, P., Beljaars, A., Holtslag, A. A. M., Vilà-Guerau de Arellano, J., and van de Wiel, B. J. H.: Fifty Years of Atmospheric Boundary-Layer Research at Cabauw Serving Weather, Air Quality and Climate, *Boundary-Layer Meteorol.*, 177, 583–612, 905 <https://doi.org/10.1007/s10546-020-00541-w>, 2020.
- Bou-Zeid, E., Anderson, W., Katul, G. G., and Mahrt, L.: The Persistent Challenge of Surface Heterogeneity in Boundary-Layer Meteorology: A Review, *Boundary Layer Meteorology*, 177, 227–245, 2020.
- Bougeault, P., Clar, A., Bénech, B., Carissimo, B., Pelon, J., and Richard, E.: Momentum budget over the Pyrenees - The PYREX experiment, 910 *Bulletin of the American Meteorological Society*, 71, 806–818, 1990.
- Browning, K. A. and Wexler: The determination of kinematic properties of a wind field using Doppler radar, *J. App. Met.*, 7, 105–113, 1968.
- Båserud, L., Reuder, J., Jonassen, M. O., Kral, S. T., Paskyabi, M. B., and Lothon, M.: Proof of concept for turbulence measurements with the RPAS SUMO during the BLLAST campaign, *Atmos. Meas. Tech.*, 9, 4901–4913, 2016.
- Båserud, L., Reuder, J., Jonassen, M. O., Bonin, T. A., Chilson, P. B., Jiménez, M. A., and Durand, P.: Potential and Limitations in Estimating Sensible-Heat-Flux Profiles from Consecutive Temperature Profiles Using Remotely-Piloted Aircraft Systems, *Boundary-Layer Meteorology*, 174, 145–177, 2020.
- Calmer, R., Roberts, G. C., Sanchez, K. J., Sciare, J., Sellegri, K., Picard, D., Vrekoussis, M., and Pikridas, M.: Aerosol–cloud closure study on cloud optical properties using remotely piloted aircraft measurements during a BACCHUS field campaign in Cyprus, *Atmos. Chem. Phys.*, 19, 13 989–14 007, <https://doi.org/10.5194/acp-19-13989-2019>, 2019.
- 920 Calmer, R., Roberts, G. C., Preissler, J., Sanchez, K. J., Derrien, S., and O’Dowd, C.: Vertical wind velocity measurements using a five-hole probe with remotely piloted aircraft to study aerosol–cloud interactions, *Atmos. Meas. Tech.*, 11, 2583–2599, 2018.
- Campistron, B., Long, A. B., and Hugin, A. W.: A method of retrieving turbulence parameters from volume processing of single-Doppler radar measurements, *J. Oc. atmos. Tech.*, 8, 491–505, 1991.



- 925 Campistron, B., Point, Y., Lohou, F., and Pages, J.-P.: Aspect sensitivity of VHF radar echoes observed in the middle and upper troposphere during the passage of a cut-off low, *Radio Science*, 34, 667–679, 1999.
- Canut, G., Calvet, J. C., Maurel, W., and Paci, A.: Seven years (2012–2018) of continuous observation of the surface energy budget and of soil moisture and temperature profiles in a peri-urban area, EMS2019-687-1, 2019, Copenhagen, 16, 2019.
- Canut, G., Couvreur, F., Lothon, M., Legain, D., Pignatelli, B., Lampert, A., Maurel, W., and Moulin, E.: Turbulence fluxes and variances measured with a sonic anemometer mounted on a tethered balloon, *Atmos. Meas. Tech.*, 9, 4375–4386, 2016.
- 930 Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J.-P.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, *Atmos. Meas. Tech.*, 3, 79–89, <https://doi.org/10.5194/amt-3-79-2010>, 2010.
- Chambers, S. D., Zahorowski, W., Williams, A. G., Crawford, J., and Griffiths, A. D.: Identifying tropospheric baseline air masses at Mauna Loa Observatory between 2004 and 2010 using Radon-222 and back trajectories, *Journal of Geophysical Research: Atmospheres*, 118, 992–1004, <https://doi.org/10.1029/2012jd018212>, <https://doi.org/10.1029/2012jd018212>, 2013.
- 935 Chambers, S. D., Williams, A. G., Conen, F., Griffiths, A. D., Reimann, S., Steinbacher, M., Krummel, P. B., Steele, L. P., van der Schoot, M. V., Galbally, I. E., Molloy, S. B., and Barnes, J. E.: Towards a Universal “Baseline” Characterisation of Air Masses for High- and Low-Altitude Observing Stations Using Radon-222, *Aerosol and Air Quality Research*, 16, 885–899, <https://doi.org/10.4209/aaqr.2015.06.0391>, <http://dx.doi.org/10.4209/aaqr.2015.06.0391>, 2016.
- Chevalier, A., Gheusi, F., Delmas, R., Ordonez, C., Sarrat, C., Zbinden, R., Thouret, V., Athier, G., and Cousin, J.-M.: Influence of altitude 940 on ozone levels and variability in the lower troposphere: a ground-based study for western Europe over the period 2001–2004, *Atmos. Chem. Phys.*, 7, 4311–4326, <https://doi.org/10.5194/acp-7-4311-2007>, 2007.
- Chiriaco, M., Dupont, J.-C., Bastin, S., Badosa, J., Lopez, J., Haefelin, M., Chepfer, H., and Guzman, R.: ReOBS: a new approach to synthesize long-term multi-variable dataset and application to the SIRTAs supersite, *Earth Syst. Sci. Data*, 10, 919–940, <https://doi.org/10.5194/essd-10-919-2018>, 2018.
- 945 Chow, J. C., Watson, J. G., Antony Chen, L. W., Oliver Chang, M. C., Robinson, N. F., Trimble, D., and Kohl, S.: The IMPROVE_A Temperature Protocol for Thermal/Optical Carbon Analysis: Maintaining Consistency with a Long-Term Database, *Journal of the Air and Waste Management Association*, 57, 1014–1023, <https://doi.org/10.3155/1047-3289>, 2007.
- Collaud Coen, M., Weingartner, E., Furger, M., Nyeki, S., Prévôt, A. S. H., Steinbacher, M., and Baltensperger, U.: Aerosol climatology and planetary boundary influence at the Jungfrauoch analyzed by synoptic weather types, *Atmospheric Chemistry and Physics*, 11, 950 5931–5944, <https://doi.org/10.5194/acp-11-5931-2011>, <https://acp.copernicus.org/articles/11/5931/2011/>, 2011.
- Collaud Coen, M., Andrews, E., Aliaga, D., Andrade, M., Angelov, H., Bukowiecki, N., Ealo, M., Fialho, P., Flentje, H., Hallar, A. G., Hooda, R., Kalapov, I., Krejci, R., Lin, N.-H., Marinoni, A., Ming, J., Nguyen, N. A., Pandolfi, M., Pont, V., Ries, L., Rodríguez, S., Schauer, G., Sellegri, K., Sharma, S., Sun, J., Tunved, P., Velasquez, P., and Ruffieux, D.: Identification of topographic features influencing aerosol observations at high altitude stations, *Atmospheric Chemistry and Physics*, 18, 12 289–12 313, <https://doi.org/10.5194/acp-18-12289-2018>, <https://acp.copernicus.org/articles/18/12289/2018/>, 2018.
- 955 Courtier, P. and Geleyn, J.-F.: The AROME-France Convective-Scale Operational Model, *Q. J. R. Meteorol. Soc.*, 114, 1321–1346, 1988.
- Couvreur, F., Bazile, E., Canut, G., Seity, Y., Lothon, M., Lohou, F., Guichard, F., and Nilsson, E.: Boundary-layer turbulent processes and mesoscale variability represented by numerical weather prediction models during the BLLAST campaign, *Atmos. Phys. Chem.*, 16, 8983–9002, 2016.



- 960 Cuxart, J., Wrenger, B., Martínez-Villagrasa, D., Reuder, J., Jonassen, M. O., Jiménez, M. A., Lathon, M., Lohou, F., Hartogensis, O.,
Dünnemann, J., Conangla, L., and Garai, A.: Estimation of the advection effects induced by surface heterogeneities in the surface energy
budget, *Atmos. Phys. Chem.*, 16, 9489–9504, 2016.
- Darbieu, C., Lohou, F., Lathon, M., de Arellano, J. V.-G., Couvreux, F., Durand, P., Pino, D., Patton, E. G., Nilsson, E., Blay-Carreras, E.,
and Gioli, B.: Turbulence vertical structure of the boundary layer during the afternoon transition, *ATMOSPHERIC CHEMISTRY AND*
965 *PHYSICS*, 15, 10 071–10 086, 2015.
- De Wekker, S.: Observational and Numerical Evidence of Depressed Convective Boundary Layer Heights near a Mountain Base, *J. App.*
Met. Clim., 47, 1017–1026, <https://doi.org/10.1175/2007JAMC1651.1>, 2008.
- Dessens, J. and Bücher, A.: Changes in minimum and maximum temperature at the Pic du Midi in relation with humidity and cloudiness,
1882-1984, *Atmospheric Research*, 37, 147–162, 1995.
- 970 El Guernaoui, O., Reuder, J., Esau, I., Wolf, T., and Maronga, B.: Scaling the Decay of Turbulence Kinetic Energy in the Free-Convective
Boundary Layer, *Boundary Layer Meteorology*, 173, 79–97, 2016.
- Elvidge, A. D. and Renfrew, I. A.: The causes of foehn warming in the lee of mountains, *Bulletin of the American Meteorological Society*,
97, 455–466, <https://doi.org/10.1175/BAMS-D-14-00194.1>, 2016.
- Enrico, M., Le Roux, G., Maruszczak, N., Heimbürger, L.-E., Claustres, A., Fu, X., Sun, R., and Sonke, J.: Atmospheric mer-
975 cury transfer to peat bogs dominated by gaseous elemental mercury dry deposition, *Environmental Science & Technology*,
<https://doi.org/10.1021/acs.est.5b06058>, <https://doi.org/10.1021/acs.est.5b06058>, 2016.
- Evrard, O., Van Beek, P., Gateuille, D., Pont, V., Lefèvre, I., Lansard, B., and P., B.: Evidence of the radioactive fallout in France due to the
Fukushima nuclear accident, *J. Env. Radioactivity*, 114, 54–60, <https://doi.org/10.1016/j.jenvrad.2012.01.024>, 2012.
- Filippo, G., Coppola, E., Solmon, F., Mariotti, L., Sylla, M. B., Bi, X., and Elguindi, N. e. a.: RegCM4: model description and preliminary
980 tests over multiple CORDEX domains, *Climate Research*, 52, 7–29, <https://doi.org/10.3354/cr01018>, 2012.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C., Luijkx, I. T.,
Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P.,
Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra,
N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser,
985 T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, O., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C.,
Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D.,
Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot,
N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K. M., Olsen, A., Omar,
A. M., Ono, T., Paulsen, M., Pierrot, D., Pockock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C.,
990 Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao,
S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-
Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth System*
Science Data, 15, 5301–5369, <https://doi.org/10.5194/essd-15-5301-2023>, <https://essd.copernicus.org/articles/15/5301/2023/>, 2023.
- Fu, X., Maruszczak, N., Heimbürger, L., Sauvage, B., Gheusi, F., and Sonke, J.: Atmospheric mercury speciation dynamics at the high-altitude
995 Pic du Midi Observatory, southern France, *Atmospheric Chemistry and Physics*, 16, 5623–5639, <https://doi.org/10.5194/acp-16-5623-2016>,
<https://doi.org/10.5194/acp-16-5623-2016>, 2016a.



- Fu, X., Maruschak, N., Wang, X., Gheusi, F., and Sonke, J.: The isotopic composition of total gaseous mercury in the free troposphere of the Pic du Midi Observatory (2877 m a.s.l, France, Environmental Science & Technology, <https://doi.org/10.1021/acs.est.6b00033>, 2016b.
- 1000 Fu, X., Jiskra, M., Yang, X., Maruschak, N., Enrico, M., Chmeleff, J., Heimbürger-Boavida, L.-E., Gheusi, F., and Sonke, J.: Mass-Independent Fractionation of Even and Odd Mercury Isotopes during Atmospheric Mercury Redox Reactions, *Environ. Sci. Technol.*, 55, 10 164–10 174, <https://doi.org/10.1021/acs.est.1c02568>, 2021.
- Füllekrug, M., Mareev, E. A., and Rycroft, M. J.: Sprites, elves and intense lightning discharges, NATO Science Series II. In: Mathematics, physics and chemistry, Springer, Berlin, 225, 2006.
- GAW: Guidelines for the measurement of atmospheric carbon monoxide - GAW report no.192, Tech. rep., World Meteorological Organization (WMO), https://library.wmo.int/doc_num.php?explnum_id=9447, 2010.
- 1005 GAW: Guidelines for the measurement of atmospheric carbon monoxide - GAW report no.209, Tech. rep., World Meteorological Organization (WMO), https://library.wmo.int/doc_num.php?explnum_id=7814, 2013.
- Gheusi, F., Ravetta, F., Delbarre, H., Tsamalis, C., Chevalier-Rosso, A., Leroy, C., Augustin, P., Delmas, R., Ancellet, G., Athier, G., Bouchou, P., Campistron, B., Cousin, J.-M., Fourmentin, M., and Meyerfeld, Y.: Pic 2005, a field campaign to investigate low-tropospheric ozone variability in the Pyrenees, *Atmospheric Research*, 101, 640–665, <https://doi.org/10.1016/j.atmosres.2011.04.014>, <https://doi.org/10.1016/j.atmosres.2011.04.014>, 2011.
- 1010 Gomez Kuri, Z.: Analyse des couplages électrodynamiques transitoires à partir d’observations terrestres et spatiales, vol. 45, PhD Thesis, University of Toulouse, Toulouse, France, 2021.
- Gomez Kuri, Z., Soula, S., Neubert, T., Mlynarczyk, J., and Köhn, C.: Converging Luminosity in Column-Sprite Filaments, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2020GL090364>, 2021.
- 1015 Gomez-Pelaez, A. J., Ramos, R., Cuevas, E., Gomez-Trueba, V., and Reyes, E.: Atmospheric CO₂, CH₄, and CO with the CRDS technique at the Izaña Global GAW station: instrumental tests, developments, and first measurement results, *Atmospheric Measurement Techniques*, 12, 2043–2066, <https://doi.org/10.5194/amt-12-2043-2019>, <https://amt.copernicus.org/articles/12/2043/2019/>, 2019.
- Gueffier, J.: Régimes de temps et composition atmosphérique associée à la Plateforme Pyrénéenne d’Observation de l’Atmosphère : caractérisation par classification hiérarchique de 5 ans de données, PhD Thesis, Université Toulouse III, 2023.
- 1020 Gueffier, J., Gheusi, F., Lothon, M., Pont, V., Philibert, A., Lohou, F., Derrien, S., Bezombes, Y., Athier, G., Meyerfeld, Y., Vial, A., and Leclerc, E.: Weather regimes and the related atmospheric composition at a Pyrenean observatory characterized by hierarchical clustering of a 5-year data set, *Atmos. Phys. Chem.*, 24, 287–316, <https://doi.org/10.5194/acp-24-287-2024>, 2024.
- Guerova, G., Jones, J., Douša, J., Dick, G., de Haan, S., Pottiaux, E., Bock, O., Pacione, R., Elgered, G., Vedel, H., and Bender, M.: Review of the state of the art and future prospects of the ground-based GNSS meteorology in Europe, *Atmospheric Measurement Techniques*, 9, 5385–5406, <https://doi.org/10.5194/amt-9-5385-2016>, 2016.
- 1025 Guillaume, B., Lioussé, C., Rosset, R., Cachier, H., Van Velthoven, P., Bessagnet, B., and Poisson, N.: ORISAM-TM4: a new global sectional multi-component aerosol model including SOA formation — focus on carbonaceous BC and OC aerosols, *Tellus B*, 59, 283–302, 2007.
- Guillaume, B., Lioussé, C., Galy-Lacaux, C., Rosset, R., Gardrat, E., Cachier, H., Bessagnet, B., and Poisson, N.: Modeling exceptional high concentrations of carbonaceous aerosols observed at Pic du Midi in spring–summer 2003: Comparison with Sonnblick and Puy de Dôme, *Atmospheric Environment*, 42, 5140–5149, <https://doi.org/10.1016/j.atmosenv.2008.02.024>, 2008.
- 1030 Hadad, D., Baray, J.-L., Montoux, N., Van Baelen, J., Fréville, P., Pichon, J.-M., Bossert, P., Ramonet, M., Yver Kwok, C., Bègue, N., and Duflot, V.: Surface and Tropospheric Water Vapor Variability and Decadal Trends at Two Supersites of CO-PDD (Cézeaux and Puy de Dôme) in Central France, *Atmosphere*, 9, <https://doi.org/10.3390/atmos9080302>, 2018.



- 1035 Haeffelin, M., Barthès, L., Bock, O., Boitel, C., Bony, S., Bouniol, D., Chepfer, H., Chiriaco, M., Cuesta, J., Delanoë, J., Drobinski, P., Dufresne, J.-L., Flamant, C., Grall, M., Hodzic, A., Hourdin, F., Lapouge, F., Lemaître, Y., Mathieu, A., Morille, Y., Naud, C., Noël, V., O'Hirok, W., Pelon, J., Pietras, C., Protat, A., Romand, B., Scialom, G., and Vautard, R.: SIRTa, a ground-based atmospheric observatory for cloud and aerosol research, *Ann. Geophys.*, 23, 253–275, 2005.
- Hattenberger, G., Verdu, T., Maury, N., Narvor, P., Couvreur, F., Bronz, M., Lacroix, S., Cayez, G., and Roberts, G. C.: Field report: Deployment of a fleet of drones for cloud exploration, *Int. J. Micro Air Vehicles*, 14, 17568293211070 830, 2022.
- 1040 Heiskanen, J., Bruemmer, C., Buchmann, N., Calfapietra, C., Chen, H., Gielen, B., Gkritzalis, T., Hammer, S., Hartman, S., and Herbst, M.: The Integrated Carbon Observation System in Europe, *Bull. Amer. Met. Soc.*, 103, E855–E872, 2022.
- Heo, B.-H., Jacoby-Koaly, S., Kim, K.-E., Campistron, B., Bénech, B., and Jung, E.-S.: Use of the Doppler spectral width to improve the estimation of the convective boundary layer height from UHF wind profiler observations, *J. Oc. Atmos. Tech.*, 20, 408–424, 2003.
- 1045 Herrmann, E., Weingartner, E., Henne, S., Vuilleumier, L., Bukowiecki, N., Steinbacher, M., Conen, F., Collaud Coen, M., Hammer, E., Jurányi, Z., Baltensperger, U., and Gysel, M.: Analysis of long-term aerosol size distribution data from Jungfraujoch with emphasis on free tropospheric conditions, cloud influence, and air mass transport, *Journal of Geophysical Research: Atmospheres*, 120, 9459–9480, <https://doi.org/10.1002/2015JD023660>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JD023660>, 2015.
- Hulin, M., Gheusi, F., Lathon, A. M., Pont, V., Lohou, F., Ramonet, M., Delmotte, M., Derrien, S., Athier, G., Meyerfeld, Y., Bezombes, Y., Augustin, P., and Ravetta, F.: Observations of thermally driven circulations in the pyrenees: Comparison of detection methods and impact on atmospheric composition measured at a mountaintop, *Journal of Applied Meteorology and Climatology*, 58, 717–740, <https://doi.org/10.1175/JAMC-D-17-0268.1>, 2019.
- 1050 Jacoby-Koaly, S., Campistron, B., Bernard, S., Bénech, B., Girard-Ardhuin, F., Dessens, J., Dupont, E., and Carissimo, B.: Turbulent Dissipation Rate In The Boundary Layer Via UHF Wind Profiler Doppler Spectral Width Measurements, *Boundary Layer Meteorol.*, 103, 361–389, 2002.
- 1055 Keeling, C. D., Bacastow, R. B., Bainbridge, A. E., Ekdahl, C. A., Guenther, P. R., Waterman, L. S., and Chin, J. F. S.: Atmospheric carbon dioxide variations at Mauna Loa Observatory, Hawaii, *Tellus*, 28, 538–551, <https://doi.org/10.1111/j.2153-3490.1976.tb00701.x>, 1976.
- Kim, K.-E., Jung, E.-S., Campistron, B., and Heo, B.-H.: A physical examination of tropopause height and stratospheric air intrusion - A case study, *J. Met. Soc. Jap.*, 79, 1093–1103, 2001.
- 1060 Kooijmans, L. J. and Hartogensis, O. K.: Surface-Layer Similarity Functions for Dissipation Rate and Structure Parameters of Temperature and Humidity Based on Eleven Field Experiments, *Boundary Layer Meteorology*, 160, 501–527, 2016.
- Kotthaus, S., Haeffelin, M., Drouin, M.-A., Dupont, J.-C., Grimmond, S., Haeefe, A., Hervo, M., Poltera, Y., and Wiegner, M.: Tailored Algorithms for the Detection of the Atmospheric Boundary Layer Height from Common Automatic Lidars and Ceilometers (ALC), *Remote Sens.*, 12, 3259, <https://doi.org/10.3390/rs12193259>, 2020.
- 1065 Lac, C., Chaboureaud, J.-P., Masson, V., Pinty, J.-P., Tulet, P., Escobar, J., Leriche, M., Barthe, C., Aouizerats, B., Augros, C., Aumond, P., Auguste, F., Bechtold, P., Berthet, S., Bielli, S., Bosseur, F., Caumont, O., Cohard, J.-M., Colin, J., Couvreur, F., Cuxart, J., Delautier, G., Dauhut, T., Ducrocq, V., Filippi, J.-B., Gazen, D., Geoffroy, O., Gheusi, F., Honnert, R., Lafore, J.-P., Lebeaupin Brossier, C., Libois, Q., Lunet, T., Mari, C., Maric, T., Mascart, P., Mogé, M., Molinié, G., Nuissier, O., Pantillon, F., Peyrillé, P., Pergaud, J., Perraud, E., Pianezze, J., Redelsperger, J.-L., Ricard, D., Richard, E., Riette, S., Rodier, Q., Schoetter, R., Seyfried, L., Stein, J., Suhre, K., Taufour, M., Thouron, O., Turner, S., Verrelle, A., Vié, B., Visentin, F., Vionnet, V., and Wautelet, P.: Overview of the Meso-NH model version 5.4 and its applications, *Geoscientific Model Development*, 11, 1929–1969, <https://doi.org/10.5194/gmd-11-1929-2018>, <https://gmd.copernicus.org/articles/11/1929/2018/>, 2018.



- Legain, D., Bousquet, O., Douffet, T., Tzanos, D., Moulin, E., Barrie, J., and Renard, J.-B.: High-frequency boundary layer profiling with reusable radiosondes, *Atmos. Meas. Tech.*, 6, 2195–2205, 2013.
- 1075 Leuchner, M., Gubo, S., Schunk, C., Wastl, C., Kirchner, M., Menzel, A., and Plass-Dülmer, C.: Can positive matrix factorization help to understand patterns of organic trace gases at the continental Global Atmosphere Watch site Hohenpeissenberg?, *Atmospheric Chemistry and Physics*, 15, 1221–1236, <https://doi.org/10.5194/acp-15-1221-2015>, <https://acp.copernicus.org/articles/15/1221/2015/>, 2015.
- Lohou, F., Lathon, M., Canut, G., Bastin, S., Cheruy, F., Cohard, J.-M., Brut, A., Zouzoua, M., and Jomé, M.: The MOSAI project: Models and Observations for surface Atmosphere Interactions, Submitted to which journal, 2024.
- 1080 Lathon, M., Lohou, F., Pino, D., Couvreux, F., Pardyjak, E. R., Reuder, J., Vilà-Guerau de Arellano, J., Durand, P., Hartogensis, O., Legain, D., Augustin, P., Gioli, B., Lenschow, D. H., Faloon, I., Yagüe, C., Alexander, D. C., Angevine, W. M., Bargain, E., Barrié, J., Bazile, E., Bezombes, Y., Blay-Carreras, E., van de Boer, A., Boichard, J. L., Bourdon, A., Butet, A., Campistron, B., de Coster, O., Cuxart, J., Dabas, A., Darbieu, C., Deboudt, K., Delbarre, H., Derrien, S., Flament, P., Fourmentin, M., Garai, A., Gibert, F., Graf, A., Groebner, J., Guichard, F., Jiménez, M. A., Jonassen, M., van den Kroonenberg, A., Magliulo, V., Martin, S., Martinez, D., Mastroiello, L., Moene, 1085 A. F., Molinos, F., Moulin, E., Pietersen, H. P., Pignatelli, B., Pique, E., Román-Cascón, C., Rufin-Soler, C., Saïd, F., Sastre-Marugán, M., Seity, Y., Steeneveld, G. J., Toscano, P., Traullé, O., Tzanos, D., Wacker, S., Wildmann, N., and Zaldei, A.: The BLLAST field experiment: Boundary-Layer Late Afternoon and Sunset Turbulence, *Atmos. Phys. Chem.*, 14, 10931–10960, 2014.
- Lathon, M., Barnéoud, P., Gabella, O., Lohou, F., Derrien, S., Rondi, S., Chiriacco, M., Bastin, S., Dupont, J.-C., Haeffelin, M., Badosa, J., Pascal, N., and Montoux, N.: ELIFAN, an algorithm for the estimation of cloud cover from sky imagers, *Atmos. Meas. Tech.*, 12, 1090 5519–5534, 2019.
- Marenco, A.: Variations of CO and O₃ in the troposphere: evidence of O₃ photochemistry, *Atmos. Env.*, 20(5), 911–918, [https://doi.org/10.1016/0004-6981\(86\)90275-1](https://doi.org/10.1016/0004-6981(86)90275-1), 1986.
- Marenco, A., Gouget, H., Nédélec, P., Pagès, J., and Karcher, F.: Evidence of a long-term increase in tropospheric ozone from Pic du Midi data series - consequences - positive radiative forcing, *JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES*, 99, 16 617–16 632, 1095 <https://doi.org/10.1029/94JD00021>, 1994.
- Maruscak, N., Sonke, J., Fu, X., and Jiskra, M.: Tropospheric GOM at the Pic du Midi Observatory-Correcting Bias in Denuder Based Observations, *Environmental Science & Technology*, 51, 863–869, <https://doi.org/10.1021/acs.est.6b04999>, <https://doi.org/10.1021/acs.est.6b04999>, 2017.
- Mather, J. H. and Voyles, J. W.: The arm climate research facility: A review of structure and capabilities, *Bulletin of the American Meteorological Society*, 94, 377–392, <https://doi.org/10.1175/BAMS-D-11-00218.1>, 2013.
- Maury, N., Roberts, G. C., Couvreux, F., Verdu, T., Narvor, P., Lacroix, S., and Hattenberger, G.: Quantifying the mixing of trade-wind cumulus during the NEPHELAE-EUREC4A field campaign with remotely piloted aircraft, *Quarterly Journal of the Royal Meteorological Society*, pp. 1–22, <https://doi.org/https://doi.org/10.1002/qj.4430>, 2023.
- Milroy, C., Martucci, G., Lolli, S., Loac, S., Sauvage, L., Xueref-Remy, I., Lavrič, J. V., Ciais, P., Feist, D. G., Biavati, G., and O’Dowd, 1105 C. D.: An Assessment of Pseudo-Operational Ground-Based Light Detection and Ranging Sensors to Determine the Boundary-Layer Structure in the Coastal Atmosphere, *Advances in Meteorology*, 2012, 929 080, <https://doi.org/10.1155/2012/929080>, 2012.
- Necki, J., Schmidt, M., Rozanski, K., Zimnoch, M., Korus, A., Lasa, J., Graul, R., Levin, I., et al.: Six-year record of atmospheric carbon dioxide and methane at a high-altitude mountain site in Poland, *Tellus B: Chemical and Physical Meteorology*, 55, 94–104, <https://doi.org/10.3402/tellusb.v55i2.16763>, <https://doi.org/10.3402/tellusb.v55i2.16763>, 2003.



- 1110 Neubert, T., Rycroft, M., Farges, T., Blanc, E., Chanrion, O., Arnone, E., Odzimek, A., Arnold, N., Enell, C. F., Turunen, E., Bösinger, T., Mika, Á., Haldoupis, C., Steiner, R. J., van der Velde, O., Soula, S., Berg, P., Boberg, F., Thejll, P., Christiansen, B., Ignaccolo, M., Füllekrug, M., Verronen, P. T., Montanya, J., and Crosby, N.: Recent results from studies of electric discharges in the mesosphere, *Surveys in Geophysics*, 29, 71–137, <https://doi.org/10.1007/s10712-008-9043-1>, 2008.
- 1115 Nilsson, E., Lohou, F., Lothon, M., Pardyjak, E., Mahrt, L., and Darbieu, C.: Turbulence kinetic energy budget during the afternoon transition - Part 1: Observed surface TKE budget and boundary layer description for 10 intensive observation period days, *ATMOSPHERIC CHEMISTRY AND PHYSICS*, 16, 8849–8872, 2016a.
- Nilsson, E., Lothon, M., Lohou, F., Pardyjak, E., Hartogensis, O., and Darbieu, C.: Turbulence kinetic energy budget during the afternoon transition - Part 2: A simple TKE model, *ATMOSPHERIC CHEMISTRY AND PHYSICS*, 16, 8873–8898, 2016b.
- Ólafsson, H. and Bougeault, P.: The effect of rotation and surface friction on orographic drag, *Journal of the Atmospheric Sciences*, 54, 193–210, [https://doi.org/10.1175/1520-0469\(1997\)054<0193:TEORAS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054<0193:TEORAS>2.0.CO;2), 1997.
- 1120 Pandolfi, M., Cusack, M., Alastuey, A., and Querol, X.: Variability of aerosol optical properties in the Western Mediterranean Basin, *Atmospheric Chemistry and Physics*, 11, 8189–8203, <https://doi.org/10.5194/acp-11-8189-2011>, <https://acp.copernicus.org/articles/11/8189/2011/>, 2011.
- Pandolfi, M., Ripoll, A., Querol, X., and Alastuey, A.: Climatology of aerosol optical properties and black carbon mass absorption cross section at a remote high-altitude site in the western Mediterranean Basin, *Atmospheric Chemistry and Physics*, 14, 6443–6460, <https://doi.org/10.5194/acp-14-6443-2014>, <https://acp.copernicus.org/articles/14/6443/2014/>, 2014.
- Pappalardo, G., Nicolae, D., Makoto, A., Vassilis, A., Balis, D., Behrendt, A., Comeron, A., Gibert, F., Landulfo, E., McCormick, M. P., and Senff, C.: ACTRIS Aerosol, Clouds and Trace Gas Research Infrastructure, 28th International laser radar conference, Book Series EPJ Web of Conferences, 176, num. 09 004, doi:10.1051/epjconf/201817609 004, 2018.
- 1130 Philibert, A., Lothon, M., Campistron, B., Vial, A., Derrien, S., Amestoy, J., Meslin, Y., Canut, G., and Saïd, F.: CALOTRITON: a convective boundary layer height estimation algorithm from ultra-high-frequency (UHF) wind profiler data, *Atmos. Meas. Tech.*, 17, 1679–1701, 2024.
- Pietersen, H. P., de Arellano, J. V.-G., Augustin, P., van de Boer, A., de Coster, O., Delbarre, H., Durand, P., Fourmentin, M., Gioli, B., Hartogensis, O., Lohou, F., Lothon, M., Ouwersloot, H. G., Pino, D., and Reuder, J.: Study of a prototypical convective boundary layer observed during BLLAST: contributions by large-scale forcings, *ATMOSPHERIC CHEMISTRY AND PHYSICS*, 15, 4241–4257, 2015.
- 1135 Prados-Roman, C., Fernández, M., Gómez-Martín, L., Cuevas, E., Gil-Ojeda, M., Maruszczak, N., Puentedura, O., Sonke, J., and Saiz-Lopez, A.: Atmospheric formaldehyde at El Teide and Pic du Midi remote high-altitude sites, *Atmospheric Environment*, 234, 117618, <https://doi.org/10.1016/j.atmosenv.2020.117618>, 2020.
- RESIF: RESIF-RENAG French national Geodetic Network, <https://doi.org/10.15778/resif.rg>, 2017.
- 1140 Reuder, J., Båserud, L., Jonassen, M. O., Kral, S. T., and Müller, M.: Exploring the potential of the RPA system SUMO for multipurpose boundary-layer missions during the BLLAST campaign, *Atmos. Meas. Tech.*, 9, 2675–2688, 2016.
- Román-Cascón, C., Yagüe, C., Arrillaga, J. A., Lothon, M., Pardyjak, E. R., Lohou, F., Inclán, R. M., Sastre, M., Maqueda, G., Derrien, S., Meyerfeld, Y., Hang, C., Campargue-Rodríguez, P., and Turki, I.: Comparing mountain breezes and their impacts on CO₂ mixing ratios at three contrasting areas, *Atmospheric Research*, 221, 111–126, <https://doi.org/10.1016/j.atmosres.2019.01.019>, <https://doi.org/10.1016/j.atmosres.2019.01.019>, 2019.
- 1145 Román-Cascón, C., Yagüe, C., Mahrt, L., Sastre, M., Steeneveld, G.-J., Pardyjak, E., van de Boer, A., and Hartogensis, O.: Scaling the Decay of Turbulence Kinetic Energy in the Free-Convective Boundary Layer, *Atmos. Phys. Chem.*, 15, 9031–9047, 2015.



- Rotach, M. W., Wohlfahrt, G., Hansel, A., Reif, M., Wagner, J., and Gohm, A.: The World is Not Flat: Implications for the Global Carbon Balance, *Bulletin of the American Meteorological Society*, 95, 1021–1028, <https://doi.org/10.1175/BAMS-D-13-00109.1>, <https://journals.ametsoc.org/view/journals/bams/95/7/bams-d-13-00109.1.xml>, 2014.
- 1150 Roudier, T., Maiherbe, J.-M., Rozelot, J.-P., Mein, P., and Muller, R.: Five decades of solar research at the Pic du Midi Turret-Dome (1960–2010).i. Part I: Overview of instrumentation and observations, *J. History of Astronomy*, 24, 585–606, 2021.
- Saiz-Lopez, A., Sitkiewicz, S., Roca-Sanjuán, D., Oliva-Enrich, J., Dávalos, J., Notario, R., Jiskra, M., Xu, Y., Wang, F., Thackray, C., Sunderland, E., Jacob, D., Travnikov, O., Cuevas, C., Acuña, A., Rivero, D., Plane, J., Kinnison, D., and Sonke, J.: Photoreduction of gaseous oxidized mercury changes global atmospheric mercury speciation, transport and deposition, *Nature Communications*, 9, 4796, <https://doi.org/10.1038/s41467-018-07075-3>, 2018.
- 1155 Scorer, R. S.: Theory of airflow over mountains. Part IV: Separation of flow from the surface, *Quart. J. Roy. Meteorol. Soc.*, pp. 340–350, 1955.
- Seity, Y., Brousseau, P., Malardel, S., Hello, G., Bénard, P., Bouttier, F., Lac, C., and Masson, V.: The AROME-France Convective-Scale Operational Model, *Monthly Weather Review*, 139, 976–991, <https://doi.org/10.1175/2010MWR3425.1>, 2011.
- 1160 Smith, R. B.: On severe downslope winds, *J. Atmos. Sci.*, 42, 2597–2603, [https://doi.org/10.1175/1520-0469\(1985\)042<OSDW>3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1985)042<OSDW>3E2.0.CO;2), 1985.
- Soula, S., Defer, E., Füllekrug, M., van der Velde, O., Montanyà, J., Bousquet, O., Mlynarczyk, J., Coquillat, S., Pinty, J.-P., Rison, W., Krehbiel, P., Thomas, R., and Pedebay, S.: Dancing sprites above a Lightning Mapping Array—An analysis of the storm and flash/sprite developments, *J. Geophys. Res. Atmos.*, 120, 552–11,574, <https://doi.org/10.1002/2015JD023894>, 2015.
- 1165 Soula, S., Mlynarczyk, J., Füllekrug, M., Pineda, N., Georgis, J. F., van der Velde, O., Montanyà, J., and Fabró, F.: Dancing sprites: Detailed analysis of two case studies, *Journal of Geophysical Research*, 122, 3173–3192, <https://doi.org/10.1002/2016JD025548>, 2017.
- Steinacker, R.: Area-height distribution and its relation to the valley wind, *Beitr. Phys. Atmos.*, 57, 64–71, 1984.
- Stevens, B., Farrell, D., Hirsch, L., Jansen, F., Nuijens, L., Serikov, I., Brüggmann, B., Forde, M., Linne, H., Lonitz, K., and Prospero, J. M.: The Barbados cloud observatory: Anchoring investigations of clouds and circulation on the edge of the itcz, *Bulletin of the American Meteorological Society*, 97, 735–754, <https://doi.org/10.1175/BAMS-D-14-00247.1>, 2016.
- 1170 Suess, E., Aemisegger, F., Sonke, J., Sprenger, M., Wernli, H., and Winkel, L.: Marine versus Continental Sources of Iodine and Selenium in Rainfall at Two European High-Altitude Locations, *Env. Sci. Tech.*, 53, 1905–1917, <https://doi.org/10.1021/acs.est.8b05533>, <https://doi.org/10.1021/acs.est.8b05533>, 2019.
- 1175 Sun, J., Hermann, M., Yuan, Y., Birmili, W., Collaud Coen, M., Weinhold, K., Madueño, L., Poulain, L., Tuch, T., Ries, L., Sohmer, R., Couret, C., Frank, G., Brem, B. T., Gysel-Beer, M., Ma, N., and Wiedensohler, A.: Long-term trends of black carbon and particle number concentration in the lower free troposphere in Central Europe, *Environmental Sciences Europe*, 33, 47, <https://doi.org/10.1186/s12302-021-00488-w>, 2021.
- Tinorua, S., Denjean, C., Nabat, P., Bourriane, T., Pont, V., Gheusi, F., and Leclerc, E.: Two years measurements of black carbon properties at the high-altitude mountain site of Pic du Midi Observatory in the French Pyrenees, *Atmos. Chem. Phys. Discuss.*, 23, 2023.
- 1180 Tomicic, M., Soula, S., Defer, E., Prieur, S., Mlynarczyk, J., and Farges, T.: Dancing sprites above a Lightning Mapping Array—An analysis of the storm and flash/sprite developments, *J. Geophys. Res. Atmos.*, 126, D035 059, <https://doi.org/10.1029/2021JD035059>, 2021.
- Tsamalis, C., Ravetta, F., Gheusi, F., Delbarre, H., and Augustin, P.: Mixing of free-tropospheric air with the lowland boundary layer during anabatic transport to a high altitude station, *Atmos. Res.*, 143, 425–437, <https://doi.org/10.1016/j.atmosres.2014.03.011>, 2014.



- 1185 Tsikerdekis, A., Zanis, P., Steiner, A. L., Solmon, F., Amiridis, V., Marinou, E., Katragkou, E., Karacostas, T. S., and Foret, G.: Impact of dust size parameterizations on aerosol burden and radiative forcing in RegCM4, *Atmos. Chem. Phys.*, 17, 769–791, <https://doi.org/10.5194/acp-17-769-2017>, 2017.
- Whiteman, C. D.: Observations of thermally developed wind systems in mountainous terrain, vol. 45 of *Atmospheric processes in complex terrain, Meteorological Monographs*, American Meteorological Society, 1990.
- 1190 Whittlestone, S. and Zahorowski, W.: Baseline radon detectors for shipboard use: Development and deployment in the First Aerosol Characterization Experiment (ACE 1), *Journal of Geophysical Research: Atmospheres*, 103, 16 743–16 751, <https://doi.org/10.1029/98JD00687>, 1998.
- Yang, X., Jiskra, M., and Sonke, J.: Experimental rainwater divalent mercury speciation and photoreduction rates in the presence of halides and organic carbon, *Science of The Total Environment*, 697, 133821, <https://doi.org/10.1016/j.scitotenv.2019.133821>, 2019.
- 1195 Yver-Kwok, C., Laurent, O., Guemri, A., Philippon, C., Wastine, B., Rella, C., Vuillemin, C., Truong, F., Delmotte, M., Kazan, V., Darding, M., Lebegue, B., Kaiser, C., Xueref-Remy, I., and Ramonet, M.: Comprehensive laboratory and field testing of cavity ring-down spectroscopy analyzers measuring H₂O, CO₂, CH₄ and CO, *Atmos. Meas. Tech.*, 8, 3867–3892, <https://doi.org/10.5194/amt-8-3867-2015>, 2015.
- Zouzoua, M., Bastin, S., Chiriaco, M., Lohou, F., Lothon, M., Jomé, M., Mallet, C., and Barthes, L.: Using supervised neural networks to better evaluate surface turbulent heat fluxes in weather and climate numerical models: a demonstration study, Submitted to which journal, 2024.
- 1200