

Dear Editor,

*Referee #1's comments are in italic and our answers to all comments of reviewer #1 are embedded in red. Bold text are manuscript additions and ~~striketrough-text~~ are manuscript deletions.*

*This manuscripts deals with airborne-based remote sensing data to characterize NO<sub>2</sub> and HONO pumes from the Sheridan Fire. It uses auxiliary data also coming from airborne-based data, but also from satellite-based data. The exponentially modified gaussian (EMG) approach is used to extract the lifetimes and the emission rates from the NO<sub>2</sub> and HONO emissions. Simulations regarding different plume cases are used to understand the limitations of the EMGs. A new methodology based on this very same approach is proposed, which results in a more consistent fit to the plume line densities. Comparison to the daily mean emissions of HONO and NO<sub>2</sub> are also carried out. An important point of this manuscript is definitely the transfer of techniques developed in this work to geostationary space-based instruments such as TEMPO and GEMS, which can provide data with a very high temporal resolution.*

*I want to congratulate the authors for the enormous amount of work related to this manuscript. I also think the English is good, with almost no typos. Regarding the text, there are several comments on this document asking for more clarity. Some times it was difficult to keep up with the text, which made it difficult to read at some points. This also alludes to the science part, where there are some key concepts that were difficult for me to follow.*

*I consider this manuscript as an added-value to science that meets most quality standards. However, I believe that this work needs some corrections before publication, mostly for clarification. I have separated these corrections into 'major comments' and 'minor comments'.*

### **Major comments**

- *Tests have been done using an airborne instrument. However, TEMPO and GEMS are space-based instruments. These instruments appear in the abstract and conclusions, and sometimes in the rest of the test. Regarding their importance for the future, it would be suitable to briefly explain how features as a different spatial resolution would affect the methodologies explain in this manuscript.*

*There is not a substantial difference in the methodology if using space-based or airborne instruments. The methodology requires the retrieval of VCDs. Each instrument will have a different sampling resolution and thus different grid resolution of these final products. That is the only difference concerning the application of this methodology. They employ different procedures to retrieve VCDs, but an instrument comparison of VCD retrievals is beyond the scope of this paper.*

*We have added the following text to the "Analysis methods" section and the "Conclusions" section of the manuscript, separately, expanding upon how the methodologies would differ for the space-based instruments:*

***"...For use with an hourly space-based instrument like TEMPO, only the regridding resolution would differ in this procedure (0.02° for TEMPO). This lower resolution in grid size results in a lower resolution in line density, where the fine structures seen with the GCAS instrument will not be as resolved..."***

“...With the launch of ~~the Tropospheric Emissions: Monitoring of Pollution (TEMPO)~~ instrument in early April 2023, hourly daytime measurements of air pollutants are possible in the North American continent. **Even though the spatial resolution of TEMPO is coarser than GCAS, the loss of the fine structures in the sampled line densities is not critical to the techniques described in this paper. The Monte Carlo 1-D model output itself is smooth and lacks the noisy, fine features found in the GCAS line densities. This coarse representation may make finding the best model parameters easier with RMSE...**”


- *Other methodologies for the estimation of emission rate the NO<sub>2</sub> and HONO plumes can provide better results? Please, consider (for example) the IME method and the Cross Sectional Flux method. Maybe these methodology can suffer from incomplete plumes. However, due to the larger swath from space-based instruments, these instruments will be capable to capture entire plumes. As one of the most important points of this study is show the implementation of methodologies on airborne data to transfer them to satellite-based data in the future, it is important to deal with these questions.*

We have included a brief discussion of a few other emission estimations methods, such as the IME and Cross-Sectional Flux methods. As this paper only evaluates the EMG method and improves upon it, it is beyond the scope of this paper to evaluate two more methodologies. The primary point of this paper was to respond to two papers using TROPOMI and the EMG method to make emissions estimates. The discussion in the manuscript is presented below:


**“The EMG is fit to a plume and the enhancement factor from the fitting procedure is directly linked to the emissions. Other methods are also used to estimate emissions from satellites, such as the integrated mass enhancement (IME) method, which multiplies the wind speed by the integrated vertical column densities (VCDs), the cross-sectional flux method (CFM), which estimates emissions by averaging the flux through multiple cross sections perpendicular to the plume direction, and Chemical Transport Models (CTMs), which can predict emissions more accurately but are computationally expensive. While these other techniques are proven to be useful, this paper focuses on dissecting the EMG approach.** Forming conclusions using daily observations of fires, while a good starting point, does not capture the diurnal variability that fires exhibit (Wiggins et al., 2020).”

- *L375-383: A fit with higher R<sup>2</sup> guarantees the most accurate values of lifetime and emission? I think it has more to do with the shape of the line density. If it has a gaussian shape it would be very suitable for the fit, but if not, then you can only try to make the best you can. What is then the meaning of the values with the maximum R<sup>2</sup>? Please, consider to develop more this concept.* Upon further review of the use of R<sup>2</sup> and other reviewer feedback, we have changed the evaluation metric to be root mean square error (RMSE). This evaluation metric is optimal for Gaussian errors (Hodson, 2002).


The new Figure 6, Figure 7, and Figure D1 are below. Note: the best model runs did not change when using RMSE as the evaluation metric:



“Figure 6: The RMSE value between 100 PECANS models with varied total emission rate and lifetime model input parameters and the observed GCAS HONO VCD line density for (a) track 12 and (b) track 14. The emission rate is plotted by multiplying the total emission rate by the normalized FRP at the time of each track. A filled contour map creates a 2D interpolation of the randomly sampled models (circles/dots). (c) and (d) Similar to (a) and (b), but for GCAS NO<sub>2</sub> VCD.”



“Figure 7: Daily mean emissions of (a) HONO and (b) NO<sub>2</sub> from the Monte Carlo diurnal 1-D model method (MC diurnal 1-D) using the  $R^2 = 0.9$  contour lowest RMSE model for each track to define the minimum and maximum emissions, and the following biomass burning emission inventories: FINNv2.5 SAPRC, FINNv2.5 MOZART, GFED4s, FINNv2.5, QFEDv2.5, and GFASv1.2.”



“Figure D1: Observed GCAS VCD line densities (orange) and best Monte Carlo diurnal 1-D model line densities (blue) for (a) HONO track 12, (b) HONO track 14, (c)  $\text{NO}_2$  track 12, and (d)  $\text{NO}_2$  track 14. Model lifetimes and emission rates are reported in the subfigures.”

The following text in the manuscript has been updated:

“...The ~~coefficient of determination~~ **root mean square error,  $R^2$  (RMSE)**, between the model and the satellite **GCAS** observations for **tracks 12 and 14** is ~~are~~ shown in Fig. 6 for HONO and  $\text{NO}_2$ . The absolute ~~maximum  $R^2$~~  **minimum RMSE** provides an indication of the most likely lifetime and emissions rate combination(s)...”

“...In Fig. 6, **RMSE is approximately constant in a combination of lifetime and emission rate values that follow an inverse relationship**~~the emission rate has an inverse relationship with lifetime~~, which is expected since general mass balance constitutes that concentration is proportional to emissions and inversely proportional to loss rate. In Figs. 6a and 6b, the ~~largest  $R^2$  (above 0.9)~~ **smallest RMSE** between the HONO observed and model derived line densities occur with a lifetime between 15 and 27 minutes and an emissions rate of 0.6 to  ~~$1.3575 \times 10^{25}$~~   **$1.19 \times 10^{25}$**   $\text{molec s}^{-1}$ . In Figs. 6c and 6d,  $\text{NO}_2$  has a larger viable range of lifetimes and total emissions rates, which range between ~~15~~ **20** and 45 minutes and 2.0 to  ~~$5.25 \times 10^{25}$~~   **$4.15 \times 10^{25}$**   $\text{molec s}^{-1}$ , respectively. Track 18 was excluded from this analysis as measurements occurred after the Sheridan Fire had ceased its high emissions for the day. The model runs with the ~~largest  $R^2$~~  **smallest RMSE** values are shown in Fig. D1...”

“...As shown in Fig. 7a, FINNv2.5 HONO emissions for the SAPRC and MOZART mechanisms underestimate the ~~minimum and maximum~~ **track 12 and track 14** daily mean emission rate of our method (MC diurnal 1-D) by a factor of ~~5~~ **9.4** for the ~~minimum~~ **track 12** and a factor of ~~10~~ **7.2** for the ~~maximum~~ **track 14**...”

Response Reference:

Hodson, T. O.: Root-mean-square error (RMSE) or mean absolute error (MAE): when to use them or not, *Geosci. Model Dev.*, 15, 5481–5487, <https://doi.org/10.5194/gmd-15-5481-2022>, 2022.

## Minor comments

### 1. General comments

- a. *Please, check the acronym definitions and once the acronym is defined, used it. There are several cases in the text that are lately defined or where the acronym is not used.*

We have corrected our acronym usage in some instances. However, we believe it is useful to define an acronym that is commonly used in an area of research. This makes searching through text easier for those readers trying to find key words.

- b. *Please, make sure that the references to websites are correctly cited.*

All websites in the main text of the manuscript have been replaced with citations to the software user manuals.

- c. *Please, avoid unnecessary repetitions along the text.*

We have read through the text and have removed unnecessary repetitions. However, some repetitions have remained to benefit the reader.

### 2. Abstract

- a. *L22: Are these magnitudes really comparable? See comments later in the text.*

These quantities are comparable. We have clarified in the abstract what we are comparing by adding the following text:

“We assess the validity of a range of emission rate and lifetime combinations for both HONO and NO<sub>2</sub> as the fire evolves by comparing the resulting line density predictions to the **line density** observations.”

### 3. Introduction

- a. *L35: Is this study only important for the US? If not, please consider to include a more global view in this sentence.*

This study is not only important for the US. To express that this study has global implications, we have amended the manuscript with the following text:

“As the intensity of fires and burned area from fires ~~are predicted to increase~~ in the United States **and globally have had increasing trends over the past few decades and are predicted to increase** in the future (Barbero et al., 2015; Burton et al., 2024; Cunningham et al., 2024; Dennison et al., 2014), it is important to understand how wildfire emissions change with fire properties and how these emissions impact local, regional, and global atmospheric composition.”

The following references were added to the References section:

**“Barbero, R., Abatzoglou, J. T., Larkin, N. K., Kolden, C. A., and Stocks, B.: Climate change presents increased potential for very large fires in the contiguous United States, *Int. J. Wildland Fire*, 24, 892, <https://doi.org/10.1071/WF15083>, 2015.**

**Burton, C., Lampe, S., Kelley, D. I., Thiery, W., Hantson, S., Christidis, N., Gudmundsson, L., Forrest, M., Burke, E., Chang, J., Huang, H., Ito, A., Kou-Giesbrecht, S., Lasslop, G., Li, W., Nieradzik, L., Li, F., Chen, Y., Randerson, J., Reyer, C. P. O., and Mengel, M.: Global burned area increasingly explained by climate change, *Nat. Clim. Chang.*, 14, 1186–1192, <https://doi.org/10.1038/s41558-024-02140-w>, 2024.**

**Cunningham, C. X., Williamson, G. J., and Bowman, D. M. J. S.: Increasing frequency and intensity of the most extreme wildfires on Earth, *Nat Ecol Evol*, 8, 1420–1425, <https://doi.org/10.1038/s41559-024-02452-2>, 2024.**

**Dennison, P. E., Brewer, S. C., Arnold, J. D., and Moritz, M. A.: Large wildfire trends in the western United States, 1984–2011, *Geophys. Res. Lett.*, 41, 2928–2933, <https://doi.org/10.1002/2014GL059576>, 2014.”**

- b. *L39: Regarding GFED and other inventories, the burned area is obtained from satellite-based measurements. Typically, a bottom-up approach is associated with ground-based, local, or detailed data collection at specific sites or sources (such as factories, vehicles, etc.). However, satellite data is more commonly associated with a top-down approach, where broad, large-scale observations (e.g., of emissions, land cover, or atmospheric concentrations) are made, and then models or algorithms are used to estimate emissions, trends, or impacts at finer scales. Please, consider checking the terminology of these approaches.*

We have checked the terminology of top-down and bottom-up approaches as applied to the fire emissions inventories. Multiple sources cited in our manuscript use the same terminology (see Darmenov and da Silva, 2015; Wiedinmyer et al., 2011; and Wiedinmyer et al., 2023). Further, the bottom-up approach is defined by using fuel accounting to determine emissions, while the top-down approach is defined by using energy to determine emissions (see Wiggins et al., 2021).

Response Reference:

Wiggins, E. B., Anderson, B. E., Brown, M. D., Campuzano-Jost, P., Chen, G., Crawford, J., Crosbie, E. C., Dibb, J., DiGangi, J. P., Diskin, G. S., Fenn, M., Gallo, F., Gargulinski, E. M., Guo, H., Hair, J. W., Halliday, H. S., Ichoku, C., Jimenez, J. L., Jordan, C. E., Katich, J. M., Nowak, J. B., Perring, A. E., Robinson, C. E., Sanchez, K. J., Schueneman, M., Schwarz, J. P., Shingler, T. J., Shook, M. A., Soja, A. J., Stockwell, C. E., Thornhill, K. L., Travis, K. R., Warneke, C., Winstead, E. L., Ziemba, L. D., and Moore, R. H.: Reconciling Assumptions in Bottom-Up and Top-Down Approaches for Estimating Aerosol Emission Rates From Wildland Fires Using Observations From FIREX-AQ, *JGR Atmospheres*, 126, e2021JD035692, <https://doi.org/10.1029/2021JD035692>, 2021.

- c. *L40: ‘as a proxy for amount of material burned’... Please, try to avoid repetition for a more compact writing.*

We have removed the repetition and amended the text as follows:

~~“There are two distinct approaches for estimating global fire emissions within biomass burning emission inventories: (1) a bottom-up approach that uses burned area as a proxy for amount of material burned and (2) a top-down approach that uses fire radiative power (FRP) as a proxy proxies for amount of material burned.”~~

- d. L55-56: Please, include references related to this statement.

We have amended the text as follows to include references:

~~“Recent advances in remote sensing and in situ observations have enabled improved assessments of wildfire emissions of reactive nitrogen (Gkatzelis et al., 2024; Lindaas et al., 2021; Theys et al., 2020).”~~

The following references have been added to the References section (Theys already included):

~~“Gkatzelis, G. I., Coggon, M. M., Stockwell, C. E., Hornbrook, R. S., Allen, H., Apel, E. C., Bela, M. M., Blake, D. R., Bourgeois, I., Brown, S. S., Campuzano-Jost, P., St. Clair, J. M., Crawford, J. H., Crouse, J. D., Day, D. A., DiGangi, J. P., Diskin, G. S., Fried, A., Gilman, J. B., Guo, H., Hair, J. W., Halliday, H. S., Hanisco, T. F., Hannun, R., Hills, A., Huey, L. G., Jimenez, J. L., Katich, J. M., Lamplugh, A., Lee, Y. R., Liao, J., Lindaas, J., McKeen, S. A., Mikoviny, T., Nault, B. A., Neuman, J. A., Nowak, J. B., Pagonis, D., Peischl, J., Perring, A. E., Piel, F., Rickly, P. S., Robinson, M. A., Rollins, A. W., Ryerson, T. B., Schueneman, M. K., Schwantes, R. H., Schwarz, J. P., Sekimoto, K., Selimovic, V., Shingler, T., Tanner, D. J., Tomsche, L., Vasquez, K. T., Veres, P. R., Washenfelder, R., Weibring, P., Wennberg, P. O., Wisthaler, A., Wolfe, G. M., Womack, C. C., Xu, L., Ball, K., Yokelson, R. J., and Warneke, C.: Parameterizations of US wildfire and prescribed fire emission ratios and emission factors based on FIREX-AQ aircraft measurements, Atmos. Chem. Phys., 24, 929–956, <https://doi.org/10.5194/acp-24-929-2024>, 2024.~~

~~Lindaas, J., Pollack, I. B., Garofalo, L. A., Pothier, M. A., Farmer, D. K., Kreidenweis, S. M., Campos, T. L., Flocke, F., Weinheimer, A. J., Montzka, D. D., Tyndall, G. S., Palm, B. B., Peng, Q., Thornton, J. A., Permar, W., Wielgasz, C., Hu, L., Ottmar, R. D., Restaino, J. C., Hudak, A. T., Ku, I., Zhou, Y., Sive, B. C., Sullivan, A., Collett, J. L., and Fischer, E. V.: Emissions of Reactive Nitrogen From Western U.S. Wildfires During Summer 2018, Geophys Res Atmos, 126, <https://doi.org/10.1029/2020JD032657>, 2021.”~~

- e. L78-79: Please, consider rewording this sentence for more clarity.

We have reworded this sentence as follows for clarity:

~~“We use a simple one-dimensional (1-D) horizontal model to evaluate the assumptions within and the accuracy of the EMG approach on wildfire smoke plumes. We evaluate the assumptions inherent to the EMG approach to assess the EMG’s utility at deriving emission rates and lifetimes of wildfire smoke plumes. We use a simple one-dimensional (1-D) horizontal model to perform this evaluation.”~~

#### 4. Data and methods

#### 5. FIREX-AQ

- a. *The section name refers to the FIREX-AQ campaign. However, the GCAS technical features, the AMF concept, the retrieval methodology and the a priori data are also explained. Please, consider to use different sections to describe the campaign and the methodology-related topics.*

We have rearranged the sections to include a new section, 2.2 GCAS Methodology. The two paragraphs concerning the AMF concept, the retrieval methodology, and the a priori data are now under this section.

- b. *L90: Is the second also measuring 'absolute nadir radiance'? Please, clarify.*

Yes, it is. We have amended the text as follows:

“The GCAS instrument is composed of two push-broom spectrometers: the first records the spectrum as absolute nadir radiance in the ultraviolet to visible (UV-Vis), from 300 to 490 nm, and the second records the spectrum **as absolute nadir radiance** in the visible to near-infrared (Vis-NIR) from 480 to 900 nm (Kowalewski and Janz, 2014).”

- c. *L93: A reference here for these instruments would be suitable.*

References have been added to the text as follows:

“The GCAS instrument shares similar design specifications with the Tropospheric Monitoring Instrument (TROPOMI; **Veefkind et al., 2012**) and Tropospheric Emissions: Monitoring of Pollution (TEMPO; **Chance et al., 2013**) instruments.”

The references below have been added to the References section:

“**Chance, K., Liu, X., Suleiman, R. M., Flittner, D. E., Al-Saadi, J., and Janz, S. J.: Tropospheric emissions: monitoring of pollution (TEMPO), SPIE Optical Engineering + Applications, San Diego, California, United States, 88660D, <https://doi.org/10.1117/12.2024479>, 2013.**”

**Veefkind, J. P., Aben, I., McMullan, K., Förster, H., De Vries, J., Otter, G., Claas, J., Eskes, H. J., De Haan, J. F., Kleipool, Q., Van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sensing of Environment*, **120**, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.”**

- d. *L93: Please, explain briefly in which aspects these instruments are similar.*

We have added the following sentence to explain the instrumental similarity:

“**TROPOMI operates in the 310 to 405 nm and 405 to 500 nm bands with a spectral resolution of 0.55 nm and in the 675 to 725 nm and 725 to 775 nm bands with a spectral resolution of 0.5 nm (Veefkind et al., 2012). TEMPO operates in the 290 to 490 nm and 540 to 740 nm bands with a spectral resolution of 0.57 nm (Zoogman et al., 2017).**”

The following references were added to the References section (Veefkind was added in response to the previous comment):

**“Zoogman, P., Liu, X., Suleiman, R. M., Pennington, W. F., Flittner, D. E., Al-Saadi, J. A., Hilton, B. B., Nicks, D. K., Newchurch, M. J., Carr, J. L., Janz, S. J., Andraschko, M. R., Arola, A., Baker, B. D., Canova, B. P., Chan Miller, C., Cohen, R. C., Davis, J. E., Dussault, M. E., Edwards, D. P., Fishman, J., Ghulam, A., González Abad, G., Grutter, M., Herman, J. R., Houck, J., Jacob, D. J., Joiner, J., Kerridge, B. J., Kim, J., Krotkov, N. A., Lamsal, L., Li, C., Lindfors, A., Martin, R. V., McElroy, C. T., McLinden, C., Natraj, V., Neil, D. O., Nowlan, C. R., O’Sullivan, E. J., Palmer, P. I., Pierce, R. B., Pippin, M. R., Saiz-Lopez, A., Spurr, R. J. D., Szykman, J. J., Torres, O., Veefkind, J. P., Veihelmann, B., Wang, H., Wang, J., and Chance, K.: Tropospheric emissions: Monitoring of pollution (TEMPO), *Journal of Quantitative Spectroscopy and Radiative Transfer*, 186, 17–39, <https://doi.org/10.1016/j.jqsrt.2016.05.008>, 2017.”**

- e. *L98: The GCAS instrument measures radiance, from which the differential absorption spectra from a gas is deduced. Therefore, the different absorption spectra is not measured, but deduced. Then, I would consider removing the term 'measured'.*

We have removed the term ‘measured’ as demonstrated in the following amendment:

**“By fitting the differential cross-sections of trace gases to the ~~measured~~ differential absorption spectra, the trace gas concentrations along the light path can be determined via the Beer-Lambert law.”**

- f. *L100: A figure showing an example of transmissivity spectra from NO<sub>2</sub> and HONO would be illustrative to justify the selection of these fitting windows. Another option could be cite other works that used them.*

We have cited other works that illustrate either the fitting window’s use in previous work or works that demonstrate the absorptivity of the compound in the UV-Vis band. Our amended text is as follows:

**“The fitting windows for NO<sub>2</sub> and HONO are 425 to 460 nm and 345 to 390 nm, respectively. The NO<sub>2</sub> fitting window is within the standard TROPOMI NO<sub>2</sub> product window of 405 to 465 nm (van Geffen et al., 2022). HONO absorbs in the UV-Vis spectrum at wavelengths 342 nm, 354 nm, and 368 nm (Stutz et al., 2000).”**

The new references are added to the References section:

**“van Geffen, J. H. G. M., Eskes, H. J., Boersma, K. F., and Veefkind, J. P.: TROPOMI ATBD of the total and tropospheric NO<sub>2</sub> data products, 2022.**

**Stutz, J., Kim, E. S., Platt, U., Bruno, P., Perrino, C., and Febo, A.: UV-visible absorption cross sections of nitrous acid, *J. Geophys. Res.*, 105, 14585–14592, <https://doi.org/10.1029/2000JD900003>, 2000.”**

- g. *L105: AMF is not only dependent of the wavelength and altitude (first sentence), but also dependent of the solar zenith angle and the viewing zenith angle, among others. You refer to these parameters by mentioning the observation geometry (second sentence).*

However, consider rewording these sentences to build a more compact sentence where the dependencies of the AMF are not split in 2 sentences. It is not wrong, but I think this correction can provide more consistency in the AMF description.

We have rewritten these sentences about AMF as follows:

“An AMF is a ~~wavelength–and–altitude–dependent~~ quantity representing the effect that the light’s path has on retrieval. An AMF depends on **wavelength, altitude**, observation geometry, surface reflectivity, a-priori vertical profiles, aerosols, and other factors that affect the measurement sensitivity.”

- h. L108: Please, include a reference for VLIDORT.

The reference for VLIDORT has now been included in the text:

“In this study, AMFs are calculated using the vector linearized discrete ordinate radiative transfer code (VLIDORT), **version 7.2 (Spurr, 2006).**”

The full reference is below:



“**Spurr, R. J. D.: VLIDORT: A linearized pseudo-spherical vector discrete ordinate radiative transfer code for forward model and retrieval studies in multilayer multiple scattering media, Journal of Quantitative Spectroscopy and Radiative Transfer, 102, 316–342, <https://doi.org/10.1016/j.jqsrt.2006.05.005>, 2006.**”

- i. L132: Which is the range of the plume altitude? Is there such a big difference that can have an impact on the results or the selected value is well-suited for this purpose? For example, at different altitude values from the same plume, is there a significant difference in the pressure values? Please, justify the assumption taken regarding the altitude.



The range of the plume’s altitude is from 4 km to 5.25 km, equivalent to 630 mb to 544 mb. The selection of the plume’s altitude/pressure level impacts the wind speed taken from ERA5. From 550 to 600 mb, there is a shift in wind as depicted below on the first and second images. From 600 mb to 650 mb, there is no shift in the wind direction as depicted below in the second and third images. However, adding the GCAS plume data, we see that the selection of 588 mb, where the DC-8 sampled the plume, is appropriate, as the plume direction follows the wind vectors, in the fourth image. I have clarified in the text that the DC-8 sampling altitude of the plume determined the pressure level chosen:

“Altitude was converted to a pressure level by recording the DC-8 aircraft data at a time when the DC-8 was flying through the smoke plume, ~~which resulted in~~. **The DC-8 plane’s altitude and the plume’s altitude were approximated to be** at a pressure level of 588 mb.”


ERA5 Winds at 2019-08-16 22 UTC and 550 mb



ERA5 Winds at 2019-08-16 22 UTC and 600 mb



ERA5 Winds at 2019-08-16 22 UTC and 650 mb



## 6. ERA5

- a. L138: Please, consider to show the pressure ranges presenting first the lower values and then the higher values.

We have rewritten this sentence to reorder the pressure ranges as follows:

“We collected zonal and meridional wind speeds at pressures every 25 mb from 1000 mb to 750 mb and every 50 mb from 750 mb to 400 mb and every 25 mb from 750 mb to 1000 mb.”

## 7. GOES FRP

- a. L145: When reading this sentence, it seems that the main goal from these satellites are to locate fires. Please, consider rewording.

The satellites have multiple goals and purposes for their operation. Specifically relating to fires and of interest to this manuscript, these satellites are used to locate fires and retrieve fire characteristics, which was stated in the main text. The sentence structure remains the same, but we now emphasize their retrieval of the fire properties:

“On both satellites, the Advanced Baseline Imager (ABI) uses visible and infrared spectral bands to locate fires and ~~provide~~**retrieve** fire characteristics.”

- b. L149: *What is GeoMAC? Please, provide some clarification or at least a reference.*

GeoMAC is an internet-based mapping tool that has been collecting and storing data on wildfire perimeters in the contiguous 48 states of the United States and Alaska since 2000. We have expanded on the main text to clarify what GeoMAC is:

“A diurnal profile of the Sheridan Fire’s sum-FRP, a sum of all fire pixel FRPs associated with the Sheridan Fire, was generated by selecting GOES data within 4 km of the final fire perimeter from the Geospatial Multi-Agency Coordination (GeoMAC). **GeoMAC is an internet-based mapping tool that stores wildfire perimeters in the contiguous 48 states of the United States and Alaska since 2000.**”

## 8. Emission inventories

- a. L159: *In previous text, the angular degree unit was deg. Please, use just one of these units for consistency.*

We thank the author for catching this inconsistency. We have selected the unit to be ° throughout the text.

- b. L160: *This sentences is something misleading. Then, this model can provide emission data for NO and HONO or only for NO? Regarding the way it is written, it seems that it says that HONO is a NO<sub>x</sub>, but it is not.*

We agree that the way the text is currently written, there is some ambiguity in whether HONO is part of NO<sub>x</sub>. We have amended the text as follows:

“The spatial resolution is 0.25° × 0.25° and provides emission data for NO<sub>x</sub> (as NO), ~~but not~~ **Emissions data is not provided for NO<sub>2</sub> or HONO.**”

- c. L165: *Which are the E units? Please, clarify.*

We have added units for all variables, as demonstrated by the additions below:

“...where *i* is a specific compound, *E* is the emissions (**g**), *A* is the area burned (**m<sup>2</sup>**), *B* is the amount of biomass (**kg m<sup>-2</sup>**), *FB* is the fraction of biomass burned (**unitless**), and *EF* is the emission factor with units of mass of *i* per mass biomass burned (**g kg<sup>-1</sup>**).”

- d. L170-173: *Please, consider a simpler description.*

We have rephrased these couple of sentences to simplify the description:

“**FINNv2.5 can also derive other emission products from total Total non-methane organic gases (NMOG) in FINNv2.5 can also be mapped to species found in using** three commonly used chemical mechanisms: Statewide Air Pollution Research Center Mechanism (SAPRC99; **Carter, 1999**), Model for Ozone and Related chemical Tracers

(MOZART; Emmons et al., 2020), and Goddard Earth Observing System with Chemistry (GOES-Chem; Bey et al., 2001). FINNv2.5 can derive HONO from SAPRC and MOZART, but not from GOES-Chem. NMOG can be mapped to HONO for use in SAPRC and MOZART; there is no map to HONO for GOES-Chem.”

- e. L185: Which are the data sources from GFASv1.2? All the data sources from the previous inventories were described.

We have amended the text to include GFASv1.2 data sources as follows:

“GFASv1.2 is also an FRP-derived emissions inventory using the MOD14 and MYD14 products from the MODIS instruments on the Terra and Aqua satellites, respectively.”

## 9. Analysis methods

### 10. Calculation of HONO and NO<sub>2</sub> line densities

- a. L196: This can work for gaussian plume shapes as the ones you show in this study. However, if the direction of the plume is extracted from the plume shape, it will not always align with the wind direction. Please, consider this for the discussion.

We have added the following discussion point to the main text, as shown below:

“Subsequently, we rotated the grids by the angle of the ~~time-averaged~~ NO<sub>2</sub> VCD plume **averaged over time**, ~~representing~~ **acting as a proxy of** the mean wind direction (Fig. A1). **In general, the plume direction may not always align with the wind direction. We applied this rotation methodology to the Sheridan Fire because of its near-ideal plume characteristics, a linear plume shape.**”

- b. L196: Time averaged VCD plume? Please, clarify.

We have rewritten this phrase as follows:

“Subsequently, we rotated the grids by the angle of the ~~time-averaged~~ NO<sub>2</sub> VCD plume **averaged over time,...**”

- c. Figure 1: 2 of these panels are already shown in Figure 2. You could consider remove them from here or from Figure 1. If you keep the panels from this figure, you can allude to Figure 1 to refer to the track 14 results.

We believe it is kinder to the reader to provide a repetition of panels (b) and (e) for two reasons. First, Figure 1 clearly delineates the steps made to produce the line densities in panels (c) and (f). Without panels (b) and (e), a casual reader may think the integration occurred over panels (a) and (d) from north to south. Second, the side-by-side comparisons of tracks 12, 14, and 18 in Figure 2 make it easier for the reader to compare the VCD magnitudes all in one place.

- d. Figure 1: Please, consider a more compact caption.

The Figure 1 caption has been revised to the following:

“Figure 1: Image of (a) gridded and (b) rotated track 14 GCAS HONO VCD. (c) HONO line density from track 14. ~~Image of (d)-(f) gridded and (e) rotated track 14 GCAS-NO<sub>2</sub>-VCD. (f) NO<sub>2</sub> line density from track 14~~ **Similar to (a)-(c), but for NO<sub>2</sub>.**”

e. L206: trackS

There is already the letter “s” in “tracks” to denote the multiple tracks (12, 14, and 18).

f. Appendix A: In my opinion, it is very difficult to understand this. Please, consider rewording to solve this.

We have added some background text to Appendix A to help prime the reader for Fig. A1. This text is shown in response to the reviewer’s next comment. We have amended the Fig. A1 caption as follows:

“Figure A1: Time-averaged NO<sub>2</sub> VCD plume **averaged over GCAS sampling time, where plume pixels were more than 3 times greater than the NO<sub>2</sub> upwind background with a plume mask applied.** The black line is a **weighted** linear fit of the plume pixels, where the standard deviation at each location was the difference between the maximum NO<sub>2</sub> VCD and the location’s NO<sub>2</sub> VCD where the plume edges have the higher weight. The angle of the line is displayed.”

g. Appendix A: Please, consider to add some text in this appendix for clarification.

We have added the following text to clarify the concept depicted in Fig. A1:

“To calculate the line densities of a sampled smoke plume, the smoke plume needs to be rotated such that the plume axis is parallel to the x-axis. Fig. A1 below summarizes our methodology of determining the plume axis and its rotation angle for all the Sheridan Fire smoke plumes. First, each data point of the regridded NO<sub>2</sub> VCDs from all GCAS-sampled tracks (tracks 10-14 and 17-20) are summed and then divided by the number of valid cells (non-NaN cells), producing a temporal average of the plume shape, since the averaging occurs over the different timestamps of the sampled tracks. Second, we calculate the background NO<sub>2</sub> VCD by averaging all cells upwind of the fire. Third, we create a plume mask, where only NO<sub>2</sub> VCDs that are greater than three times the background NO<sub>2</sub> VCD remain. These are the colored pixels in Fig. A1. Finally, we perform a linear regression on the plume NO<sub>2</sub> VCD, weighting the regression with the difference between the NO<sub>2</sub> VCDs and the maximum NO<sub>2</sub> VCD plume pixel. This weighting prioritizes the plume’s edges, which define the plume shape. The angle of this linear fit becomes the plume rotation angle we apply to all the sampled plume tracks.”

## 11. Exponentially Modified Gaussian (EMG)

a. L217: NO<sub>x</sub> = NO + NO<sub>2</sub>. Then, NO<sub>2</sub> repeated?

That is correct. NO<sub>x</sub> and NO<sub>2</sub> are different quantities. Modelers are interested in quantifying NO<sub>x</sub> emissions for their models, but satellites can only retrieve NO<sub>2</sub>. The Griffin et al. (2021) paper communicated their results for NO<sub>2</sub> solely and for NO<sub>x</sub>, the sum of NO<sub>2</sub> and NO.

## 12. Results and discussion

### 13. HONO and NO<sub>2</sub> plume structure in the Sheridan Fire

a. Why there is a better resolution for NO<sub>2</sub> maps? Please, clarify.

There is a better resolution for NO<sub>2</sub> maps because the signal-to-noise ratio of HONO is lower than that of NO<sub>2</sub>. The maximum NO<sub>2</sub> is about four times greater than maximum

HONO. However, HONO and NO<sub>2</sub> have the same average minimum sensitivities of  $0.7 \times 10^{15}$  molec cm<sup>-2</sup>.

We amend our text to explain this concept after Fig. 1 is displayed, which is the first time the resolution difference can be noticed:

**“The resolution of the HONO image is lower than the NO<sub>2</sub> image because the magnitude of NO<sub>2</sub> VCD is approximately four times greater than HONO. This results in a higher signal-to-noise ratio for NO<sub>2</sub>.”**

- b. L264-265: ‘... in all three overpasses, HONO and NO<sub>2</sub> share local maxima, plume edges, and plume shape’. Please, consider a discussion about this result. What does this mean? We have added a small discussion about this result as follows:

**“Overall, in all three overpasses, HONO and NO<sub>2</sub> share local maxima, plume edges, and plume shape. This means that both chemicals shared similar emission characteristics, were exposed to a consistent dispersal pattern, and traveled together under the same atmospheric conditions.”**

#### **14. EMG emission rates and lifetimes from the Sheridan Fire**

- a. L274: I would avoid to use the term 'step function' when you are referring to the extracted line densities as they are not extracted according to a known function. Please, consider to use other terms as 'drop sharply' or similar.

We agree that the use of the term “step function” is not appropriate as applied in this paragraph. We have amended the text as follows:

**“In Fig. 3e, track 12 appears to have a ~~step function~~ decrease sharply in integrated NO<sub>2</sub> VCD around 25 km away from the fire center...”**

**“HONO appears to have the ~~step function~~ sharp declines in integrated HONO VCD that NO<sub>2</sub> demonstrates...”**

- b. L291-293: This sentence is somewhat confusing in expressing the timeline of events. Please, consider to reword this.

We have rewritten the sentence as follows:

**“Before analyzing the track 18 line densities, it is crucial to note that ~~In Fig. 3a, we see that by the time the ER-2 sampled the Sheridan Fire for track 18, sum FRP at track 18 is nearly at~~ back to the background value before the Sheridan Fire grew around 21 UTC.”**

- c. L294: You could avoid to refer to Fig. 3f if you are already alluded to 'track 14'.

We agree. We have removed referring to Fig. 3f as shown below:

**“The NO<sub>2</sub> line density maximum for track 18 in Fig. 3g is nearly four times smaller than that of track 14 ~~in Fig. 3f.~~”**

- d. L301: If you show the ratio of emission rates, you should first calculate them individually. Please, consider to include this calculation in the text (or if not, in an appendix).

The emission rates come from the EMG fits like those shown in Fig. 3 and described in Eq. 5,  $E_{EMG}$ . We have included the calculation in the text as follows:

“...we can take the ratio of the emissions rates of HONO and NO<sub>2</sub> ( $E_{EMG,HONO}/E_{EMG,NO_2}$ ) **from all flight tracks** to explore how reactive nitrogen in wildfire smoke is partitioned and processed.”

- e. L301: *What do you mean with 'similar sampling biases'? Please, clarify.*

We mean to say that we acknowledge that HONO and NO<sub>2</sub> have the same sampling biases as described in lines 204-207. The sweeping bowtie flight pattern will produce the same incomplete and asymmetric samplings in HONO as in NO<sub>2</sub> for every flight track, not just the three tracks mentioned earlier. We have amended the text as follows to make this connection for the reader:

~~“Assuming~~**Acknowledging** that HONO and NO<sub>2</sub> have ~~similar~~**the same** sampling biases, **meaning that each GCAS sample of the plume shares the same sampling orientation of the sweeping bowtie**, we can take the ratio...”

- f. L303: *'... less HONO is being emitted than NO<sub>2</sub>...' Is this correct? As a I understand, it makes more sense to me 'a relative decrease of HONO in reference to previous values' instead of 'less HONO is being emitted than NO<sub>2</sub>'. 'less HONO is being emitted than NO<sub>2</sub>' is always true as the ratio is lower than 1. Please, clarify.*

We thank the reviewer for catching this error. We agree with your interpretation and have amended the text as follows:

~~“A decrease in the emission ratio indicates that over time, less HONO is being emitted than NO<sub>2</sub>”~~**there is a relative decrease in HONO in reference to previous values.”**

- g. L306: *As I understand, constant emission factor ratios should lead to constant emission ratios. To make this clear, I would change here 'emission factor ratio' to 'emission ratio'.* We agree. We have amended the text as follows:

~~“If this were true, HONO and NO<sub>2</sub> should have a constant emission factor ratios over both time and sum FRP.”~~

- h. *Figure 4: How you can extract the emission rates of HONO and NO<sub>2</sub> from 7 points (Figure 4) regarding that you only have 3 plumes (3 tracks) (L204-207)*

As stated earlier in the manuscript, only three tracks capture the smoke plume core: tracks 12, 14, and 18. However, in this section, we are extracting and reporting emission ratios. Even though the remaining tracks capture the smoke plume at an angle and are asymmetric, both the HONO and NO<sub>2</sub> VCDs share the exact same sampling bias, meaning that they are both asymmetric in the same way. While the emission rates themselves will not be accurate, both emission rates were derived from the same sampling bias. Thus, taking their ratio eliminates this sampling bias. We have amended the text as follows to clarify our methods:

~~“Assuming~~**Acknowledging** that HONO and NO<sub>2</sub> have ~~similar~~**the same** sampling biases, **meaning that each GCAS sample of the plume shares the same sampling orientation of**

**the sweeping bowtie, we can take the ratio of the emissions rates of HONO and NO<sub>2</sub> ( $E_{EMG,HONO}/E_{EMG,NO_2}$ ) from all flight tracks to explore how reactive nitrogen in wildfire smoke is partitioned and processed. By taking a ratio, the sampling biases in HONO and NO<sub>2</sub> VCDs and EMG emission rates cancel out.**

**15. An improved EMG methodology: Monte Carlo diurnal 1-D models**

- a. L353-355: Please, add a reference related to this idea. Why an increase of thermal output lead to an underestimation of the emission rate? Please, clarify.

We have included a second reference to Wiggins et al. (2020), where Fig. 1 shows the average diurnal cycle of fires from the FIREX-AQ campaign. Many of these fires have peak thermal output, measured as FRP, after 1:30 PM local time. Thus, a satellite like TROPOMI would not observe the maximum thermal output, and would underestimate the emission rate. We have amended the text as follows:

“Third, wildfires have been observed to increase their thermal output after the 1:30 PM local time that TROPOMI and some other polar-orbiting satellites observe at (Wiggins et al., 2020), and thus daily EMG estimates from ~~the EMG method~~ polar-orbiting satellites may underestimate maximum emission rate.”

- b. L378: From which time? Please, clarify.

The summation and division was conducted from 16 August 7 UTC to 17 August 7 UTC, or midnight to 11:59 PM local time. We have added this clarification to the text as follows:

“~~The First, the~~ diurnal FRP shape is obtained by dividing the summed GOES-16 and GOES-17 5-minute FRP product by the GOES total daily FRP **for times 16 August 7 UTC to 17 August 7 UTC, equivalent to 16 August 00:00 to 23:59 local time.**”

- c. L375-383: As I understand, here you are referring to the 4th PECAN 1-D configuration. Please, clarify this in the text.

We have clarified this in the text as follows:

“We run ~~the~~ **this fourth configuration of the** 1-D PECANS model one hundred times...”

- d. L378: See above where? Please, clarify.

We tried to refer to our discussion in the last sentence of the previous paragraph. We see that this is confusing to the reader and have amended the text as follows:

“**Second, EMG functions are fit to all appropriate satellite GCAS observations (see above greater than forty minutes after fire start and before fire intensity diminishes)...**”

- e. L379: Where are these ranges coming from exactly?

These ranges come from the population of generated EMG fit lifetimes and total emission rates from the second step. The text has been amended as follows:

“...**greater than forty minutes after fire start and before fire intensity diminishes) to create a population of EMG fit emission rates and lifetimes and establish sampling ranges.**”

- f. L3979: As written here, it seems that the lifetime is also included in the division. Please, consider to reword this sentence to better readability.

We have broken up this long sentence into shorter, step-by-step sentences to expand on the process and improve readability, as demonstrated by our last four responses. The last sentence of this process is as follows:

**“Third, the ranges of EMG fit emission rate divided by the fractional FRP (total daily emission rate) lifetimes and lifetime total-emission rate divided by the fractional FRP are used to create sampling distributions for a Monte Carlo simulation.”**

- g. L375-383: Please, specify at which data this exercise is going to be applied.

This exercise will be applied to GCAS data for track 12 and 14. We have amended the text as follows:

**“The coefficient of determination root mean square error,  $R^2$  (RMSE), between the model and the satellite GCAS observations for tracks 12 and 14 is are shown in Fig. 6 for HONO and  $\text{NO}_2$ .”**

- h. L381: PECANS

The text has been modified to, **“We run the this fourth configuration of the 1-D PECANS model...”**

- i. L375-L383: This paragraph needs further improvement as it deals with complex concepts that need some clarification.

We believe that the responses to the past 7 comments clarifies these complex concepts.

- j. L385: Here, the plotted value is  $R^2$ . You could say that, regarding a  $R^2$  value, you could find that it is approximately constant in a combination of lifetime and emission rate values that follow an inverse relationship. However, you cannot say that 'the emission rate has an inverse relationship with lifetime'. Please, reword this sentence.

We agree with the reviewer. We have amended the text as follows:

**“In Fig. 6, RMSE is approximately constant in a combination of lifetime and emission rate values that follow an inverse relationship the-emission rate has an inverse relationship with lifetime...”**


- k. Figure 6: Why are two y-axis showing the emission rate in two different units? Please, consider just using one and write to the relation between these 2 units in the text (if needed).

There are two y-axes showing the emission rate in two different units for the benefit of the reader. There are three kinds of audiences that may be interested in this article. First, there is the satellite retrieval group. They inherently work with molecules. Second, there is the emission inventory and modeling group. They inherently work with grams. Third, the audience group that is not within the first or second audience groups. We are not favoring either of the first or second groups by providing an axis for both. We will be keeping both y-axes.

- I. Figure 6: There are 2 colorbars that are representing the same value. Please, consider to keep only one.

We see how the presence of 2 colorbars is busy, distracting, and slightly repetitive. We have removed the coloring of the randomly sampled models and likewise removed the second color bar. As noted by the reviewer and others, we have remade the figure using root mean square error (RMSE) as the fitting metric.

The updated Fig. 6 is now in the manuscript and is shown below:



- m. L400: Which are the input values from the TUV Quick Calculator? Please, clarify.

Upon looking at the previous inputs for the TUV Quick Calculator, we find that there was room for improvement of the inputs used to calculate HONO's lifetime from photolysis. We have selected a new set of values and provided explanations for our selections. The table found below is now included in "Appendix E: TUV Quick Calculator Inputs and Outputs". The following text has been amended in the manuscript:

"The lifetime of HONO from photolysis alone just above the smoke plume, estimated from the TUV Quick Calculator (**Madronich, 2016** <https://www.acom.ucar.edu/Models/TUV/Interactive-TUV/>), would be ~~13~~ **11** min and ~~15~~ **13** min during the overpasses of track 12 and 14 respectively, which is shorter than those lifetimes found in Figs. 6a and 6b. **The inputs and outputs of the TUV Quick Calculator can be found in Tables E1 and E2.**"

The input and output values from the TUV Quick Calculator are as follows in the tables below and in the manuscript:

**Table E1. TUV Quick Calculator Inputs**

<b>Inputs</b>	<b>Values</b>	<b>Justification</b>
<b>Wavelength Start (nm)</b>	<b>280</b>	<b>Default</b>
<b>Wavelength End (nm)</b>	<b>700</b>	<b>Default</b>
<b>Wavelength Increments (-)</b>	<b>420</b>	<b>Default</b>
<b>Latitude (°)</b>	<b>34.68</b>	<b>Sheridan Fire latitude</b>
<b>Longitude (°)</b>	<b>-112.89</b>	<b>Sheridan Fire longitude</b>
<b>Date (YYMMDD)</b>	<b>20190816</b>	<b>Analysis Date</b>
<b>Time (hh:mm:ss, GMT)</b>	<b>22:45:15 and 23:20:30</b>	<b>Track 12 and track 14 ER-2 overpass times</b>
<b>Overhead ozone column (du)</b>	<b>300</b>	<b>Default</b>
<b>Surface Albedo (0-1)</b>	<b>0.15</b>	<b>Default albedo of forests</b>
<b>Ground Elevation (km asl)</b>	<b>1.55</b>	<b>Sheridan Fire elevation</b>
<b>Measurement Altitude (km asl)</b>	<b>6.25</b>	<b>4.7 km altitude</b>
<b>Clouds Optical Depth (-)</b>	<b>0</b>	<b>No clouds</b>
<b>Clouds Base (km asl)</b>	<b>N/A</b>	<b>No clouds</b>
<b>Clouds Top (km asl)</b>	<b>N/A</b>	<b>No clouds</b>
<b>Aerosols Optical Depth (-)</b>	<b>0.5</b>	<b>DC-8 DIAL measurement</b>
<b>Aerosols Single Scattering Albedo (SSA; -)</b>	<b>0.7</b>	<b>Default smoke SSA; 0.37 – 0.95 are valid values (Lewis et al., 2008; Liu et al., 2014)</b>
<b>Aerosols Alpha</b>	<b>1.63</b>	<b>Saleh et al. (2013)</b>
<b>Sunlight Direct beam (-)</b>	<b>1</b>	<b>Default</b>
<b>Sunlight Diffuse down (-)</b>	<b>1</b>	<b>Default</b>
<b>Sunlight Diffuse up (-)</b>	<b>1</b>	<b>Default</b>

**Table E2. TUV Quick Calculator Outputs**

<b>Output Times</b>	<b>HONO photolysis frequency (s<sup>-1</sup>)</b>	<b>HONO photolysis lifetime (min)</b>
<b>22:45:15</b>	<b>1.467E-03</b>	<b>11.36</b>
<b>23:20:30</b>	<b>1.290-03</b>	<b>12.92</b>

Here are the new references added to the manuscript:

“Lewis, K., Arnott, W. P., Moosmüller, H., and Wold, C. E.: Strong spectral variation of biomass smoke light absorption and single scattering albedo observed with a novel dual-wavelength photoacoustic instrument, *J. Geophys. Res.*, 113, 2007JD009699, <https://doi.org/10.1029/2007JD009699>, 2008.

Liu, S., Aiken, A. C., Arata, C., Dubey, M. K., Stockwell, C. E., Yokelson, R. J., Stone, E. A., Jayarathne, T., Robinson, A. L., DeMott, P. J., and Kreidenweis, S. M.: Aerosol single scattering albedo dependence on biomass combustion efficiency: Laboratory and field studies, *Geophysical Research Letters*, 41, 742–748, <https://doi.org/10.1002/2013GL058392>, 2014.

Saleh, R., Hennigan, C. J., McMeeking, G. R., Chuang, W. K., Robinson, E. S., Coe, H., Donahue, N. M., and Robinson, A. L.: Absorptivity of brown carbon in fresh and photochemically aged biomass-burning emissions, *Atmos. Chem. Phys.*, **13**, 7683–7693, <https://doi.org/10.5194/acp-13-7683-2013>, 2013.”

- n. L424-426: *What does this mean? A presence of a large source of RO<sub>2</sub> is realistic? If it is not, what you can say about it? Looking at the plume from Figure A1, it can be seen that there is some influence from the surface. Can the surface have an impact on the retrieval and therefore in the lifetime? Please, clarify.*

Yes, the presence of a large source of RO<sub>2</sub> is realistic. A previous chemical modeling study based on actual observations of wildfire smoke indicated that there was a missing sink of NO<sub>x</sub> and a missing source of organic nitrates (see text added to the manuscript below).

Regarding the surface influence in Figure A1, we believe the reviewer is either talking about the linear artifacts in the image or the pixels outside of the core plume. For the first point, these linear artifacts are the result of overlapping swath samples with some lingering edge processing effects and averaging the swaths over different time periods. For the second point, we believe that the isolated pixels outside of the plume edge are primarily because we chose a conservative plume mask, which was three times the background value. With a less conservative plume mask, the disconnected pixels rejoin the plume.

However, the surface can impact the retrieval by affecting SCDs and AMFs through surface reflectivity, as mentioned earlier in the manuscript. We cannot determine which way the surface reflectivity impacts the lifetime without performing a larger separate analysis, but generally higher surface albedo causes higher AMFs. However, by dividing the SCDs by the AMF, the VCDs now consider the effect that surface reflectivity has on the slant column retrieval. We would anticipate minimal surface impacts on the retrieval and lifetime determination.

We have added the following discussion to the manuscript:

“However, a simple chemical box model applied to the Sheridan Fire indicates that a large source of organic peroxy radicals (RO<sub>2</sub>) is needed to drive the NO<sub>2</sub> chemical lifetime to the short values we infer from applying the 1-D plume model (with no chemical mechanism) to the GCAS observations. **This is not dissimilar to a hypothesized missing RO<sub>2</sub> source seen in the Taylor Creek Fire from 2018 (Peng et al., 2021). In that study, a baseline box model simulation of the Taylor Creek Fire overestimated NO<sub>x</sub> and underestimated organic nitrates, therefore missing a NO<sub>x</sub> to organic nitrate reaction pathway. By incorporating proxy peroxy radicals with organic nitrate formation pathways, their model better reflected the observed NO<sub>x</sub> decay. This has crucial implications for understanding the chemical evolution of wildfire smoke.**”

The following reference has been added to the References section:

“Peng, Q., Palm, B. B., Fredrickson, C. D., Lee, B. H., Hall, S. R., Ullmann, K., Campos, T., Weinheimer, A. J., Apel, E. C., Flocke, F., Permar, W., Hu, L., Garofalo, L. A., Pothier, M. A., Farmer, D. K., Ku, I.-T., Sullivan, A. P., Collett, J. L., Fischer, E., and Thornton, J. A.: Observations and Modeling of NO<sub>x</sub> Photochemistry and Fate in Fresh Wildfire Plumes, *ACS Earth Space Chem.*, 5, 2652–2667, <https://doi.org/10.1021/acsearthspacechem.1c00086>, 2021.”

**16. Biomass burning emission inventories underestimate observationally constrained emission estimates**

- a. L434: Please, consider previous comments regarding the bottom-up and top-down approaches.

Please see our response to your comment on manuscript L39. We use bottom-up and top-down appropriately.

- b. Figure 7: In this section, you deal with the minimum and maximum values of emission rate extracted from track #12 and #14 as representative of the whole day. You called them as 'daily emission rate', which is not true because it does not consider the emission rate at different times during the day. Please, consider to change the terminology and to develop a discussion with this change. Stating that there are biases in reference to daily mean emission from inventories can be misleading.

In this section, we deal with the minimum and maximum values of daily mean emissions. In each iteration of the 1-D PECANS model, we extract the total daily emission rate and the lifetime. The total daily emission rate is converted to a daily mean emission rate by multiplying the total daily emission rate by the normalized daily FRP to get a diurnal profile of instantaneous emission rates. This diurnal profile is then averaged to produce the daily mean emission rate plotted in Fig. 7.

We have added text to clarify this conversion as follows:

“We compare our Monte Carlo diurnal 1-D model HONO and NO<sub>2</sub> emissions with those reported in other biomass burning emissions inventories. **We first convert the total emission rate from the lowest RMSE diurnal 1-D model to a daily mean emission rate. The total emission rate is multiplied by the normalized diurnal FRP profile to get a diurnal profile of instantaneous emission rates. This diurnal profile is then averaged to produce the daily mean emission rate.**”