Retrieval of NO₂ profiles from three years of Pandora MAX-DOAS measurements in Toronto, Canada

Ramina Alwarda^{1,2}, Kristof Bognar³, Xiaoyi Zhao², Vitali Fioletov², Jonathan Davies², Sum Chi Lee²,

5 Debora Griffin², Alexandru Lupu², Udo Frieß⁴, Alexander Cede^{5,6}, Yushan Su⁷, Kimberly Strong¹

¹Department of Physics, University of Toronto, Toronto, Canada
 ²Air Quality Research Division, Environment and Climate Change Canada, Toronto, Canada
 ³3vGeomatics, Vancouver, Canada
 ⁴Institute of Environmental Physics, Heidelberg University, Heidelberg, Germany
 ⁵NASA Goddard Space Flight Center, Greenbelt, USA
 ⁶LuftBlick, Innsbruck, Austria

⁷Ontario Ministry of the Environment, Conservation and Parks, Toronto, Canada

Correspondence to: Ramina Alwarda (ramina.alwarda@mail.utoronto.ca)

Abstract. The purpose of this work is to derive new NO2 vertical profiling data products from Pandora spectrometers and

- 15 investigate the factors contributing to the bias of this dataset relative to established ground-based and spaceborne datasets. Possible applications of the NO₂ vertical profile dataset include air quality monitoring and satellite validation studies. We explore the application of the optimal estimation method to Pandora multi-axis differential optical absorption spectroscopy (MAX-DOAS) measurements to retrieve vertical profile information for nitrogen dioxide (NO₂). We use the Heidelberg Profile (HeiPro) retrieval algorithm to derive, for the first time, NO₂ profiles and partial columns (0–4 km) from Pandora MAX-DOAS
- 20 measurements from 2018–2020 at Downsview, a suburban neighbourhood in the north end of Toronto, Canada that is subject to local traffic emissions and urban influences. Validation of the new dataset was done via comparison with official Pandora direct-Sun measurements, in situ observations, satellite data, and an air quality forecasting model. We find that, for tropospheric partial column comparisons, the HeiPro dataset has a positive mean relative bias to Pandora direct-Sun (61% ± 9.7%) and TROPOMI (37% ± 51%) observations, as well as the GEM-MACH model output (67% ± 7.1%), with similar
- 25 seasonal and diurnal cycles in the bias with Pandora direct-Sun and GEM-MACH. Contributing factors to the large bias of HeiPro to Pandora direct-Sun were investigated, and NO₂ heterogeneity, combined with differences between direct-Sun and multi-axis viewing geometries, was found to contribute a maximum of 52% of the total relative bias during morning measurement times. For surface NO₂ comparisons, we find that HeiPro measurements capture the magnitude and diurnal variability of surface NO₂ reasonably well (mean relative bias to in situ surface NO₂: -<u>9.7% + 7.5%</u>) but are low-biased
- 30 compared to GEM-MACH (mean relative bias: −3⁷/₂% ± 2.4%). Compared to HeiPro, the GEM-MACH model profiles are high-biased in the lower boundary layer and low-biased in the free troposphere.

Deleted: 41		
Deleted: 47		
Deleted: 1		
Deleted: 5		
Formatted: Not Highligh	ıt	
Formatted: Not Highligh	nt	
Deleted: 8.7		
Deleted: %		
Deleted: 5		
Deleted: 3		
Deleted: 6		

1 Introduction

Nitric oxide (NO) and nitrogen dioxide (NO₂), together comprising NO_x, play important roles in air quality, tropospheric chemistry, and stratospheric ozone (O₃) chemistry. In the troposphere, NO_x is primarily emitted from fossil fuel combustion and nitrogen fertilizers and has a near-surface peak in the profile shape in polluted regions; additionally, it is emitted from natural sources such as lightning and biomass burning (e.g., ECCC 2016; Murray, 2016). The photolysis of NO₂ rapidly contributes to the production of O₃ and therefore photochemical smog formation. NO₂ also plays a more direct role in air quality as it has toxic biological effects at high concentrations and has been linked to negative health outcomes such as asthma exacerbations (ECCC, 2016). Tropospheric NO₂ has a large spatiotemporal variability and this can be attributed to its short lifetime, variability of emissions, and meteorological variations (Beirle et al., 2003).

50

45

Due to its role in various atmospheric processes and impact on air quality, it is important to continuously monitor NO₂ and its spatiotemporal variability. Surface NO₂ is measured using in situ photolytic conversion/chemiluminescence as part of Canada's Air Quality Health Index (AQHI, Stieb et al., 2008) along with surface O_3 and surface fine particulate matter with diameter < 2.5 microns (PM2.5). While in situ instruments provide surface measurements with high temporal resolution,

- 55 such measurements are spatially limited both in geographic extent and altitude. Ground-based remote sensing measurements address the latter limitation by measuring NO₂ throughout the troposphere and stratosphere and are used in air quality and satellite validation studies (e.g., Ma et al., 2013; Griffin et al., 2019; Zhao et al., 2019, 2020; Verhoelst et al., 2021). Non-geostationary satellite instruments, on the other hand, address spatial limitations from ground-based monitoring and provide global coverage. A suite of UV-visible satellite instruments has monitored NO₂ columns since 1995, beginning with the Global
- 60 Ozone Monitoring Experiment (GOME, Burrows et al., 1999) and more recently, the TROPOspheric Monitoring Instrument (TROPOMI) onboard the European Space Agency's Sentinel-5 Precursor (S5P) satellite. While such satellite instruments provide global coverage, the temporal frequency of at any location is limited, which hinders insight into the spatiotemporal variation of NO₂ on large scales. The recent additions of geostationary satellite instruments such as the Tropospheric Emissions: Monitoring of Pollution (TEMPO, Zoogman et al., 2017) and the Geostationary Environmental Monitoring
- 65 Spectrometer (GEMS, Kim et al., 2020) address this limitation of low-Earth satellite orbits. Thus, in situ, ground-based, and spaceborne measurements all contribute to our understanding of the spatiotemporal variability of NO₂.

Ground-based UV-visible spectra are usually analysed using Differential Optical Absorption Spectroscopy (DOAS, Platt and Stutz, 2008). These measurements include zenith-scattered-light DOAS (ZSL-DOAS), which is most sensitive to the root stratospheric column, multi-axis DOAS (MAX-DOAS), which is most sensitive to the tropospheric column, and direct-Sun (DS), which measures the total column with equal sensitivity to the troposphere and stratosphere. One such ground-based instrument that performs routine direct-Sun, zenith-sky, and multi-axis measurements of the atmosphere is the Pandora UVvisible spectrometer (Herman et al., 2009). Pandora instruments are distributed globally as part of the Pandonia Global

Network (PGN), an international collaboration led by the National Aeronautics and Space Administration (NASA), European Space Agency (ESA), the US Environmental Protection Agency (US-EPA), and the Korean National Institute of Environmental Research (NIER), which provides standardized high-quality datasets, such as direct-Sun NO₂ total columns, for various purposes including satellite validation and air quality assessments (e.g., Zhao et al., 2022).

- Additionally, Pandora MAX-DOAS spectra can be used to retrieve vertical profiles of trace gases in the lower troposphere, for example by using the optimal estimation method (OEM), a maximum a posteriori profile inversion algorithm which is based on Bayesian statistics and provides a robust error analysis (Rodgers, 2000; Frieß et al., 2019). Utilizing a MAX-DOAS NO₂ dataset is advantageous because it provides sensitivity to tropospheric absorbers, profile information, increased temporal resolution compared to spaceborne measurements, horizontal distribution of NO₂ surrounding a measurement site (Dimitropoulou et al., 2020), and a horizontal resolution that is more suited for assessing model performance compared to in
- 85 situ data (Blechschmidt et al., 2020). NO₂ profiles from MAX-DOAS measurements are also a useful tool for aiding in satellite retrieval algorithms and validation; previous studies have utilized NO₂ profiles derived from MAX-DOAS data as inputs for the TROPOMI retrieval algorithm to reduce the systematic underestimation in the satellite data compared to ground-based data (e.g., Dimitropoulou et al., 2020). Partial columns of NO₂ from MAX-DOAS data have also been used in ground-based validation of spaceborne measurements, whereby satellite tropospheric NO₂ is typically biased low to ground-based MAX-
- 90 DOAS NO₂ in polluted regions (e.g., Pinardi et al., 2020; Verhoelst et al., 2021). Satellite measurements rely on ground-based data to continuously improve satellite algorithms and understand the bias that exists between spaceborne and ground-based measurements as a function of location, especially due to the difficulty in implementing location-specific a priori NO₂ profiles in satellite retrieval algorithms (Verhoelst et al., 2021). In addition to OEM, there are various other retrieval techniques that convert MAX-DOAS measurements to NO₂ profiles and subsequent data products. For example, Pandora MAX-DOAS spectra
- 95 can be used to retrieve NO₂ vertical profiles, tropospheric columns, and surface values using an L2 Air-Ratio Sky Algorithm developed by Elena Spinei and the PGN (see PGN software manual v1.8 available at https://www.pandonia-globalnetwork.org/home/documents/manuals/ for further information, last access 13 September 2024). Comparisons of these PGN sky algorithm data products with other datasets at the measurement site in this study will be the subject of a future study.
- 100 While OEM has been applied to Pandora MAX-DOAS data in a study by Nowlan et al. (2018) that compared aircraft retrievals of NO₂ to ground-based MAX-DOAS and direct-Sun Pandora measurements for an eleven-day period, it has not yet been applied to Pandora MAX-DOAS observations for extended periods of time. A longer intercomparison period would enable investigations of annual and seasonal trends compared to other NO₂ datasets, especially comparing Pandora MAX-DOAS to direct-Sun observations. Previous comparisons between NO₂ observations of MAX-DOAS to direct-Sun are limited but would be useful in addressing the difference in the relative biases of multi-axis and direct-Sun observations to satellite measurements. For example, Pinardi et al. (2020) found a bias of 10–15% in the ground-based MAX-DOAS NO₂ columns
 - compared to direct-Sun tropospheric columns from the same station, but such comparisons were during the overpass times of 3

the two satellites in the study. The relative bias was attributed to NO₂ heterogeneity as well as the differences in airmasses measured by the direct-Sun and multi-axis geometries. Performing NO₂ comparisons of MAX-DOAS to direct-Sun throughout the day would be an important consideration for the validation of geostationary satellite NO₂ measurements, since both are

used for the validation of spaceborne tropospheric NO₂ (e.g., Verhoelst et al., 2021).

Therefore, due to the advantages of obtaining NO₂ profiles, as well as the limited studies comparing MAX-DOAS to direct-Sun NO₂, the aim of this work is to derive NO₂ profiles from Pandora MAX-DOAS measurements using OEM and to
assess the quality of the dataset by comparison to established datasets. We present, for the first time, a three-year dataset of 0–4 km NO₂ profiles from 2018–2020 at Downsview, a Pandora measurement site in the north part of Toronto, Ontario, Canada, located in a suburban area subject to local traffic emissions and urban influences. To our knowledge, this work is the first study to apply the optimal estimation method to Pandora MAX-DOAS data to retrieve a multi-year NO₂ profile dataset. We assess the biases and contributing factors to the biases between the OEM-based Pandora dataset versus (1) standard measurements
comprising Pandora direct-Sun (Pandora-DS) and TROPOMI tropospheric columns as well as in situ surface NO₂, and (2) profiles, partial columns, and surface NO₂ from the Global Environmental Multiscale–Modelling Air-quality and Chemistry (GEM-MACH) high-resolution regional air quality forecast model. Where possible, we quantify the contributing factors to the biases and assess the impact of spatial heterogeneity on the biases. The resulting OEM-based Pandora NO₂ profile dataset can used to assess spatiotemporal NO₂ heterogeneity around the measurement site as well as any biases between spaceborne and

125 ground-based NO2 at the measurement site.

110

130

140

The paper is organized as follows: Section 2 describes the various instruments, algorithms, and models used in this study, as well as the data filtering and co-location criteria that were applied. In Section 3, the quality of the OEM-based Pandora dataset is assessed by comparing it with Pandora <u>DS</u> and <u>TROPOMI</u> tropospheric columns as well as <u>GEM-MACH</u> partial columns. Section 4 presents the surface NO₂ volume mixing ratio (VMR) comparisons among the OEM-based Pandora dataset, in situ data from the National Air Pollution Surveillance (NAPS) monitoring program, and <u>GEM-MACH</u> data, as well as the comparisons between Pandora OEM and <u>GEM-MACH</u> NO₂ profiles. Seasonal and diurnal trends are examined where possible, and the causes of differences among the measurements are investigated. Finally, Section 5 summarizes the implications and conclusions.

135 2 Methods: Instruments, datasets, and models

An overview of the instruments and corresponding datasets used in this study is presented in Fig. <u>1</u>, Section 2.1 describes the <u>Pandora spectrometer and Sections 2.1.1–2.1.3 describe the direct-Sun, MAX-DOAS, and profiling retrieval algorithms, respectively. Sections 2.2 and 2.3 describe the <u>TROPOML</u> and photolytic conversion/chemiluminescence in situ <u>measurements, respectively</u>. Section 2.4 describes the models used in this work, both auxiliary models used in producing other</u>





Deleted: Lon. Figure 1: A map (© Google Maps) of the measurement site (Downsview, white circle) and surrounding Greater Toronto Area masked by the TROPOMI pixel-averaged tropospheric NO₂ field from 2018–2020. The color bar indicates the NO₂ tropospheric column in molec cm². Solid and dashed lines represent the minimum (yellow) and maximum (red) solar azimuth angle (SAA) during summer and winter, respectively. The solid black line represents the Pandora instrument's multiaxis azimuth viewing angle (AVA) of 255°.

Deleted:	direct-Sun
----------	------------

Deleted: Retrieval Algorithms
Deleted: M
Formatted: Space Before: 18 pt, After: 0 pt
Deleted: 2
Deleted: three instruments: the Pandora spectrometer,
Deleted: ,
Deleted: the
Deleted: sensor
Deleted: 2 discusses the retrieval algorithms that were used to

Convert Pandora direct-Sun spectra to tropospheric columns and Pandora MAX-DOAS data to profiles

Deleted: . Lastly, Section 2.3 focuses on



2.1 Pandora UV-visible spectrometer

Pandora instruments utilize a temperature-stabilized Czerny-Turner spectrometer with a grating of 1200 gr/mm and a back-thinned charge-coupled device (CCD) detector to record spectra between 280–530 nm with a spectral resolution of 0.6 nm. All Pandora NO₂ data products presented in this work were obtained from Pandora no. 103 (P103) located on the rooftop of the Environment and Climate Change Canada (ECCC) five-storey building at Downsview (43.78 °N, 79.47 °W, 186 m a.s.l.) in Toronto, Canada. This building is located in a suburban region with several nearby roads. P103 operates in direct-Sun, zenith-sky, and multi-axis viewing geometries during the sunlit period. Figure 2 shows the May 2018 – December 2020 TROPOMI tropospheric NO₂ field at the measurement site and surrounding region, as well as the differences in viewing

185 directions between the direct-Sun and multi-axis measurements. Direct-Sun and multi-axis data products from P103 from February 2018 to June 2020 were used in this study. Further details on the <u>retrieval algorithms used on the Pandora data are</u> discussed in Sections 2.1, 1 (direct-Sun), 2.1.2 (multi-axis), and 2.1.3 (profiling),

2.1, 1, Pandora-DS retrievals,

The standard <u>Pandora-DS</u> total column NO₂ data product is obtained using Total Optical Absorption Spectroscopy (TOAS), as implemented by PGN's BlickP software (Cede, 2019). Direct-Sun spectra in the 400–440 nm range are fitted with cross-sections of NO₂ (at an effective temperature of 254.5 K, Vandaele et al., 1998), O₃ (at an effective temperature of 255 K, Brion et al., 1993, 1998; Daumont et al., 1992), and a fourth-order polynomial to produce SCDs of NO₂ with a clear-sky precision of 2.7×10¹⁴ molec cm⁻² (Herman et al., 2009). A synthetic reference spectrum is used in the analysis and is obtained by taking an average of several measured spectra which are corrected for their total optical depth, Following this, NO₂ SCDs

- 195 are converted to vertical column densities (VCDs or total columns) using geometric AMFs. <u>The Pandora-DS NO₂ VCD has an absolute accuracy of 1.3×10¹⁵ molec cm⁻² (Herman et al., 2009). The Pandora-DS NO₂ VCDs used in this study are obtained from the PGN (https://data.pandonia-global-network.org/Downsview/Pandora103s1/, file version: rnvs1p1-7, last access: 16 April 2024) and only high-quality data (i.e., L2 flags of 0 and 10) were used. Because the focus of this work is on tropospheric</u>
- NO₂ intercomparisons, the resulting NO₂ VCDs were converted to tropospheric NO₂ columns by subtracting the stratospheric
 NO₂ columns obtained from the Ozone Monitoring Instrument (OMI, NASA standard product version 3.0 level 2, Krotkov et al., 2017) onboard the Aura satellite. Due to the diurnal variation of NO₂, and the satellite's overpass time of 13:30 local time (LT), a photochemical box model (Pratmo, discussed further in Section 2.3.1) was used to calculate stratospheric NO₂ at various Pandora measurement times throughout the day. <u>The stratospheric portion that was removed accounted for 34% ± 2.8% of the Pandora-DS NO₂ total columns. The Pandora-DS tropospheric NO₂ data were averaged to obtain hourly means for comparison with the other datasets.
 </u>

Deleted: 2.1.1 Pandora UV-Visible Spectrometer

Deleted: direct-Sun retrieval technique are

Deleted: 2

Deleted: provided be	while a preface to the multi-axis viewing geometry is slow prior to being further discussed in Section 2.2.2.
Deleted:	1
Deleted:	2
Deleted:	1
Deleted:	Direct-Sun
Deleted:	PGN
Deleted:	from Pandora direct-Sun spectra
Deleted:	, Cede, 2021
Deleted:	and
Deleted:	
Deleted:	
Deleted:	(

Deleted: The Pandora direct-Sun NO₂ VCDs are obtained from the PGN (https://data.pandonia-globalnetwork.org/Downsview/Pandora103s1/, file version: rnvs1p1-7, last access: 16 April 2024) and only high-quality data (i.e., L2 flags of 0 and 10) were used.

Deleted: Pandora direct-Sun (from here on,

Deleted:)



230

235

Figure 2: A map (© Google Maps) of the measurement site (Downsview, white circle) and surrounding Greater Toronto Area masked by the TROPOMI pixel-averaged tropospheric NO₂ field from May 2018 – December 2020. The color bar indicates the NO₂ tropospheric column in molec cm⁻². Solid and dashed lines represent the minimum (yellow) and maximum (red) solar azimuth angle (SAA) during summer and winter, respectively. The solid black line represents the Pandora instrument's multi-axis azimuth viewing angle (AVA) of 255°.

2.1.2 Differential slant column density (dSCD) retrievals

The multi-axis viewing geometry can provide vertical information from tropospheric absorbers by measuring scattered sunlight at various elevation viewing angles (EVAs) of the instrument. P103 performs both 'long' and 'short' <u>multiaxis</u> scans in an alternating manner, with an azimuth viewing angle of 255° (see Fig. 2). Long scans consist of EVAs of 1°, 2°, 3°, 5°, 8°, 10°, 15°, 20°, 30°, 40°, 50°, and 90° and short scans consist of EVAs of 1°, 2°, 15°, 30°, and 90°. The resulting multiaxis spectra from P103 (i.e., corrected Level 1 data, data file version: smca1c1p1-7) were obtained from the PGN. Utilizing the DOAS technique and fitting parameters outlined in Appendix A of Kreher et al. (2020), spectral fitting to the ratio of two spectra, with one spectrum at each EVA and the other at a 90° reference EVA, is performed to retrieve the differential slant column density (dSCD) of the oxygen collision complex (O4) and NO₂ at each EVA.<u>This is achieved using the QDOAS</u> software (https://uv-vis.aeronomie.be/software/QDOAS/, last access: 16 April 2024), developed by the Royal Belgian Institute for Space Aeronomy (Danckaert et al., 2017). O4 and NO₂ dSCDs were retrieved in both the ultraviolet (UV, 338–370 nm,

only measurements with the UV band pass filters) and visible (vis, 425–490 nm) windows. Differential cross-sections of NO2
 at 294 and 220 K for both windows (Vandaele et al., 1998), O4 at 293 K for both windows (Thalman and Volkamer, 2013), O3 at 223 and 243 K for UV and 223 K for vis (Serdyuchenko et al., 2014), BrO at 223 K for UV only (Fleischmann et al., 2004),

7

Deleted: 2.2

Deleted: 1

Moved (insertion) [1]

HCHO at 297 K for UV only (Meller and Moortgat, 2000), and H2O for vis only (Rothman et al., 2010) were convolved using
 the instrument slit function and the nominal wavelength calibration file from PGN. A fifth-degree polynomial, linear offset, and first-order shift and stretch were used in both windows (Kreher et al., 2020). Following the QDOAS analysis, O4 and NO2 dSCDs with fitting residual root-mean-square (RMS) values ≥ 3×10⁻³ were discarded (only 5% of the data, mostly at SZA > 80°). This filter was selected to improve the quality of the fits and discard noisier fits from the analysis. Next, we discuss the Heidelberg Profile Retrieval algorithm (HeiPro, Frieß et al., 2011; Yilmaz, 2012), which was used to produce aerosol extinction and NO2 profiles.

2.1.3 HeiPro retrievals

HeiPro was used to convert Pandora O4 and NO2 dSCDs of at various EVAs to aerosol extinction and NO2 profiles.
respectively, from 0–4 km. HeiPro is a maximum a posteriori optimal estimation method (Rodgers, 2000) that utilizes a forward model (a radiative transfer model or RTM, here being SCIATRAN version 2.2, Rozanov et al., 2005), to simulate dSCDs
based on a priori NO2 profile information and additional parameters such as aerosol extinction, aerosol optical properties, surface albedo, and temperature. Exponentially decreasing aerosol extinction and NO2 profiles with scale heights of 1 km and column values of 0.18 (aerosol optical depth, AOD) and 9×10¹⁵ molec cm⁻² (NO2) were used as a priori information in the HeiPro simulation. Additional a priori inputs to HeiPro were daily pressure and temperature profiles from ERA5 reanalysis data at the grid box nearest to the measurement site (discussed further in Section 2.4, 2). For the aerosol retrievals, a single
scattering albedo value of 0.92 and asymmetry factor of 0.68 were used. For all retrievals, a surface albedo value of 0.06 was used, and correlation matrices with a variance of 50% and correlation length of 200 m were used.

Given these settings, a cost function is used to assess the difference between the dSCDs simulated by the RTM and*
the measured dSCDs while accounting for additional a priori constraints. The Levenberg–Marquardt algorithm is then used to
iteratively find the minimum of the cost function. At this minimum, the most probable atmospheric state or the maximum *a posteriori* estimate (MAP) is then taken to be the NO₂ profile. HeiPro utilizes a two-step approach: prior to each NO₂ profile
retrieval, aerosol extinction profiles are retrieved based on O₄ dSCDs; these are subsequently used as inputs for the NO₂ profile
retrievals. Profiles were retrieved separately for UV and visible scans, with RTM wavelengths of 360 nm for O₄-UV and NO₂-UV, 477 nm for O₄-vis, and 460 nm for NO₂-vis. Full details <u>on</u> the HeiPro algorithm can be found in Frieß et al. (2006,
2011). The profile grid is from 0–4 km, with a vertical resolution of 200 m. Therefore, the NO₂ concentration for the 0–200 m grid level represents the average throughout this range, i.e., the concentration at 100 m. The HeiPro 'surface' concentration reported in this study is then obtained by extrapolating the VMR at the 0–200 m grid level to 0 m for each individual profile. The linear extrapolation method used in this study produces NO₂ surface values that are, on average, 1 ppbv larger than the NO₂ value at 100 m. Therefore, this extrapolation method may underestimate the HeiPro surface NO₂ values that we report.

Moved up [1]: This is achieved using the QDOAS software (https://uv-vis.aeronomie.be/software/QDOAS/, last access: 16 April 2024), developed by the Royal Belgian Institute for Space Aeronomy (Danckaret et al., 2017).

(Deleted: 2	
(Deleted: 2	\mathbb{D}
(Deleted: NO ₂	\Box
(Deleted: an	\Box

Deleted: information	
Deleted: utilized	
Deleted: by	
Deleted: 3	

Deleted: about

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1, English (UK)

285

300 Additionally, only the long multi-axis scans (see Section 2.1.2 for list of EVAs) recorded in the UV window, together termed the 'long UV' scans, were incorporated in this work as they have the most reliable results. We found the HeiPro long UV results to be better suited for this study for the following reasons: (1) as expected, the degrees of freedom for signal (DOFS) for each profile, which represents the number of independent pieces of information obtained from the measurements, are greater for the long scans (2.35 and 3.13 for UV and vis, respectively) compared to the short scans (2.03 and 2.75 for UV and vis).

- 305 vis, respectively), see Fig. A1 (in Appendix A). (2) The NO₂ partial column errors are smaller for the long scans (5.45×10¹⁴ and 2.99×10¹⁴ molec/cm² for UV and vis) compared to the short scans (7.51×10¹⁴ and 3.49×10¹⁴ molec/cm² for UV and vis, respectively), see Fig. A1. (3) Lastly, the long UV scans show improved regression slopes, intercept, and correlation coefficient, to the NAPS in situ surface VMRs compared to the long vis scans (see Fig. A2). This is likely due to the fact that scans in the visible window cover a greater horizontal region than in the UV (Ortega et al., 2015), and so UV scans may better capture the
- 310 NO₂ field localized to the in situ instrument inlet. Therefore, only HeiPro retrievals with long UV scans have been used in this study for the comparisons with <u>Pandora-DS and</u> TROPOMI tropospheric columns, GEM-MACH partial columns and surface <u>NO₂</u>, and NAPS in situ surface NO₂. <u>Example MAX-DOAS dSCDs of O₄ and NO₂, along with averaging kernels for aerosol extinction and NO₂ profile retrievals, are demonstrated in Fig. A3.</u>
- 315 Lastly, retrievals for which both the NO₂ profiles and aerosol extinction profiles had DOFS < 1 were excluded from the analysis for quality control purposes (e.g., Vlemmix et al., 2015) and represented 19% of the dataset. Such retrievals contain limited information from the measurements and are more influenced by the a priori profile. The mean DOFS values before and after filtering are 1.61 ± 0.68 and 1.88 ± 0.42 for aerosol extinction, respectively, and 2.35 ± 0.49 and 2.39 ± 0.40 for NO₂, respectively. In addition to providing NO₂ profiles from 0–4 km, HeiPro also provides an integrated value from 0–4 km,
- 320 thereby providing a partial NO₂ column for the lower troposphere. All HeiPro data products presented here are hourly averages. From here on, surface NO₂ as well as 0–4 km partial columns and profiles retrieved from Pandora MAX-DOAS measurements using HeiPro will be described as HeiPro data products.

2,2 TROPOMI

The TROPOMI instrument measures nadir solar UV-visible radiance in a sun-synchronous orbit at an altitude of 817 325 km and a local overpass time of 13:30 LST. The TROPOMI instrument provides spaceborne NO₂ measurements at a horizontal resolution of 3.5 km × 5.5 km (3.5 km × 7.5 km prior to August 2019). The DOAS technique is used to convert UV-visible spectra to NO₂ slant column densities (SCDs). Following this, a chemical transport model (TM5-MP) assimilates the SCDs to NO₂ vertical profiles, from which a stratospheric SCD is determined. Finally, air mass factors (AMFs, which are based on surface albedo, NO₂ profile shape, etc.) are used to obtain a tropospheric column. Details about the instrument and NO₂

330 retrieval scheme can be found in van Geffen et al. (2020). The TROPOMI tropospheric columns used in this work are from processor version 2.3.1 (van Geffen et al., 2022). We used tropospheric NO₂ data product quality assurance values (qa_value)

9

Deleted: 1

Deleted: associated with the retrieval of each profile,

• Deleted: root-mean-square (RMS) residuals are reduced

Deleted: RMS:

Deleted: a reduced multiplicative bias relative

Deleted:

Deleted: retrieved in the visible window

Deleted: (zero-intercept slopes of 0.94 and 0.82 for UV and vis, respectively,

(Deleted: and Pandora-DS

Deleted: in

Formatted: Font: Not Italic, Font color: Text 1

Deleted: to ensure that profile information was not mostly coming from the a priori NO₂ profile.

Deleted: 1.

 \geq 0.75 to exclude retrievals with very cloudy scenes. Only TROPOMI pixels that were within 10 km of the Pandora measurement site, i.e., the ECCC building at Downsview, and within ±10 minutes of the Pandora MAX-DOAS measurement time (Zhao et al., 2020) were included in the comparisons.

	2,3 In situ measurements	Deleted: 1.
350	The in situ instrument (Thermo 42i) at the Toronto North site, i.e., the Pandora measurement site, monitors surface NO ₂ under the NAPS Program and is located near a major road. Horizontally, it is located 100 m away from the Pandora	Deleted: M
	instrument; vertically, it has an air intake inlet that is 4 m above ground level and 11 m below the Pandora instrument. It utilizes	
	the photolytic conversion/chemiluminescence technique to measure NO_2 with a precision of 0.4 ppb. The NO_2 data collected	
	at the Toronto North site was provided on an hourly-averaged timescale and had undergone final validation by the Ontario	
355	Ministry of the Environment, Conservation and Parks	Deleted: 1
	2.4 Models	Deleted: 3
	2.41 Pratmo	Deleted: 3
	Pratmo is a photochemical box model with detailed stratospheric chemistry that is of relevance to NO2 (McLinden et	
	al., 2000; Brohede et al., 2008; Lindenmaier et al., 2011). Because the Pandora-DS observations provide NO2 total columns	
360	and we are interested in only the tropospheric portion, the stratospheric column must be removed. However, sun-synchronous	
	satellite observations of stratospheric NO2 cannot solely be used to do this because they provide a single observation per day	
	whereas stratospheric NO2 exhibits diurnal variability and Pandora measures throughout the day. Therefore, Pratmo provides	
	a modelled ratio of stratospheric NO_2 at the Pandora measurement time and the OMI overpass time. The measured OMI $% \mathcal{O}_{\mathcal{O}}$	
	stratospheric columns can then be multiplied by this modelled ratio to obtain stratospheric NO_2 columns at Pandora	
365	$measurement times. \ The \ OMI \ stratospheric \ NO_2 \ measurements used in this \ study \ are \ from \ version \ 3.0 \ level \ 2 \ (SPv3.0, Krotkov)$	
	et al., 2017) of NASA's standard product (SP).	
	2.4.2 ERA5	Deleted: 3
	Various data products from the European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis v5	
	(ERA5) were utilized in this work (Hersbach et al., 2023). As previously mentioned, pressure and temperature profiles were	
370	used as a priori information in the HeiPro algorithm that retrieves NO_2 profiles from Pandora MAX-DOAS spectra. ERA5	

Deleted: data

pressure and temperature profiles from 0.1–30 km at the grid box nearest to the Downsview site were averaged at timestamps of 11:00, 14:00, 17:00, 20:00, and 23:00 UTC to produce daily a priori temperature and pressure profiles. Secondly, hourly ERA5 2 m temperature and surface pressure data at the grid box nearest to the Downsview site were used to calculate a 15 m NO₂ column (see Section 3 and Equation 1) to account for the rooftop location of the Pandora instrument. Additionally,

planetary boundary layer (PBL) height was obtained from ERA5 at the Downsview site to investigate how the bias among datasets varies as a function of PBL height.

2.4.3 GEM-MACH

385

GEM-MACH is ECCC's operational regional air quality model, which provides predictions of pollutants over North America at $a_10 \text{ km} \times 10 \text{ km}$ horizontal resolution on an hourly timescale for a 72 h period, and is run twice per day at 0 and 12 UTC (Moran et al., 2009; Pavlovic et al., 2016; Pendlebury et al., 2018). The model provides output for 80 vertical levels from the surface to approximately 0.1 hPa. In this work, NO₂ profiles and partial columns from ~0–5.0 km within the model grid box nearest to the Pandora measurement site were extracted from 2018 to 2020, with a 200 m vertical grid. GEM-MACH

390 utilizes emissions generated from inventories from the United States Environmental Protection Agency Air Emissions Modeling Platform, from Canada's Air Pollutant Emission Inventory, and from Mexico's National Emissions Inventory. The model includes physical and chemical processes such as tropospheric gas- and aqueous-phase chemistry, inorganic heterogeneous chemistry, and wet and dry deposition. It should be noted that the operational version of GEM-MACH does not currently include NO_x sources in the free troposphere, such as lightning and aircraft emissions.

395 3 HeiPro nartial column comparisons

In this section, we compare the HeiPro partial columns of NO₂ to (i) Pandora-DS tropospheric columns, (ii) TROPOMI tropospheric columns, and (iii) GEM-MACH partial columns. It is important to note that the comparisons are not entirely equivalent because HeiPro provides an integrated partial column from 0–4 km while TROPOMI and Pandora-DS provide tropospheric columns and GEM-MACH provides 0–5 km partial columns. Figure 3 displays the following scatter plots: HeiPro partial columns. Three types of linear regressions are presented in Fig. 3: the first is the York linear fit (York et al., 2004) in which the uncertainties in both datasets are incorporated into the regression, the second is the zero-intercept linear regression, and the third is the ordinary least squares linear regression (used only for the HeiPro vs. GEM-MACH regression since error estimates were not available for GEM-MACH data). The multiplicative bias is obtained from the zero-intercept slope and is presented as a deviation, in %, from a zero-intercept slope of 1 (100 × (zero-intercept slope-1)). The mean relative bias is (100 × (HeiPro-X)/X), where X is Pandora-DS, TROPOMI, or GEM-MACH.

3.1 HeiPro vs. Pandora-DS

Figure 3a shows the correlation between the HeiPro partial columns and Pandora-DS tropospheric columns. This comparison provides a good opportunity to study the differences between scattered sunlight and direct-Sun measurements by the same instrument. Although the direct-Sun and MAX-DOAS retrieval wavelength ranges are different due to the varying standard protocols for each, it is worthwhile to note that the HeiPro long vis versus long UV NO₂ partial column comparisons showed good agreement with one another, with a zero-intercept slope of 0.97 ±0.004 and mean relative bias of 0.7% ±5.9%

11

Deleted: 3

	-	- 1			-	-1	-	
1		0	0	т.	_	п	•	-
۰.	~	-				ч		

Deleted: P Deleted: C Deleted: C

Formatted: Font: Not Italic, Font color: Text 1	
Formatted: Font: Not Italic, Font color: Text 1	
Formatted: Font: Not Italic, Font color: Text 1	
Formatted: Font: Not Italic, Font color: Text 1	

(results not shown). We therefore do not expect the choice of retrieval window to significantly impact the HeiPro long UV partial column comparisons to Pandora-DS (see Table A1 for the HeiPro long vis partial column NO₂ comparisons). We find

- 420 that HeiPro partial columns, although measuring a smaller altitude range, are larger than Pandora-DS tropospheric columns, with a multiplicative bias of 51% ± 0.8% and a mean relative bias of 61% ± 9.7%. Additionally, the multiplear Pandora dataset presented here provides an opportunity to investigate seasonal differences between HeiPro and Pandora-DS. Monthly and hourly box-and-whisker plots are displayed in Fig. 4a and b, respectively. The top and bottom of each box represent the 75th and 25th percentiles, respectively, the horizontal line within is the median value, and the whiskers are the most extreme non-outlier data points, where outliers are values that are 1.5 times greater or less than the interquartile range. The mean relative
- bias between the datasets for each month of the year or hour of the day are depicted by the circled markers. Both HeiPro <u>(0-4</u> <u>km)</u> and Pandora-DS tropospheric column median values display typical seasonal and diurnal trends for tropospheric NO₂, whereby NO₂ is greater during the winter months and morning hours from 6–9 a.m. LT due to various chemical and dynamical factors such as increased emissions, shallow PBL, and photochemistry. The HeiPro partial columns have even larger seasonal
- 430 and diurnal trends compared to Pandora-DS tropospheric columns. This is further exemplified by the mean relative bias between the datasets, which exhibits a strong seasonal variability, with the largest values observed in the winter (84–114%) and some of the smallest values observed in the summer (39–51%). The larger relative bias during the winter months is also shown by a larger zero-intercept slope value during the winter (1.60 ± 0.02) compared to summer (1.38 ± 0.02), as well as a greater York linear fit intercept value during the winter (2.17×10¹⁵ molec cm⁻²) compared to summer (0.38×10¹⁵ molec cm⁻²).







Deleted: 77 Deleted: three

Deleted: partial

Formatted: Font: Not Italic, Font color: Text 1

445 regression (dashed red line), zero-intercept linear regression (dashed blue line), ordinary least squares regression (dashed magenta line), and the 1:1 line (dashed green line) are depicted. The color bar indicates the normalized density of the data points.

For reference, HeiPro (0–4 km) vs. Pandora-DS total columns are compared in panel (a) of Fig. A4, which shows that
HeiPro partial columns exhibit a positive multiplicative bias of 16% ± 0.7% and a mean relative bias of 6.1% ± 4.8%. Not surprisingly, there is better agreement here as compared to Fig. 3a (i.e., HeiPro vs. Pandora-DS tropospheric NO₂) since the Pandora-DS total columns are larger. The TROPOMI vs. Pandora-DS NO₂ total and tropospheric column comparisons are shown in Fig. A4b–c, respectively. Pandora-DS and TROPOMI show good agreement with one another for both total column (multiplicative bias: -12% ± 1.9%; mean relative bias: 0.1% ± 21%) and tropospheric NO₂ (multiplicative bias: -4.4% ±

- 455 3.5% ; mean relative bias: -0.9% ± 34%). Note that the large uncertainties are due to the relatively large TROPOMI total column and tropospheric NO₂ errors. Additionally, the tropospheric NO₂ agreement in panel (c) provides more confidence in the stratospheric-tropospheric separation method that was used in the study (i.e., Pratmo-OMI data). <u>TROPOMI total column</u> NO₂ at this measurement site has been studied and validated in Zhao et al. (2020). Using the version 1 data product, Zhao et al. (2020) found that TROPOMI vs. Pandora-DS total column NO₂ had a zero-intercept slope of 0.70 and correlation coefficient of 0.75. The version 2.3 data product used in this work showed an improvement from version 1, with a zero-intercept slope of 0.89 and correlation coefficient of 0.81. The time period of the study in which version 1 was used (March 2018 to March 2019)
- 0.89 and correlation coefficient of 0.81. The time period of the study in which version 1 was used (March 2018 to March 2019) was similar to that of this study (May 2018 to June 2020). Comparisons and validation of the newer version 2.3 TROPOMI data products are outside the scope of this work.

Mext, we explore various contributing factors to the generally large bias of HeiPro to Pandora-DS. We investigate how the PBL height contributes to the bias because the Pandora-DS measurements are missing the first ~15 m of the vertical column due to the rooftop instrument location while the HeiPro measurements attempt to include the first 15 m since the multi-axis viewing geometry has some sensitivity to lower layers and the HeiPro algorithm extrapolates the profile to ground level. This difference in detection may further be amplified by shallower PBL heights during winter months and morning hours, when the PBL height is smaller due to lower surface temperatures and less boundary layer dynamics. This leads to less vertical mixing of pollutants, with NO2 accumulating near the surface (e.g., Lin and McElroy, 2010; Chan et al., 2018; Schreier et al., 2019), where it is not captured by the Pandora-DS measurements in the first 15 m. We additionally investigate the effects of the SAA on the bias due to the spatiotemporal heterogeneity of NO2 as well as the varying airmasses measured by the multi-axis and direct-Sun viewing geometries. Lastly, while NO2 increases during the wintertime due to greater anthropogenic emissions from heat sources (e.g., Meng et al., 2018) and increased lifetimes due to decreased solar radiation, it is possible that increased emissions can contribute to the bias if there are more NO2 emissions coming from the multi-axis azimuth viewing direction of 255° compared to the various direct-Sun viewing angles.

Formatted: Font color: Text 1

Deleted:

(Deleted: since shallower PBL heights correspond to
(Deleted: , which
~(Deleted: do not capture
(Deleted: since
~(Deleted: emissions
(Deleted: increased
\mathcal{I}	Deleted: (
Ý	Deleted: and shallower PBL heights





Figure 4: (a) Monthly and (b) hourly box-and-whisker plots of HeiPro 0-4 km (orange) and Pandora-DS tropospheric (brown) NO₂ columns as well as the mean relative bias between the two (green circles). (c) Monthly and (d) hourly 490 box-and-whisker plots of HeiPro 0-4 km (orange) and GEM-MACH 0-5 km (blue) NO₂ columns as well as mean relative bias between the two (green circles). The top and bottom of each box represent the 75th and 25th percentiles, respectively, the horizontal line within is the median value, and the whiskers are the most extreme non-outlier data points, where outliers are values that are 1.5 times greater or less than the interquartile range.

495

The impact of the above factors on the bias is illustrated in Fig. 5a and b, which shows the zero-intercept slope, i.e., multiplicative bias, and mean relative bias, respectively, for HeiPro partial versus Pandora-DS tropospheric NO2 columns under the following scenarios versus the SAA range, with 20° bins per SAA range: (1) HeiPro partial vs. Pandora-DS standard tropospheric NO₂ columns, (2) HeiPro partial vs. Pandora-DS 'modified' tropospheric NO₂ columns, whereby a 0-15 m column is added to the Pandora-DS column, see Section 3.1.1 for further details, (3) HeiPro partial vs. Pandora-DS modified 500 tropospheric NO2 columns during summer and winter, (4) HeiPro partial vs. Pandora-DS standard tropospheric NO2 during winter only. In Sections 3.1.1–3.2.2, we quantify the contributions to the multiplicative and mean relative biases (see markers in Fig. 5c and d, respectively) between HeiPro and Pandora-DS NO2 from various factors and indicate the scenarios that lead to the best and worst agreement between the two at this site. The relative contribution is calculated as the absolute value of the percent change in the bias after incorporating a factor (e.g., the PBL contribution is calculated as: $100 \times (1)$ bias of HeiPro vs.

Pandora-DS modified – bias of HeiPro vs. Pandora-DS standard|)/ bias of HeiPro vs. Pandora-DS standard, at constant SAA range), and so any decreases in the bias when incorporating a factor such as PBL are reported as positive numbers and therefore as contributions to the bias.





515

520

Figure 5: (a) Zero-intercept slope and (b) mean relative bias for HeiPro partial vs. Pandora-DS (PDS) tropospheric NO₂ columns under various scenarios (see legend), with relative contributions (%) from various factors (see legend) to the (c) zero-intercept slope and (d) mean relative bias. The dashed yellow lines represent the Pandora azimuth viewing angle of 255° and the dashed blue lines represent the mean SAA during the TROPOMI overpass time of 204°. Each marker represents a 20° bin of SAA, with a range of values that precede and include the marker value.

3.1.1 PBL height effects

The effect of the missing 0–15 m in the Pandora-DS tropospheric column can be enhanced when the PBL height is shallow. Overall, we found that this partially contributed to why the HeiPro partial NO₂ columns were larger than the Pandora-DS tropospheric columns but was not able to fully account for the biases observed. This was investigated by producing an approximate 0–15 m column, which was calculated using the NAPS in situ surface NO₂ value taken as an indication of the

first 15 m of the NO2 profile at the measurement location, calculated as:

15

Deleted: H

Deleted: E

$$0 - 15m \ column = \frac{h \times VMR \times n_A \times P}{R \times T \times 10^9 \ molec. \ air} \qquad Equation (1)$$

where *h* is the height of the column (15 m), *VMR* is the volume mixing ratio of NO₂ in ppbv $\left(\frac{X \text{ molecules of NO}_2}{10^9 \text{ molecules of air}}\right)$, n_A is Avogadro's number, *P* is the surface pressure in *atm*, *R* is the ideal gas constant (8.21 × 10⁻⁵m³ · atm · K⁻¹ · mol⁻¹), and *T* is the surface temperature in *K*. Surface pressure and temperature values were obtained from ERA5 reanalysis data for each

- 530 hour corresponding to the in situ measurement of NO₂ VMR. This column was then added to the Pandora-DS tropospheric column to create a 'modified' column and the biases between HeiPro and Pandora-DS were re-examined using the modified column. The effects of the missing 15 m and the PBL height on the bias can be quantified by looking at how the multiplicative bias and mean relative bias change from the HeiPro partial versus Pandora-DS standard tropospheric columns to the HeiPro partial versus Pandora-DS modified tropospheric columns. The most notable reductions in the biases are seen when the SAA
- 535 range is 66°–85°, although large multiplicative biases and mean relative differences of > 50% remain at this SAA range. Such reductions in the multiplicative and relative biases (HeiPro partial column vs. Pandora-DS tropospheric column HeiPro partial column vs. Pandora-DS modified tropospheric column) are depicted by the circles in Fig. 5c and d and termed the "PBL contribution", respectively. The largest reductions of 8.4% in the multiplicative bias and 18% in the mean relative bias occur at the SAA range of 66°–85°, which corresponds to the morning hours of 6–8 a.m. LT. This is unsurprising since the PBL
- height is shallow during this time, which presents a greater opportunity for any NO₂ accumulating in the first 15 m to be missed by the Pandora-DS measurements. The impact of accounting for the missing 15 m then drops to negligible amounts throughout the day as the SAA increases, presumably due to a more well-mixed PBL, Because the 0–15 m column derivation assumes a constant NO₂ VMR from 0 to 15 m, the estimations provided here of the PBL contribution to the bias represent upper limits (aside from cases with lofted plumes below 15 m). In cases where the NO₂ VMR at 15 m <<u>NO₂ VMR at 0 m</u>, the contribution 545 of the 0–15 m column would be lower, and therefore, the PBL contribution to the bias would be lower.
- --

3.1.2 NO₂ heterogeneity contribution

Another potential factor contributing to the large bias of HeiPro to Pandora-DS is the three-dimensional heterogeneity of the NO₂ field combined with the fact that multi-axis and direct-Sun viewing geometries inherently measure different airmasses. Importantly, the direct-Sun and multi-axis viewing geometries point to different azimuth angles, with direct-Sun tracking the Sun and multi-axis constantly pointing to 255°, which also contributes to differences in the airmasses being probed due to the spatial heterogeneity of NO₂. To investigate how differences in viewing angles contributes to the bias, the zerointercept slope and mean relative bias between HeiPro partial NO₂ columns versus Pandora-DS modified tropospheric NO₂ columns at each SAA range were compared to the bias at 246°–265°, i.e., Pandora-DS measurements within ±10° of the MAX-DOAS HeiPro measurements. This bias difference is represented by the diamond markers in Fig. 5c and d and termed the Formatted: Font: Not Italic, Font color: Text 1
Formatted: Font: Not Italic, Font color: Text 1

Deleted: H

Formatted: Font color: Text 1

Deleted: C

16

"SAA contribution". The SAA contribution was not calculated for the SAA ranges of 266°–285° since the HeiPro bias relative to Pandora-DS in these ranges is smaller than the bias at the reference SAA range of 246°–265°.

- 560 The largest zero-intercept slope and mean relative biases (1.85 and 91%, respectively) are seen when the direct-Sun and multi-axis viewing angles are farthest apart during the measurement day in the hours from 6–8 a.m. LT. The zero-intercept slope then drops from 1.85 at the 66°–85° SAA range to 1.36 at the 246°–265° SAA range, while the mean relative bias drops from 91% to 44% for these ranges, representing 27% and 52% reductions in the multiplicative and mean relative biases, respectively. These large contributions are during the morning hours when the direct-Sun and multi-axis viewing geometries
- 565 are farther apart, and the SAA contribution to the bias decreases to negligible values as the SAA increases. It is important to note that even when Pandora-DS measurements are within ±10° of the HeiPro measurements, the biases are still large, with a zero-intercept slope of 1.30 and mean relative bias of 44%. One contribution to this bias may be the 10° differences between direct-Sun and multi-axis viewing angles, which may still exhibit NO₂ heterogeneity, as well as the different horizontal sensitivities between the direct-Sun and multi-axis viewing geometries.

570

Additionally, since the mean relative bias between HeiPro partial versus Pandora-DS tropospheric columns was at a maximum during winter months and minimum during summer months (see Fig. 4a), seasonal effects on the bias were investigated. The zero-intercept slope and mean relative bias between HeiPro partial versus Pandora-DS modified tropospheric columns were plotted against the SAA range for summer months (JJA) and winter months (DJF, see Fig. 5a and b), and are

- 575 both larger during the winter months compared to summer months across all SAA ranges, which is consistent with the mean relative bias shown in Fig. 4a. The differences in the biases of HeiPro partial versus Pandora-DS tropospheric columns between winter and summer at each SAA range are depicted by the crosses in Fig. 5c and d and termed the "seasonal contribution". Since there are no Pandora wintertime measurements for the first two SAA ranges and the last SAA range, seasonal contributions cannot be assessed for those hours of the day (at 6 a.m. and onwards from 6 p.m., LT). The seasonal contribution
- 580 to (i) the zero-intercept slope ranges from 16% to 39% and to (ii) the mean relative bias ranges from 25% to 85%, depending on the SAA range. The seasonal contribution peaks when the SAA range is between 126°–165°, and then decreases as the SAA increases. Due to increased NO₂ emissions during the winter, it is possible that the seasonal contribution stems from increased emissions from the multi-axis azimuth viewing direction of 255°, and so HeiPro measurements, and subsequently the biases towards Pandora-DS, are even larger during winter months.
- 585

590

Next, we investigate the spatiotemporal NO₂ heterogeneity around Downsview and assess whether we can observe the heterogeneity that supports the biases observed (i.e., large differences in the NO₂ columns between the azimuth viewing angle and the various SAAs, and even larger differences between the two from 6–9 a.m. LT and winter months). To do this, the NO₂ field around Downsview was explored using two datasets from 2018–2020: (i) TROPOMI tropospheric NO₂ columns and (ii) GEM-MACH (0–5 km) NO₂ columns. The TROPOMI pixel-averaged (Fioletov et al., 2011; Sun et al., 2018) NO₂

field was used to investigate the larger bias in winter, while the mean GEM-MACH NO₂ field was used to investigate the larger bias in the early morning hours since TROPOMI only provides data at 13:30 LST. We found that using such datasets to demonstrate that there was NO₂ heterogeneity between the multi-axis and direct-Sun viewing directions was insufficient because (i) TROPOMI tends to underestimate NO₂ in polluted regions and is not as suited to capture small local NO₂ enhancements as MAX-DOAS measurements are (Verhoelst et al., 2021, see Section 3.2 for further discussion), and (ii) the

GEM-MACH model resolution of 10 km \times 10 km may also not capture local enhancements and it is difficult to use model inventories as an interpretation of actual conditions since the model utilizes inventories from the year 2013. See Fig. A⁵₂ and A⁶₄ in the Appendix for the TROPOMI and GEM-MACH NO₂ fields, respectively, and for further discussion.

- To summarize, the smallest biases in the dataset are observed during the summer months, when using Pandora-DS modified tropospheric columns and when the SAA > 125°, which exhibit multiplicative biases of $7.2\% \pm 3.4\%$ to $31\% \pm 2.0\%$ and mean relative biases of $14\% \pm 9.0\%$ to $35\% \pm 11\%$. We term these combined conditions the "best-case scenario", which constitutes the following: a well-mixed PBL, accounting for the missing 15 m, smaller viewing angle differences, and summer months with less NO₂ emissions. Therefore, when conditions are suitable for agreement, HeiPro partial columns show
- 605 good agreement with Pandora-DS tropospheric NO₂ columns. Conversely, the largest biases are observed when using Pandora-DS standard columns (i) during the winter months throughout the measurement day (see HeiPro partial column bias to Pandora-DS standard columns in Fig. 5a and b), which exhibit multiplicative and mean relative biases mostly greater than 50% and 70%, respectively, and (ii) during the early morning hours across all seasons (see HeiPro partial column bias to Pandora-DS standard columns when the SAA ≤ 105° in Fig. 5a and b), which exhibit multiplicative biases of 80% ± 2.6% to 102% ±
- 610 3.1% and mean relative biases of 86% ± 10% to 110% ± 12%. We term the conditions outlined in (i) and (ii) as the "worst-case scenario". Therefore, some portion of the large bias of HeiPro to Pandora-DS may be explained by the following environmental conditions: shallow PBL, the missing 15 m in the Pandora-DS measurements, larger viewing angle differences, and increased NO₂ heterogeneity during the winter. Scatter plots for the best-case (multiplicative bias: 19% ± 1.3%; mean relative bias: 24% ± 9.7%) and worst-case (multiplicative bias: 72% ± 1.7%; mean relative bias: 101% ± 9.4%) subsets of
- 615 the data are presented in Fig. A 7 and b, respectively. Although the biases are minimized in the best-case scenario, nonzero biases remain. This is shown in Fig. 5a and b by the offset from the dotted grey line of the HeiPro partial column bias towards Pandora-DS modified tropospheric columns during the summer months when the SAA > 125°). Possible contributing factors to the remaining bias are discussed in Section 5.

3.2 HeiPro vs. TROPOMI

620

595

Figure 3b shows the scatter plot for HeiPro partial versus TROPOMI tropospheric NO₂ columns, with HeiPro exhibiting multiplicative and mean relative biases of $17\% \pm 4.0\%$ and $37\% \pm 51\%$, respectively, compared to TROPOMI measurements. The relatively large uncertainty in the mean relative bias can be attributed to the larger retrieval errors in the TROPOMI tropospheric NO₂ measurements. Our findings regarding the differences between HeiPro and TROPOMI are in

18

Deleted: 3

Deleted: 4

Deleted: 5

Deleted: 41

Deleted: 47

accordance with a study by Verhoelst et al. (2021), who categorized the TROPOMI bias relative to ground-based MAX-DOAS
NO2 into the following regimes: (1) regions with low pollution levels can have median relative differences (100 × (TROPOMI–MAX-DOAS)/MAX-DOAS) of up to -27%, (2) regions with moderate pollution levels exhibit biases between -15% to -56%, and (3) for extremely polluted regions, differences of -37% to -74% are seen. The pollution level categories were based on the median MAX-DOAS ground-based tropospheric columns. As Toronto is a moderately polluted region with a median HeiPro 0–4 km partial column of 7.71×10¹⁵ molec cm⁻², our findings are in accordance with the bias range in the moderately polluted categorization of Verhoelst et al. (2021), i.e., the median bias (TROPOMI–HeiPro) within this dataset is -1.4 ×10¹⁵ molec cm⁻², with a median relative bias of -27%, as per the equation used in the study. Factors contributing to the discrepancy between TROPOMI and MAX-DOAS NO₂, and therefore also the limitations of using the TROPOMI NO₂ field to address the HeiPro overestimation of Pandora-DS, are discussed next.

- 640 Firstly, MAX-DOAS measurements can capture more local enhancements in NO₂, while the satellite retrievals provide a smoothed pixel representation. For example, satellite underestimation of MAX-DOAS measurements can occur if the footprint of the emission source is smaller than the satellite footprint of 3.5 km × 5.5 km for TROPOMI (3.5 km × 7.5 km prior to August 2019, Verhoelst et al., 2021). Additionally, the TROPOMI retrieval algorithm is sensitive to the a priori NO₂ profile shape, and the use of low-resolution a priori NO₂ profiles in the TROPOMI retrieval algorithm can contribute to the 645 underestimation of ground-based MAX-DOAS measurements, as shown in various studies (Zhao et al., 2020; Dimitropoulou et al., 2020). For example, Zhao et al. (2020) showed a 10% reduction in the bias between TROPOMI and Pandora-DS NO₂ total columns when replacing the standard a priori profile in the TROPOMI retrieval algorithm with GEM-MACH, a profile from a high-resolution regional air quality forecast model.
- 650 The TROPOMI overpass time of 13:30 LST occurs at a time of day such that the HeiPro biases relative to Pandora-DS are at the lower end of the bias range (see where markers intersect with the dashed line at 204° in Fig. 5a and b). It would be interesting to observe how the bias of geostationary NO₂ measurements (e.g., TEMPO) relative to ground-based direct-Sun and multi-axis measurements changes throughout the day, given the large seasonal and diurnal dependency of the HeiPro bias relative to Pandora-DS. The results of the HeiPro comparisons to Pandora-DS in this study can therefore aid in future ground-
- 655 based direct-Sun and MAX-DOAS validation studies of TEMPO by providing possible explanations for differences in the bias between TEMPO versus MAX-DOAS and TEMPO versus direct-Sun at this measurement site.

3.3 HeiPro vs. GEM-MACH

Here we compare the HeiPro NO₂ partial columns to GEM-MACH. Figure 3c shows the regression between HeiPro 660 (0-4 km) and GEM-MACH (0-5 km) partial columns, While the GEM-MACH standard dataset includes partial columns from 0-5 km, we did not generate a 0-4 km partial column because we found the GEM-MACH NO₂ VMRs from 4-5 km to be very

Formatted: Font color: Text 1 Formatted: Font: Not Italic, Font color: Text 1



	small, and that the integrated NO ₂ from 4–5 km constituted only 0.3% of the GEM-MACH (0–5 km) partial columns, HeiPro
	is greater than GEM-MACH with a multiplicative bias of $\frac{12}{5} \pm \frac{12}{5}$ and a mean relative bias of $\frac{67\%}{5} \pm 7.1\%$. Figure 4c
	and d display monthly and hourly, respectively, box-and-whisker plots of both datasets as well as the mean relative bias
665	between them. As with comparisons of HeiPro to Pandora-DS, the HeiPro dataset exhibits a stronger annual cycle than GEM-
	MACH, with a larger interquartile range and whiskers during the winter months. The seasonal trend appears weaker in the
	GEM-MACH box-and-whisker plots, which do not seem to capture the seasonal pattern in NO2. Such differences in seasonal
	patterns between GEM-MACH and HeiPro are further reflected in the monthly mean relative biases (see circled markers), in
	which the winter months exhibit some of the largest percent differences (95-145%) compared to the summer months (38-
670	53%). Also similar to the HeiPro and Pandora-DS hourly comparisons, the HeiPro versus GEM-MACH hourly comparisons
	exhibit the largest (107-162%) mean relative biases during the morning hours (6-9 a.m. LT), with values generally decreasing
	throughout the day. The large bias of HeiPro to GEM-MACH NO2 can partially be explained by the NO2 inventories used in
	the GEM-MACH model. The model inventories account for surface NO2 emissions in the PBL, but do not include lightning
	and aircraft emissions of NO2 in the free troposphere. This is further demonstrated in the NO2 profile and surface comparisons
675	in Section 4

Due to the similarly large biases in HeiPro partial columns relative to Pandora-DS and to GEM-MACH, as well as similar seasonal and diurnal patterns in these biases, a comparison was done between Pandora-DS and GEM-MACH (see Appendix, Fig. A. The scatter plot (Fig. A. (a)) shows a zero-intercept slope of 1.00 (i.e., no apparent multiplicative bias), indicating that there is good agreement between Pandora-DS tropospheric and GEM-MACH (0–5 km) partial NO₂ columns, although a positive mean relative bias (100 × (GEM-MACH– Pandora-DS)/Pandora-DS) of 20% exists. The positive mean relative bias, indicating that GEM-MACH values are greater than Pandora-DS, is also evident in the mean monthly and hourly box-and-whisker plots in Fig. A. (a) and c, respectively, which show that GEM-MACH is greater than Pandora-DS, particularly in the afternoon hours with an approximate mean relative bias of 30%.

685 4 HeiPro surface NO2 & profile comparisons

This section presents the surface NO₂ comparisons between HeiPro versus (i) NAPS in situ and (ii) GEM-MACH, as well as NO₂ profile comparisons between HeiPro and GEM-MACH. The surface comparisons are presented in Section 4.1 while the profile comparisons are presented in Section 4.2. GEM-MACH is used for the profile comparison as it is the only source of NO₂ profile data available at this site. The multiplicative bias is as described in Section 3 and the mean relative bias 690 is (100 × (HeiPro-X)/X), where X is NAPS in situ or GEM-MACH.

4.1 Surface NO₂ comparisons

Figure 6 shows the scatter plots of HeiPro versus (a) NAPS in situ surface NO₂ and (b) GEM-MACH surface NO₂, whereby HeiPro surface NO₂ exhibits negative multiplicative biases of $-\frac{5}{2}\frac{80}{2} \pm 0.7\%$ and $-\frac{41}{2}\% \pm 0.5\%$ to NAPS in situ

Formatted. Fort. Not Italic, Fort Color. Text I
Formatted: Font color: Text 1
Deleted: 0
Deleted: 84
Deleted: 1
Deleted: 5

. . .

Net Italia Cont

Deleted: 74	
Deleted: 32	
Deleted: 23	
Deleted: 6	
Deleted: 99	
Deleted: 75	

Formatted: Font: Italic, Font color: Red, English (CAN)

Deleted: 6 Deleted: 6

Deleted: 7

-(Deleted: S	
•(Deleted: P	
4	Deleted: C	_

Deleted: C	
Deleted: 4	
Deleted: 4	
Deleted: 1	
Deleted: 0	
Deleted: 4	
Deleted: 7	

2	A
2	υ

and GEM-MACH, respectively. Likewise, HeiPro exhibits mean relative biases of $-9.7\% \pm 7.5\%$ and $-37\% \pm 2.4\%$ to NAPS in situ and GEM-MACH, respectively. Figure 6c–f shows box-and-whisker plots of the seasonal diurnal trends of HeiPro, GEM-MACH, and NAPS in situ surface NO₂ (box-and-whisker plots are as defined in Section 3.1). All datasets

- 720 display peaks in surface NO₂ around 6–8 a.m. LT and decrease throughout the day, capturing the diurnal trend of surface NO₂, which is shaped by emission intensity, PBL dynamics, and photochemistry. Additionally, the small negative bias between HeiPro and NAPS in situ is evident in the diurnal plots throughout most of the measurement day and across all seasons. On the other hand, the larger negative bias of HeiPro to GEM-MACH is evident in the diurnal plots throughout the measurement day, particularly during the morning hours.
- 725

Across all seasons, the largest surface NO₂ values are from the GEM-MACH data from 6–9 a.m. LT. The GEM-MACH surface NO₂ is likely to be overestimated due to an outdated NO₂ inventory being used in the model, which does not account for reduced NO₂ emissions over the years or reduced commutes during the COVID-19 lockdown periods in 2020 (Zhao et al., 2022). Additionally, all three datasets capture a stronger diurnal trend on weekdays (Fig. 6g) compared to the

730 diurnal trend on weekends (Fig. 6h), which shows smaller median values as well as interquartile and whisker ranges. It is important to note that the HeiPro seasonal diurnal surface NO₂ values, as well as weekday versus weekend trends, are more closely aligned to NAPS in situ surface NO₂ than GEM-MACH is to NAPS in situ surface NO₂. HeiPro provides a reasonable estimate of surface NO₂ values as well as the seasonal diurnal pattern of surface NO₂ but tends to underestimate the intensity during the evening hours. Contributions to the discrepancy between HeiPro and NAPS in situ surface NO₂ are discussed next.

Deleted: 8.7	
Deleted: 5	
Deleted: 3	
Deleted: 6	





740 Figure 6: Comparisons of surface NO₂ VMR (2018–2020) for HeiPro vs. (a) NAPS in situ, and (b) GEM-MACH. The dashed lines and color bar are as indicated in Fig. 3. (c)–(f) Seasonal diurnal and (e), (f) weekday, weekend box-and-whisker plots of HeiPro (orange), GEM-MACH (blue), and NAPS in situ (black) surface NO₂ VMRs. Box-and whisker values are as defined in Fig. 4.

745

750

Due to the vertical offset of 11 m between the Pandora and in situ instruments (the former on the rooftop at 15 m and the latter at 4 m above ground level) and the heterogeneity of the NO₂ field, PBL height was investigated as a potential source of discrepancy between HeiPro and NAPS in situ surface NO₂. The relative bias of HeiPro to NAPS in situ (100 × (HeiPro–NAPS in situ)/NAPS in situ) versus PBL height range is presented in the box-and-whisker plots of Fig. 7. The boxand-whisker values are as defined in Section 3.1. The PBL range were chosen to ensure that the number of data points in each bin were in the same order of magnitude. For the shallowest PBL range from 0–0.40 km, the upper and lower extents of the boxes and whiskers indicate that, at times, HeiPro is greater than NAPS in situ surface NO₂, while at other times, HeiPro is

smaller than NAPS in situ. Although the HeiPro profiles have some sensitivity to the 15 m of the atmospheric column below the instrument altitude, the vertical offset of the Pandora instrument may impact its sensitivity to those 15 m. It is not known

- 755 how the HeiPro surface value would change if the instrument were on ground level or had negative elevation viewing angles (and therefore more sensitivity to the first 15 m). Nonetheless, it appears that the shallowest PBL range exhibits a larger range of differences but smaller (less negative) median values (e.g., for summer, whisker range: -128% to 195%; median value: -10%) compared to the largest PBL range (e.g., for summer, whisker range: -119% to 22%; median value: -51%), and so the vertical offset between the instruments may contribute to the larger range of differences observed when the PBL height is
- 760 shallow and there is more vertical NO₂ heterogeneity. In addition to the narrower whisker range for PBL heights > 0.90 km, this PBL range consistently also has median values of approximately -50% across all seasons (see horizontal lines within each box in Fig. 7). At this PBL range, which corresponds to the afternoon periods when surface NO₂ is at a minimum, the airmass measured by both the Pandora and NAPS instruments is more uniform, which may contribute to a more consistent difference between the two, hence the narrower whisker range. Additionally, the NAPS instrument measures a very localized
- airmass due to the nature of the in situ technique, while the HeiPro measurements stem from an instrument field-of-view of
 1.6° (Herman et al., 2009) and a photon effective pathlength between 5 and 10 km for UV measurements (Ortega et al., 2015).
 These varying horizontal sensitivities between HeiPro and NAPS in situ, combined with spatiotemporal NO₂ heterogeneity, can also contribute to differences in the airmasses being measured.

(Deleted:



770



4.2 Profile comparisons

Figure 8 shows the seasonal median NO₂ profiles from HeiPro, GEM-MACH, and GEM-MACH smoothed by the
HeiPro averaging kernel. The shaded regions represent the 90th percentile (right of solid line) and 10th percentile (left of solid line) values for HeiPro and GEM-MACH but are not shown for GEM-MACH smoothed for visual clarity purposes. The seasonal median NAPS in situ surface NO₂ values are also displayed for reference. As stated previously, the surface values for HeiPro and GEM-MACH were extrapolated from the midpoint of the grid level closest to the surface (e.g., 0–200 m for HeiPro), for each available profile. Due to the presence of some lofted layers in the HeiPro profiles, in which the 0–200 m grid level NO₂ VMR is less than that of the 200–400 m grid level, the extrapolated surface VMR for such cases is smaller than that of the grid level closest to the surface does not have a sharper gradient towards the surface (see HeiPro profiles in Fig. 8), although this is only because the median values are plotted; aside from profiles with lofted layers, each individual profile has a larger increase from the 100 m layer to the surface were reference. NO₂

Deleted: C

Deleted: then

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Deleted: When plotting the median profiles, as was done in Fig. 8, this appears to skew the profile shape (see HeiPro profiles near 0 m in Fig. 8), and it appears that the extrapolation was not linear with respect to the near-surface values, though it is the case that each profile extrapolation to the surface was linear from the grid level closest to the surface.

Formatted: Font color: Text 1

values during the winter months (5.7, 7.8, and 10 ppbv, respectively) compared to spring (4.2, 5.4, and 7.7 ppbv), summer (2.7, 4.0, and 6.5 ppbv), and autumn (4.3, 5.7, and 8.1 ppbv). This is in accordance with winter conditions that increase surface
NO₂ such as larger anthropogenic emissions, meteorological conditions, etc.



Figure 8: HeiPro (solid orange line), GEM-MACH (solid blue line), and GEM-MACH smoothed (dotted blue line) median NO₂ profiles for each season. The shaded regions represent the 90th percentile (right of line) and 10th percentile (left of line) values for HeiPro and GEM-MACH. The black triangles indicate the seasonal median NAPS in situ surface NO₂ values.

800

Across all seasons, the HeiPro NO2 median profiles from 0-200 m underestimate the unsmoothed GEM-MACH median values, while from 1.5-4 km, the HeiPro median profiles then overestimate the unsmoothed GEM-MACH median values. For the 0-200 m layer, the mean relative bias of HeiPro towards GEM-MACH decreases from -37% (unsmoothed) to -6.1% (smoothed). Note that these biases are representative of the integrated 0-200 m layer and may differ slightly from the surface values reported in Table 2. The most significant changes occur in the layer from 1.5-4 km, where the HeiPro bias towards GEM-MACH decreases from > 1000% (unsmoothed) to 2.6% (smoothed). The HeiPro surface underestimation and free tropospheric overestimation of the unsmoothed GEM-MACH profiles can probably be explained by the NO2 inventories used in the GEM-MACH model, which, respectively, (i) utilize older inventories that do not account for reduced emissions over the years, and (ii) do not account for free tropospheric NO2 sources while the a priori NO2 profile contains free

Formatted: Font color: Text 1
Formatted: Font color: Text 1, Subscript
Formatted: Font color: Text 1
Formatted: Font color: Text 1
Deleted: surface and near-surface NO ₂
Deleted: from 0–0.80 km
Deleted: . At an altitude of approximately
Deleted: 0.80
Deleted: and above
Deleted:
Formatted: Font color: Text 1
Formatted: Font color: Text 1
Formatted: Font color: Text 1
Deleted: profiles
Deleted: .
Deleted: As stated previously, this is likely due to
Formatted: Font color: Text 1
Deleted:

25

tropospheric NO2. For the layers above 200 m and below 1 km, there is no clear pattern between HeiPro and GEM-MACH.

Although it is difficult to assess the accuracy of the HeiPro NO₂ profiles without the availability of in situ NO₂ profiles at this site, it is worthwhile to note that HeiPro seems to provide a conservative estimate of the NAPS in situ value at the surface, where NO₂ VMRs are typically the largest. Additionally, while the GEM-MACH surface NO₂ VMRs are appropriately larger in the winter, this trend is not similarly captured for the GEM-MACH partial columns (see box-and-whisker plots in Fig. 4c).

On the other hand, both the HeiPro surface NO₂ and partial columns display increased values during winter months.

5 Conclusions

825

This study produced a three-year dataset of NO₂ profiles and partial columns from 0-4 km at Downsview, Ontario by applying HeiPro, an optimal estimation profile retrieval algorithm, to Pandora UV MAX-DOAS measurements. Table 2

- 830 provides a summary of all the intercomparisons made using the HeiPro NO₂ data products at this measurement site. The HeiPro partial NO₂ columns obtained from the profiles were found to overestimate the partial columns from Pandora-DS, TROPOMI, and GEM-MACH, with multiplicative biases of 51%, 17%, and 12%, respectively, and mean relative biases of 61%, 37%, and 67%, respectively. The seasonal and diurnal trends in the mean relative biases between HeiPro and Pandora-DS as well as HeiPro and GEM-MACH were similar, with larger mean relative biases during the winter months and morning hours from 6–
- 835 8 a.m. LT. Additionally, HeiPro partial columns exhibit larger variability, as evidenced by the larger box-and-whisker ranges in Fig. 4a and b, compared to Pandora-DS; this is consistent with a study by Pinardi et al. (2020), who found that MAX-DOAS measurements tend to depict a fuller range of NO₂ variability due to their ability to measure under partially cloudy conditions, while direct-Sun measurements require clear skies. Although Fig. 4 presents hourly-averaged coincident measurements between HeiPro and Pandora-DS, the measurements are not perfectly coincident during that hour that is averaged, which can
- 840 also contribute to the differences between the direct-Sun and MAX-DOAS datasets. Seasonal and diurnal comparisons to TROPOMI were not possible due to limited datapoints during winter months and there being one measurement per day, respectively. The TROPOMI bias to HeiPro reported in this study (-27%) matched the bias range reported by Verhoelst et al. (2021) for a moderately polluted region, and it would be interesting to investigate how this satellite bias relative to groundbased MAX-DOAS measurements changes throughout the day using a geostationary satellite data product such as that of TEMPO (Terement et al. 2017).
- 845 TEMPO (Zoogman et al., 2017).

 Table 2. Summary of the multiplicative biases and mean relative biases (± uncertainties) for the HeiPro

 comparisons to partial columns and surface NO2.

HeiPro (0-4 km) vs. TROPOMI tropospheric NO2 51% ± 0.8% 61% ± 9.7% Deleted: 7 HeiPro (0-4 km) vs. TROPOMI tropospheric NO2 17% ± 4.0% 37% ± 51% Deleted: 7 HeiPro (0-4 km) vs. GEM-MACH (0-5 12% + 1.2% 67% + 7.1% Deleted: 7	Datasets compared	Multiplicative Bias (%)	Mean Relative Bias (%)	Eormatted Table
HeiPro (0-4 km) vs. Pandora-DS 51% ± 0.8% 61% ± 9.7% Deleted: 7 HeiPro (0-4 km) vs. TROPOMI 17% ± 4.0% 37% ± 51% 17% ± 4.0% 17% ± 4.0% 17% ± 51% leiPro (0-4 km) vs. GEM-MACH (0-5 12% ± 1.2% 67% ± 7.1% 12% ± 1.2%		Munipicative Dias (70)	incun renuive Blus (70)	ronnatted Table
tropospheric NO2 17% ± 4.0% 37% ± 51% HeiPro (0-4 km) vs. GEM-MACH (0-5 12% + 1.2% 67% + 7.1%	HeiPro (0-4 km) vs. Pandora-DS	$51\% \pm 0.8\%$	$61\% \pm 9.7\%$	Deleted: 7
Heirro (0-4 km) vs. 1 KOPOWI $17\% \pm 4.0\%$ $37\% \pm 51\%$ tropospheric NO ₂ $12\% \pm 1.2\%$ $67\% \pm 7.1\%$	tropospneric NO ₂	-		
tropospieric NO2 HeiPro (0-4 km) vs. GEM-MACH (0-5 12% + 1.2% 67% + 7.1%	HeiPro (0-4 km) vs. 1 ROPOMI	$17\% \pm 4.0\%$	$37\% \pm 51\%$	
Heirro (0-4 km) vs. GEM-MACH (0-5 $12\% + 1.2\%$ $67\% + 7.1\%$	Uropospheric NO ₂			
	HeiPro (0-4 km) vs. GEM-MACH (0-5	$12\% \pm 1.2\%$	$67\% \pm 7.1\%$	

26

Deleted: HeiPro measurements have some sensitivity to such layers, and (ii) utilize older inventories that do not account for reduced emissions over the years.

Deleted: that discrepancies between HeiPro and GEM-MACH profiles can be explained by the model inventories, and

Deleted: While the unsmoothed GEM-MACH profile is biased high/low near the surface/above 1 km compared to HeiPro due to the model inventory, the smoothed GEM-MACH profile shows better agreement with HeiPro as it accounts for the MAX-DOAS measurement limitations and vertical sensitivity.

Deleted: 1	
Deleted: 9.0	
Deleted: 41	
Deleted: 1	

HeiPro vs. NAPS in situ surface NO2	$-5.8\% \pm 0.7\%$	- <u>9.7</u> % ± 7.5%
HeiPro vs. GEM-MACH surface NO2	$-4\underline{1}\% \pm 0.\underline{5}\%$	$-3\underline{7}\% \pm 2.4\%$
HeiPro vs. GEM-MACH-smoothed surface NO ₂	$-\underline{3}0\% \pm \underline{0.8}\%$	$-6.0\% \pm 3.4\%$

- 865 We found that the HeiPro bias relative to Pandora-DS can partially be explained by several factors, which vary in their contributions to the bias throughout the day. The PBL height, combined with the missing 0–15 m partial column in the Pandora-DS measurements, contributed a maximum of 8.4% of the multiplicative bias and 18% of the mean relative bias in the morning hours, with these values declining to < 5% in the evening hours. The differences between the direct-Sun and multi-axis azimuthal viewing angles throughout the day, combined with the spatiotemporal heterogeneity of the NO₂ field, contributed a maximum of 27% and 52% of the multiplicative and mean relative biases, respectively, when the two viewing angles were farthest apart, and declined throughout the day as the direct-Sun azimuthal viewing angle approached the multi-axis one. Lastly, the maximum seasonal contributions to the multiplicative and mean relative biases ranged from 39% and 85% respectively, with systematically larger biases during the winter months. We were not able to assess seasonal contributions at certain SAA ranges during the wintertime since there were no measurements during these hours, i.e., at 6 a.m. and onwards
- 875 from 6 p.m. LT. We utilize best-case (multiplicative bias: 19%; mean relative bias: 24%) and worst-case (multiplicative bias: 72%; mean relative bias: 101%) scenarios to exemplify how these contributing factors affect the bias, and note that when these factors are minimized, the bias decreases and the agreement between HeiPro and Pandora-DS significantly improves.
- There are various possible contributions and sources of uncertainty to the remaining bias. (1) The HeiPro and Pandora-880 DS comparisons use hourly averages and are not perfectly temporally coincident. This, combined with NO₂ variability, may contribute to the scatter within the data. (2) The HeiPro data stems from NO2 retrieved in the UV window (338-370 nm) while the Pandora-DS data stems from NO₂ retrieved in the 400-440 nm range, so differences in the spectroscopic analysis may be a factor but we do not expect this to be a large contribution due to the similar agreement of HeiPro long UV vs. long vis partial columns (see Table A1). (3) Uncertainties in the Pratmo model that was used to calculate OMI stratospheric NO2 throughout 885 the day can overestimate or underestimate Pandora-DS tropospheric NO2, thereby underestimating or overestimating the bias, respectively. We do not believe that the use of OMI stratospheric NO2 itself contributed to the large bias, since, for example, TROPOMI stratospheric NO2 was greater than OMI stratospheric NO2 at this site. (4) The AOD retrieved from HeiPro is used as a parameter for the radiative transfer model of the trace gas profile retrieval as it helps to constrain the atmospheric light path. Inaccuracies in this retrieved AOD can therefore lead to errors in the NO2 retrievals. (5) The ERA5 temperature profile 890 data utilized in the HeiPro algorithm may underestimate measurements, which can contribute 3% to the remaining bias (as demonstrated by a sensitivity test that was performed with HeiPro and variable temperature inputs, results not shown). (6) The a priori NO2 profile used in the retrievals may also contribute to the HeiPro bias towards Pandora-DS. Although a conservative

Deleted: for

Deleted: Although not quantified,	
Deleted: t	
Deleted: is likely	
Deleted: contributing	

Deleted:

surface VMR of \sim 3.5 ppbv was used for the a priori profile, it is difficult to know if the upper layers of the a priori profile are 27 900 conservative estimates, given the limited in situ profile information at this site. It is possible that the VMR values in the upper layers of the a priori profile are larger than the true values, which would contribute to the larger HeiPro bias relative to Pandora-DS.

Additionally, we investigated whether the NO₂ heterogeneity that we believe is contributing to the HeiPro bias to 905 Pandora-DS can be supported by TROPOMI and GEM-MACH NO₂ fields at the measurement site. Overall, we were not able to utilize these datasets to show strong NO₂ heterogeneity between the two viewing geometries during the early morning hours and during the winter months. This suggests that satellite measurements and model output may not capture small-scale NO₂ enhancements: TROPOMI provides a smoothed representation of the NO₂ field and lacks sensitivity to the boundary layer due to the use of low-resolution a priori profiles, while GEM-MACH may not accurately represent real-time NO₂ gradients and 910 intensities. Pandora MAX-DOAS measurements may therefore provide a better tool for probing lower tropospheric NO₂ and its heterogeneity around a measurement site since the sensitivity and temporal coverage of satellites are limited, in situ measurements are spatially limited, and the direct-Sun measurement viewing geometry is subject to the Sun's position and

- 915 While HeiPro NO₂ partial columns are larger than Pandora-DS, GEM-MACH, and TROPOMI, HeiPro surface NO₂ agrees reasonably well with NAPS in situ, with a mean relative bias of -<u>9.7% ± 7.5%</u>, and appears to underestimate GEM-MACH, with a mean relative bias of -<u>3.7% ± 2.4%</u>. This underestimation of GEM-MACH was consistent across all measurement hours of the day and all seasons, while HeiPro more closely matched the diurnal and seasonal trends of NAPS in situ surface NO₂. In comparing the HeiPro and GEM-MACH NO₂ profiles, <u>HeiPro</u> generally <u>underestimates GEM-MACH</u>.
- 920 in the 0–200 m layer with a mean relative bias of –37%, and overestimates GEM-MACH in the 1.5–4 km layer with a significantly large mean relative bias > 1000%. These discrepancies as a function of altitude can probably be explained by the GEM-MACH model inventories. On the other hand, the smoothed GEM-MACH profiles more closely match, the HeiPro profiles across all seasons (mean relative bias from 0–200 m: –6.1%; mean relative bias from 1.5–4 km; 2.6%) as the measurement limitations and vertical sensitivity are removed in the comparison. In summary, the three-year NO₂ profile dataset presented in this study provides information about the spatiotemporal vertical distribution of NO₂ at the Downsview
- measurement site and can be used to assess discrepancies between spaceborne and ground-based NO₂ measurements.

930

clear-sky conditions.

Deleted	: 8
Deleted	:7
Deleted	: 5
Deleted	: 3
Deleted	: 6
Deleted	: GEM-MACH
Deleted	• overestimates
Deleted	: HeiPro
Formatt	t ed: Not Highlight
Deleted bias of 61 more stee with a me	across all seasons from 0–0.80 km with a mean relative %, but above 0.80 km, the GEM-MACH profiles decrease ply with altitude, and GEM-MACH underestimates HeiPro an relative bias of –81%.
Deleted	: be
Deleted	: T
Deleted	es

Deleted: 0.80 Deleted: k Deleted: 34 Deleted: , Deleted: above 0.80 Deleted: -8 Deleted: 0%

Deleted: 1







Figure A1: (a) Histogram of the normalized frequency of degrees of freedom for signal (DOFS) for the <u>HeiPro</u> retrieved <u>NO2</u> profiles from each type of Pandora multi-axis scan, where the me<u>an</u>,values are indicated in brackets. (b) Same as (a) but for <u>NO2</u> partial column <u>errors</u>.







970 Figure A2, Scatter plots for HeiPro surface NO2 vs. NAPS in situ surface NO2 VMRs for scan types and retrieval windows of (a) long UV, (b) short UV, (c) long vis, and (d) short vis. Only the long UV scans were incorporated in the results of this work. The dashed lines and color bar are as indicated in Fig. 3.

Deleted: the

Formatted: Font: Bold



975

Figure A3: MAX-DOAS dSCDs (UV, 338–370 nm) of (a) NO₂ and (c) O₄ retrieved on January 28, 2020. The right panels show the averaging kernels and respective DOFS for <u>a single HeiPro profile retrieval of (b) NO₂ and (d)</u> aerosol extinction, <u>The dSCDs used in the NO₂ and aerosol extinction</u>, profile retrievals are indicated by the red dashed lines in panels (a) and (c), <u>respectively</u>.

980

Table A1. Multiplicative biases and mean relative biases (±, uncertainties) of HeiPro towards NO2 partial columns from Pandora-DS, TROPOMI, and GEM-MACH, for both the HeiPro long UV and long vis results.

NO2 Partial Column Comparison		HeiPro Scan Type	
		long UV	long vis
HeiPro vs. Pandora-DS	Multiplicative Bias	51% ± 0.8%	49% ± 0.9%
partial columns	Mean Relative Bias	$61\%\pm9.7\%$	$61\%\pm6.8\%$
H-: Day of TRODOMI	Multiplicative Bias	$17\% \pm 4.0\%$	13% ± 4.6%
	Mean Relative Bias	37% ± 51%	$40\% \pm 45\%$
HeiDreine CEM MACH	Multiplicative Bias	$12\% \pm 1.2\%$	13% ± 1.3%
Helfro vs. GEMI-MACH	Mean Relative Bias	67% ± 7.1%	64% ± 2.4%

Deleted.	
Deleted: profile retrievals using HeiPro	
Deleted: , whereby the	
Deleted: se	
Formatted: Subscript	
Deleted: .	
Formatted: Font color: Text 1	
Formatted: Font: Bold, Font color: Text 1	
Formatted Table	
Deleted: L	
Deleted: L	
Deleted: 7	
Deleted: 1	

Formatted: Font color: Text 1



Figure A4. Comparisons (2018–2020) of (a) HeiPro (0–4 km) NO₂ partial columns vs. Pandora-DS NO₂ total columns, (b) TROPOMI vs. Pandora-DS NO₂ total columns, and (c) TROPOMI vs. Pandora-DS tropospheric NO₂ columns. The dashed lines and color bar are as indicated in Fig. 3.

- In Figs. A5, and A6, various measurement lines of sight are depicted. Figure A5a-b displays the TROPOMI pixelaveraged NO₂ field from 2018–2020 for summer and winter months, respectively. The MAX-DOAS azimuth viewing angle (255°) and direct-Sun viewing angle (average during summer or winter time periods) are shown. A marker indicates the horizontal extent of the multi-axis path length, i.e., the effective path length. For UV MAX-DOAS measurements, this value is in the range of 5–10 km, and so a horizontal path length of 7.5 km is indicated in the figures (Ortega et al., 2015). During both summer and winter, there does not appear to be an obvious difference in NO₂ between the Pandora MAX-DOAS azimuth
- 1005 viewing angle and the direct-Sun viewing angle at this time of day. However, there are limitations to using the TROPOMI NO₂ field to investigate the HeiPro bias to Pandora-DS, since the TROPOMI data itself underestimates HeiPro and may be insufficient to address how NO₂ heterogeneity from local emissions contributes to the HeiPro overestimation (discussed in Section 3.2). The single temporal datapoint from TROPOMI (1:30 LST) also prevents an investigation as to how spatiotemporal NO₂ heterogeneity around Downsview is impacting the HeiPro overestimation. Alternatively, the GEM-1010 MACH dataset allows an exploration of how the NO₂ heterogeneity changes throughout the day and is discussed next.
- Deleted: 3 Deleted: 4 Deleted: 3



Figure A5, TROPOMI pixel-averaged NO2 field: (a) 2018–2020, summer only; (b) 2018–2020, winter only, around Downsview (white circle), with the MAX-DOAS azimuth viewing angle depicted by the black line and the mean direct-Sun viewing angle during the TROPOMI overpass time (13:30 LT) depicted by the magenta line. The average MAX-DOAS effective path length of 7.5 km is depicted by the black marker along the line of sight.



Deleted: In Fig. A4a-c, the 2018–2020 mean NO₂ field from GEM-MACH is displayed for 7 a.m., 1 p.m., and 5 p.m. LT, respectively, with the Pandora-DS viewing angles shown for each time of day. From the 7 a.m. field, it is not evident that a spatially heterogeneous NO₅ field, combined with differences in viewing geometries, is contributing to the larger bias of HeiPro to Pandora-DS during the early moming hours, since the direct-Sun viewing angle at this time of day faces a similarly polluted region to the multi-axis viewing direction. Although Fig. 5 shows that the bias decreases throughout the day as the sampling directions approach one another, the difference in NO₂ levels between the two sampling directions in the GEM-MACH NO₂ field does not reflect that, possibly due to

Deleted: 4

Deleted: 3 Formatted: Font: Bold

Formatted: Font: Bold

Figure A₆ GEM-MACH averaged NO₂ field from 2018–2020 at (a) 7 a.m. LT, (b) 1 p.m. LT, and (c) 5 p.m. LT around Downsview (white circle), with the MAX-DOAS azimuth viewing angle and average direct-Sun viewing angle during the corresponding times depicted by the black and magenta lines, respectively. The MAX-DOAS effective path length (7.5 km) is depicted by the black marker along the line of sight.

1025

In Fig. A6a–c, the 2018–2020 mean NO₂ field from GEM-MACH is displayed for 7 a.m., 1 p.m., and 5 p.m. LT, respectively, with the Pandora-DS viewing angles shown for each time of day. From the 7 a.m. field, it is not evident that a spatially heterogeneous NO₂ field, combined with differences in viewing geometries, is contributing to the larger bias of HeiPro to Pandora-DS during the early morning hours, since the direct-Sun viewing angle at this time of day faces a similarly polluted region to the multi-axis viewing direction. Although Fig. 5 shows that the bias decreases throughout the day as the sampling directions approach one another, the difference in NO₂ levels between the two sampling directions in the GEM-MACH NO₂ field does not reflect that, possibly due to limitations in using model data to capture local enhancements in NO₂.







Figure A& (a) Scatter plot for the GEM-MACH (0–5 km) vs. Pandora-DS tropospheric NO₂. The dashed lines and color bar are as indicated in Fig. 3. (b) Monthly and (c) hourly box-and-whisker plots of <u>Pandora-DS tropospheric</u> (brown) and GEM-MACH (0–5 km, blue) NO₂ columns as well as the mean relative bias between the two (green circles). Box-and-whisker values are as defined in Fig. 4.

C	
 Deleted: 6	

Formatted: Font: Bold

Formatted: Font: Bold

1065

060

Data availability

Pandora data (L1 multi-axis spectra and L2 direct-Sun NO2 total columns) are available from the Pandonia network 1075 (https://data.pandonia-global-network.org/Downsview/Pandora103s1/; Pandonia Global Network, 2024). OMI NO2 SPv3.1 data are available from https://doi.org/10.5067/Aura/OMI/DATA2017 (Krotkov et al., 2019), TROPOMI L2 NO2 data are available on the Copernicus Open Access Hub at https://scihub.copernicus.eu (ESA, 2024). The NAPS in situ data can be downloaded from https://data-donnees.az.ec.gc.ca/data/air/monitor/national-air-pollution-surveillance-naps-program/, last accessed on 10 September 2024. The HeiPro data (O4 and NO2 dSCDs that went into the retrievals as well as the subsequent profile retrievals) are made available at https://borealisdata.ca/dataset.xhtml?persistentId=doi:10.5683/SP3/J8PDHW.

1080

Author contribution

RA, XZ, KB, and KS conceived this project. RA was responsible for conducting the data analysis, creating all the figures and plots, and drafting the paper. XZ, KB, and KS provided significant inputs throughout the study, and XZ, DG, VF, UF, KB, 1085 and KS provided significant edits to the paper. JD, VF, XZ, and SCL operated and maintained the Pandora103 instrument. DG and AL provided the GEM-MACH data products. UF developed the HeiPro algorithm, YS provided the NAPS in situ data. AC managed the Pandonia network and provided vital technical support for the Downsview site and the ensuing data analysis. All coauthors provided feedback on the paper.

1090 **Competing interests**

At least one of the authors is a member of the editorial board of Atmospheric Measurement Techniques.

Acknowledgments

- Ramina Alwarda was supported by the ECCC Research Affiliate Program and funding from the University of Toronto Faculty 1095 of Arts & Science and would like to express her gratitude for this support. We would like to thank the ECCC Grants & Contributions Program (contribution agreement no. GCXE22S069) for their financial support, which also made this research possible. We acknowledge the NASA Earth Science Division for providing the OMI NO2 SPv3.0 data as well as the Netherlands Space Office (NSO) and the European Space Agency (ESA) for providing the Sentinel-5 Precursor TROPOMI Level 2 product and appreciate the efforts of all those involved in the collection, processing, and distribution of the satellite
- 1100 data. We are grateful to the Pandonia Global Network team and collaborators for their invaluable contributions and ongoing support in the operation and maintenance of the Pandora spectrometer network. The PGN is a bilateral project supported with funding from NASA and ESA. We also thank Vitali Fioletov and staff for their effort in establishing and maintaining Pandora103 at the Downsview site. We acknowledge the European Centre for Medium-Range Weather Forecasts (ECMWF) for providing the ERA5 reanalysis data used in this study, which is publicly available through the Copernicus Climate Change
- 105 Service (C3S). The authors would like to thank the anonymous reviewers for providing constructive feedback, which helped improve the quality of this manuscript,

36

Deleted:

Deleted: https://disc.gsfc.nasa.gov/ Deleted: (NASA, 2020)

Deleted:

References

Alwarda, Ramina; Kristof Bognar; Xiaoyi Zhao; Vitali Fioletov; Jonathan Davies; Sum Chi Lee; Debora Griffin; Alexandru Lupu; Udo Frieß; Alexander Cede; Yushan Su; Kimberly Strong, 2024, "Replication Data for: Retrieval of NO₂ profiles from three years of Pandora MAX-DOAS measurements in Toronto, Canada", https://doi.org/10.5683/SP3/J8PDHW, Borealis, V1

1115 Beirle, S., Platt, U., Wenig, M., and Wagner, T.: Weekly cycle of NO₂ by GOME measurements: a signature of anthropogenic sources, Atmos. Chem. Phys., 3, 2225–2232, 2003.

Blechschmidt, A.-M., Arteta, J., Coman, A., Curier, L., Eskes, H., Foret, G., Gielen, C., Hendrick, F., Marécal, V., Meleux, F., Parmentier, J., Peters, E., Pinardi, G., Piters, A. J. M., Plu, M., Richter, A., Segers, A., Sofiev, M., Valdebenito, Á. M., Van Roozendael, M., Vira, J., Vlemmix, T., and Burrows, J. P.: Comparison of tropospheric NO₂ columns from MAX-DOAS retrievals and regional air quality model simulations, Atmos. Chem. Phys., 20, 2795–2823, https://doi.org/10.5194/acp-20-

Brion, J., Chakir, A., Daumont, D., Malicet, J., and Parisse, C.: High-resolution laboratory absorption cross section of O ₂ .	(

Brion, J., Chakir, A., Charbonnier, J., Daumont, D., Parisse, C., and Malicet, J.: Absorption Spectra Measurements for the 25 Ozone Molecule in the 350–830 nm Region. J. Atmos. Chem., 30, 291–299. https://doi.org/10.1023/a:1006036924364, 1998.

Brohede, S., McLinden, C. A., Urban, J., Haley, C. S., Jonsson, A. I., and Murtagh, D.: Odin stratospheric proxy NOy measurements and climatology, Atmos. Chem. Phys., 24, 2008.

Burrows, J. P., Weber, M., Buchwitz, M., Rozanov, V., Ladstätter-Weißenmayer, A., Richter, A., DeBeek, R., Hoogen, R., Bramstedt, K., Eichmann, K.-U., Eisinger, M., and Perner, D.: The Global Ozone Monitoring Experiment (GOME): Mission
Concept and First Scientific Results, J. Atmos. Sci., 56, 151–175, https://doi.org/10.1175/1520-0469(1999)056<0151:TGOMEG>2.0.CO;2, 1999.

Cede, A.: Manual for Blick Software Suite 1.8, available at: https://www.pandonia-global-network.org/wpcontent/uploads/2021/09/BlickSoftwareSuite_Manual_v1-8-4.pdf, last accessed: 16 April 2024,

Chan, K. L., Wiegner, M., Wenig, M., and Pöhler, D.: Observations of tropospheric aerosols and NO₂ in Hong Kong over 5 1135 years using ground based MAX-DOAS, Science of The Total Environment, 619–620, 1545–1556, https://doi.org/10.1016/j.scitotenv.2017.10.153, 2018

Choi, S., Lamsal, L. N., Follette-Cook, M., Joiner, J., Krotkov, N. A., Swartz, W. H., Pickering, K. E., Loughner, C. P., Appel, W., Pfister, G., Saide, P. E., Cohen, R. C., Weinheimer, A. J., and Herman, J. R.: Assessment of NO₂ observations during DISCOVER-AQ and KORUS-AQ field campaigns, Atmos. Meas. Tech., 13, 2523–2546, https://doi.org/10.5194/amt-13-2523-2020, 2020.

Danckaert, T., Fayt, C., Roozendael, M. V., Smedt, I. D., Letocart, V., Merlaud, A., and Pinardi, G.: QDOAS Software user manual, available at: http://uv-vis.aeronomie.be/software/QDOAS/QDOAS manual.pdf, last access: 16 April 2024

Daumont, D., Brion, J., Charbonnier, J., and Malicet, J.: Ozone UV spectroscopy I: Absorption cross-sections at room temperature, J. Atmos. Chem., 15, 145–155, https://doi.org/10.1007/bf00053756, 1992

1145 Dimitropoulou, E., Hendrick, F., Pinardi, G., Friedrich, M. M., Merlaud, A., Tack, F., De Longueville, H., Fayt, C., Hermans, C., Laffineur, Q., Fierens, F., and Van Roozendael, M.: Validation of TROPOMI tropospheric NO₂ columns using dual-scan Formatted: Subscript

Formatted: Normal, Space After: 12 pt Formatted: English (UK)

Deleted:

Formatted: English (US)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Space After: 12 pt

Deleted: 1

Deleted:

1150	multi-axis differential optical absorption spectroscopy (MAX-DOAS) measurements in Uccle, Brussels, Atmos. Meas. Tech., 13, 5165–5191, https://doi.org/10.5194/amt-13-5165-2020, 2020.	
	ECCC Canadian Environmental Sustainability Indicators: Air Quality; Environment and Climate Change Canada: Dufferin St North York, ON, Canada, 2016; ISBN 978-0-660-06016-3.	
1155	Eskes, H. J. and Eichmann, KU.: S5P Mission Performance Centre Nitrogen Dioxide [L2 NO2] Readme, 2019.	
	Fioletov, V. E., McLinden, C. A., Krotkov, N., Moran, M. D., and Yang, K.: Estimation of SO ₂ emissions using OMI retrievals, Geophys. Res. Lett., 38, https://doi.org/10.1029/2011GL049402, 2011.	
1160	Fleischmann, O. C., Hartmann, M., Burrows, J. P., and Orphal, J.: New ultraviolet absorption cross-sections of BrO at atmospheric temperatures measured by time-windowing Fourier transform spectroscopy, J. Photoch. Photobio. A, 168, 117–132, https://doi.org/10.1016/j.jphotochem.2004.03.026, 2004,	Formatted: Normal, Space After: 12 pt Formatted: English (UK)
I	Frieß, U., Monks, P. S., Remedios, J. J., Rozanov, A., Sinreich, R., Wagner, T., and Platt, U.: MAX-DOAS O ₄ measurements: A new technique to derive information on atmospheric aerosols: 2. Modeling studies, J. Geophys. Res., 111, D14203, https://doi.org/10.1029/2005JD006618, 2006.	
1165	Frieß, U., Sihler, H., Sander, R., Pöhler, D., Yilmaz, S., and Platt, U.: The vertical distribution of BrO and aerosols in the Arctic: Measurements by active and passive differential optical absorption spectroscopy, J. Geophys. Res., 116, D00R04, https://doi.org/10.1029/2011JD015938, 2011.	
1170	Frieß, U., Beirle, S., Alvarado Bonilla, L., Bösch, T., Friedrich, M. M., Hendrick, F., Piters, A., Richter, A., van Roozendael, M., Rozanov, V. V., Spinei, E., Tirpitz, JL., Vlemmix, T., Wagner, T., and Wang, Y.: Intercomparison of MAX-DOAS vertical profile retrieval algorithms: studies using synthetic data, Atmos. Meas. Tech., 12, 2155–2181, https://doi.org/10.5194/amt-12-2155-2019, 2019.	
ļ	van Geffen, J., Boersma, K. F., Eskes, H., Sneep, M., ter Linden, M., Zara, M., and Veefkind, J. P.: S5P TROPOMI NO ₂ slant column retrieval: method, stability, uncertainties and comparisons with OMI, Atmos. Meas. Tech., 13, 1315–1335, https://doi.org/10.5194/amt-13-1315-2020, 2020.	Deleted: Herman, J., Cede, A., Spinei, E., Mount, G., Tzortziou, M., and Abuhassan, N.: NO ₂ column amounts from ground-based Pandora and MFDOAS spectrometers using the direct-sun DOAS technique: Intercomparisons and application to OMI validation, J.
1175	van Geffen, J., Eskes, H., Compernolle, S., Pinardi, G., Verhoelst, T., Lambert, JC., Sneep, M., ter Linden, M., Ludewig, A., Boersma, K. F., and Veefkind, J. P.: Sentinel-5P TROPOMI NO ₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data, Atmos. Meas. Tech., 15, 2037–2060, https://doi.org/10.5194/amt-15-2037-2022, 2022.	Geophys. Res., 114, D13307, https://doi.org/10.1029/2009JD011848, 2009.¶
1180	Griffin, D., Zhao, X., McLinden, C. A., Boersma, F., Bourassa, A., Dammers, E., Degenstein, D., Eskes, H., Fehr, L., Fioletov, V., Hayden, K., Kharol, S. K., Li, S., Makar, P., Martin, R. V., Mihele, C., Mittermeier, R. L., Krotkov, N., Sneep, M., Lamsal, L. N., Linden, M. ter, Geffen, J. van, Veefkind, P., and Wolde, M.: High-Resolution Mapping of Nitrogen Dioxide With TROPOMI: First Results and Validation Over the Canadian Oil Sands, Geophys. Res. Lett., 46, 1049–1060, https://doi.org/10.1029/2018GL081095, 2019.	
	Herman, J., Cede, A., Spinei, E., Mount, G., Tzortziou, M., and Abuhassan, N.: NO2 column amounts from ground-based	

Herman, J., Spinei, E., Fried, A., Kim, J., Kim, J., Kim, W., Cede, A., Abuhassan, N., and Segal-Rozenhaimer, M.: NO₂ and HCHO measurements in Korea from 2012 to 2016 from Pandora spectrometer instruments compared with OMI retrievals and

Pandora and MFDOAS spectrometers using the direct-sun DOAS technique: Intercomparisons and application to OMI validation, J. Geophys. Res., 114, D13307, https://doi.org/10.1029/2009JD011848, 2009.

1185

1195 with aircraft measurements during the KORUS-AQ campaign, Atmos. Meas. Tech., 11, 4583–4603, https://doi.org/10.5194/amt-11-4583-2018, 2018

1200

Deleted:

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J-N.: ERA5 hourly data on pressure levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), 10.24381/cds.bd0915c6 (Accessed on 21-Jan-2024), 2023.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J-N.: ERA5 hourly data on single levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), 10.24381/cds.adbb2d47 (Accessed on 21-Jan-2024), 2023.

Kim, J., Jeong, U., Ahn, M.-H., Kim, J. H., Park, R. J., Lee, H., Song, C. H., Choi, Y.-S., Lee, K.-H., Yoo, J.-M., Jeong, M.-J., Park, S. K., Lee, K.-M., Song, C.-K., Kim, S.-W., Kim, Y. J., Kim, S.-W., Kim, M., Go, S., Liu, X., Chance, K., Chan Miller, C., Al-Saadi, J., Veihelmann, B., Bhartia, P. K., Torres, O., Abad, G. G., Haffner, D. P., Ko, D. H., Lee, S. H., Woo,

- 1210 J.-H., Chong, H., Park, S. S., Nicks, D., Choi, W. J., Moon, K.-J., Cho, A., Yoon, J., Kim, S., Hong, H., Lee, K., Lee, H., Lee, S., Choi, M., Veefkind, P., Levelt, P. F., Edwards, D. P., Kang, M., Eo, M., Bak, J., Baek, K., Kwon, H.-A., Yang, J., Park, J., Han, K. M., Kim, B.-R., Shin, H.-W., Choi, H., Lee, E., Chong, J., Cha, Y., Koo, J.-H., Irie, H., Hayashida, S., Kasai, Y., Kanaya, Y., Liu, C., Lin, J., Crawford, J. H., Carmichael, G. R., Newchurch, M. J., Lefer, B. L., Herman, J. R., Swap, R. J., Lau, A. K. H., Kurosu, T. P., Jaross, G., Ahlers, B., Dobber, M., McElroy, C. T., and Choi, Y.: New Era of Air Quality
- 1215 Monitoring from Space: Geostationary Environment Monitoring Spectrometer (GEMS), Bulletin of the American Meteorological Society, 101, E1–E22, https://doi.org/10.1175/BAMS-D-18-0013.1, 2020.

Kreher, K., Van Roozendael, M., Hendrick, F., Apituley, A., Dimitropoulou, E., Frieß, U., Richter, A., Wagner, T., Lampel, J., Abuhassan, N., Ang, L., Anguas, M., Bais, A., Benavent, N., Bösch, T., Bognar, K., Borovski, A., Bruchkouski, I., Cede, A., Chan, K. L., Donner, S., Drosoglou, T., Fayt, C., Finkenzeller, H., Garcia-Nieto, D., Gielen, C., Gómez-Martín, L., Hao,

- 1220 N., Henzing, B., Herman, J. R., Hermans, C., Hoque, S., Irie, H., Jin, J., Johnston, P., Khayyam Butt, J., Khokhar, F., Koenig, T. K., Kuhn, J., Kumar, V., Liu, C., Ma, J., Merlaud, A., Mishra, A. K., Müller, M., Navarro-Comas, M., Ostendorf, M., Pazmino, A., Peters, E., Pinardi, G., Pinharanda, M., Piters, A., Platt, U., Postylyakov, O., Prados-Roman, C., Puentedura, O., Querel, R., Saiz-Lopez, A., Schönhardt, A., Schreier, S. F., Seyler, A., Sinha, V., Spinei, E., Strong, K., Tack, F., Tian, X., Tiefengraber, M., Tirpitz, J.-L., van Gent, J., Volkamer, R., Vrekoussis, M., Wang, S., Wang, Z., Wenig, M., Wittrock, F.,
- 1225 Xie, P. H., Xu, J., Yela, M., Zhang, C., and Zhao, X.: Intercomparison of NO₂, O₄, O₃ and HCHO slant column measurements by MAX-DOAS and zenith-sky UV-visible spectrometers during CINDI-2, Atmos. Meas. Tech., 13, 2169–2208, https://doi.org/10.5194/amt-13-2169-2020, 2020.

Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., Marchenko, S. V., Bucsela, E. J., Chan, K. L., Wenig, M., and Zara, M.: The version 3 OMI NO² standard product, Atmos. Meas. Tech., 10, 3133–3149, https://doi.org/10.5194/amt-10-3133-2017, 2017.

Lin, J.-T. and McElroy, M. B.: Impacts of boundary layer mixing on pollutant vertical profiles in the lower troposphere: Implications to satellite remote sensing, Atmospheric Environment, 44, 1726–1739, https://doi.org/10.1016/j.atmosenv.2010.02.009, 2010.

Lindenmaier, R., Strong, K., Batchelor, R. L., Bernath, P. F., Chabrillat, S., Chipperfield, M. P., Daffer, W. H., Drummond, J.
 R., Feng, W., Jonsson, A. I., Kolonjari, F., Manney, G. L., McLinden, C., Ménard, R., and Walker, K. A.: A study of the Arctic NOy budget above Eureka, Canada, J. Geophys. Res., 116, n/a-n/a, https://doi.org/10.1029/2011JD016207, 2011.

Ma, J. Z., Beirle, S., Jin, J. L., Shaiganfar, R., Yan, P., and Wagner, T.: Tropospheric NO₂ vertical column densities over Beijing: results of the first three years of ground-based MAX-DOAS measurements (2008-2011) and satellite validation, 1240 Atmos. Chem. Phys., 13, 1547–1567, https://doi.org/10.5194/acp-13-1547-2013, 2013.

McLinden, C. A., Olsen, S. C., Hannegan, B., Wild, O., Prather, M. J., and Sundet, J.: Stratospheric ozone in 3-D models: A simple chemistry and the cross-tropopause flux, J. Geophys. Res., 105, 14653-14665, https://doi.org/10.1029/2000JD900124, 2000.

Meller, R. and Moortgat, G. K.: Temperature dependence of the absorption cross sections of formaldehyde between 223 and 323 K in the wavelength range 225-375 nm, J. Geophys. Res., 105, 7089-7101, https://doi.org/10.1029/1999JD901074. 2000. 1245

Meng, K., Xu, X., Cheng, X., Xu, X., Qu, X., Zhu, W., Ma, C., Yang, Y., and Zhao, Y.: Spatio-temporal variations in SO2 and NO₂ emissions caused by heating over the Beijing-Tianjin-Hebei Region constrained by an adaptive nudging method with OMI data, Science of The Total Environment, 642, 543-552, https://doi.org/10.1016/j.scitotenv.2018.06.021, 2018.

Moran, M. D., Menard, S., Talbot, D., Huang, P., Makar, P. A., Gong, W., Landry, H., Gong, S., Gravel, S., Crevier, L.-P., and Kallaur, A.: Particulate-matter forecasting with GEM-MACH15, a new Canadian operational air quality forecast model, 1250 in 30th NATO/SPS ITM on Air Pollution Modelling and Its Application, 289-293, Springer, San Francisco, 2009.

Murray, L. T.: Lightning NO_x and Impacts on Air Quality, Curr Pollution Rep. 2, 115-133, https://doi.org/10.1007/s40726-016-0031-7, 2016.

- 1255 Nowlan, C. R., Liu, X., Janz, S. J., Kowalewski, M. G., Chance, K., Follette-Cook, M. B., Fried, A., González Abad, G., Herman, J. R., Judd, L. M., Kwon, H.-A., Loughner, C. P., Pickering, K. E., Richter, D., Spinei, E., Walega, J., Weibring, P., and Weinheimer, A. J.: Nitrogen dioxide and formaldehyde measurements from the GEOstationary Coastal and Air Pollution Events (GEO-CAPE) Airborne Simulator over Houston, Texas, Atmos. Meas. Tech., 11, 5941-5964, https://doi.org/10.5194/amt-11-5941-2018, 2018.
- Ortega, I., Koenig, T., Sinreich, R., Thomson, D., and Volkamer, R.: The CU 2-D-MAX-DOAS instrument Part 1: Retrieval 1260 of 3-D distributions of NO2 and azimuth-dependent OVOC ratios, Atmos. Meas. Tech., 8, 2371-2395, https://doi.org/10.5194/amt-8-2371-2015, 2015.

Pavlovic, R., Chen, J., Anderson, K., Moran, M. D., Beaulieu, P.-A., Davignon, D., and Cousineau, S.: The FireWork air quality forecast system with near-real-time biomass burning emissions: Recent developments and evaluation of performance 1265 for the 2015 North American wildfire season, Journal of the Air & Waste Management Association, 66, 819-841, https://doi.org/10.1080/10962247.2016.1158214, 2016.

Pendlebury, D., Gravel, S., Moran, M. D., and Lupu, A.: Impact of chemical lateral boundary conditions in a regional air quality forecast model on surface ozone predictions during stratospheric intrusions, Atmospheric Environment, 174, 148-170, https://doi.org/10.1016/j.atmosenv.2017.10.052, 2018.

Pinardi, G., Van Roozendael, M., Hendrick, F., Thevs, N., Abuhassan, N., Bais, A., Boersma, F., Cede, A., Chong, J., Donner, 1270 S., Drosoglou, T., Dzhola, A., Eskes, H., Frieß, U., Granville, J., Herman, J. R., Holla, R., Hovila, J., Irie, H., Kanaya, Y., Karagkiozidis, D., Kouremeti, N., Lambert, J.-C., Ma, J., Peters, E., Piters, A., Postylyakov, O., Richter, A., Remmers, J., Takashima, H., Tiefengraber, M., Valks, P., Vlemmix, T., Wagner, T., and Wittrock, F.: Validation of tropospheric NO2 column measurements of GOME-2A and OMI using MAX-DOAS and direct sun network observations, Atmos. Meas. Tech., 1275 13, 6141-6174, https://doi.org/10.5194/amt-13-6141-2020, 2020.

Platt, U. and Stutz, J.: Differential optical absorption spectroscopy: principles and applications, Springer, Berlin, 597 pp., 2008.

Rodgers, C. D.: Inverse methods for atmospheric sounding: Theory and practice, vol. 2, World Scientific, Singapore, ISBN 920 978-981-02-2740-1, 256 pp., 2000.

1280 Rothman, L., Gordon, I., Barber, R., Dothe, H., Gamache, R., Goldman, A., Perevalov, V., Tashkun, S., and Tennyson, J.: HITEMP, the high-temperature molecular spectroscopic database, J. Quant. Spectrosc. Ra., 111, 2139–2150, https://doi.org/10.1016/j.jqsrt.2010.05.001, 2010.

Rozanov, A., Rozanov, V., Buchwitz, M., Kokhanovsky, A., and Burrows, J. P.: SCIATRAN 2.0 – A new radiative transfer model for geophysical applications in the 175–2400 nm spectral region, Advances in Space Research, 36, 1015–1019, https://doi.org/10.1016/j.asr.2005.03.012, 2005.

Schreier, S. F., Richter, A., and Burrows, J. P.: Near-surface and path-averaged mixing ratios of NO₂ derived from car DOAS zenith-sky and tower DOAS off-axis measurements in Vienna: a case study, Atmos. Chem. Phys., 19, 5853–5879, https://doi.org/10.5194/acp-19-5853-2019, 2019.

Serdyuchenko, A., Gorshelev, V., Weber, M., Chehade, W., and Burrows, J. P.: High spectral resolution ozone absorption crosssections – Part 2: Temperature dependence, Atmos. Meas. Tech., 7, 625–636, https://doi.org/10.5194/amt-7-625-2014, 2014.

Stieb, D. M., Burnett, R. T., Smith-Doiron, M., Brion, O., Shin, H. H., and Economou, V.: A New Multipollutant, No-Threshold Air Quality Health Index Based on Short-Term Associations Observed in Daily Time-Series Analyses, Journal of the Air & Waste Management Association, 58, 435–450, https://doi.org/10.3155/1047-3289.58.3.435, 2008.

1295 Sun, K., Zhu, L., Cady-Pereira, K., Chan Miller, C., Chance, K., Clarisse, L., Coheur, P.-F., González Abad, G., Huang, G., Liu, X., Van Damme, M., Yang, K., and Zondlo, M.: A physics-based approach to oversample multi-satellite, multispecies observations to a common grid, Atmos. Meas. Tech., 11, 6679–6701, https://doi.org/10.5194/amt-11-6679-2018, 2018.

Thalman, R. and Volkamer, R.: Temperature dependent absorption cross-sections of O₂–O₂ collision pairs between 340 and 630 nm and at atmospherically relevant pressure, Phys. Chem. Chem. Phys., 15, 15371–15381, https://doi.org/10.1039/C3CP50968K, 2013

Thermo Scientific. (2015). Model 42i instruction manual: Chemiluminescence NO-NO₂-NO_x analyzer (Part No. 101350-00, 25JUL2015). Thermo Fisher Scientific.

Vandaele, A., Hermans, C., Simon, P., Carleer, M., Colin, R., Fally, S., Mérienne, M., Jenouvrier, A., and Coquart, B.:
 Measurements of the NO₂ absorption cross-section from 42 000 cm-1 to 10 000 cm-1 (238–1000 nm) at 220 K and 294 K, J. Quant. Spectrosc. Ra., 59, 171–184, https://doi.org/10.1016/S0022- 4073(97)00168-4, 1998.

Verhoelst, T., Compernolle, S., Pinardi, G., Lambert, J.-C., Eskes, H. J., Eichmann, K.-U., Fjæraa, A. M., Granville, J., Niemeijer, S., Cede, A., Tiefengraber, M., Hendrick, F., Pazmiño, A., Bais, A., Bazureau, A., Boersma, K. F., Bognar, K., Dehn, A., Donner, S., Elokhov, A., Gebetsberger, M., Goutail, F., Grutter de la Mora, M., Gruzdev, A., Gratsea, M., Hansen,

- 1310 G. H., Irie, H., Jepsen, N., Kanaya, Y., Karagkiozidis, D., Kivi, R., Kreher, K., Levelt, P. F., Liu, C., Müller, M., Navarro Comas, M., Piters, A. J. M., Pommereau, J.-P., Portafaix, T., Prados-Roman, C., Puentedura, O., Querel, R., Remmers, J., Richter, A., Rimmer, J., Rivera Cárdenas, C., Saavedra de Miguel, L., Sinyakov, V. P., Stremme, W., Strong, K., Van Roozendael, M., Veefkind, J. P., Wagner, T., Wittrock, F., Yela González, M., and Zehner, C.: Ground-based validation of the Copernicus Sentinel-5P TROPOMI NO₂ measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandonia global
- 1315 networks, Atmos. Meas. Tech., 14, 481-510, https://doi.org/10.5194/amt-14-481-2021, 2021.

300

41

Formatted: Normal, Space After: 12 pt

Deleted: 1

Deleted:

Vlemmix, T., Hendrick, F., Pinardi, G., De Smedt, I., Fayt, C., Hermans, C., Piters, A., Wang, P., Levelt, P., and Van Roozendael, M.: MAX-DOAS observations of aerosols, formaldehyde and nitrogen dioxide in the Beijing area: comparison of two profile retrieval approaches, Atmos. Meas. Tech., 8, 941–963, https://doi.org/10.5194/amt-8-941-2015, 2015.

Yilmaz, S.: Retrieval of atmospheric aerosol and trace gas vertical profiles using multi-axis differential optical absorption spectroscopy, University of Heidelberg, 2012.

York, D., Evensen, N. M., Martínez, M. L., and De Basabe Delgado, J.: Unified equations for the slope, intercept, and standard errors of the best straight line, American Journal of Physics, 72, 367–375, https://doi.org/10.1119/1.1632486, 2004.

1325 Zhao, X., Griffin, D., Fioletov, V., McLinden, C., Davies, J., Ogyu, A., Lee, S. C., Lupu, A., Moran, M. D., Cede, A., Tiefengraber, M., and Müller, M.: Retrieval of total column and surface NO₂ from Pandora zenith-sky measurements, Atmos. Chem. Phys., 19, 10619–10642, https://doi.org/10.5194/acp-19-10619-2019, 2019.

Zhao, X., Griffin, D., Fioletov, V., McLinden, C., Cede, A., Tiefengraber, M., Müller, M., Bognar, K., Strong, K., Boersma, F., Eskes, H., Davies, J., Ogyu, A., and Lee, S. C.: Assessment of the quality of TROPOMI high-spatial-resolution NO₂ data products in the Greater Toronto Area, Atmos. Meas. Tech., 13, 2131–2159, https://doi.org/10.5194/amt-13-2131-2020, 2020.

Zhao, X., Fioletov, V., Alwarda, R., Su, Y., Griffin, D., Weaver, D., Strong, K., Cede, A., Hanisco, T., Tiefengraber, M., McLinden, C., Eskes, H., Davies, J., Ogyu, A., Sit, R., Abboud, I., and Lee, S. C.: Tropospheric and Surface Nitrogen Dioxide Changes in the Greater Toronto Area during the First Two Years of the COVID-19 Pandemic, Remote Sensing, 14, 1625, https://doi.org/10.3390/rs14071625, 2022.

- 1335 Zoogman, P., Liu, X., Suleiman, R. M., Pennington, W. F., Flittner, D. E., Al-Saadi, J. A., Hilton, B. B., Nicks, D. K., Newchurch, M. J., Carr, J. L., Janz, S. J., Andraschko, M. R., Arola, A., Baker, B. D., Canova, B. P., Chan Miller, C., Cohen, R. C., Davis, J. E., Dussault, M. E., Edwards, D. P., Fishman, J., Ghulam, A., González Abad, G., Grutter, M., Herman, J. R., Houck, J., Jacob, D. J., Joiner, J., Kerridge, B. J., Kim, J., Krotkov, N. A., Lamsal, L., Li, C., Lindfors, A., Martin, R. V., McElroy, C. T., McLinden, C., Natraj, V., Neil, D. O., Nowlan, C. R., O'Sullivan, E. J., Palmer, P. I., Pierce, R. B., Pippin, M.
- 1340 R., Saiz-Lopez, A., Spurr, R. J. D., Szykman, J. J., Torres, O., Veefkind, J. P., Veihelmann, B., Wang, H., Wang, J., and Chance, K.: Tropospheric emissions: Monitoring of pollution (TEMPO), Journal of Quantitative Spectroscopy and Radiative Transfer, 186, 17–39, https://doi.org/10.1016/j.jqsrt.2016.05.008, 2017.